


Essential Mathematics Core for the Victorian Curriculum Year 9
ISBN 978-1-009-59315-1
© Greenwood et al. 2025
Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

CAMBRIDGE
UNIVERSITY PRESS & ASSESSMENT

Shaftesbury Road, Cambridge CB2 8EA, United Kingdom

One Liberty Plaza, 20th Floor, New York, NY 10006, USA

477 Williamstown Road, Port Melbourne, VIC 3207, Australia

314–321, 3rd Floor, Plot 3, Splendor Forum, Jasola District Centre, New Delhi – 110025, India

103 Penang Road, #05–06/07, Visioncrest Commercial, Singapore 238467

Cambridge University Press & Assessment is a department of the University of Cambridge.

We share the University's mission to contribute to society through the pursuit of education, learning and research at the highest international levels of excellence.

www.cambridge.org

First edition © David Greenwood, Sara Woolley, Jenny Goodman and Jennifer Vaughan 2021
Second edition © David Greenwood, Sara Woolley, Jenny Goodman and Jennifer Vaughan 2025

This publication is in copyright. Subject to statutory exception and to the provisions of relevant collective licensing agreements, no reproduction of any part may take place without the written permission of Cambridge University Press & Assessment.

First published 2021
Second Edition 2025

20 19 18 17 16 15 14 13 12 11 10 9 8 7 6 5 4 3 2 1

Cover design by Sardine Design
Typeset by diacriTech
Printed in China by C & C Offset Printing Co., Ltd.

A catalogue record for this book is available from the National Library of Australia at www.nla.gov.au

ISBN 978-1-009-59315-1

Additional resources for this publication at www.cambridge.edu.au/GO

Reproduction and Communication for educational purposes
The Australian *Copyright Act 1968* (the Act) allows a maximum of one chapter or 10% of the pages of this publication, whichever is the greater, to be reproduced and/or communicated by any educational institution for its educational purposes provided that the educational institution (or the body that administers it) has given a remuneration notice to Copyright Agency Limited (CAL) under the Act.

For details of the CAL licence for educational institutions contact:

Copyright Agency Limited
Level 12, 66 Goulburn Street
Sydney NSW 2000
Telephone: (02) 9394 7600
Facsimile: (02) 9394 7601
Email: memberservices@copyright.com.au

Reproduction and Communication for other purposes
Except as permitted under the Act (for example a fair dealing for the purposes of study, research, criticism or review) no part of this publication may be reproduced, stored in a retrieval system, communicated or transmitted in any form or by any means without prior written permission. All inquiries should be made to the publisher at the address above.

Cambridge University Press & Assessment has no responsibility for the persistence or accuracy of URLS for external or third-party internet websites referred to in this publication and does not guarantee that any content on such websites is, or will remain, accurate or appropriate. Information regarding prices, travel timetables and other factual information given in this work is correct at the time of first printing but Cambridge University Press & Assessment does not guarantee the accuracy of such information thereafter.

Cambridge University Press & Assessment acknowledges the Aboriginal and Torres Strait Islander peoples of this nation. We acknowledge the traditional custodians of the lands on which our company is located and where we conduct our business. We pay our respects to ancestors and Elders, past and present. Cambridge University Press & Assessment is committed to honouring Aboriginal and Torres Strait Islander peoples' unique cultural and spiritual relationships to the land, waters and seas and their rich contribution to society.

Contents

			Strand and content description
	About the authors	*viii*	
	Acknowledgements	*ix*	
	Introduction	*x*	
	Guide to the working programs	*xi*	
	Guide to this resource	*xii*	
1	**Reviewing number**	2	Number, Algebra, Measurement
	Warm-up quiz	4	
1A	Adding and subtracting integers CONSOLIDATING	5	
1B	Multiplying and dividing integers CONSOLIDATING	9	
1C	Rounding decimals and significant figures	14	
1D	Rational and irrational numbers CONSOLIDATING	19	
1E	Adding and subtracting fractions CONSOLIDATING	25	
	Progress quiz	30	
1F	Multiplying and dividing fractions CONSOLIDATING	31	
1G	Ratios CONSOLIDATING	36	
1H	Rates and best buys	40	
	Maths@Work: Cooks and chefs	46	
	Modelling	48	
	Technology and computational thinking	50	
	Puzzles and games	52	
	Chapter summary and checklist	53	
	Chapter review	57	
2	**Financial mathematics**	60	Number, Algebra, Measurement
	Warm-up quiz	62	
2A	Percentages CONSOLIDATING	63	
2B	Applying percentages CONSOLIDATING	68	
2C	Percentage increase and decrease CONSOLIDATING	71	
2D	Profits and discounts CONSOLIDATING	76	
2E	Income	81	
	Progress quiz	86	
2F	Taxation	87	
2G	Simple interest	91	
2H	Applications of simple interest	96	
	Maths@Work: Facebook cake-decorating business	101	
	Modelling	103	
	Technology and computational thinking	105	
	Puzzles and games	107	
	Chapter summary and checklist	108	
	Chapter review	111	

Essential Mathematics Core for the Victorian Curriculum Year 9 ISBN 978-1-009-59315-1 © Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3 Expressions and equations 114 — Number, Algebra

	Warm-up quiz	116
3A	Algebraic expressions CONSOLIDATING	117
3B	Adding and subtracting algebraic expressions CONSOLIDATING	123
3C	Multiplying and dividing algebraic expressions CONSOLIDATING	127
3D	Expanding algebraic expressions	131
3E	Solving linear equations CONSOLIDATING	136
3F	Solving linear equations involving fractions	140
	Progress quiz	145
3G	Solving equations with brackets	146
3H	Solving equations with pronumerals on both sides	149
3I	Solving word problems ★	154
3J	Solving linear inequalities	159
3K	Using formulas ★	164
	Maths@Work: Plumber	169
	Modelling	171
	Technology and computational thinking	173
	Puzzles and games	175
	Chapter summary and checklist	176
	Chapter review	179

4 Pythagoras' theorem and trigonometry 182 — Number, Measurement, Space

	Warm-up quiz	184
4A	Exploring Pythagoras' theorem	185
4B	Finding the length of the hypotenuse	189
4C	Finding the lengths of the shorter sides	194
4D	Applying Pythagoras' theorem ★	198
4E	Trigonometric ratios	203
4F	Finding side lengths	210
	Progress quiz	215
4G	Solving for the denominator ★	217
4H	Finding an angle	222
4I	Applying trigonometry ★	227
	Maths@Work: Carpenter	232
	Modelling	234
	Technology and computational thinking	236
	Puzzles and games	238
	Chapter summary and checklist	239
	Chapter review	242

5 Linear relationships 246 — Algebra, Measurement

	Warm-up quiz	248
5A	Introduction to linear relationships	249
5B	Finding x- and y-intercepts	256
5C	Graphing straight lines using intercepts	262
5D	Lines with one intercept	266

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5E	Gradient	272
5F	Gradient and direct proportion	279
5G	Gradient–intercept form	284
	Progress quiz	291
5H	Finding the equation of a line ★	293
5I	Midpoint and length of a line segment	298
5J	Linear modelling ★	303
	Maths@Work: Trading in foreign currencies	307
	Modelling	309
	Technology and computational thinking	311
	Puzzles and games	313
	Chapter summary and checklist	314
	Chapter review	318

Semester review 1 322

6 Measurement 328 — Number, Measurement

	Warm-up quiz	330
6A	Errors and accuracy in measurement	331
6B	Length and perimeter CONSOLIDATING	336
6C	Circumference of a circle CONSOLIDATING	341
6D	Area CONSOLIDATING	346
6E	Area of a circle CONSOLIDATING	352
6F	Composite shapes	356
6G	Surface area of prisms	361
	Progress quiz	366
6H	Surface area of a cylinder	368
6I	Volume	373
6J	Volume of a cylinder	379
	Maths@Work: Vegetable and fruit growers	383
	Modelling	385
	Technology and computational thinking	387
	Puzzles and games	389
	Chapter summary and checklist	390
	Chapter review	394

7 Indices 398 — Number, Algebra, Measurement

	Warm-up quiz	400
7A	Index notation CONSOLIDATING	401
7B	Index laws for multiplication and division	409
7C	Index laws for power of a power and the zero index	415
7D	Index laws for brackets and fractions	420
7E	Negative indices ★	426
	Progress quiz	431
7F	Scientific notation	432

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

7G	Scientific notation using significant figures	437
	Maths@Work: Lab technician	442
	Modelling	444
	Technology and computational thinking	446
	Puzzles and games	448
	Chapter summary and checklist	449
	Chapter review	452

8 Geometry 454 — Measurement, Space

	Warm-up quiz	456
8A	Angles and triangles CONSOLIDATING	457
8B	Parallel lines CONSOLIDATING	464
8C	Quadrilaterals CONSOLIDATING	470
8D	Polygons ★	475
8E	Congruent triangles	480
	Progress quiz	486
8F	Enlargement and similar figures	487
8G	Similar triangles	493
8H	Applying similar triangles	499
	Maths@Work: Animator	503
	Modelling	505
	Technology and computational thinking	507
	Puzzles and games	509
	Chapter summary and checklist	510
	Chapter review	514

9 Algebraic techniques 518 — Algebra

	Warm-up quiz	520
9A	Reviewing algebra CONSOLIDATING	521
9B	Expanding binomial products	526
9C	Expanding perfect squares	530
9D	Forming a difference of two squares	534
9E	Factorising algebraic expressions	537
	Progress quiz	541
9F	Factorising the difference of two squares	542
9G	Factorising by grouping in pairs	546
9H	Factorising monic quadratic trinomials	550
9I	Simplifying algebraic fractions: multiplication and division ★	554
9J	Simplifying algebraic fractions: addition and subtraction ★	559
	Maths@Work: Automotive technology	563
	Modelling	565
	Technology and computational thinking	567
	Puzzles and games	569
	Chapter summary and checklist	570
	Chapter review	573

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10 Statistics and probability 576 — Statistics, Probability

Warm-up quiz 578
10A Review of probability CONSOLIDATING 579
10B Venn diagrams and two-way tables 586
10C Using arrays for two-step experiments 595
10D Using tree diagrams 602
10E Using relative frequencies to estimate probabilities 607
Progress quiz 612
10F Data and sampling 614
10G Summarising data: range and measures of centre 620
10H Interpreting data from tables and graphs 625
10I Stem-and-leaf plots 634
10J Grouped data 641
Maths@Work: Personal trainer 648
Modelling 650
Technology and computational thinking 652
Puzzles and games 654
Chapter summary and checklist 655
Chapter review 659

11 Introduction to quadratic equations and graphs 664 — Algebra

Warm-up quiz 666
11A Quadratic equations 667
11B Solving $ax^2 + bx = 0$ and $x^2 = d$ 672
11C Solving $x^2 + bx + c = 0$ 677
11D Using quadratic equations to solve problems 681
Progress quiz 685
11E The parabola 686
11F Sketching $y = ax^2$ with dilations and reflections 695
11G Sketching translations of $y = x^2$ 703
11H Sketching parabolas using intercept form ★ 711
Maths@Work: Fencing 718
Modelling 719
Technology and computational thinking 721
Puzzles and games 723
Chapter summary and checklist 724
Chapter review 727

Semester review 2 732

12 Algorithmic thinking (online only) 830

Activity 1: Algorithms for number patterns and financial maths 833
Activity 2: Minimising and maximising 837
Activity 3: Sorting, simulations and sampling 839

Glossary ***742***
Answers ***748***

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

About the Authors

David Greenwood is the Head of Mathematics at Trinity Grammar School in Melbourne and has 30+ years teaching mathematics from Year 7 to 12. He is the lead author for the Cambridge Essential series and has authored more than 80 titles for the Australian Curriculum and for the syllabuses of the states and territories. He specialises in analysing curriculum and the sequencing of course content for school mathematics courses. He also has an interest in the use of technology for the teaching of mathematics.



Sara Woolley was born and educated in Tasmania. She completed an Honours degree in Mathematics at the University of Tasmania before completing her education training at the University of Melbourne. She has taught mathematics from Years 7 to 12 since 2006 and is currently a Head of Mathematics. She specialises in lesson design and creating resources that develop and build understanding of mathematics for all students.



Jenny Goodman has taught in schools for over 28 years and is currently teaching at a selective high school in Sydney. Jenny has an interest in the importance of literacy in mathematics education, and in teaching students of differing ability levels. She was awarded the Jones Medal for education at Sydney University and the Bourke Prize for Mathematics. She has written for *CambridgeMATHS NSW* and was involved in the *Spectrum* and *Spectrum Gold* series.



Jennifer Vaughan has taught secondary mathematics for over 30 years in New South Wales, Western Australia, Queensland and New Zealand and has tutored and lectured in mathematics at Queensland University of Technology. She is passionate about providing students of all ability levels with opportunities to understand and to have success in using mathematics. She has had extensive experience in developing resources that make mathematical concepts more accessible; hence, facilitating student confidence, achievement and an enjoyment of maths.



Stuart Palmer was born and educated in New South Wales. He is a fully qualified high school mathematics teacher with more than 25 years' experience teaching students from all walks of life in a variety of schools. He has been Head of Mathematics in two schools. He is very well known by teachers throughout the state for the professional learning workshops he delivers. Stuart also assists thousands of Year 12 students every year as they prepare for their HSC Examinations. At the University of Sydney, Stuart spent more than a decade running tutorials for pre-service mathematics teachers.



Essential Mathematics Core for the Victorian Curriculum Year 9 ISBN 978-1-009-59315-1 © Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Acknowledgements

The author and publisher wish to thank the following sources for permission to reproduce material:

Cover: © Getty Images / Westend61.

Images: © Getty Images / Maskot, Chapter 1 Opener / RyersonClark, p.13 / Gri-spb, p.14 / Caia Image, p.18(1) / Busà Photography, p.24 / Hans Neleman, p.29(1) / bergserg, p.29(2) / Annabelle Breakey, p.30 / Gaelle Beller Studio, p.31 / EujarimPhotography, p.34 / Nicola Tree, p.35 / ArtMarie, p.36 / Betsie Van der Meer, p.39(1) / Halfdark, p.39(2) / GlobalP, p.40(1–12) / aluxum, p.43 / Michael Hall, p.44(1) / Connect Images, p.44(2) / Jakob Helbig, p.45 / Claudia Totir, p.46 / andresr, p.49 / Jeffrey Coolidge, p.50 / Jacobs Stock Photography Ltd, p.59 / Alfexe, Chapter 2 Opener / irwan rosidi / 500px, p.66 / yasinguneysu, p.74 / clubfoto, p.77 / Firdausiah Mamat, p.91 / Betsie Van der Meer, p.94 / Grafissimo, p.99 / Aleksandr Zubkov, p.102 / courtneyk, p.111 / Busà Photography, Chapter 3 Opener / Yulia Naumenko, p.122 / Maria Graziani, p.126 / Westend61, p.135 / Nigel Stripe, p.158 / ebrind, p.172 / Enrique Díaz / 7cero, p.174 / MediaProduction, p.181 / FluxFactory, Chapter 4 Opener / ZU_09, p.185 / Courtneyk, p.193 / Ed Freeman, p.208 / Frisco, p.209 / Monty Rakusen, p.221 / Matt Carr, p.235 / Amanda Caroline da Silva, Chapter 5 Opener / Hispanolistic, p.249 / © Jackie Bale, p.255 / Santiago Urquijo, p.261 / EyeEm Mobile GmbH, p.265 / Stephen Frink, p.271 / slowmotiongli, p.283 / Iuliia Burmistrova, Chapter 6 Opener / Paopano, p.331 / yalcinsonat1, p.334 / Group4 Studio, p.335 / AleksandarGeorgiev, p.340 / Achim Thomae, p.346 / mtreasure, p.355 / MirageC, p.361 / Tiina & Geir, p.365 / Thankful Photography, p.372 / gilaxia, p.386 / EvgeniyShkolenko, Chapter 7 Opener / Westend61, p.408 / Jennifer A Smith, p.425 / Photo by Mike Kline (notkalvin), p.43 / LEONELLO CALVETTI, p.435 / ClaudioVentrella, p.441 / Jose A. Bernat Bacete, p.445 / chuchart duangdaw, Chapter 8 Opener / mbbirdy, p.468 / nazar_ab, p.480 / wilatlak villette, p.491 / Santypan, Chapter 9 Opener / Mint Images, p.526 / Marcia Straub, p.533 / Lingbeek, p.566 / Witthaya Prasongsin, p.568 / sarayut Thaneerat, Chapter 10 Opener / Zero Creatives, p.579 / itsabreeze photography, p.593 / Tetra Images, p.595 / Rafe Swan, p.599 / wragg, p.609 / Tom Werner, p.618 / Noel Hendrickson, Chapter 11 Opener / Juan Maria Coy Vergara, p.680 / Mikael Vaisanen, p.685 / RICOWde, p.701.

Every effort has been made to trace and acknowledge copyright. The publisher apologises for any accidental infringement and welcomes information that would redress this situation.

The Victorian Curriculum F-10 content elements are © VCAA, reproduced by permission. The VCAA does not endorse or make any warranties regarding this resource. The Victorian Curriculum F-10 and related content can be accessed directly at the VCAA website.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025
Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Introduction

The second edition of *Essential Mathematics CORE for the Victorian Curriculum* has been significantly revised and updated to suit the teaching and learning of Version 2.0 of the Victorian Curriculum. Many of the established features of the series have been retained, but there have been some substantial revisions, improvements and new elements introduced for this edition across the print, digital and teacher resources.

New content and some restructuring

New content has been added at all year levels. In **Year 7**, there is new content on ratios and proportions, volume of triangular prisms, nets of solids and measurement relating to circles. All geometry topics are now contained in a single chapter (Chapter 7). In **Year 8**, there is new content on order of operations, 3D-coordinates, operations with negative fractions, areas of sectors and composite shapes, Pythagoras' theorem, inequalities, similar figures, two-step experiments and tree diagrams. For **Year 9**, there is new content on errors in measurement, inequalities, factorisation, sampling and proportion, quadratics expressions and parabolas. In **Year 10**, there is new content on composite solids, errors in measurement, networks and logarithmic scales.

Version 2.0 places increased emphasis on **investigations** and **modelling**, and this is covered with new Modelling activities at the end of chapters and revised downloadable Investigations. There are also many new elaborations covering **First Nations Peoples' perspectives** on mathematics, ranging across all six content strands of the curriculum. These are covered in a suite of specialised investigations provided in the Online Teaching Suite.

Other new features

- **Technology and computational thinking** activities have been added to the end of every chapter to address the curriculum's increased focus on the use of technology and the understanding and application of algorithms.
- **Targeted skillsheets** – downloadable and printable – have been written for every lesson in the series, with the intention of providing additional practice for students who need support at the basic skills covered in the lesson, with questions linked to worked examples in the book.
- **Editable PowerPoint lesson summaries** are also provided for each lesson in the series, with the intention of saving the time of teachers who were previously creating these themselves.

Diagnostic assessment tool

Also new for this edition is a flexible, comprehensive diagnostic assessment tool, available through the Online Teaching Suite. This tool, featuring around $10,000$ new questions, allows teachers to set diagnostic tests that are closely aligned with the textbook content, view student performance and growth via a range of reports, set follow-up work with a view to helping students improve, and export data as needed.

Essential Mathematics Core for the Victorian Curriculum Year 9 ISBN 978-1-009-59315-1 © Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Guide to the working programs


Essential Mathematics CORE for the Victorian Curriculum 2.0 contains working programs that are subtly embedded in the exercises. The suggested working programs provide two pathways through the book to allow differentiation for Building and Progressing students.

Each exercise is structured in subsections that match the Victorian Curriculum 2.0 proficiency strands (with Problem-solving and Reasoning combined into one section to reduce exercise length), as well as 'Gold star' (★). The questions* suggested for each pathway are listed in two columns at the top of each subsection.

- The left column (lightest shade) shows the questions in the Building working program.
- The right column (darkest shade) shows the questions in the Progressing working program.

Gradients within exercises and proficiency strands

The working programs make use of two gradients that have been carefully integrated into the exercises. A gradient runs through the overall structure of each exercise – where there's an increasing level of sophistication required as a student progresses through the proficiency strands and then on to the 'Gold Star' question(s) – but also within each proficiency strand; the first few questions in Fluency are easier than the last few, for example, and the first few Problem-solving and reasoning questions are easier than the last few.



The right mix of questions

Questions in the working programs have been selected to give the most appropriate mix of types of questions for each learning pathway. Students going through the Building pathway are given extra practice at the Understanding and basic Fluency questions and only the easiest Problem-solving and reasoning questions. The Progressing pathway, while not challenging, spends a little less time on basic Understanding questions and a little more on Fluency and Problem-solving and reasoning questions. The Progressing pathway also includes the 'Gold star' question(s).

Choosing a pathway

There are a variety of ways of determining the appropriate pathway for students through the course. Schools and individual teachers should follow the method that works best for them. If required, the Warm-up quiz at the start of each chapter can be used as a diagnostic tool. The following are recommended guidelines:

- A student who gets 40% or lower should heavily revise core concepts before doing the Building questions, and may require further assistance.
- A student who gets between 40% and 75% should do the Building questions.
- A student who gets 75% and higher should do the Progressing questions.

For schools that have classes grouped according to ability, teachers may wish to set either the Building or Progressing pathways as the default pathway for an entire class and then make individual alterations depending on student need. For schools that have mixed-ability classes, teachers may wish to set a number of pathways within the one class, depending on previous performance and other factors.

* The nomenclature used to list questions is as follows:

- 3, 4: complete all parts of questions 3 and 4
- 10(½): complete half of the parts from question 10 (a, c, e, ... or b, d, f, ...)
- 4(½), 5: complete half of the parts of question 4 and all parts of question 5
- 1–4: complete all parts of questions 1, 2, 3 and 4
- 2–4(½): complete half of the parts of questions 2, 3 and 4
- – : complete none of the questions in this section.

Essential Mathematics Core for the Victorian Curriculum Year 9
ISBN 978-1-009-59315-1
© Greenwood et al. 2025
Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Guide to this resource

PRINT TEXTBOOK FEATURES

1 NEW **New lessons:** authoritative coverage of new topics in the Victorian Curriculum 2.0 in the form of new, road-tested lessons throughout each book.

2 **Victorian Curriculum 2.0:** content strands, sub-strands and content descriptions are listed at the beginning of the chapter (see the teaching program for more detailed curriculum documents)

3 **In this chapter:** an overview of the chapter contents

4 NEW **Quick reference:** Multiplication, prime number, fraction wall and divisibility rules tables at the back of the book

5 **Chapter introduction:** sets context for students about how the topic connects with the real world and the history of mathematics

6 **Warm-up quiz:** a quiz for students on the prior knowledge and essential skills required before beginning each chapter

7 **Sections labelled to aid planning:** All non-core sections are labelled as 'Consolidating' (indicating a revision section) or with a gold star (indicating a topic that could be considered challenging) to help teachers decide on the most suitable way of approaching the course for their class or for individual students.


8 **Learning intentions:** sets out what a student will be expected to learn in the lesson


9 **Lesson starter:** an activity, which can often be done in groups, to start the lesson

10 **Key ideas:** summarises the knowledge and skills for the section

11 **Worked examples:** solutions and explanations of each line of working, along with a description that clearly describes the mathematics covered by the example. Worked examples are placed within the exercise so they can be referenced quickly, with each example followed by the questions that directly relate to it.

12 **Now you try:** try-it-yourself questions provided after every worked example in exactly the same style as the worked example to give students immediate practice







Essential Mathematics Core for the Victorian Curriculum Year 9 ISBN 978-1-009-59315-1 © Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

13 **Working programs:** differentiated question sets for two ability levels in exercises

14 **Puzzles and games:** in each chapter provide problem-solving practice in the context of puzzles and games connected with the topic


15 **Gentle start to exercises:** the exercise begins at Understanding then Fluency with the first question always linked to the first worked example in the lesson






16 **Chapter checklist:** a checklist of the learning intentions for the chapter, with example questions

17 **Chapter reviews:** with short-answer, multiple-choice and extended-response questions; questions that are 'Gold Star' are clearly signposted





ISBN 978-1-009-59315-1
© Greenwood et al. 2025
Photocopying is restricted under law and this material must not be transferred to another party.

18 **Maths@Work:** a set of extended questions across two pages that give practice at applying the mathematics of the chapter to real-life contexts

19 NEW **Technology and computational thinking:** activity in each chapter addresses the curriculum's increased focus on the use of different forms of technology, and the understanding and implementation of algorithms

20 NEW **Modelling activities:** an activity in each chapter gives students the opportunity to learn and apply the mathematical modelling process to solve realistic problems


INTERACTIVE TEXTBOOK FEATURES

21 NEW **Targeted Skillsheets**, one for each lesson, focus on a small set of related Fluency–style skills for students who need extra support, with questions linked to worked examples

22 **Workspaces:** almost every textbook question – including all working-out – can be completed inside the Interactive Textbook by using either a stylus, a keyboard and symbol palette, or uploading an image of the work

23 **Self-assessment:** students can then self-assess their own work and send alerts to the teacher. See the Introduction on page x for more information

24 **Interactive question tabs** can be clicked on so that only questions included in that working program are shown on the screen




25 **HOTmaths resources:** a huge catered library of widgets, HOTsheets and walkthroughs seamlessly blended with the digital textbook

26 **Desmos graphing calculator**, scientific calculator and geometry tool are always available to open within every lesson


27 **Scorcher:** the popular competitive game

28 **Worked example videos:** every worked example is linked to a high-quality video demonstration, supporting both in-class learning and the flipped classroom



29 A revised set of **differentiated auto-marked practice quizzes** per lesson with saved scores

30 **Auto-marked maths** literacy activities test students on their ability to understand and use the key mathematical language used in the chapter



31 **Auto-marked prior knowledge pre-test** (the 'Warm-up quiz' of the print book) for testing the knowledge that students will need before starting the chapter

32 **Auto-marked progress quizzes and chapter review questions** in the chapter reviews can be completed online

ISBN 978-1-009-59315-1
© Greenwood et al. 2025
Photocopying is restricted under law and this material must not be transferred to another party.

DOWNLOADABLE PDF TEXTBOOK

33 In addition to the Interactive Textbook, a **PDF version of the textbook** has been retained for times when users cannot go online. PDF search and commenting tools are enabled.

ONLINE TEACHING SUITE

34 NEW **Diagnostic Assessment Tool** included with the Online Teaching Suite allows for flexible diagnostic testing, reporting and recommendations for follow-up work to assist you to help your students to improve

35 NEW **PowerPoint lesson** summaries contain the main elements of each lesson in a form that can be annotated and projected in front of class

36 **Learning Management System** with class and student analytics, including reports and communication tools

37 **Teacher view of students' work and self-assessment** allows the teacher to see their class's workout, how students in the class assessed their own work, and any 'red flags' that the class has submitted to the teacher


38 **Powerful test generator** with a huge bank of levelled questions as well as ready-made tests

39 **Revamped task manager** allows teachers to incorporate many of the activities and tools listed above into teacher-controlled learning pathways that can be built for individual students, groups of students and whole classes


40 **Worksheets, Skill and drill, maths literacy worksheets**, and **two differentiated chapter tests in every chapter**, provided in editable Word documents

41 **More printable resources:** all Pre-tests and Progress quizzes and Applications and problem-solving tasks are provided in printable worksheet versions


34



34



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.



1 Reviewing number

Essential mathematics: why number skills are important

Number skills are fundamental in many aspects of daily life and various professions. Understanding how to work with integers, decimals, fractions, ratios, and rates can make everyday tasks easier and open opportunities in numerous fields.

Retail workers use numerical skills for various tasks including:

- **Cash handling**, where they are responsible for calculating totals, making change, and tracking sales.
- **Pricing and discounts**, when they might apply discounts and adjust prices.
- **Financial reporting**, where they may be summarising daily transactions and reconciling cash registers.

Gamers use numerical skills throughout their gameplay:

- In-game economies require currency conversions and budgeting.
- **Strategic planning** involves calculating damage outputs and optimising time management.
- **Maths puzzles** and pattern recognition often appear in games.

Bakers rely on precise ratios for consistent results:

- The $1 : 2 : 3$ ratio for pie crusts (one part water, two parts fat, three parts flour) ensures the right texture.
- A $1 : 1$ ratio of flour to fat is crucial for shortbread's crumbly consistency.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.



In this chapter

1A Adding and subtracting integers (Consolidating)
1B Multiplying and dividing integers (Consolidating)
1C Rounding decimals and significant figures
1D Rational and irrational numbers (Consolidating)
1E Adding and subtracting fractions (Consolidating)
1F Multiplying and dividing fractions (Consolidating)
1G Ratios (Consolidating)
1H Rates and best buys

Victorian Curriculum 2.0

This chapter covers the following content descriptors in the Victorian Curriculum 2.0:

NUMBER

VC2M9N01

ALGEBRA

VC2M9A06

MEASUREMENT

VC2M9M05

Please refer to the curriculum support documentation in the teacher resources for a full and comprehensive mapping of this chapter to the related curriculum content descriptors.

© VCAA

Online resources

A host of additional online resources are included as part of your Interactive Textbook, including HOTmaths content, video demonstrations of all worked examples, auto-marked quizzes and much more.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Warm-up quiz

1 Arrange the following mathematical terms under four headings: 'Addition', 'Subtraction', 'Multiplication' and 'Division'.

a Sum **b** Total **c** Less than **d** Lots of
e Product **f** Into **g** Take away **h** Difference
i Add **j** Times **k** Minus **l** More than
m Quotient

2 Without using a calculator, find an answer to each of the following.

a 16 less 12 **b** 24 more than 8
c the difference between 12 and 8 **d** increase 45 by 7
e the total of 40, 34 and 0 **f** 9 into 45
g the quotient of 63 and 7 **h** 480 shared between 12

3 Evaluate the following.

a $9 + 47$ **b** $135 - 35$ **c** $19 - 19$ **d** $56 + 89 - 12$
e 9×7 **f** $320 \div 4$ **g** 17×60 **h** $200 - 47 - 100$

4 Use a number line to find:

a $-5 - 7$ **b** $12 - 15$ **c** $-6 + 9$ **d** $-12 + 12$
e $16 - 17$ **f** $-4 + 3$ **g** $-7 + 4 + 3$ **h** $-4 - 4 - 4$

5 Copy and complete each of the following statements.

a $5 + 5 + 5 = \square \times 5$ **b** $-6 - 6 - 6 = \square \times (-6)$
c $9 - (+10) = 9 \square 10$ **d** $12 - (-2) = 12 \square 2$

6 The population of Australia in 2050 is projected to be 26 073 258. Round this number to the nearest:

a ten **b** hundred **c** thousand **d** million

7 Write down the place value of the 5 in each of the following numbers.

a 1256 **b** 345 **c** 5049 **d** 0.56 **e** 0.15 **f** 9.005

8 Arrange the numbers in each of the following sets in descending order (largest to smallest).

a 2.645, 2.654, 2.465 and 2.564 **b** 0.456, 0.564, 0.0456 and 0.654

9 Evaluate each of the following.

a $4.26 + 3.73$ **b** $3.12 + 6.99$ **c** $10.89 - 3.78$

10 Evaluate:

a 7×0.2 **b** 0.3×0.2 **c** 2.3×1.6
d 4.2×3.9 **e** $14.8 \div 4$ **f** $12.6 \div 0.07$

11 Evaluate each of the following.

a 0.345×100 **b** $3.74 \times 100\,000$ **c** $37.54 \div 1000$ **d** $3.7754 \div 100\,000$

12 Complete these equivalent fractions.

a $\frac{1}{2} = \frac{\square}{12}$ **b** $\frac{3}{4} = \frac{\square}{16}$ **c** $\frac{5}{6} = \frac{25}{\square}$ **d** $\frac{\square}{9} = \frac{2}{18}$

13 Find the lowest common denominator for these pairs of fractions.

a $\frac{1}{3}$ and $\frac{1}{5}$ **b** $\frac{1}{6}$ and $\frac{1}{4}$ **c** $\frac{1}{5}$ and $\frac{1}{10}$

14 Find:

a $\frac{3}{7} + \frac{2}{7}$ **b** $2 - \frac{3}{4}$ **c** $4 \div \frac{1}{2}$ **d** $\frac{3}{4} \times \frac{1}{2}$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

1A Adding and subtracting integers

CONSOLIDATING

Learning intentions

- To be able to use a number line for addition and subtraction of integers
- To review the rules for adding and subtracting negative numbers
- To be able to add and subtract a negative integer

Key vocabulary: integer, positive, negative

Integers are the set of positive and negative whole numbers, as well as the number zero.

Being able to work with whole numbers is very important, since whole numbers are used every day for counting, ordering, calculating, measuring and computing.

Landscapers use whole numbers every day to measure garden areas to determine the amount of materials needed.



Lesson starter: Naming groups

Here are some groups of numbers. In groups of two or three, use the correct mathematical terms to describe each group. (Suggestions include: 'multiples of', 'factors of', 'integers', 'squares' and 'cubes'.)

- $2, 4, 6, 8, \ldots$
- $1, 4, 9, 16, \ldots$
- $1, 3, 5, 7, 9, \ldots$
- $1, 2, 3, 4, 5, 6, \ldots$
- $-1, -2, -3, -4, -5, \ldots$
- $1, 8, 27, 64, \ldots$
- $1, 2, 3, 4, 6$ and 12

Key ideas

- **Integers** are the **positive** and **negative** whole numbers, including zero:
  $\ldots, -3, -2, -1, 0, 1, 2, 3, \ldots$
- To add a negative number, you subtract its opposite: $a + (-b) = a - b$

  e.g. $5 + (-7) = 5 - 7$

  $-6 + (-2) = -6 - 2$
- To subtract a negative number, you add its opposite: $a - (-b) = a + b$

  e.g. $5 - (-7) = 5 + 7$

  $-6 - (-2) = -6 + 2$
- Adding or subtracting zero leaves a number unchanged.
  $a + 0 = a$ e.g. $5 + 0 = 5$
  $a - 0 = a$ e.g. $5 - 0 = 5$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Exercise 1A

Understanding

1–3 | 3

1 Match each of the following sentences to the correct setting out on the right.

a	The sum of 5 and 7	**i**	$5+(-7)$
b	The total of negative 5 and 7	**ii**	$5-(-7)$
c	The difference between negative 5 and 7	**iii**	$5+7$
d	The sum of 5 and negative 7	**iv**	$7-(-5)$
e	The difference between 5 and negative 7	**v**	$-5+7$

2 Match each question on the left to an expression or answer on the right.

a	$10+(-7)$	**i**	$-10-7$
b	$-10-(-7)$	**ii**	$-10+7$
c	$10-(-7)$	**iii**	0
d	$-10+(-7)$	**iv**	$10-7$
e	$-10+10$	**v**	$10+7$



Hint for Q2: Recall:
$a+(-b)=a-b$
and
$a-(-b)=a+b$

3 True or false?

a $18+0=18$

b $6-(-4)=6+4$

c $4-(-2)=4-2$

Fluency

4–8(½) | 5–8(½)

Example 1 Using a number line for addition and subtraction of integers

Use a number line to find:

a $-10+8$ **b** $-3-5$


Solution

a



$-10+8=-2$

b



$-3-5=-8$

Explanation

Draw a number line showing -10 on it.

Start at -10 and count up (for addition) eight places to finish at -2.

Draw a number line showing -3 on it.

Count down (for subtraction) five places to -8.


Now you try

Use a number line to find:

a $-7+4$ **b** $-6-4$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

4 Use the number line below to help you find the answers to the following.





Hint for Q4: Move right for addition and left for subtraction.

a $-6+4$ **b** $-6+8$ **c** $0-3$
d $-1+5$ **e** $6-10$ **f** $-2-3$
g $-1-1$ **h** $-3+3$ **i** $-3-3$

5 Use a number line to evaluate the following.

a $9-6$ **b** $9-7$ **c** $9-8$
d $9-9$ **e** $9-10$ **f** $-9-0$
g $-9+8$ **h** $-9-8$ **i** $-6-4$
j $-3+12$ **k** $-8+6$ **l** $-1+12$
m $-12+5$ **n** $-10-8$ **o** $18-19$

6 Find:

a $4+8-7$ **b** $5+6+9$ **c** $-19-1$
d $-9+8+1$ **e** $-12-3+8$ **f** $8+3-5$
g $-12-12$ **h** $15+5-15$ **i** $-6-5-4$



Hint for Q6: Work from left to right.

Example 2 Adding a negative integer

Find $17+(-12)$.

Solution

$17+(-12)=17-12$
$=5$

Explanation

Adding a negative is the same as subtraction:
$17+(-12)=17-12$

Now you try

Find $12+(-8)$.

7 Find:

a $9+(-5)$ **b** $12+(-16)$ **c** $3+(-7)$
d $15+(-24)$ **e** $-9+(-23)$ **f** $-13+(-25)$
g $-100+(-89)$ **h** $56+(-80)$ **i** $-9+(-9)$
j $18+(-18)$ **k** $-245+(-560)$ **l** $98+(-155)$
m $-89+(-78)$ **n** $145+(-3)$ **o** $-567+(-237)$



Hint for Q7:
$9+(-5)=9-5$
$-9+(-23)=-9-23$

Example 3 Subtracting a negative integer

Find $-13-(-9)$.

Solution

$-13-(-9)=-13+9$
$=-4$

Explanation

Subtracting a negative is the same as addition:
$-13-(-9)=-13+9$

Now you try

Find $-18-(-10)$.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

1A

8 Find:

a $-8-(-6)$ **b** $-5-(-9)$ **c** $7-(-6)$
d $2-(-7)$ **e** $12-(-12)$ **f** $-34-(-34)$
g $-35-(-7)$ **h** $-90-(-9)$ **i** $90-(-90)$
j $-68-(-70)$ **k** $-90-(-87)$ **l** $234-(-6)$
m $670-(-85)$ **n** $-6-(-100)$ **o** $-230-(-240)$



Hint for Q8: $-5-(-9)=-5+9$

Problem-solving and reasoning

9, 10 | 9–10(½), 11, 12

9 Simplify each of these using the previous rules.

a $-9-9-9$ **b** $-23-(-8)+12$
c $50-46-(-6)$ **d** $24-(-8)+(-6)$
e $-20-(-5)-(-15)$ **f** $-18-(-6)+(-12)$
g $-90-(-89)-90$ **h** $125-35-(-35)$



Hint for Q9: Work from left to right.

10 Copy and complete each of the following statements.

a $\square+6=8$ **b** $-6+\square=5$
c $12-\square=15$ **d** $-9+\square=-10$
e $13-\square=20$ **f** $10-\square=-1$



Hint for Q10: Try using a number line.

11 What must be added to each of the following to obtain a final result of zero?

a 8 **b** $-7+3$ **c** $-8-5$
d -124 **e** 19 **f** 0
g $12-8+18$ **h** -98 **i** $12-(-14)$

12 The temperature on a mountain top reaches a maximum of -2°C during the day. By night it has dropped 10°C. What is the night-time temperature?

Magic squares with negatives

— | 13

13 In a magic square, each row, column and diagonal add to the same number.

a Copy and complete this magic square.

−8		
	−2	−6
		4



Hint for Q13: What does the diagonal add up to?

b Arrange these 9 integers into a 3 by 3 magic square.
$-13, -10, -7, -4, -1, 2, 5, 8$ and 11

c Complete this 4 by 4 magic square.

5			
0	10		6
−1	4	2	8
		7	−4

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

1B Multiplying and dividing integers

CONSOLIDATING

Learning intentions

- To know the rules for multiplying and dividing integers with the same or different signs
- To be able to multiply and divide with negative integers
- To be able to find the square and cube of negative integers
- To be able to apply order of operations to problems involving negative integers

Key vocabulary: integer, square, cube

Multiplication is a way of writing repeated addition. It is possible to develop rules for multiplying and dividing with negative integers.

Lesson starter: Repeated addition

1 Write each of the following as a multiplication before finding the answer.

a $4+4+4$
b $(-4)+(-4)+(-4)$
c $5+5+5+5$
d $(-5)+(-5)+(-5)+(-5)$
e $(-7)+(-7)+(-7)+(-7)+(-7)$
f $(-8)+(-8)$

2 Use your results from above to answer these divisions.

a $12 \div 4$
b $-12 \div (-4)$
c $-12 \div 3$
d $-20 \div 4$
e $-20 \div (-5)$
f $20 \div 5$
g $-35 \div (-7)$
h $-35 \div 5$
i $-16 \div (-8)$

3 What can you conclude about dividing a negative number by:

a a positive number?
b a negative number?

Key ideas

- When multiplying or dividing two numbers with the same sign, the answer is a positive number.
  $+ \times + = +$ $+ \div + = +$
  $- \times - = +$ $- \div - = +$
  e.g. $-5 \times (-4) = 20$ and $-18 \div (-6) = 3$
- When multiplying or dividing two numbers with different signs, the answer is a negative number.
  $+ \times - = -$ $+ \div - = -$
  $- \times + = -$ $- \div + = -$
  e.g. $-5 \times 4 = -20$ and $18 \div (-6) = -3$
- The **square** of a number is the number multiplied by itself, e.g. $5^2 = 5 \times 5$. The **cube** of a number is $a^3 = a \times a \times a$, e.g. 7^3 is $7 \times 7 \times 7$.
- To square a negative number, use brackets.
  For example, $(-7)^2 = -7 \times (-7) = 49$
  Note that $-7^2 = -49$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

1B

- For cube numbers $-2^3 = (-2)^3 = -8$ since $-2 \times (-2) \times (-2) = -8$
- Order of operations
  - Deal with brackets first
  - Do multiplication and division next, working from left to right
  - Do addition and subtraction last, working from left to right

Exercise 1B

Understanding

1–3 | 3

1 Complete the following statements.

a A negative number multiplied by a negative number equals a _____ number.

b A negative number multiplied by a positive number equals a _____ number.

c A negative number divided by a negative number equals a _____ number.

d A positive number divided by a negative number equals a _____ number.

e A negative number divided by a positive number equals a _____ number.

2 Without actually finding the answer to any of these questions, decide whether the answer to each would be a positive or a negative number.

a $2 \times (-8)$ **b** $78 \times (-1)$

c $56 \div (-2)$ **d** $-90 \div (-10)$

e 8×12 **f** $-18 \times (-9)$

3 Answer true (T) or false (F) to the following.

a $(-3)^2 = 9$ **b** $-3^2 = 9$ **c** $-2^3 = (-2)^3$



Hint for Q3: $(-3)^2 = -3 \times (-3)$, whilein -3^2 square 3 first.

Fluency

4–7(½) | 4–7(½)

Example 4 Multiplying with negatives

Find the value of:

a $-6 \times (-7)$ **b** $8 \times (-12)$

Solution

a $-6 \times (-7) = 42$

Explanation

The product of two negative numbers gives a positive answer.

$- \times - = +$

b $8 \times (-12) = -96$

The product of a positive number and a negative number results in a negative answer.

$+ \times - = -$

Now you try

Find the value of:

a $-8 \times (-4)$ **b** -5×7

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

4 Find these products. Some may need to be done with a calculator.

a -8×6	**b** -10×2	**c** -9×8
d $-9 \times (-8)$	**e** $-15 \times (-3)$	**f** $45 \times (-2)$
g $100 \times (-6)$	**h** $-9 \times (-9)$	**i** -12×12
j -13×11	**k** $11 \times (-13)$	**l** $34 \times (-1)$
m $-24 \times (-3)$	**n** $-18 \times (-5)$	**o** $36 \times (-3)$



Hint for Q4: $+ \times + = +$
$- \times - = +$
$+ \times - = -$
$- \times + = -$

Example 5 Finding squares and cubes of negative numbers

Find the value of:

a $(-5)^2$ **b** $(-6)^3$

Solution	Explanation
a $(-5)^2 = 25$	$(-5)^2$ is the same as $-5 \times (-5)$ The product of two negative numbers gives a positive answer. $- \times - = +$
b $(-6)^3 = -216$	$(-6)^3 = -6 \times (-6) \times (-6)$ Multiply the first two numbers together: $-6 \times (-6) = 36$ Now: $36 \times (-6) = -216$

Now you try

Find the value of:

a $(-8)^2$ **b** $(-3)^3$

5 Find the answers to each of the following. Use a calculator to check your answers.

a $(-6)^2$	**b** $(-8)^2$	**c** $(-12)^2$
d $(-3)^2$	**e** $(-10)^2$	**f** $(-13)^2$
g $(-2)^3$	**h** $(-1)^3$	**i** $(-4)^3$



Hint for Q5:
$(-6)^2 = -6 \times (-6)$
$(-2)^3 = -2 \times (-2) \times (-2)$

Example 6 Dividing with a negative number

Find the value of:

a $-24 \div 8$ **b** $-45 \div (-3)$

Solution	Explanation
a $-24 \div 8 = -3$	The division of two numbers with different signs gives a negative answer. $- \div + = -$
b $-45 \div (-3) = 15$	The division of two numbers with the same sign gives a positive answer. $- \div - = +$

Now you try

Find the value of:

a $-30 \div 6$ **b** $-28 \div (-7)$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

1B

6 Perform the following divisions. Use a calculator if necessary.

a $18 \div (-2)$	**b** $-36 \div (-6)$	**c** $-100 \div (-10)$
d $-50 \div 5$	**e** $-6 \div (-2)$	**f** $-10 \div (-2)$
g $-50 \div (-25)$	**h** $16 \div (-16)$	**i** $24 \div (-3)$
j $-100 \div (-4)$	**k** $312 \div (-3)$	**l** $-45 \div (-9)$
m $185 \div (-5)$	**n** $-428 \div 2$	**o** $-156 \div (-12)$



Hint for Q6: $+ \div + = +$
$- \div - = +$
$+ \div - = -$
$- \div + = -$

Example 7 Applying the order of operations

Find the value of $9 \times (-8) + (5 - 12)$.

Solution	Explanation
$9 \times (-8) + (5 - 12)$	Work out the value of the brackets first: $(5 - 12) = (-7)$
$= 9 \times (-8) + (-7)$	Multiplication is next: $9 \times (-8) = -72$
$= -72 + (-7)$	The addition of a negative is the same as subtraction: $-72 + (-7) = -72 - 7$
$= -79$	

Now you try

Find the value of $8 \div (-2) \times (-8 + 5)$.

7 Follow the order of operations and evaluate:

a $(7 - 3) \times 4$	**b** $(4 + 9) \times (-2)$
c $(6 - 2) \times (-3)$	**d** $-6 + 4 \times (-2)$
e $10 \div (-2) \times 4$	**f** $(-5 - 2) \times 4$
g $6 \times (-3) \times 4$	**h** $-21 \div 7 + (-5)$
i $-45 - 9 \times 5$	**j** $(4 - 6) \times (7 - 12)$
k $(15 - 9) \times (3 - 7)$	**l** $9 + 9 \times (-3)$
m $-(3 - 12)$	**n** $10 - 9 \times 4$



Hint for Q7: The order of operations:
- brackets first, following the order of operations within them
- then $\times$ and $\div$, working from left to right
- then $+$ and $-$, working from left to right.

Problem-solving and reasoning 8, 9 10–12

8 Copy and complete:

a $-5 \times \square = -35$	**b** $\square \div (-4) = 8$	**c** $-10 \times \square = 200$
d $17 \times \square = -68$	**e** $34 \div \square = -34$	**f** $-6 \times \square = -36$
g $\square \div 9 \times (-3) = 3$	**h** $15 \div \square \div 3 = -1$	**i** $-15 \times \square = 225$

9 The sum of two numbers is -3 and their product is -10. What are the two numbers?



Hint for Q9: If the product is negative, what do we know about the 2 numbers?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10 Give the value of two different numbers that when squared each produce an answer of:

a 144

b 64

c 10 000

11 Explain why $-5 \times 4 \div (-3)$ produces a positive answer.

12 Decide whether each of the following would produce a positive or a negative answer.

a $-6 \times (-4) \times 5$

b $-12 \div 4 \times 9 \div (-3)$

c $-8 \div (-2) \times 5 \times (6 - 9)$

d $(-3)^4$

e $(-1)^{201}$

f $-(-5)^3$

Substitutions involving negatives — 13

13 Evaluate these expressions by substituting $a = -2$, $b = 6$ and $c = -3$. Check that you can get the same answer with a calculator.

a $a^2 - b$

b $a - b^2$

c $2c + a$

d $b^2 - c^2$

e $a^3 + c^2$

f $3b + ac$

g $c - 2ab$

h $abc - (ac)^2$



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

1C Rounding decimals and significant figures

Learning intentions

- To know how to round a number to a required number of decimal places
- To understand what is meant by significant figures
- To be able to round a number to a required number of significant figures

Key vocabulary: decimal places, critical digit, significant figure

Numbers with and without decimal places can be rounded. The time for a 100 m sprint race, for example, might be 9.94 seconds, correct to two decimal places.

Another way of rounding numbers is to use significant figures. The number of cubic metres of gravel required for a road, for example, might be calculated as 3485 but is rounded to 3500. This number is written using two significant figures.



To ensure sufficient gravel for road construction, 3485 cubic metres is rounded up to 3500 cubic metres.


Lesson starter: In the middle

For each number given on the left, decide which of the two numbers on the right it is closest to.

a	84	80 or 90
b	856	800 or 900
c	856	850 or 860
d	0.599	0.5 or 0.6
e	1.2099	1.2 or 1.3
f	1.2099	1.20 or 1.21
g	89 543	89 000 or 90 000
h	0.035	0.03 or 0.04

Key ideas

- To round a number to a required number of **decimal places** (number of digits after the decimal point):
  - Locate the digit in the required decimal place.
  - Round down (leave as is) if the next digit (**critical digit**) is 4 or less.
  - Round up (increase by 1) if the next digit is 5 or more.



For example:
- To two decimal places, 1.543 rounds to 1.54 and 32.9283 rounds to 32.93.
- To one decimal place, 0.248 rounds to 0.2 and 0.253 rounds to 0.3.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

- To round a number to a required number of **significant figures**:
  - Locate the first *non-zero* digit counting from left to right.
  - From this first significant digit, count out the number of digits, including zeros.
  - Stop at the required number of digits and round this last digit.
  - Replace any non-significant digits to the left of a decimal point with a zero.

  For example, these numbers are all rounded to 3 significant figures:
  $2.5391 \approx 2.54$ $0.002713 \approx 0.00271$ $568\,810 \approx 569\,000$

Exercise 1C

Understanding

1–4 | 4

1 How many decimal places do each of these numbers have?
a 1.467 **b** 146.1 **c** 0.08 **d** 47 900

2 How many significant figures do each of these numbers have?
a 0.46 **b** 21.46 **c** 0.08 **d** 4906

Hint for Q2: Count the digits starting from the first non-zero digit. These are the significant ones.



3 In the following, the critical digit is circled. Use this to state if you round the previous digit down (leave as is) or up.
a 2.75(4)37 **b** 0.87(7)2 **c** 21.5(5)4

4 Choose the correct answer if the first given number is rounded to three significant figures.
a 32 124 is rounded to 321, 3210 or 32 100
b 431.92 is rounded to 431, 432 or 430
c 5.8871 is rounded to 5.887, 5.88 or 5.89
d 0.44322 is rounded to 0.44, 0.443 or 0.44302
e 0.0019671 is rounded to 0.002, 0.00197 or 0.00196

Fluency

5–6(½), 7, 8 | 5-6(½), 7, 8, 9(½)

Example 8 Rounding to a number of decimal places

Round each of these to two decimal places.
a 256.1793 **b** 0.04459 **c** 4.8972

Solution	Explanation
a $256.1793 \approx 256.18$	$256.\underline{17}93$ The number after the second decimal place is 9, so round up (increase the 7 by 1).
b $0.04459 \approx 0.04$	$0.\underline{04}459$ The number after the second decimal place is 4, so round down.

Continued on next page

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

1C

c $4.8972 \approx 4.90$

$4.8\underline{9}72$

The number after the second decimal place is 7, so round up 89 rounds to 90

Now you try

Round each of these to two decimal places.

a 107.3874 **b** 0.0321 **c** 2.7956

5 Round each of the following numbers to two decimal places.

a 17.962 **b** 11.082
c 72.986 **d** 47.859
e 63.925 **f** 23.807
g 804.5272 **h** 500.5749
i 821.2749 **j** 5810.2539
k 1004.9981 **l** 2649.9974



Hint for Q5: The critical digit will be the third decimal place.

Example 9 Rounding to a number of significant figures

Round each of these numbers to two significant figures.

a 2567 **b** 23 067.453 **c** 0.04059

Solution **Explanation**

a $2567 \approx 2600$

2567
significant digits

The next digit is 6, so round up.

Replace last 2 digits with zeros to maintain place value.

b $23\,067.453 \approx 23\,000$

23067.453
significant digits

The next digit is 0, so round down.

Replace the remaining digits, in front of the decimal point, with zeros.

c $0.04059 \approx 0.041$

Locate the first non-zero digit and count two digits:

0.04059
significant digits

The next digit is 5, so round up.

No extra zeros are needed, as they are after the decimal point.

Now you try

Round each of these numbers to two significant figures.

a 3472 **b** 1792.41 **c** 0.00226

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

6 Round each of these numbers to two significant figures.

a 2436	**b** 35 057.4	**c** 0.06049
d 34.024	**e** 107 892	**f** 0.00245
g 2.0745	**h** 0.7070	**i** 4706
j 59 134	**k** 0.4567	**l** 1.0631



Hint for Q6: Start at the first non-zero digit on the left.

7 Copy and complete this table.

	Number	Rounded to two decimal places	Rounded to two significant figures
a	1.4638		
b	0.0936		
c	23.7124		
d	0.00783		
e	100.465		

8 Round these numbers to the nearest whole number.

a 6.814 **b** 73.148 **c** 129.94 **d** 36 200.49

9 Round these numbers to one significant figure.

a 32 000	**b** 194.2	**c** 0.0492
d 0.0006413	**e** 4793	**f** 890
g 0.89	**h** 0.000304	**i** 0.95



Hint for Q9: $48.06 = 50$ to one significant figure. $4730 = 5000$ to one significant figure. $0.638 = 0.6$ to one significant figure.

Problem-solving and reasoning

10–12 | 11–14

Example 10 Estimating using significant figures

Estimate the answer to $27 + 1329.5 \times 0.0064$ by rounding each number in the problem to one significant figure. Use your calculator to check how reasonable your answer is.

Solution

$27 + 1329.5 \times 0.0064$
$\approx 30 + 1000 \times 0.006$
$= 30 + 6$
$= 36$

The estimated answer is reasonable.

Explanation

Round each number to one significant figure and evaluate.
Recall multiplication occurs before addition.

By calculator (to 1 d.p.):
$27 + 1329.5 \times 0.0064 = 35.5$

Now you try

Estimate the answer to $32 - 117 \div 5.2$ by rounding each number in the problem to one significant figure. Use your calculator to check how reasonable your answer is.

10 Estimate the answers to the following by rounding each number in the problem to one significant figure. Check how reasonable your answer is with a calculator.


a $567 + 3126$	**b** $795 - 35.6$	**c** 97.8×42.2
d $965.98 + 5321 - 2763.2$	**e** $4.23 - 1.92 \times 1.827$	**f** $17.43 - 2.047 \times 8.165$
g $0.0704 + 0.0482$	**h** 0.023×0.98	**i** $0.38 \div 1.9$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

1C

11 An electronic timer records the time for a running relay between two teams, A and B. Team A's time is 54.283 seconds and team B's time is 53.791 seconds. Find the difference in the times for teams A and B if the times were written down using:

a one decimal place
b four significant figures
c two significant figures
d one significant figure



12 One tonne (1000 kg) of soil is to be equally divided between 7 garden beds. How much soil does each garden bed get? Write your answer in tonnes rounded to the nearest kilogram.

13 A scientific experiment uses very small amounts of magnesium (0.0025 g) and potassium (0.0062 g). Why does it make sense to use two significant figures instead of two decimal places when recording numbers in a situation like this?



14 Should 2.14999 be rounded down or up if it is to be rounded to one decimal place? Give reasons.

*n*th decimal place and π — 15, 16

15 $\frac{2}{11}$ can be written as 0.18181818 correct to eight decimal places.

a Using the decimal pattern described, find the digit in the:
i 20th decimal place **ii** 45th decimal place **iii** 1000th decimal place

b Express $\frac{1}{7}$ as a decimal correct to 13 decimal places.

c Using the decimal pattern from part **b** find the digit in the:
i 20th decimal place **ii** 45th decimal place **iii** 1000th decimal place

16 π is a decimal that is non-terminating and has no pattern. $\pi = 3.141592653\ldots$

a How many decimal places can you remember?
b Quiz yourself and your classmates.
c Use the internet to find π correct to 100 decimal places.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

1D Rational and irrational numbers

CONSOLIDATING

Learning intentions

- To know the difference between rational and irrational numbers
- To be able to convert between fractions and decimals
- To know the notation for recurring decimals
- To be able to compare fractions using a common denominator

Key vocabulary: real numbers, rational numbers, irrational numbers, numerator, denominator, proper fraction, improper fraction, mixed number, highest common factor, recurring decimal, equivalent fractions


Rational numbers are any numbers that can be written as a fraction in the form $\frac{a}{b}$, where a and b are integers and $b \neq 0$. Pythagoras and his followers discovered around 500 BCE that not all numbers are rational. Pi (π) and $\sqrt{2}$ are examples of these numbers. When written as decimals, they do not terminate or repeat. We call them irrational numbers.

This is $\sqrt{2}$ to 100 decimal places:

1.41421356237309504880168872420969807856967187537694807317667973799073247846210703885038753432764157273


Lesson starter: The real number system

Copy the diagram on the right and insert these words correctly: rational, irrational, real, integers, counting.



Key ideas

- The **real numbers** (any positive or negative number or zero) can be classified as rational or irrational numbers.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025
Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

1D

- An infinite decimal is one where the decimal places continue forever.
- **Equivalent fractions** have the same value. For example: $\frac{2}{3} = \frac{6}{9}$
- Fractions can be simplified by dividing the **numerator** and **denominator** by their **highest common factor** (HCF).
- If $\frac{a}{b}$ is a **proper fraction**, then $a < b$. For example: $\frac{2}{7}$
- If $\frac{a}{b}$ is an **improper fraction**, then $a \geq b$. For example: $\frac{10}{3}$
- A **mixed number** is written as a whole number plus a proper fraction. For example: $2\frac{3}{5}$

  As a mixed number $\frac{11}{3}$ is $3\frac{2}{3}$ and as an improper fraction $4\frac{1}{5}$ is $\frac{21}{5}$
- Fractions can be compared using a common denominator. This should be the lowest common multiple of both denominators.
- A dot or bar is used to show a pattern in a **recurring decimal** number.
  For example: $\frac{1}{6} = 0.16666... = 0.1\dot{6}$ or $\frac{3}{11} = 0.272727... = 0.\overline{27}$

Exercise 1D

Understanding

1–3 | 1

1 Answer the following as true or false.

a $\sqrt{3}$ is an irrational number.

b $\sqrt{4}$ is an irrational number.

c $0.\dot{3}$ is a rational number.

d $\frac{4}{5}$ is a rational number.



Hint for Q1: An irrational number as a decimal is never-ending and has no pattern.

2 State whether each of the following is a proper fraction, improper fraction or mixed number.

a $\frac{2}{3}$

b $4\frac{1}{7}$

c $\frac{11}{6}$

d $\frac{1}{5}$

3 Write each of these in the form $\frac{a}{b}$.

a 0.7 **b** 0.3

c 0.23 **d** 0.09



Hint for Q3: $0.1 = \frac{1}{10}$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Fluency

4, 5, 6–9(½) | 4, 5, 6–9(½)

Example 11 Identifying rational numbers

Choose the rational numbers in the following list.

$\sqrt{9}, \sqrt{2}, 0.6, \pi, -\frac{3}{4}$

Solution

$\sqrt{9}, 0.6$ and $-\frac{3}{4}$ are rational numbers.

Explanation

$\sqrt{9} = 3 = \frac{3}{1}$ $\therefore$ rational (a fraction)

$0.6 = \frac{6}{10}$ $\therefore$ rational

$-\frac{3}{4}$ is also rational.

π and $\sqrt{2}$ are infinite non-recurring decimals.

Now you try

Choose the rational numbers in the following list.

$2\pi, \frac{5}{8}, 0.32, \sqrt{3}, -\sqrt{25}$

4 Which of the following are rational numbers?

a 0.1	**b** 20%	**c** $6\frac{1}{4}$
d $\frac{2}{5}$	**e** $\frac{2}{3}$	**f** 0.001
g $\sqrt{64}$	**h** $\sqrt{7}$	**i** 4^2
j π	**k** $0.\dot{2}$	**l** $\sqrt{11}$



Hint for Q4: Rational numbers can be written as a fraction.

Example 12 Converting between mixed numbers and improper fractions

Express

a $\frac{11}{4}$ as a mixed number

b $2\frac{3}{5}$ as an improper fraction

Solution

a $\frac{11}{4} = 2\frac{3}{4}$

b $2\frac{3}{5} = \frac{13}{5}$

Explanation

4 divides into 11, 2 whole times. $2 \times 4 = 8$, leaving a remainder of 3.

To obtain the numerator $2 \times 5 + 3 = 13$.

Now you try

Express

a $\frac{17}{5}$ as a mixed number

b $2\frac{1}{4}$ as an improper fraction

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

1D

5 Complete the following conversions.

a Express each of the following improper fractions as mixed numbers.

i $\frac{12}{5}$ **ii** $\frac{17}{6}$ **iii** $\frac{23}{4}$ **iv** $\frac{9}{8}$

b Express each of the following mixed numbers as improper fractions.

i $3\frac{1}{5}$ **ii** $6\frac{2}{7}$ **iii** $4\frac{4}{5}$ **iv** $10\frac{3}{8}$

Example 13 Writing fractions as decimals

Write $4\frac{3}{8}$ as a decimal.

Solution

$$8\overline{)3.^{3}0^{6}0^{4}0}\quad 0.\ 3\ 7\ 5$$

$4\frac{3}{8} = 4.375$

Explanation

Find a decimal for $\frac{3}{8}$ by dividing 8 into 3 using the short division algorithm.

Now you try

Write $2\frac{5}{16}$ as a decimal.

6 Write these fractions as decimals.

a $\frac{11}{4}$ **b** $\frac{7}{20}$ **c** $3\frac{2}{5}$ **d** $\frac{15}{8}$

e $2\frac{5}{8}$ **f** $3\frac{4}{5}$ **g** $\frac{37}{16}$ **h** $\frac{7}{28}$

Hint for Q6: Use short division. $\frac{11}{4}$ means $11 \div 4$



Example 14 Writing recurring decimals from fractions

Write $\frac{5}{13}$ as a recurring decimal.

Solution

$$13\overline{)5.^{5}0^{11}0^{6}0^{8}0^{2}0^{7}0^{5}0}\quad 0.\ 3\ \ 8\ 4\ 6\ 1\ 5\ 3$$

$\frac{5}{13} = 0.\overline{384615}$

Explanation

Divide 13 into 5 and continue until the pattern repeats.
Add a bar over the repeating pattern.

Writing $0.\dot{3}84615\dot{3}$ is an alternative.

Now you try

Write $\frac{5}{12}$ as a recurring decimal.

7 Write these fractions as recurring decimals.

a $\frac{3}{11}$ **b** $\frac{7}{9}$ **c** $\frac{9}{7}$ **d** $\frac{7}{12}$

e $\frac{10}{9}$ **f** $3\frac{5}{6}$ **g** $7\frac{4}{15}$ **h** $\frac{29}{11}$

Hint for Q7: Check using your calculator.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Example 15 Writing decimals as fractions

Write these decimals as fractions.

a 0.24 **b** 2.385

Solution

a $0.24 = \frac{24}{100}$

$= \frac{6}{25}$

b $2.385 = \frac{2385}{1000}$

$= \frac{477}{200}$

$= 2\frac{77}{200}$

OR

$2\frac{385}{1000}$

$= 2\frac{77}{200}$

Explanation

Write as a fraction using the smallest place value (hundredths) and then simplify by dividing by the HCF of 4.

The smallest place value is thousandths.

Simplify to an improper fraction or a mixed number, using the HCF of 5.

Now you try

Write these decimals as fractions.

a 0.45 **b** 3.225

8 Write these decimals as fractions.

a 0.35	**b** 0.06	**c** 3.7
d 0.56	**e** 1.07	**f** 0.075
g 3.32	**h** 7.375	**i** 2.005
j 10.044	**k** 6.45	**l** 2.101



Hint for Q8: Divide by 100 when there are two decimal places.

Example 16 Comparing fractions

Decide which is the larger fraction of the following.

$\frac{7}{12}$ or $\frac{8}{15}$

Solution

LCM of 12 and 15 is 60.

$\frac{7}{12} = \frac{35}{60}$ and $\frac{8}{15} = \frac{32}{60}$

$\therefore \frac{7}{12} > \frac{8}{15}$

Explanation

Find the lowest common multiple of the two denominators (lowest common denominator).

Write each fraction as an equivalent fraction using the common denominator. Then compare numerators (i.e. $35 > 32$) to determine the larger fraction.

Now you try

Decide which is the larger fraction of the following.

$\frac{4}{9}$ or $\frac{5}{12}$

9 Decide which is the larger fraction in the following pairs.

a $\frac{3}{4}, \frac{5}{6}$	**b** $\frac{13}{20}, \frac{3}{5}$	**c** $\frac{7}{10}, \frac{8}{15}$	**d** $\frac{5}{12}, \frac{7}{18}$
e $\frac{7}{16}, \frac{5}{12}$	**f** $\frac{26}{35}, \frac{11}{14}$	**g** $\frac{7}{12}, \frac{19}{30}$	**h** $\frac{7}{18}, \frac{11}{27}$



Hint for Q9: First write both fractions using a common denominator.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

1D

Problem-solving and reasoning 10–12 11, 13, 14

10 Express the following quantities as simplified fractions.

a \$45 out of \$100 **b** 12 kg out of 80 kg

c 64 baskets out of 90 shots in basketball **d** 115 mL out of 375 mL

11 These sets of fractions form a pattern. Find the next two fractions in the pattern.

a $\frac{1}{3}, \frac{5}{6}, \frac{4}{3}, \frac{\square}{\square}, \frac{\square}{\square}$ **b** $\frac{6}{5}, \frac{14}{15}, \frac{2}{3}, \frac{\square}{\square}, \frac{\square}{\square}$ **c** $\frac{2}{3}, \frac{3}{4}, \frac{5}{6}, \frac{\square}{\square}, \frac{\square}{\square}$ **d** $\frac{1}{2}, \frac{4}{7}, \frac{9}{14}, \frac{\square}{\square}, \frac{\square}{\square}$

12 The 'Weather forecast' website says there is a 0.45 chance that it will rain tomorrow. The 'Climate control' website says that the chance of rain is $\frac{14}{30}$. Which website gives the least chance that it will rain?



Hint for Q12: Express both values as decimals or as fractions with a common denominator



13 A jug has 400 mL of $\frac{1}{2}$ strength orange juice. The following amounts of full strength juice are added to the mix. Find a fraction to describe the strength of the orange drink after the full strength juice is added.

a 100 mL **b** 50 mL **c** 120 mL **d** 375 mL

14 If x is an integer, determine the values that x can take in the following.

a The fraction $\frac{x}{3}$ is a number between (and not including) 10 and 11.

b The fraction $\frac{x}{7}$ is a number between (and not including) 5 and 8.

c The fraction $\frac{34}{x}$ is a number between 6 and 10.



Hint for Q14: You may have more than one answer.

Converting recurring decimals to fractions — 15

15 Here are two examples of how to convert recurring decimals to fractions.

$0.\dot{6} = 0.6666\ldots$

Let $x = 0.6666\ldots$ (1)

$10x = 6.6666\ldots$ (2)

(2) − (1) $9x = 6$

$x = \frac{6}{9} = \frac{2}{3}$

$\therefore \quad 0.\dot{6} = \frac{2}{3}$

$1.\overline{27} = 1.272727\ldots$

Let $x = 1.272727\ldots$ (1)

$100x = 127.2727\ldots$ (2)

(2) − (1) $99x = 126$

$x = \frac{126}{99}$

$\therefore \quad 1.\overline{27} = \frac{126}{99} = 1\frac{27}{99} = 1\frac{3}{11}$

Convert these recurring decimals to fractions using the above method.

a $0.\dot{8}$ **b** $1.\dot{2}$ **c** $0.\overline{81}$ **d** $3.\overline{43}$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

1E Adding and subtracting fractions

CONSOLIDATING

Learning intentions

- To understand that common denominators are needed to add or subtract fractions
- To be able to find the lowest common denominator to add or subtract fractions
- To be able to add and subtract mixed numbers

Key vocabulary: equivalent fractions, numerator, denominator, lowest common denominator


As you will remember, fractions represent parts of a whole.


Fractions, like other numbers, can be added and subtracted.


To add or subtract fractions they need to be the same type; that is, they both need to be eighths, quarters, thirds or tenths etc. They need to have a common denominator.


Lesson starter: Shade the fraction


Shade the fraction suggested by each of these additions and subtractions.


a $\frac{1}{4}+\frac{1}{4}=$ 

b $\frac{1}{4}+\frac{1}{2}=$ 

c $\frac{3}{4}-\frac{1}{2}=$ 


d $\frac{5}{8}+\frac{1}{4}=$ 

e $\frac{1}{2}+\frac{1}{3}=$ 

f $\frac{1}{3}-\frac{1}{4}=$ 

Key ideas

- Equivalent fractions are created by multiplying or dividing both the numerator and the denominator by the same factor.
  For example:



- To add or subtract fractions, the denominators need to be the same.
  - If the denominators are the same, add/subtract the numerators, keeping the denominator unchanged.
    For example: $\frac{3}{5}+\frac{1}{5}=\frac{4}{5}$
  - If the denominators are different, find the **lowest common denominator** (LCD) using equivalent fractions. Then add/subtract the numerators, remembering to simplify your answer where possible.
    For example: $\frac{1}{2}+\frac{2}{3}=\frac{3}{6}+\frac{4}{6}$ The LCD of halves and thirds is sixths.

$$=\frac{7}{6}$$

$$=1\frac{1}{6}$$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Exercise 1E

Understanding

1–4 | 4

1 Copy and complete these equivalent fractions.

a $\frac{3}{4}=\frac{\square}{16}$ **b** $\frac{3}{5}=\frac{\square}{10}$ **c** $\frac{5}{6}=\frac{\square}{18}$

d $\frac{3}{7}=\frac{21}{\square}$ **e** $\frac{4}{5}=\frac{12}{\square}$ **f** $\frac{5}{7}=\frac{25}{\square}$

Hint for Q1: What you do to the numerator must also be done to the denominator, and vice versa.



2 True or false?

a $\frac{7}{9}+\frac{1}{9}=\frac{8}{9}$ **b** $\frac{6}{9}-\frac{1}{3}=\frac{5}{6}$

3 Copy and complete.

a $\frac{3}{10}+\frac{2}{5}$

$=\frac{3}{10}+\frac{\square}{10}$

$=\frac{\square}{10}$

b $\frac{1}{2}-\frac{3}{8}$

$=\frac{\square}{8}-\frac{3}{8}$

$=\frac{\square}{8}$

c $1\frac{1}{2}+\frac{1}{4}$

$=1\frac{2}{\square}+\frac{1}{4}$

$=1\frac{\square}{\square}$



4 What is the LCD needed if these fractions are to be added or subtracted?

a $\frac{1}{2}, \frac{1}{3}$ **b** $\frac{3}{7}, \frac{5}{9}$ **c** $\frac{3}{10}, \frac{8}{15}$

d $\frac{1}{2}, \frac{3}{8}$ **e** $\frac{9}{11}, \frac{4}{33}$ **f** $\frac{5}{12}, \frac{7}{30}$

Hint for Q4: The LCD is the lowest common multiple of both denominators.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Fluency

5–7(½) 5–7(½)

Example 17 Adding and subtracting with same denominators

Find:

a $\frac{5}{11}+\frac{8}{11}$ b $1\frac{3}{8}-\frac{5}{8}$

Solution

a $\frac{5}{11}+\frac{8}{11}=\frac{13}{11}$

$=1\frac{2}{11}$

b $1\frac{3}{8}-\frac{5}{8}=\frac{11}{8}-\frac{5}{8}$

$=\frac{6}{8}$

$=\frac{3}{4}$

Explanation

Denominators are the same (elevenths).

Keep the denominator and add the numerators.

Write as a mixed number.

Write $1\frac{3}{8}$ as an improper fraction.

Denominators are the same (eighths). Subtract the numerators.

Simplify the answer by dividing by the HCF (2).

Now you try

Find:

a $\frac{3}{7}+\frac{5}{7}$ b $1\frac{1}{5}-\frac{4}{5}$

5 Find:

a $\frac{2}{5}+\frac{1}{5}$ b $\frac{3}{8}+\frac{2}{8}$ c $\frac{9}{10}-\frac{2}{10}$ d $\frac{2}{7}+\frac{2}{7}$

e $\frac{13}{17}-\frac{8}{17}$ f $\frac{3}{5}-\frac{2}{5}$ g $\frac{11}{6}-\frac{1}{6}$ h $\frac{13}{8}-\frac{3}{8}$

i $\frac{4}{5}+\frac{3}{5}$ j $1\frac{1}{5}+\frac{3}{5}$ k $2\frac{2}{3}-\frac{1}{3}$ l $1\frac{1}{8}-\frac{5}{8}$

m $1\frac{1}{10}-\frac{7}{10}$ n $3\frac{1}{4}+1\frac{1}{4}$ o $2\frac{3}{10}-\frac{9}{10}$ p $3\frac{7}{8}+2\frac{5}{8}$



Hint for Q5: Writing mixed numbers as improper fractions may help.

Example 18 Adding with different denominators

Evaluate $\frac{1}{2}+\frac{3}{5}$.

Solution

$\frac{1}{2}+\frac{3}{5}$

$=\frac{5}{10}+\frac{6}{10}$

$=\frac{11}{10}$ or $1\frac{1}{10}$

Explanation

The lowest common multiple of the denominators of 2 and 5 is 10. Rewrite as equivalent fractions using a denominator of 10: $\frac{1\times5}{2\times5}=\frac{5}{10}$ and $\frac{3\times2}{5\times2}=\frac{6}{10}$.

Add the numerators.

Now you try

Evaluate $\frac{4}{7}+\frac{1}{3}$.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

1E

6 Evaluate the following.

a $\frac{1}{2}+\frac{1}{8}$ **b** $\frac{1}{2}-\frac{1}{6}$ **c** $\frac{3}{10}+\frac{1}{5}$ **d** $\frac{4}{3}-\frac{5}{6}$

e $\frac{3}{2}-\frac{3}{5}$ **f** $\frac{1}{3}+\frac{1}{4}$ **g** $\frac{3}{5}+\frac{1}{10}$ **h** $\frac{3}{4}-\frac{3}{5}$

i $\frac{5}{6}-\frac{3}{8}$ **j** $\frac{2}{3}+\frac{1}{2}$ **k** $\frac{5}{3}-\frac{2}{7}$ **l** $\frac{3}{8}+\frac{1}{3}$

m $\frac{2}{9}+\frac{1}{12}$ **n** $\frac{9}{21}+\frac{1}{7}$ **o** $\frac{27}{50}-\frac{7}{20}$ **p** $\frac{11}{30}-\frac{7}{45}$



Hint for Q6: First express each fraction using the LCD.

Example 19 Adding and subtracting mixed numbers

Evaluate:

a $1\frac{2}{3}+4\frac{5}{6}$ **b** $3\frac{2}{5}-2\frac{3}{4}$

Solution

a $1\frac{2}{3}+4\frac{5}{6}=\frac{5}{3}+\frac{29}{6}$

$=\frac{10}{6}+\frac{29}{6}$

$=\frac{39}{6}$

$=\frac{13}{2}$ or $6\frac{1}{2}$

Explanation

Change each mixed number to an improper fraction.

Remember that the lowest common denominator of 3 and 6 is 6. Change $\frac{5}{3}$ to an equivalent fraction with denominator 6 then add the numerators and simplify.

b $3\frac{2}{5}-2\frac{3}{4}=\frac{17}{5}-\frac{11}{4}$

$=\frac{68}{20}-\frac{55}{20}$

$=\frac{13}{20}$

Convert to improper fractions and then express as equivalent fractions with the same denominator.

Subtract the numerators.

Now you try

Evaluate:

a $2\frac{3}{4}+1\frac{5}{8}$ **b** $4\frac{1}{3}-3\frac{4}{5}$

7 Evaluate the following.

a $1\frac{1}{3}+2\frac{1}{2}$ **b** $1\frac{1}{3}-\frac{3}{4}$ **c** $5\frac{1}{2}-2\frac{7}{10}$ **d** $2\frac{1}{3}+\frac{1}{4}$

e $4\frac{1}{2}+2\frac{1}{4}$ **f** $6\frac{1}{3}-2\frac{1}{2}$ **g** $1\frac{3}{8}-1\frac{1}{4}$ **h** $2\frac{3}{4}-1\frac{1}{3}$

i $4\frac{1}{10}-2\frac{2}{5}$ **j** $1\frac{7}{12}-\frac{2}{3}$ **k** $3\frac{1}{5}+1\frac{1}{2}$ **l** $2\frac{5}{6}-1\frac{1}{4}$



Hint for Q7: First rewrite using improper fractions.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Problem-solving and reasoning 8–10 9–12

8 To remove impurities a mining company filters $3\frac{1}{2}$ tonnes of raw material. If $2\frac{5}{8}$ tonnes are removed, what quantity of material remains?





Hint for Q8: Decide which mathematical operation is involved first.

9 When a certain raw material is processed it produces $3\frac{1}{7}$ tonnes of mineral and $2\frac{3}{8}$ tonnes of waste. How many tonnes of raw material were processed?

10 The ingredients in a punch recipe were: $2\frac{1}{4}$ L of apple juice, $1\frac{1}{2}$ L of guava juice and $1\frac{1}{5}$ L of lemonade.

a How much punch is produced?

b If a cup holds 150 mL, how many cups can this punch serve?



11 Clarissa owns $\frac{3}{10}$ of a company, Sally owns another $\frac{1}{4}$ of the company and Keith owns $\frac{2}{5}$ of it. The bank owns the rest. How much of the company does the bank own?

12 Here is an example involving the subtraction of fractions where improper fractions are not used.

$$2\frac{1}{2} - 1\frac{1}{3} = 2\frac{3}{6} - 1\frac{2}{6}$$

$$= 1\frac{1}{6}$$

Try this technique on the following problem and explain the difficulty that you find.

$2\frac{1}{3} - 1\frac{1}{2}$

Which fractions do you choose? — 13

13 The six fractions listed below are each used exactly once in part **a**.

$\frac{1}{8}, \frac{5}{12}, \frac{1}{2}, \frac{2}{3}, \frac{3}{4}, \frac{5}{6}$

a Arrange the six fractions correctly so that each equation produces the required answer. Use each fraction only once.

i $\square + \square = \frac{7}{8}$ **ii** $\square - \square = \frac{1}{4}$ **iii** $\square - \square = \frac{1}{3}$

b Which of the fractions, when added, produce an answer of 2?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Progress quiz

1A **1** Find the value of:

a $-15+7$ **b** $-10-4$ **c** $18-(-7)$

d $12+(-4)$ **e** $-8+5-(-7)$ **f** $-9-(-6)$

1B **2** Find the value of:

a $-7\times(-6)$ **b** $12\times(-5)$ **c** $-56\div 7$

d $-48\div(-6)$ **e** $(-4)^3$ **f** $(-9)^2$

1B **3** Apply the order of operations to find the value of:

a $3\times(4-9)+6$ **b** $7-3\times(-4)$ **c** $(6-10)\div(-2)-9$

1C **4** **a** Round each of the following to two decimal places.

i 24.187 **ii** −6.143988 **iii** 7.595699

b Round each of the following to two significant figures.

i 3.268 **ii** 237 **iii** 7932.8 **iv** 0.0078812

1C **5** Estimate the following using one significant figure rounding of each number. Check how reasonable your answer is with a calculator.

a $242-121$ **b** $93.4-9.7\times 0.0234$

1D **6** Write these fractions as decimals, parts **c** and **d** are recurring decimals.

a $\frac{5}{8}$ **b** $3\frac{8}{25}$ **c** $\frac{7}{12}$ **d** $2\frac{7}{11}$

1D **7** Write these decimals as simplified fractions.

a 0.8 **b** 0.36 **c** 0.004 **d** 4.205

1D **8** Arrange these fractions in ascending (smallest to largest) order.

$\frac{11}{24}, \frac{3}{8}, \frac{5}{12}$

1E **9** Evaluate the following.

a $\frac{3}{8}+\frac{2}{8}$ **b** $\frac{2}{3}-\frac{4}{7}$ **c** $2\frac{3}{10}+1\frac{1}{4}$ **d** $4\frac{1}{5}-2\frac{3}{4}$

1E **10** A cake recipe requires $\frac{3}{4}$ cup of sugar, $2\frac{1}{3}$ cups of flour and $\frac{1}{2}$ cup of cocoa. How many cups of ingredients are needed in total?



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

1F Multiplying and dividing fractions

CONSOLIDATING

Learning intentions

- To review how to multiply fractions
- To know to cancel any common factors between numerators and denominators before multiplying
- To know that 'of' means multiply
- To be able to find the reciprocal of a fraction
- To know how to divide by a fraction by multiplying by its reciprocal
- To know to convert mixed numbers to improper fractions before multiplying or dividing

Key vocabulary: numerator, denominator, highest common factor, reciprocal, proper fraction, improper fraction, mixed number

A series of steps can be followed to make multiplying and dividing fractions easier. This includes cancelling, which reduces the size of the numbers inside the fractions.

Lesson starter: How much is left?

How much is left for Kimberly to eat if:

- Tom eats half the tart
- then Sarah eats half of what's left
- then Zara eats a third of the remaining section?

To determine Kimberley's share of the tart, each step involves multiplying the remaining portion by a fraction.



Key ideas

- To multiply fractions (proper or improper), multiply the numerators together and multiply the denominators together.
  - In general:

$$\frac{a}{b} \times \frac{c}{d} = \frac{a \times c}{b \times d}$$

For example:

$$\frac{2}{5} \times \frac{3}{7} = \frac{2 \times 3}{5 \times 7} = \frac{6}{35}$$

- To divide a number by a fraction, multiply by its reciprocal.
  - In general:

$$\frac{a}{b} \div \frac{c}{d} = \frac{a}{b} \times \frac{d}{c} = \frac{a \times d}{b \times c}$$

For example:

$$\frac{2}{3} \div \frac{5}{6} = \frac{2}{3} \times \frac{6}{5} = \frac{2}{\cancel{3}_{1}} \times \frac{\cancel{6}^{2}}{5} = \frac{4}{5}$$

- Cancel any common factors in the numerator with any common factors in the denominator before multiplying.

$$\frac{\cancel{15}^{5}}{\cancel{8}_{4}} \times \frac{\cancel{12}^{1}}{\cancel{15}_{3}} = \frac{1}{4} \times \frac{1}{3} = \frac{1}{12}$$

- *Of* means multiply.
  For example:

$$\frac{1}{3} \text{ of } 24 = \frac{1}{3} \times 24$$

- To find the **reciprocal** of a fraction, you simply invert it (that is, flip the proper or improper fraction upside down).
  - The reciprocal of 2 is $\frac{1}{2}$ since $2 = \frac{2}{1}$.
  - The reciprocal of $\frac{3}{4}$ is $\frac{4}{3}$.
  - The reciprocal of $1\frac{1}{5}$ $\left(\text{or } \frac{6}{5}\right)$ is $\frac{5}{6}$.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Exercise 1F

Understanding

1–3 | 3

1 State if the following are true (T) or false (F).

a $\frac{2}{5} \times \frac{1}{3} = \frac{2}{15}$

b $\frac{1}{6}$ of $30 = 6 \times 30$

c $\frac{3}{7} \div \frac{2}{5} = \frac{3}{7} \times \frac{5}{2}$

2 Find:

a $\frac{1}{2}$ of 24 **b** $\frac{1}{3}$ of 12

c $\frac{3}{4}$ of 16 **d** $\frac{9}{10}$ of 200

3 Find the reciprocal of:

a $\frac{3}{4}$ **b** $\frac{1}{7}$ **c** 6 **d** $1\frac{2}{3}$



Hint for Q3: Whole numbers can be written with a denominator of $1. \; 6 = \frac{6}{1}$

Fluency

4–7(½) | 4–7(½)

Example 20 Multiplying with proper fractions

Evaluate the following.

a $\frac{2}{3} \times \frac{5}{7}$ **b** $\frac{4}{5} \times \frac{25}{32}$

Solution

a $\frac{2}{3} \times \frac{5}{7} = \frac{10}{21}$

b $\frac{\cancel{4}^{1}}{\cancel{5}_{1}} \times \frac{\cancel{25}^{5}}{\cancel{32}_{8}} = \frac{1}{1} \times \frac{5}{8}$

$= \frac{5}{8}$

Explanation

No common factors therefore multiply the numerators together and multiply the denominators together.

$\frac{2}{3} \times \frac{5}{7} = \frac{2 \times 5}{3 \times 7}$

Cancel the common factors between the numerators and the denominators.

Multiply the numerators and then multiply the denominators.

$\frac{1 \times 5}{1 \times 8}$

Now you try

Evaluate the following.

a $\frac{3}{7} \times \frac{4}{5}$ **b** $\frac{3}{8} \times \frac{20}{27}$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

4 Evaluate the following.

a $\frac{3}{5} \times \frac{1}{4}$ **b** $\frac{2}{3} \times \frac{2}{5}$ **c** $\frac{3}{4} \times \frac{5}{7}$ **d** $\frac{1}{2} \times \frac{1}{3}$

e $\frac{1}{4} \times \frac{1}{5}$ **f** $\frac{1}{9} \times \frac{2}{3}$ **g** $\frac{5}{8} \times \frac{16}{17}$ **h** $\frac{1}{2} \times \frac{6}{7}$

i $\frac{2}{3} \times \frac{3}{5}$ **j** $\frac{8}{9} \times \frac{3}{4}$ **k** $\frac{4}{5} \times \frac{1}{2}$ **l** $\frac{5}{6} \times \frac{24}{25}$

m $\frac{6}{7} \times \frac{7}{8}$ **n** $\frac{7}{9} \times \frac{18}{21}$ **o** $\frac{8}{21} \times \frac{7}{12}$ **p** $\frac{5}{14} \times \frac{21}{40}$



Hint for Q4: First cancel any common factors, if possible.

Example 21 Multiplying with mixed numbers

Evaluate $1\frac{2}{3} \times 2\frac{1}{10}$.

Solution

$$1\frac{2}{3} \times 2\frac{1}{10} = \frac{{}^{1}\cancel{5}}{\cancel{3}_{1}} \times \frac{\cancel{21}^{7}}{\cancel{10}_{2}}$$

$$= \frac{7}{2} \text{ or } 3\frac{1}{2}$$

Explanation

Rewrite as improper fractions.

Cancel common factors between numerators and denominators and then multiply numerators and denominators.

Now you try

Evaluate $1\frac{4}{5} \times 2\frac{1}{3}$.

5 Evaluate:

a $1\frac{1}{2} \times \frac{1}{3}$ **b** $1\frac{2}{3} \times \frac{3}{4}$ **c** $2\frac{1}{3} \times \frac{9}{10}$ **d** $1\frac{1}{2} \times 1\frac{1}{3}$ **e** $2\frac{1}{3} \times 1\frac{4}{5}$

f $\frac{3}{4} \times 1\frac{1}{7}$ **g** $2\frac{1}{2} \times 2\frac{1}{2}$ **h** $4\frac{1}{2} \times \frac{8}{9}$ **i** $1\frac{3}{4} \times 1\frac{2}{3}$ **j** $3\frac{3}{4} \times \frac{8}{5}$

Example 22 Dividing fractions

Evaluate the following.

a $\frac{4}{15} \div \frac{12}{25}$ **b** $1\frac{17}{18} \div 1\frac{1}{27}$

Solution

a $\frac{4}{15} \div \frac{12}{25} = \frac{{}^{1}\cancel{4}}{\cancel{15}_{3}} \times \frac{\cancel{25}^{5}}{\cancel{12}_{3}}$

$$= \frac{5}{9}$$

b $1\frac{17}{18} \div 1\frac{1}{27} = \frac{35}{18} \div \frac{28}{27}$

$$= \frac{{}^{5}\cancel{35}}{\cancel{18}_{2}} \times \frac{\cancel{27}^{3}}{\cancel{28}_{4}}$$

$$= \frac{15}{8} \text{ or } 1\frac{7}{8}$$

Explanation

To divide by $\frac{12}{25}$ we multiply by its reciprocal $\frac{25}{12}$.

Cancel common factors between numerators and denominators then multiply fractions.

Rewrite mixed numbers as improper fractions.

Multiply by the reciprocal of the second fraction.

The reciprocal of $\frac{28}{27}$ is $\frac{27}{28}$.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

1F

Now you try

Evaluate the following.

a $\frac{5}{12} \div \frac{5}{18}$ b $1\frac{7}{8} \div 1\frac{1}{24}$

6 Evaluate the following (recall that the reciprocal of 8 is $\frac{1}{8}$).

a $\frac{4}{7} \div \frac{3}{5}$ b $\frac{3}{4} \div \frac{2}{3}$ c $\frac{5}{8} \div \frac{7}{9}$

d $\frac{3}{7} \div \frac{4}{9}$ e $\frac{3}{4} \div \frac{9}{16}$ f $\frac{4}{5} \div \frac{8}{15}$

g $\frac{8}{9} \div \frac{4}{27}$ h $\frac{15}{42} \div \frac{20}{49}$ i $15 \div \frac{5}{6}$

j $6 \div \frac{2}{3}$ k $\frac{4}{5} \div 8$ l $\frac{8}{9} \div 6$



Hint for Q6: Multiply by the reciprocal of the second fraction.

7 Evaluate the following.

a $14 \div 4\frac{1}{5}$ b $6 \div 1\frac{1}{2}$ c $1\frac{1}{3} \div 8$

d $2\frac{1}{4} \div 1\frac{1}{2}$ e $4\frac{2}{3} \div 5\frac{1}{3}$ f $7\frac{1}{5} \div \frac{4}{15}$



Hint for Q7: First change any mixed numbers to improper fractions.

Problem-solving and reasoning

8(½), 9, 10 | 8(½), 9, 11, 12

8 Find the following amounts.

a $\frac{1}{2}$ of \$20 b $\frac{1}{2}$ of \$7 c $\frac{1}{3}$ of \$18

d $\frac{3}{4}$ of \$24 e $\frac{4}{5}$ of \$8000 f $\frac{9}{10}$ of \$1



Hint for Q8: 'Of' means multiply and 20 as a fraction is $\frac{20}{1}$.

9 Evaluate these mixed-operation problems.

a $\frac{2}{3} \times \frac{1}{3} \div \frac{7}{9}$ b $\frac{4}{5} \times \frac{3}{5} \div \frac{9}{10}$ c $\frac{4}{9} \times \frac{6}{25} \div \frac{1}{150}$

10 In a $1\frac{1}{2}$ hour maths exam, $\frac{1}{6}$ of that time is allocated as reading time. How long is the reading time?

11 A car's fuel gauge shows that it has $\frac{1}{4}$ of a tank of petrol remaining. The petrol tank holds 64 litres of fuel when full. The car can travel 10 km on 1 litre. How many kilometres can you travel on the amount of petrol that remains in the tank?



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

12 Thomas, Ahn and Oscar agree to equally share the job of cleaning the house after a party. They estimate the job will take $4\frac{1}{2}$ hours to complete. How many minutes of work should they each contribute?



Electronics and reciprocals — 13

13 Reciprocals are used in the study of electronics. On the calculator, the x^{-1} button can be used to find the reciprocal of a number.

a Use this button to find reciprocals of the following numbers.

i 2 **ii** 5 **iii** 6.2 **iv** 1.5 **v** -1


b If $\frac{1}{6.4}+\frac{1}{7.2}=A$, find the value of $\frac{1}{A}$ (to two decimal places).

c If $\frac{1}{50}+\frac{1}{50}=B$, what is $\frac{1}{B}$?

d Investigate, using your calculator, the types of numbers for which the reciprocal is bigger than the original number.

e What positive number is its own reciprocal?

f Are there any numbers that do not have a reciprocal?



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

1G Ratios CONSOLIDATING

Learning intentions

- To understand how ratios are used
- To know how to express a ratio in simplest form using whole numbers with no common factor
- To know that the unitary method involves finding the value of one part
- To be able to use the unitary method to divide a quantity in a given ratio

Key vocabulary: ratio, unitary method, highest common factor

Fractions, ratios and rates are used to compare quantities. A lawn mower, for example, might require $\frac{1}{6}$ of a litre of oil to make a petrol mix of 2 parts oil to 25 parts petrol, which is an oil to petrol ratio of 2 to 25 or 2 : 25.



A 2 : 25 oil to petrol ratio means that a lawn mower needs 2 parts oil to 25 parts petrol.

Lesson starter: The lottery win

$100 000 is to be divided up for three lucky people into a ratio of 2 to 3 to 5 (2 : 3 : 5). Work out how the money is to be divided.

- Clearly write down your method and answer. There may be many different ways to solve this problem.
- Write down and discuss the methods suggested by other students in the class.

Key ideas

- **Ratios** are used to compare quantities with the same units.
  - The ratio of a to b is written $a : b$.
  - Ratios in simplest form use whole numbers that have no common factor.
- The **unitary method** involves finding the value of one part of a total.
  - Once the value of one part is found, then the value of several parts can easily be determined.

Exercise 1G

Understanding 1–3 3

1 State if each of the following ratios is in simplest form (yes or no).


a 4 : 10 **b** 3 : 7 **c** 20 : 39 **d** 48 : 64

2 Draw up a table with the following headings.


Diagram	Ratio of shaded parts to unshaded parts	Ratio of shaded parts to parts in the whole

Enter information about these diagrams to complete the table.


a



b



c



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3 A pet shop has birds, fish and kittens. At the pet shop there were five times as many birds as kittens and twice as many fish as birds. Write the ratio of:

- **a** birds to kittens
- **b** birds to fish
- **c** fish to birds
- **d** fish to kittens
- **e** birds to the total number of pets at the pet shop.



Hint for Q3: Remember that, with a ratio, the order matters.

Fluency

4–8(½) 4–8(½)

Example 23 Simplifying ratios using HCFs

Simplify the ratio 38 : 24.

Solution

$38 : 24 = 19 : 12$

Explanation

The HCF of 38 and 24 is 2 so divide both sides by 2.

Now you try

Simplify the ratio 45 : 27.

4 Simplify these ratios.

- **a** 6 : 30
- **b** 8 : 20
- **c** 40 : 50
- **d** 2 : 20
- **e** 15 : 18
- **f** 9 : 27
- **g** 18 : 6
- **h** 24 : 36
- **i** 52 : 39
- **j** 144 : 36
- **k** 48 : 96
- **l** 30 : 10
- **m** 2000 : 5600
- **n** 3 : 6 : 12
- **o** 15 : 30 : 10



Hint for Q4: Divide by the HCF.

Example 24 Simplifying ratios involving fractions and decimals

Simplify these ratios.

a $0.2 : 0.14$ **b** $2\frac{1}{2} : 1\frac{1}{3}$

Solution

a $0.2 : 0.14 = 20 : 14$
$= 10 : 7$

Explanation

Multiply by 100 to remove all the decimal places and simplify.

b $2\frac{1}{2} : 1\frac{1}{3} = \frac{5}{2} : \frac{4}{3}$
$= \frac{15}{6} : \frac{8}{6}$
$= 15 : 8$

Write as improper fractions using the same denominator.
Multiply both sides by 6 to write as whole numbers.

Now you try

Simplify these ratios.

a $0.5 : 0.32$ **b** $3\frac{1}{4} : 1\frac{7}{8}$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

1G

5 Simplify these ratios.

a 0.1 : 0.2 **b** 0.3 : 4.1 **c** 0.5 : 3.2
d 0.3 : 0.9 **e** 0.7 : 3.5 **f** 0.4 : 0.12
g 8 : 0.2 **h** 15 : 0.01 **i** 1.6 : 0.56

6 Simplify:

a $\frac{1}{4} : \frac{1}{3}$ **b** $\frac{2}{3} : \frac{1}{4}$ **c** $1\frac{1}{2} : 3\frac{1}{3}$
d $2\frac{1}{4} : 1\frac{2}{5}$ **e** $\frac{3}{8} : 1\frac{3}{4}$ **f** $1\frac{5}{6} : 3\frac{1}{4}$



Hint for Q6: A simplified ratio must include only whole numbers (with no common factors).

Example 25 Simplifying ratios involving units

Simplify the ratio of 560 metres to 2 kilometres.

Solution

560 m : 2 km
= 560 m : 2000 m
= 560 : 2000
= 7 : 25

Explanation

Write using ratio notation.
Convert the units so that both are the same:
2 km = 2000 m
Cancel the units and divide both numbers by their HCF. HCF of 560 and 2000 is 80

Now you try

Simplify the ratio of 35 cents to $2.

7 Write each of the following as a ratio in simplest form.

a 80c : $8 **b** 90c : $4.50
c 80 cm : 1.2 m **d** 0.7 kg : 800 g
e 2.5 kg : 400 g **f** 30 min : 2 hours
g 45 min : 3 hours **h** 4 hours : 50 min
i 40 cm : 2 m : 50 cm **j** 80 cm : 600 mm : 2 m
k 2.5 hours : 1.5 days **l** 0.09 km : 300 m : 1.2 km



Hint for Q7: Convert to the same units first.

Example 26 Dividing into a given ratio

$300 is to be divided into the ratio 2 : 3.
Find the value of the larger portion using the unitary method.

Solution

Total number of parts is 2 + 3 = 5
5 parts = $300

1 part = $\frac{1}{5}$ of $300
= $60

Larger portion = 3 × $60
= $180

Explanation

Use the ratio 2 : 3 to get the total number of parts.

Calculate the value of each part. ($300 ÷ 5)

Calculate the value of 3 parts.

Now you try

$600 is to be divided into the ratio 5 : 7.

Find the value of the larger portion using the unitary method.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

8 Use the unitary method to divide:

a $500 in the ratio of 1 : 4
b $36 in the ratio of 4 : 5
c 88 kg in the ratio of 3 : 8
d $96 in the ratio of 7 : 5
e $500 in the ratio of 2 : 3
f 2000 g in the ratio of 3 : 5
g $100 in the ratio of 7 : 3
h $600 in the ratio of 1 : 1
i $70 in the ratio of 2 : 7 : 1
j 420 g in the ratio of 8 : 2



Hint for Q8: First find the value of one part.

Problem-solving and reasoning 9–11 10–13



9 420 g of flour is to be divided into a ratio of 7 : 3 for two different recipes. Find the smaller amount.

10 Kirsty manages a restaurant. Each day she buys watermelons and mangoes in the ratio of 3 : 2. How many watermelons did she buy if, on one day, the total number of watermelons and mangoes was 200?

11 If a prize of $6000 was divided among Georgia, Leanne and Maya in the ratio of 5 : 2 : 3, how much did each girl get?



Hint for Q12: If 72 litres is two parts, what is one part?

12 The dilution ratio for a particular chemical with water is 2 : 3 (chemical to water). If you have 72 litres of chemical, how much water is needed to dilute the chemical?

13 Amy, Belinda, Candice and Diane invested money in the ratio of 2 : 3 : 1 : 4 in a publishing company. If the profit was shared according to their investment, and Amy's profit was $2400, find the profit each investor made.

Mixing drinks — 14

14 Four jugs of cordial have a cordial to water ratio as shown and a given total volume.

Jug	Cordial to water ratio	Total volume
1	1 : 5	600 mL
2	2 : 7	900 mL
3	3 : 5	400 mL
4	2 : 9	330 mL



a How much cordial is in:
i Jug 1? **ii** Jug 2?

b How much water is in:
i Jug 3? **ii** Jug 4?

c If Jug 1 and 2 were mixed together to give 1500 mL of drink:
i how much cordial is in the drink?
ii find the ratio of cordial to water in the drink.

d Find the ratio of cordial to water if the following jugs are mixed.
i Jug 1 and 3 **ii** Jug 2 and 3
iii Jug 2 and 4 **iv** Jug 3 and 4

e Which combination of two jugs gives the strongest cordial to water ratio?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

1H Rates and best buys

Learning intentions

- To understand what a rate represents and how it is expressed in simplest form
- To know and be able to use the speed = distance ÷ time relationship to determine an unknown
- To be able to use rates to determine best buys

Key vocabulary: rate, best buys

It is often necessary to compare two quantities with different units. When this occurs it is called a rate.

Rates are used to describe speed (m/s, km/h), pay, lap times in formula one racing, and even the prices at the supermarket. Using rates is important in our everyday lives.



Speed is important in Formula One racing, where lap speeds are measured in km/h. These lap speeds can be over 250 km/h!

Lesson starter: Fastest animals on Earth

In pairs, arrange these animals from fastest to slowest. Then see if you can write their speeds in metres per second (m/s).



Zebra 64.37 km/h



Ostrich 70 km/h



Giraffe 52 km/h



African bush elephant 40.7 km/h



Lion 80 km/h



Cheetah 120 km/h



Dog 72.4 km/h



Galapagos tortoise 0.32 km/h



Kangaroo 70.01 km/h



Wildebeest 64.05 km/h



Hyena 65 km/h




Fox 32.03 km/h

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Key ideas

- A **rate** compares related quantities with different units.
  - The rate is usually written with one quantity compared to a single unit of the other quantity. For example: 50 km per 1 hour or 50 km/h
- Rates can be used to determine best buys when purchasing products.
- Speed (s) = distance (d) ÷ time (t)
  For example:
  80 km in 2 hours = 40 km/h



$s = \frac{d}{t}$, $d = s \times t$, $t = \frac{d}{s}$

- The 'best buy' is the best value for money. This is found by comparing the cost per unit value, e.g. cost per kg.

Exercise 1H

Understanding

1–4 | 3, 4



Hint for Q1: Compare to 1 hour in part c.

1 Write the following as simple rates using the / symbol.
   a Tom travelled 60 km in 1 hour
   b $84 in 1 hour
   c 600 km in 10 hours

2 Find the cost of 1 kg if:
   a 2 kg costs $8
   b 5 kg costs $15
   c 4 kg costs $10



Hint for Q3: Use units m/s or km/h.

3 Determine the speed of the following trips
   a 100 m in 10 seconds
   b 120 km in 2 hours
   c 42 km in 3 hours

4 Which is better value for money (the best buy)?
   A 4 kg of potatoes at $6
   B 10 kg of potatoes at $14

Fluency

5–6(½), 7, 8–9(½) | 5–6(½), 8–9(½)

Example 27 Simplifying rates

Write these rates in simplest form.
a 120 km every 3 hours
b 5000 revolutions in $2\frac{1}{2}$ minutes

Solution

a 120 km per 3 hours $= \frac{120}{3}$ km/h
$= 40$ km/h

Explanation

Divide by 3 to write the rate compared to 1 hour.

Continued on next page

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

1H

b 5000 revolutions per $2\frac{1}{2}$ minutes

= 10 000 revolutions per 5 minutes

$= \frac{10\,000}{5}$ revs/min

= 2000 revs/min

First multiply by 2 to remove the fraction.

Then divide by 5 to write the rate using 1 minute.

Now you try

Write these rates in simplest form.

a \$150 in 5 hours

b 1800 revolutions in $1\frac{1}{2}$ minutes

5 Write these as rates in their simplest form.

a \$84 in 3 hours

b \$200 in 4 hours

c 12 kg every 2 minutes

d \$3.50 for $\frac{1}{2}$ kg

e 64 runs in 16 overs

f 623 points in 7 games

g 76 cm in 4 years

h 56 metres in 4 seconds

i 207 heart beats in $2\frac{1}{4}$ minutes

j 180 mL in 22.5 seconds



Hint for Q5:
Work out each per one unit.

Example 28 Determining distance, speed and time

A family took 3 hours to complete the 210 km from their home to their holiday house.

a Find their average speed for the trip.

b If they increased their average speed to 90 km/h, how many hours and minutes would the trip have taken?

Solution


a Average speed = distance ÷ time

= 210 km ÷ 3 hours

= 70 km/h

Explanation

Use the *dst* triangle and cover the *s* to provide the formula for finding speed: $s = \frac{d}{t}$ or $d \div t$



Substitute in your values and calculate the speed.

Remember to put units in your answer.

b Time = distance ÷ speed


= 210 km ÷ 90 km/h

$= 2\frac{1}{3}$ hours

= 2 hours 20 minutes

Use the *dst* triangle and cover the *t* to find the formula for time:

$t = \frac{d}{s}$ or $d \div s$



Substitute in your values and find the time:

$\frac{1}{3}$ hour $= \frac{1}{3} \times 60$ min = 20 minutes

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Now you try

A rally driver took 5 hours to complete a 300 km course.

a Find the average speed for the rally.

b If the average speed was increased to 72 km/h, how many hours and minutes would the course have taken?

6 Find the average speed (km/h) of:

a a car travelling 140 km in 2 hours

b a bike travelling 60 km in 4 hours

c a walker travelling 12 km in 3 hours

d a horse galloping 2.7 km in 3 minutes $\left(\frac{1}{20} \text{ hour}\right)$

e a truck travelling 760 km in 9.5 hours

7 Rick drives the 1040 km from Sydney to Melbourne to watch the Sydney Swans play the Western Bulldogs in the AFL. He averaged 80 km/h.

a How many hours did the trip take?

b How many hours and minutes would the trip take if the average speed was reduced to 64 km/h?



Example 29 Finding best buys

Which is better value?
5 kg of potatoes for $3.80 or 3 kg for $2.20

Solution

Price per kg.

5 kg bag.
1 kg costs $3.80 ÷ 5 = $0.76

3 kg bag.
1 kg costs $2.20 ÷ 3 = $0.73
∴ the 3 kg bag is cheaper

Explanation

Divide each price by the number of kilograms to find the price per kilogram.

Then compare and choose the cheapest.

Now you try

Which is better value?

2 L of chocolate ice cream for $4.50 or 3 L of the same ice cream for $6.45.

8 Determine the best buy in each of the following.

a 2 kg of washing powder for $11.70 or 3 kg for $16.20

b 1.5 kg of red delicious apples for $4.80 or 2.2 kg of royal gala apples for $7.92

c 2.4 litres of orange juice for $4.20 or 3 litres of orange juice for $5.40

d 0.7 GB of internet usage for $14 or 1.5 GB for $30.90 with different service providers



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

1H

9 Which is the best buy?

A 4 hours hire for $120 **B** 5 hours hire for $160 **C** 7 hours hire for $203

Problem-solving and reasoning 10–12 11–13

10 Hamish rides his bike at an average speed of 22 km/h. How far does he ride in:

a $2\frac{1}{2}$ hours? **b** $\frac{3}{4}$ hours? **c** 15 minutes?

11 Find the cost of 100 g of each product below then decide which is the best buy.

a 300 g of coffee A at $10.80 or 220 g of coffee B at $8.58.

b 600 g of pasta A for $7.50 or 250 g of pasta B for $2.35

c 1.2 kg of cereal A for $4.44 or 825 g of cereal B for $3.30

12 A shearer sheared 80 sheep in 2 hours.

a Express this as a rate.

b If the shearer is able to continue at this rate for $5\frac{1}{2}$ hours, how many sheep can be sheared?

c How long would it take, at this rate, to shear all 1000 sheep on the property?



13 Light travels at approximately 300 000 km/s.

a Express this speed as km/h.

b How far does light travel in 2 minutes?

c How long does light from the Sun take to travel the 149 million kilometres to Earth?



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

The cyclist — 14

14 A cyclist's progress over a journey of 160 km is recorded. The ratio of time (in hours) to distance (in kilometres) is 1 : 20.

a Copy and complete the table showing the progress of the cyclist.


Time in hours	0	1	2	3	4	5	6	7	8
Distance in km	0	20							

b What is the cyclist's average speed?

c Draw a set of axes like the ones shown and record the information from the table onto the axes.

d What conclusions can you make about the shape of the graph? Why is this the case?

e If the cyclist doubles their speed, on the same set of axes plot a graph that shows the journey for this new speed.





ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Maths@Work

Cooks and chefs

Being a professional cook or chef is a physically demanding job. Long hours in hot kitchens and being on your feet all day are challenging. Most chefs have a passion for food and flavours and are seen as creative people who create not just a meal but a masterpiece on a plate.

Chefs need to have good communication skills, a practical sense of measurement and excellent number skills. Their job involves ordering stock, understanding cooking temperatures and adapting recipes using ratios and direct proportion.



1 A recipe for a classic Pavlova uses 6 egg whites and 270 grams ($1\frac{1}{4}$ cups) of superfine sugar or caster sugar and serves 8. (A Pavlova also contains cornflour and white vinegar.)

- **a** What is the simplified ratio of egg whites to servings?
- **b** How many egg whites are needed for a Pavlova serving 12 people?
- **c** How many whole egg whites are needed for a Pavlova serving 6 people?
- **d** How many servings are in a Pavlova using 15 egg whites?
- **e** How many grams of sugar are equivalent to 1 cup?
- **f** How many grams of sugar are needed per egg white for this Pavlova recipe?

2 A café style banana bread recipe indicates that it would make 3 large loaves. The recipe is as follows:

9 cups of plain flour
$4\frac{1}{2}$ cups sugar
3 teaspoons of bicarb of soda
1125 grams of melted butter
9 eggs
10 mashed bananas



- **a** If each loaf can be cut into 18 single slices, how many loaves need to be baked per day for 50 single slice serves at a café?
- **b** How many cartons of a dozen eggs are needed for a week where the cook makes this recipe 8 times?
- **c** If 1 egg weighs 75 grams, what is the simplified ratio of eggs to butter in this recipe?
- **d** If 1 cup of plain flour is equivalent to 150 grams of flour, how many grams of flour is needed for this recipe?
- **e** What is the simplified ratio of flour to melted butter for this recipe?
- **f** If only one loaf is to be made, write down the amounts of ingredients required.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3 Many chefs start their careers mastering the basics such as mayonnaise.
The recipe for 1 cup of classic mayonnaise is:

2 egg yolks
2 teaspoons of lemon juice
1 teaspoon of mustard
$\frac{1}{2}$ teaspoon of salt
pinch of pepper
250 mL of olive oil



a What is the simplified ratio of:
- **i** lemon juice to mustard?
- **ii** lemon juice to olive oil (1 teaspoon = 5 mL)?
- **iii** mustard to salt?

b If a litre is the same as 4 cups, how many eggs are needed for 1 L of mayonnaise?

c If one lemon yielded 25 mL of fresh juice, how many cups of mayonnaise can be made using all of the juice in this one lemon?

4 Scaling a recipe means that the ingredient quantities are each multiplied by a conversion factor so that the recipe makes more or fewer serves. If a cake recipe makes 12 cupcakes, what conversion factor is required to scale up the recipe quantities to make 42 cupcakes?

Using technology

5 Set up an Excel spreadsheet as shown below for scaling cookie recipe quantities. Enter formulas in the shaded cells.



Hint for Q5: Format all formula cells to 'number'/one decimal place. Use $ signs to fix (i.e. anchor) a cell reference, e.g. $C$4.

	A	B	C	D
1	Bella's Bakery			
2	Chocloate chip cookies recipe			
3	Recipe number of cookies	Desired number of cookies	Conversion factor	
4	24	60		
5	**Ingredient**	**Recipe quantities**	**Unit**	**Scaled quantities**
6	Eggs	1	egg	
7	Macadamia nut oil	60	mL	
8	Apple sauce	2	teaspoon	
9	Brown sugar	160	g	
10	Vanilla extract	1	teaspoon	
11	Plain flour	1	cup	
12	Baking powder	1	teaspoon	
13	Rolled oats	1	cup	
14	Dark chocolate	100	g	

a For a recipe to make 60 cookies, state the quantities of the following ingredients:
- **i** oil
- **ii** flour
- **iii** chocolate

b Bella has a school tuckshop order for 32 bags with 3 cookies in each. State the quantities of the following ingredients that Bella will need:
- **i** eggs
- **ii** apple sauce
- **iii** sugar



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Modelling

Painting makeover

Archer and Emily are mixing their own paint to form a special colour for their loungeroom and bedroom makeover. They have tested the following ratios and feel that these two options, A and B, suit their tastes.

	White	Yellow	Red	Blue
Colour Ratio A	18	2	1	3
Colour Ratio B	14	3	2	1

Archer and Emily want to know the amount and cost of paint required to cover the walls and ceiling of the loungeroom and bedroom using either ratio A mix or ratio B mix.

The white paint costs $\$8$ per litre and the colours all cost $\$24$ per litre.
The paint covers $10\,m^2$ per litre.
The loungeroom is 6 m long, 5 m wide and 2.5 m high.
The bedroom is 5 m long, 4 m wide and 2.5 m high.

Present a report for the following tasks and ensure that you show clear mathematical workings, explanations and diagrams where appropriate.

1 Preliminary task

a Using the table above, copy and complete the colour ratios for:

i Colour Ratio A, W : Y : R : B = ☐ : ☐ : ☐ : ☐

ii Colour Ratio B, W : Y : R : B = ☐ : ☐ : ☐ : ☐

b What fraction of:

i white paint is in ratio A?

ii blue paint is in ratio A?

iii white paint is in ratio B?

iv red paint is in ratio B?

c If 36 L of paint is required in total, find the volume of the following amounts giving each answer as a mixed number.

i white paint in ratio A

ii blue paint in ratio A


iii white paint in ratio B

iv red paint in ratio B

2 Modelling task

Formulate

The problem is to compare the cost of painting the loungeroom and bedroom using the colour mixes A and B.



a Write down all the relevant information that will help solve this problem.

b For the loungeroom and bedroom

i Draw a labelled diagram of each rectangular ceiling and each room's 4 rectangular walls.

ii For each room, calculate the total area of the walls and ceiling in m^2.

iii If 2 coats of paint are applied, calculate the total area to be painted.

iv Determine the total volume of paint needed given that each litre covers $10\,m^2$.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Solve

c For Colour ratio A, determine:
- **i** The fraction that is white paint and the fraction that is coloured paint.
- **ii** The volume of white paint needed.
- **iii** The cost of white paint at $8 per litre.
- **iv** The volume of coloured paint needed.
- **v** The cost of coloured paint at $24 per litre.

d For Colour ratio B, determine:
- **i** The fraction that is white paint and the fraction that is coloured paint.
- **ii** The volume of white paint needed.
- **iii** The cost of white paint at $8 per litre.
- **iv** The volume of coloured paint needed.
- **v** The cost of coloured paint at $24 per litre.

Evaluate and verify

e Determine the total cost of the paint for colour ratio A and for colour ratio B.

f Which colour ratio will be the cheaper option and by how much?

g Compare the percentages of white and coloured paint in ratios A and B.

h Explain why one ratio produces a more expensive paint.

Communicate

i Summarise your results and describe any key findings.

3 Extension questions

Emily paints the bedroom at the same time as Archer paints the loungeroom. Emily paints at a rate of $13\,m^2$ per hour and Archer paints at a rate of $17\,m^2$ per hour.

a How long does it take for one coat of paint to be applied to each room?

b If there is a 3 hour gap between coats, find the total time needed for this painting makeover.




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Forming irrational numbers

Key technology: Spreadsheets

We know that an irrational number like pi or the square root of 2 cannot be expressed as a fraction exactly and when written as a decimal will show infinite non-recurring decimal places.



One way to generate these types of numbers is to use a sum or an infinite set of fractions called a series. An example is the number pi (π) which is equal to 3.14159265359..... with an infinite non-recurring pattern. Pi can be generated using this series with the first 6 terms showing then multiplying the result by 4..

$$\frac{1}{1}-\frac{1}{3}+\frac{1}{5}-\frac{1}{7}+\frac{1}{11}-\frac{1}{13}+\ldots\ldots\ldots$$

1 Getting started

a Do you notice any patterns in the series above? If so, describe them.

b Work out the value of the following, writing your answer as a decimal.

i $\left(\frac{1}{1}-\frac{1}{3}\right)\times 4$

ii $\left(\frac{1}{1}-\frac{1}{3}+\frac{1}{5}\right)\times 4$

iii $\left(\frac{1}{1}-\frac{1}{3}+\frac{1}{5}-\frac{1}{7}\right)\times 4$

iv $\left(\frac{1}{1}-\frac{1}{3}+\frac{1}{5}-\frac{1}{7}+\frac{1}{9}\right)\times 4$

c What do you notice about the answers to the above calculations? Which one is the best approximation to pi?

2 Using technology

We will generate the answers to the questions in part **1b** above using a spreadsheet.

a Fill in the details for row 1, 3 and 4 as shown, including the formula in cell D4.

b Enter the formulas in cells A5, B5, C5 and C6.

	A	B	C	D
1	Forming pi			
2				
3	Terms	Denominator	pi/4	pi
4	1	1	1	=C4*4
5	=A4+1	=B4+2	=C4-1/B5	=C5*4
6	=A5+1	=B5+2	=C5-1/B6	=C6*4
7	=A6+1	=B6+2	=C6-1/B7	=C7*4
8	=A7+1	=B7+2	=C7-1/B8	=C8*4

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

c Now fill down the following.
- **i** Column A using cell A5
- **ii** Column B using cell B5
- **iii** Column C using cells C5 and C6. Highlight both then fill down
- **iv** Column D using cell D4.

d Check that the numbers in column D match your answers from part **1b** above.

3 Applying an algorithm


a Fill down your spreadsheet for a total of 20 terms. Remember to highlight two cells in column C before filling down.

b How many terms are required for the first digit to be a 3 and then never change from that point onwards?

c Repeatedly extend your spreadsheet and find how many terms are required for the next digit of pi to settle on a number and never change from that point onwards. Complete this table to summarise your results. The first one is done for you which is the answer to part **b** above.

Digit	Number of terms required
1st (3)	7
2nd (1)	
3rd (4)	
4th (1)	

d Is 10 000 terms enough to ensure that your approximation of pi is accurate for the first 5 digits and never changes from that point onwards?




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Puzzles and games

1 Can you arrange the first nine counting numbers using signs (+, −, ×, ÷) and brackets to give an answer of 100?

2 In how many ways can four girls and two boys be seated in a row if the two boys are to sit at the ends?

3 Complete the number cross below.



Clues

Across

2. The sum of 100, 150 and 3
4. The product of 5 and the first prime number after 6
5. Reverse the digits of the product of 13 and 5
7. One less than the square of 10
9. Half of 6 times 8
10. 2 to the power of 7

Down

1. A palindromic number
2. $5 \times 10 - 5 \times 5$
3. A multiple of 11
6. 639 written correct to two significant figures
8. The next even integer after 90
9. A multiple of 7

4 a Perfect numbers are positive integers which are equal to the sum of all their factors, excluding the number itself.

i Show that 6 is a perfect number.

ii There is one perfect number between 20 and 30. Find the number.

b Triangular numbers are the number of dots required to form triangles as shown in this table.

i Complete this table.

Number of rows	1	2	3	4	5	6
Diagram						
Number of dots (triangular number)	1	3				

ii Find the 7th and 8th triangular numbers.

c Fibonacci numbers are a sequence of numbers where each number is the sum of the two preceding numbers. The first two numbers in the sequence are 0 and 1.

i Write down the first 10 Fibonacci numbers.

ii If the Fibonacci numbers were to be extended in the negative direction, what would the first four negative Fibonacci numbers be?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter summary

Reviewing number

Decimals
Decimals are a way of writing fractions with denominators of 10, 100, 1000.

$\frac{1}{10} = 0.1$, $\frac{1}{100} = 0.01$,

$\frac{1}{1000} = 0.001$

Rounding decimals
If the critical digit (the next one in the number) is 0, 1, 2, 3 or 4, round down. If it is 5, 6, 7, 8 or 9, round up.

4.763 = 4.8 (1 d.p.)
= 4.76 (2 d.p.)

Integers
Integers are the set of positive and negative whole numbers including zero.

0, ±1, ±2, ±3, ±4, ...

Working with negatives
- To add a negative number, you subtract.
  $a + (-b) = a - b$
- To subtract a negative number, you add.
  $a - (-b) = a + b$

× and ÷
- Two numbers of the same sign multiply or divide to give a positive answer. $-4 \times (-5) = 20$
  $-10 \div (-2) = 5$
- Two numbers with different signs multiply or divide to give a negative answer. $-4 \times 5 = -20$
  $10 \div (-2) = -5$

Rational numbers
Rational numbers can be written as a fraction.

3, -4, $\frac{2}{3}$, $0.\dot{3}$, $\sqrt{4}$, 0.416 are all rationals.

π is an example of an irrational number. It cannot be written as a fraction.

Ratios
Ratios compare two or more quantities with the same units in a definite order.

5 cents to \$1 = 5 : 100 = 1 : 20

You can simplify ratios by using the HCF.

24 : 36 = 2 : 3 (dividing both by 12)

Use whole numbers:

0.3 : 7 = 3 : 70

$\frac{3}{5} : \frac{8}{5} = 3 : 8$

Reciprocals
The reciprocal of a proper or improper fraction is the fraction turned upside down (inverted).

The reciprocal of:

6 is $\frac{1}{6}$

$1\frac{1}{2}$ is $\frac{2}{3}$

$\frac{4}{5}$ is $\frac{5}{4}$ or $1\frac{1}{4}$

Fractions
- To add or subtract fractions, you need to use the lowest common denominator.
  $\frac{3}{4} + \frac{1}{6} = \frac{9}{12} + \frac{2}{12} = \frac{11}{12}$
- To multiply fractions, multiply the numerators and multiply the denominators after cancelling common factors.
  $\frac{8}{9} \times \frac{3}{7} = \frac{8 \times \cancel{3}^{1}}{\cancel{9}^{3} \times 7} = \frac{8}{21}$
- To divide by a fraction, multiply by the reciprocal.
  $\frac{3}{2} \div \frac{4}{5} = \frac{3}{2} \times \frac{5}{4}$
- Convert mixed numbers to improper fractions to apply operations.

Fractions and decimals
- Divide to express a fraction as a decimal.
  $\frac{11}{4} = 3.75$, $\frac{8}{3} = 2.\dot{6}$
- A dot on top of a number indicates it repeats infinitely; a bar is used for more than one digit
  $0.\overline{27} = 0.272727...$
- Use a denominator of 10, 100 etc. to write a decimal as a fraction.
  $0.37 = \frac{37}{100}$, $2.4 = 2\frac{4}{10}$
  $= 2\frac{2}{5}$

Rates
Rates compare two quantities with different units.

50 km in 2 hours = 25 km/h

Best buys
3 kg of carrots for \$6.45 is \$2.15/kg

5 kg of carrots for \$10.20 is \$2.04/kg

∴ 5 kg is best buy

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter checklist

Chapter checklist

A version of this checklist that you can print out and complete can be downloaded from your Interactive Textbook.

		✔
1A	**1 I can add and subtract integers.** e.g. Find **a** $-15+5$ **b** $-4-6$	
1A	**2 I can add a negative integer.** e.g. Find $4+(-9)$	
1A	**3 I can subtract a negative integer.** e.g. Find $3-(-7)$	
1B	**4 I can multiply and divide with negatives.** e.g. Find the value of **a** $-7\times(-4)$ **b** $48\div(-6)$	
1B	**5 I can find squares and cubes of negative numbers.** e.g. Find the value of **a** $(-4)^2$ **b** $(-5)^3$	
1B	**6 I can apply order of operations.** e.g. Find the value of $3\times(-4+2)-5$	
1C	**7 I can round to a required number of decimal places.** e.g. Round each of these to two decimal places: **a** 32.2389 **b** 3.712 **c** 1.4954	
1C	**8 I can round to a number of significant figures.** e.g. Round each of these numbers to two significant figures: **a** 7842 **b** 0.0375	
1C	**9 I can estimate using significant figures.** e.g. Estimate the answer to $1067-506\times0.052$ by rounding each number in the problem to one significant figure. Use your calculator to check how reasonable your answer is.	

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter checklist

		✔
1D	**10 I can identify rational numbers.** e.g. Choose the rational numbers in the following list: $\sqrt{10}, \pi, 1.4, \sqrt{36}, -\frac{3}{5}$	
1D	**11 I can convert between mixed numbers and improper fractions.** e.g. Express **a** $\frac{18}{5}$ as a mixed number **b** $2\frac{3}{7}$ as an improper fraction	
1D	**12 I can write fractions as decimals, including recurring decimals.** e.g. Write **a** $2\frac{7}{8}$ as a decimal **b** $\frac{4}{11}$ as a recurring decimal	
1D	**13 I can write decimals as fractions.** e.g. Write these decimals as fractions: **a** 0.36 **b** 2.215	
1D	**14 I can compare fractions.** e.g. Decide which is the larger fraction of the following: $\frac{7}{9}$ or $\frac{11}{15}$	
1E	**15 I can add and subtract fractions with the same denominator.** e.g. Find **a** $\frac{8}{17} - \frac{3}{17}$ **b** $2\frac{1}{8} + \frac{5}{8}$	
1E	**16 I can add and subtract fractions with different denominators.** e.g. Evaluate $\frac{5}{6} + \frac{3}{5}$	
1E	**17 I can add and subtract mixed numbers.** e.g. Evaluate: **a** $2\frac{3}{4} + 1\frac{5}{6}$ **b** $2\frac{1}{3} - 1\frac{3}{7}$	
1F	**18 I can multiply proper fractions.** e.g. Evaluate the following: **a** $\frac{5}{6} \times \frac{1}{4}$ **b** $\frac{6}{7} \times \frac{14}{27}$	
1F	**19 I can divide proper fractions.** e.g. Evaluate $\frac{5}{8} \div \frac{15}{28}$	

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter checklist

		✔
1F	**20 I can multiply and divide with mixed numbers.** e.g. Evaluate $4\frac{4}{7} \div 2\frac{2}{3}$	
1G	**21 I can simplify ratios.** e.g. Simplify these ratios: **a** $40 : 32$ **b** $0.3 : 0.21$ **c** $\frac{5}{8} : \frac{9}{20}$	
1G	**22 I can simplify ratios by converting to common units.** e.g. Simplify the ratio of 80 cm to 5 metres	
1G	**23 I can divide into a given ratio using the unitary method.** e.g. $\$480$ is to be divided into the ratio $3 : 5$. Find the value of the smaller portion using the unitary method.	
1H	**24 I can simplify a rate.** e.g. Write these rates in simplest form **a** 120 L in 2 hours **b** 108 strokes in $1\frac{1}{2}$ minutes	
1H	**25 I can determine distance, speed and time.** e.g. A driver took 3 hours to complete a 180 km trip. What was their average speed and what was the time taken, in hours and minutes, on the return trip if the average speed was 75 km/h?	
1H	**26 I can determine a best buy.** e.g. Which is better value? 4 L of paint for $\$84$ or 7 L of the same paint for $\$126$?	

Essential Mathematics Core for the Victorian Curriculum Year 9 ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Short-answer questions

1A/B **1** Evaluate the following.

a $-4-(-12)$ **b** $18-9\times 7$ **c** $-90\times(-3)$

d $(-4)^3$ **e** $-15\div 5\times(-2)$ **f** $(10-12)\times(7-11)$

1C **2** Round these numbers to three significant figures.

a 21.483 **b** 29 130 **c** 0.15271 **d** 0.002414

1C **3** Round to two decimal places:

a 14.97683 **b** 0.7149871498… **c** 1.999

1D **4** Write these fractions as decimals.

a $2\frac{1}{8}$ **b** $\frac{5}{6}$ **c** $\frac{13}{7}$

1D **5** Write these decimals as fractions.

a 0.75 **b** 1.6 **c** 2.55

1F **6** State the reciprocal of:

a $\frac{3}{4}$ **b** $\frac{1}{5}$ **c** 8 **d** $1\frac{2}{7}$

1E/F **7** Simplify the following.

a $\frac{5}{6}-\frac{1}{3}$ **b** $1\frac{1}{2}+\frac{2}{3}$ **c** $\frac{13}{8}-\frac{4}{3}$

d $3\frac{1}{2}\times\frac{4}{7}$ **e** $5\div\frac{4}{3}$ **f** $3\frac{3}{4}\div 1\frac{2}{5}$

1G **8** Simplify these ratios.

a 30 : 12 **b** 1.6 : 0.9 **c** $7\frac{1}{2} : 1\frac{2}{5}$

1G **9** Divide 80 into the given ratio.

a 5 : 3 **b** 5 : 11 **c** 1 : 2 : 5

1H **10** Dry dog food can be bought from store A for \$18 for 8 kg or from store B for \$14.19 for 5.5 kg.

a Determine the cost per kilogram at each store and state which is the best buy.

b Determine to the nearest whole number how many grams of each brand you get per dollar.

1G **11** Share \$660 in the ratio of 2 : 3 : 5.

1H **12** Complete these rates.

a \$96 in 3 hours = \$_____/h

b 14 g in 2 minutes = _____g/min

c 32 m in 10 seconds = _____m/min

d \$7.40 in 30 minutes = \$_____/h

e 660 runs in 8 matches = _____runs/match



1H **13** Write the following speeds in km/h.

a 780 km in 5 hours **b** 90 km in $2\frac{1}{2}$ hours

1G **14** Tom and Claire share their lottery win in the ratio of 3 : 2. If Tom got \$9000, how much did Claire get?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter review

Multiple-choice questions

1D **1** $\frac{2}{7}$ written as a decimal is:

A 0.29 **B** 0.286 **C** $0.28\dot{5}$
D $0.\overline{285714}$ **E** $0.285714\dot{4}$

1C **2** 3.0456 written to three significant figures is:

A 3.04 **B** 3.05 **C** 3.045
D 3.046 **E** 3.45

1D **3** 2.25 written as a fraction in simplest form is:

A $2\frac{1}{2}$ **B** $\frac{5}{4}$ **C** $\frac{9}{4}$
D $9\frac{1}{4}$ **E** $\frac{225}{100}$

1E **4** $1\frac{1}{2} - \frac{5}{6}$ is equal to:

A $\frac{2}{3}$ **B** $\frac{5}{6}$ **C** $-\frac{1}{2}$
D $\frac{2}{6}$ **E** $\frac{1}{2}$

1F **5** $\frac{2}{7} \times \frac{3}{4}$ is equivalent to:

A $\frac{8}{11}$ **B** $\frac{3}{7}$ **C** $\frac{5}{11}$
D $\frac{8}{12}$ **E** $\frac{3}{14}$

1F **6** $\frac{3}{4} \div \frac{5}{6}$ is equivalent to:

A $\frac{5}{8}$ **B** 1 **C** 21
D $\frac{4}{5}$ **E** $\frac{9}{10}$

1G **7** Simplifying the ratio 50 cm : 4 m gives:

A $50 : 4$ **B** $8 : 1$ **C** $25 : 2$
D $1 : 8$ **E** $5 : 40$

1D **8** 0.28 as a fraction in its simplest form is:

A 0.28 **B** $\frac{28}{100}$ **C** $\frac{0.28}{100}$
D $\frac{2.8}{100}$ **E** $\frac{7}{25}$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

1G **9** $2.50 divided in the ratio of 4 : 1 is:

A 50 : 200 **B** 50c **C** $2 : 50c **D** $200 **E** $2

1G **10** A childcare centre requires the ratio of carers to children to be 1 : 5. How many carers are needed for 30 children?

A 5 **B** 6 **C** 150 **D** 30 **E** 36

1H **11** Michael earns $460 in 25 hours. His hourly rate of pay is:

A $11500/h **B** $0.05/h **C** $18.40/h
D $18.40/day **E** $5/h

1H **12** 36 km in 40 minutes is the same as:

A 40 km/h **B** 36 km/h **C** 54 km/h
D 90 km/h **E** 56 km/h

Extended-response questions

1 In a cinema with 28 patrons, the ratio of adults to children is 3 : 4.

a How many adults are in the cinema?

b Two adults leave and two more children arrive. How many adults and children are now in the cinema?

c What is the new ratio of adults to children in part **b**?

2 The Tomslin family plans to drive the 856 km from Perth to Monkey Mia in Western Australia.

a At what speed should they travel, on average, if they hope to only take 10 hours?

b The family left home at 6 a.m. and arrived at Monkey Mia at 8 p.m. How long did the trip take?

c Calculate their average speed in part **b**. Answer to one decimal place.

d How far could they have travelled at the speed in part **c** if they only drove for the original 10 hours as planned? Answer to the nearest km.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

2 Financial mathematics



Essential mathematics: why skills in financial mathematics are important

Understanding financial mathematics is crucial for managing personal finances and making informed decisions in various aspects of life. Whether you are shopping, working, or planning for the future, these skills are invaluable.

Taxpayers rely on financial numeracy for accurate tax returns. They calculate total income from various sources, including wages, investments, and rental properties. Understanding deductions and credits reduces taxable income and maximises refunds. A small business owner may also manage tax payments throughout the year, estimating quarterly taxes to avoid penalties.

By understanding simple interest calculations, borrowers can determine the total cost of loans and monthly repayments, helping them compare loan offers and choose the best option. This knowledge aids in budgeting and avoiding overborrowing. Additionally, being financially literate enables borrowers to recognize the impact of interest rates and loan terms on overall debt, empowering them to manage repayments effectively and reduce long-term financial stress. Ultimately, it promotes responsible borrowing and financial stability.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.



In this chapter

2A Percentages (Consolidating)
2B Applying percentages (Consolidating)
2C Percentage increase and decrease (Consolidating)
2D Profits and discounts (Consolidating)
2E Income
2F Taxation
2G Simple interest
2H Applications of simple interest

Victorian Curriculum 2.0

This chapter covers the following content descriptors in the Victorian Curriculum 2.0:

NUMBER

VC2M9N01

ALGEBRA

VC2M9A06

MEASUREMENT

VC2M9M05

Please refer to the curriculum support documentation in the teacher resources for a full and comprehensive mapping of this chapter to the related curriculum content descriptors.

© VCAA

Online resources

A host of additional online resources are included as part of your Interactive Textbook, including HOTmaths content, video demonstrations of all worked examples, auto-marked quizzes and much more.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025
Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Warm-up quiz

1 Simplify:

a $\frac{12}{100}$ **b** $\frac{20}{100}$ **c** $\frac{35}{100}$ **d** $\frac{75}{100}$ **e** $\frac{60}{100}$ **f** $\frac{50}{100}$

2 Multiply these decimals by 100.

a 0.99 **b** 0.58 **c** 0.9 **d** 1.22 **e** 0.08 **f** 1.5

3 Which is larger in each of the following pairs?

a $\frac{1}{2}$ or 55% **b** $\frac{3}{4}$ or 70% **c** 0.89 or 98%

4 Copy and complete.

a $61\% = \frac{\square}{100}$ **b** $9\% = \frac{\square}{100}$ **c** $37\% = \frac{\square}{100}$

d $121\% = \frac{\square}{100}$ **e** $1\% = \frac{\square}{100}$ **f** $75\% = \frac{\square}{100} = \frac{3}{\square}$

5 Copy and complete the following table.

Fraction	Decimal	Percentage
$\frac{1}{100}$		
	0.1	
	0.25	
		50%
$\frac{3}{4}$		

6 Write down 10% of these amounts.

a 100 g **b** 70 km **c** \$450 **d** \$8000 **e** \$5 **f** 90c

7 Find:

a 25% of \$400 **b** 75% of 80 m **c** 50% of \$3

d 10% of \$678 **e** 1% of 600 days **f** $\frac{1}{2}$% of 600 days

8 Complete the following.

a 1% of 60 = 60 divided by $\square$ **b** 10% of 50 = 50 divided by $\square$

c 5% of 100 = 100 divided by $\square$ **d** $33\frac{1}{3}$% of 963 = 963 divided by $\square$

e 25% of 88 = 88 divided by $\square$

9 Find $33\frac{1}{3}$% of 6300 km.

10 Which is larger: 40% of 50 or 25% of 100?

11 Copy and complete:

a 1 week = $\square$ days **b** 1 year = $\square$ days **c** 1 year = $\square$ weeks **d** 1 year = $\square$ months

12 How many hours are there from:

a 5 a.m. to 7 p.m.? **b** 9 a.m. to 3 p.m.? **c** 8:30 a.m. to 9 p.m.?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

2A Percentages CONSOLIDATING

Learning intentions

- To understand what a percentage represents
- To review how to convert between percentages and fractions or decimals
- To know common fraction, decimal and percentage conversions
- To be able to express one quantity as a percentage of another

Key vocabulary: percentage, fraction, decimal

We use percentages for many different things in our daily lives. Some examples include home loans, credit cards, sales and profits.

We know from our previous work on percentages that they represent a fraction with a denominator of 100. 'Per cent' comes from the Latin word *per centum*, and means 'out of 100'.

Lesson starter: Ordering with percentages

Ten different values are given below.

$$\frac{1}{2}, 0.8, 0.05, 15\%, \frac{7}{20}, 0.9, 9\%, \frac{3}{5}, \frac{1}{3}, 0.3$$


In pairs, decide:

a which of the numbers is the smallest **b** which of the numbers is the largest

Write the 10 numbers in ascending order.

Key ideas

- A **percentage** is the number of parts out of 100.
- Converting percentages
  - To change a **decimal** or a **fraction** into a percentage, *multiply* by 100.
  - For example: $\frac{1}{2} \times 100 = 50$ so $\frac{1}{2} = 50\%$
  - To convert a percentage into a fraction, *divide* by 100, using fraction notation. For example: $37\% = \frac{37}{100}$
  - To convert a percentage into a decimal, *divide* by 100.
    For example: $8\% = 8 \div 100$
    $= 0.08$



- Percentage composition
  - To express one quantity as a percentage of another, write them as a fraction, making sure the units are the same. Then convert this fraction to a percentage by multiplying by 100.
    For example: 8 grams out of 32 grams $= \frac{8}{32} \times 100$
    $= 25\%$
- It is useful to memorise some common fraction/decimal/percentage conversions.

Fraction	1	$\frac{1}{2}$	$\frac{1}{3}$	$\frac{1}{4}$	$\frac{1}{5}$	$\frac{1}{10}$	$\frac{1}{20}$	$\frac{1}{50}$	$\frac{1}{100}$
Decimal	1	0.5	$0.\dot{3}$	0.25	0.2	0.1	0.05	0.02	0.01
Percentage	100%	50%	$33\frac{1}{3}\%$	25%	20%	10%	5%	2%	1%

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Exercise 2A

Understanding

1–4 | 1, 4

1 Complete the following using the words multiply or divide.

a To convert a decimal into a percentage _____ by 100.

b To convert a percentage into a fraction _____ by 100.

c To convert a fraction into a percentage _____ by 100.


d To convert a percentage into a decimal _____ by 100.

2 Copy and complete this table of common percentages.

%	10%			50%			
Fraction			$\frac{1}{4}$			$\frac{1}{3}$	$\frac{2}{3}$
Decimal		0.2			0.75		

3 What percentage of each of the following diagrams has been shaded?


a




b



c



d





Hint for Q3: Consider common fractions and percentages for the shaded areas.

4 Scott scored 38 out of 50 on a maths quiz and Sarah scored 79% on the same test. Who scored the higher mark?

Fluency

5–9(½) | 5–9(½)

Example 1 Percentages and fractions

a Write $\frac{12}{25}$ as a percentage.

b Write 7.5% as a fraction.

Solution

a $\frac{12}{25} = \frac{12}{25} \times 100$

$= 48\%$

Explanation

Multiply the fraction by 100.

$\frac{12}{25} \times 100 = \frac{12}{\cancel{25}_{1}} \times \frac{\cancel{100}^{4}}{1}$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

b $7.5\% = \frac{7.5}{100}$

$= \frac{15}{200}$

$= \frac{3}{40}$

Write the percentage as a fraction, using a denominator of 100, $\frac{7.5}{100}$.

Multiply this fraction by $\frac{2}{2}$ so that we don't have a decimal in the fraction and it will be easier to simplify. $\frac{7.5}{100} \times \frac{2}{2} = \frac{15}{200}$

Simplify: $\frac{15}{200} = \frac{3}{40}$ by cancelling a common factor of 5.

Now you try

a Write $\frac{11}{20}$ as a percentage.

b Write 12.5% as a fraction.

5 Express the following fractions as percentages.

a $\frac{1}{5}$	**b** $\frac{4}{5}$	**c** $\frac{8}{10}$	**d** $\frac{3}{10}$
e $\frac{1}{4}$	**f** $\frac{1}{8}$	**g** $\frac{3}{4}$	**h** $\frac{12}{20}$
i $\frac{14}{25}$	**j** $\frac{7}{20}$	**k** $\frac{9}{100}$	**l** $\frac{3}{40}$



Hint for Q5: Multiply by 100.

6 Express the following percentages as simplified fractions.

a 19%	**b** 23%	**c** 99%	**d** 5%
e 22%	**f** 45%	**g** 74%	**h** 75%
i 2.5%	**j** 17.25%	**k** 1%	**l** 125%



Hint for Q6: Divide by 100.

Example 2 Converting between percentages and decimals

a Write 0.45 as a percentage.

b Write 25% as a decimal.

Solution

a $0.45 = 0.45 \times 100 = 45\%$

b $25\% = 25 \div 100 = 0.25$

Explanation

Multiply by 100. This moves the decimal point two places to the right.

Divide by 100. This moves the decimal point two places to the left.

Now you try

a Write 0.23 as a percentage.

b Write 48% as a decimal.

7 Express the following decimals as percentages.

a 0.78	**b** 0.95	**c** 0.65	**d** 0.48
e 0.75	**f** 1.42	**g** 0.07	**h** 0.3
i 0.03	**j** 1.04	**k** 0.12	**l** 0.1225



Hint for Q7: Move the decimal point two places to the right.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

2A

8 Express the following percentages as decimals.

a 12% **b** 83% **c** 57% **d** 88%
e 99% **f** 100% **g** 120% **h** 5%

Example 3 Writing a quantity as a percentage

Write 50c out of $2.50 as a percentage.

Solution

$$50\text{c out of } \$2.50 = \frac{\overset{1}{\cancel{50}}}{\underset{5}{\cancel{250}}} \times 100$$
$$= 20\%$$

Explanation

Convert to the same units ($2.50 = 250c) and write as a fraction. Multiply by 100, cancelling first.

Now you try

Write 90c out of $3.60 as a percentage.

9 In each of the following cases, express the first quantity as a percentage of the second.

a 5 g out of 200 g **b** 40c out of $4
c 10 km out of 200 km **d** 3 s out of 1 minute
e 200 m out of 1 km **f** 100 mL out of $\frac{1}{2}$ L
g 200c out of $1 **h** 45 marks out of a possible 60 marks



Hint for Q9: Write as a fraction and then multiply by 100.

Problem-solving and reasoning 10–12 11–14

10 Copy and complete the table of the favourite summer sports of Year 9 students.

Sport	Number of students who chose sport	Fraction of the total	Percentage of the total
Swimming	44		
Golf	12		
Volleyball	58		
Cricket	36		
Total			



11 Toni pays 31.5 cents in the dollar in tax. Express this as a percentage.

12 Bad weather stopped a cricket game for 35 minutes of a scheduled $3\frac{1}{2}$ hour match. What percentage of the scheduled time was lost?

13 Joe lost 4 kg and now weighs 60 kg. What percentage of his original weight did he lose?



Hint for Q13: What was Joe's original weight?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

14 A company claims that the apple pies it makes are 97% fat free. If the nutritional information on the side of the pack states that total fat is 7 grams of the 250 gram pie, is the claim correct?



Today's timeline — 15

15 Complete this table by filling in estimated times. In the first two columns, insert the times that you did each activity. Use that information to help you fill in the other columns. Round percentages to one decimal place.

a	Time you went to bed last night:	Time you woke up this morning:	Hours and minutes spent in bed:	Percentage of a day spent in bed:
b	Time you started breakfast today:	Time you finished breakfast today:	Minutes spent eating breakfast:	Percentage of the day you spent at breakfast:
c	Time school started today:	Time school is due to finish today:	Hours and minutes spent at school:	Percentage of the day spent at school:
d	Time this maths lesson started:	Time this maths lesson will finish:	Minutes spent in the maths lesson:	Percentage of the day spent in the maths lesson:
e	Time school will finish today:	Time you will arrive home:	Minutes spent travelling home:	Percentage of the day spent travelling home:
f	Time you started your homework yesterday:	Time you finished your homework yesterday:	Minutes spent on homework:	Percentage of your day spent on homework:
g	Time you started watching TV or playing games yesterday:	Time you finished watching TV or playing games:	Minutes spent at this activity:	Percentage of the day spent at this activity:
h	Time you woke up today:	Time you will go to bed tonight:	Hours and minutes spent awake today:	Percentage of the day spent awake:



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

2B Applying percentages CONSOLIDATING

Learning intentions

- To know how to find a percentage of an amount using a fraction or decimal
- To be able to find the original amount from a given percentage

Key vocabulary: percentage, unitary method

The media often quotes percentages in news stories and advertisements. For example:

- 90% of dentists prefer this toothbrush.
- A shirt is reduced by 45%.
- A swing of 5% towards the Liberals is expected in the next election.

These examples involve finding a percentage of a quantity or amount and this is an important part of the work we do with percentages.



Percentages are used a lot in media and advertising. Understanding how they work will help you make decisions.

Lesson starter: Today's news challenge

In pairs, go through today's newspaper or online news site and find articles and advertisements that use percentages. Choose two and explain to the class how percentages are used in the articles you choose.

Key ideas

- To find a percentage of an amount, write the percentage as a fraction or a decimal, then multiply by the amount.

  For example: 3% of $200 = \frac{3}{100} \times 200$ or $0.03 \times 200 = 6$

- To find the original amount when given a percentage, you can work backwards using the **unitary method** (finding the value of 1 unit first).
  For example: *3% of an amount is 9. What is the original amount?*
  Dividing both numbers by 3 gives 1% of the amount = 3
  To find 100%, multiply 3 by 100 so 100% of the amount is 300.

Exercise 2B

Understanding 1–3 3

1 State if the following are true or false.

a 34% of $568 = 0.34 \times 568$ **b** 6% of $81 = 0.6 \times 81$

c 17% of $50 = \frac{17}{50} \times 100$ **d** 12% of $75 = \frac{3}{25} \times 75$

2 Find 1% of the amount if:

a 5% of the amount is 60 **b** 7% of the amount is 63 **c** 20% of the amount is 120

3 **a** If 1% of an amount is $\$4$, what is 100% of the amount?

b If 1% of an amount is $\$7.50$, what is 100% of the amount?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Fluency

4–6(½) | 4–7(½)

Example 4 Finding a percentage of a quantity

Find 15% of $35.

Solution

$$15\% \text{ of } \$35 = \frac{\cancel{15}^{3}}{\cancel{100}_{20}} \times \$35$$
$$= \$5.25$$

Explanation

Write the percentage as a fraction out of 100 and multiply by $35.

Or write 0.15×35.

Note: 'of' means to 'multiply'.

Now you try

Find 18% of $60.

4 Find the following amounts.

a 10% of 20	**b** 5% of 200	**c** 20% of 40
d 15% of 50	**e** 8% of 720	**f** 5% of 680
g 15% of 8200	**h** 70% of 60	**i** 90% of 500
j 75% of 44	**k** 99% of 200	**l** 3% of 50



Hint for Q4:
$10\% \text{ of } 20 = \frac{10}{100} \times 20$

5 Find:

a 10% of $360	**b** 50% of $420	**c** 75% of 64 kg
d 12.5% of 240 km	**e** 37.5% of 40 apples	**f** 87.5% of 400 m
g $33\frac{1}{3}\%$ of 750 people	**h** $66\frac{2}{3}\%$ of 300 cars	**i** $8\frac{3}{4}\%$ of $560

Example 5 Finding the original amount given a percentage amount

Determine the original amount if 5% of the amount is $45.

Solution

5% of the amount = $45
1% of the amount = $9
100% of the amount = $900
So the original amount is $900

Explanation

To use the unitary method, find the value of 1 part or 1% then multiply by 100 to find 100%.

Alternate method: $5\% \text{ of } A = \$45$

$$0.05 \times A = \$45$$
$$A = \$45 \div 0.05$$
$$= \$900$$

Write an equation using the pronumeral A to represent the amount.

Solve for A by dividing both sides by 0.05.

Now you try

Determine the original amount if 8% of the amount is $40.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

2B

6 Determine the original amount if:

a 10% of the amount is \$12
b 6% of the amount is \$42
c 3% of the amount is \$9
d 40% of the amount is \$2.80
e 90% of the amount is \$0.18
f 6% of the amount is \$27
g 12% of the amount is \$96
h 15% of the amount is \$54



Hint for Q6: First find the value of 1%.

7 Determine the value of x in the following if:

a 10% of x is \$54
b 15% of x is \$90
c 25% of x is \$127
d 18% of x is \$225
e 105% of x is \$126
f 110% of x is \$44



Hint for Q7: x is the original amount.

Problem-solving and reasoning 8–11 8(½), 11–14

8 Without a calculator, evaluate the following.

a 10% of \$58
b 5% of \$84
c 1% of \$46
d $2\frac{1}{2}\%$ of \$20
e $33\frac{1}{3}\%$ of \$132
f $66\frac{2}{3}\%$ of \$60



Hint for Q8: $33\frac{1}{3}\% = \frac{1}{3}$

9 If $\frac{1}{3}$ of $96 = 32$, what is $66\frac{2}{3}\%$ of 96?

10 If 10% of \$800 is \$80, explain how you can use this result to find:

a 1% of \$800
b 5% of \$800
c $2\frac{1}{2}\%$ of \$800

11 About 80% of the mass of the human body is water. If Carla weighs 60 kg, how many kilograms of water make up her body weight?

12 In a class of 25 students, 40% have been to England. How many students have not been to England?



13 Explain why 10% of 24 = 24% of 10.

14 10% of 1 day is the same as x hours and y minutes. What is the value of x and y?

More than 100% — 15

15 **a** Find 120% of 60.
b Determine the value of x if 165% of $x = 1.5$.
c Write 2.80 as a percentage.
d Write 325% as a fraction.
e \$2000 in a bank account increases to \$5000 over a period of time. By how much has the amount increased as a percentage?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

2C Percentage increase and decrease

CONSOLIDATING

Learning intentions

- To understand how to form the percentage amount for a percentage increase or decrease
- To be able to increase or decrease an amount by a given percentage
- To know that percentage change is the percentage of the original amount
- To be able to calculate the percentage change

Key vocabulary: percentage

Percentages are often used to describe by how much a quantity has increased or decreased. The price of a car in the new year might be increased by 5%. On a $70 000 car, this is a $3500 increase. The price of a shirt might be marked down by 30%. If the shirt originally cost $60, this provides an $18 discount. It is important to note that the increase or decrease is calculated on the original amount.



Car sales often make use of percentages during negotiations. Understanding them will let you know if you're getting a good deal.

Lesson starter: The quicker method

Two students, Nicky and Mila, consider the question: $250 is increased by 15%. What is the final amount?

Nicky puts his solution on the board with two steps.

Step 1: 15% of $\$250 = 0.15 \times \$250$
$= \$37.50$

Step 2: Final amount $= \$250 + \$37.50$
$= \$287.50$


Mila says that the same problem can be solved with only one step using the number 1.15.

- Can you describe Mila's method? Write it down.
- What if the question was altered so that $250 is decreased by 15%. How would Nicky's and Mila's methods work in this case?
- Which of the two methods do you prefer and why?

Key ideas


- To increase an amount by a given percentage:
  - add the percentage increase to 100%
  - multiply the amount by this new percentage.

  For example: to increase by 30%, multiply by $100\% + 30\% = 130\% = 1.3$



- To decrease an amount by a given percentage:
  - subtract the percentage from 100%
  - multiply the amount by this new percentage.

  For example: to decrease by 25%, multiply by $100\% - 25\% = 75\% = 0.75$



- To find a percentage change, use:

$$\text{Percentage change} = \frac{\text{change}}{\text{original amount}} \times 100\%$$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Exercise 2C

Understanding

1–3 | 3

1 Write the missing number for these increases.

a To increase a number by 40%, multiply by _____.

b To increase a number by 26%, multiply by _____.

c To increase a number by _____, multiply by 1.6.

d To increase a number by _____, multiply by 1.21.



Hint for Q1: To increase by 40%, need 140% of amount

2 Write the missing number for these decreases.

a To decrease a number by 20%, multiply by _____.

b To decrease a number by 73%, multiply by _____.

c To decrease a number by _____, multiply by 0.94.

d To decrease a number by _____, multiply by 0.31.



Hint for Q2: To decrease by 20%, need 80% of amount

3 Find the percentage change if

a Original amount = \$120, change = \$30

b Original amount = \$35, change = \$70

Fluency

4–5(½), 6, 7 | 4–5(½), 6, 7(½), 8

Example 6 Increasing by a percentage

Increase \$70 by 15%

Solution

$100\% + 15\% = 115\%$

$= 1.15$

$\$70 \times 1.15 = \$80.50$

Explanation

First add 15% to 100%

Note that $15\% = 0.15$ and $100\% = 1$

Multiply by 1.15 to give \$70 plus the increase in one step.

Now you try

Increase \$120 by 30%

4 Complete the following.

a Increase 56 by 10%

b Increase 980 by 20%

c Increase 100 by 12%

d Increase 890 by 5%

e Increase 180 by 15%

f Increase 450 by 20%

g Increase 8 by 50%

h Increase 98 by 100%

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Example 7 Decreasing by a percentage

Decrease \$5.20 by 40%

Solution

$100\% - 40\% = 60\%$

$= 0.6$

$\$5.20 \times 0.6 = \$3.12$

Explanation

First subtract the 40% from 100% to find the percentage remaining.

Multiply by $60\% = 0.6$ to get the result.

Now you try

Decrease \$64 by 15%

5 Complete the following.

a Decrease 80 by 5%
b Decrease 600 by 10%
c Decrease 45 by 50%
d Decrease 700 by 12%
e Decrease 8000 by 8%
f Decrease 450 by 25%
g Decrease 68 by 75%
h Decrease 9000 by 1%
i Decrease 7000 by 100%
j Decrease 10 000 by 1.5%

Example 8 Finding a percentage change

a The price of a mobile phone increased from \$250 to \$280. Find the percentage increase.
b The population of a town decreases from 3220 to 2985. Find the percentage decrease and round to one decimal place.

Solution

a Increase $= \$280 - \$250$

$= \$30$

Percentage increase $= \frac{30}{250} \times \frac{100}{1}$

$= 12\%$

Explanation

First find the actual increase.

Divide the increase by the original amount and multiply by 100.

b Decrease $= 3220 - 2985$

$= 235$

Percentage decrease $= \frac{235}{3220} \times \frac{100}{1}$

$= 7.3\%$ (to 1 d.p.)

First find the actual decrease.

Divide the decrease by the original population and multiply by 100. Round as indicated.

Now you try

a The height of a plant increased from 20 cm to 28 cm. Find the percentage increase.
b The price of a washing machine decreased from \$649 to \$545. Find the percentage decrease and round to one decimal place.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

2C

6 The price of a flight increased from $125 to $150 overnight. Find the percentage increase.

7 Copy and complete the tables showing percentage change. Round to one decimal place where necessary.

a

Original amount	New amount	Increase	Percentage change
40	60		
12	16		
100	125		
24	30		
88	100		

b

Original amount	New amount	Decrease	Percentage change
90	81		
100	78		
20	15		
24	18		
150	50		

Example 9 Finding the original amount from an increase or decrease

After rain, the volume of water in a tank increased by 24% to 2200 L. How much water was in the tank before it rained? Round to the nearest litre.

Solution

$100\% + 24\% = 124\%$

124% of original value $= 2200$ L

1% of original value $= 17.74$ L

100% of original value $= 1774$ L

Alternate method: $100\% + 24\% = 124\%$

124% of $V = 2200$ L

$1.24 \times V = 2200$

$V = 2200 \div 1.24$

$= 1774$

Explanation

Write the total percentage.

The original volume is increased by 24% to give 2200 litres

Divide by 124 to find 1%

Multiply by 100 to find the original amount.

Write the total percentage.

Write an equation using the pronumeral V to represent the volume.

Divide both sides by 1.24 to solve for V.

Round to the nearest litre.

Now you try

At the start of a new year, daily public transport tickets increase by 4% to $9.20. What was the cost of a daily ticket before the increase? Round to the nearest cent.

8 Find the original cost for each of the following if:

- **a** an increase of 10% on the cost of a can of cola drink increased it to $3.30
- **b** an increase of 10% on the cost of a meal increased the cost to $88
- **c** after an increase of 5%, the cost of a pair of running shoes came to $210
- **d** a decrease of 30% made the cost of car insurance $350
- **e** a decrease of 60% brought the price of a used car down to $5000



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Problem-solving and reasoning 9–12 13–16

9 The price of a computer was decreased by 15% in a sale. What is the sale price, if the original price was $2100?

10 Plumbers on a salary of $82 570 were given a $2\frac{1}{2}\%$ pay increase. Find their new annual salary.

11 A car manufacturer intends to increase sales by 14.7% next year. If the company sold 21 390 new cars this year, how many does it expect to sell next year?



12 The length of a bike sprint race is increased from 800 m to 1200 m. Find the percentage increase.

Hint for Q12:
$$\%\text{ increase} = \frac{\text{increase}}{\text{original amount}} \times 100$$

13 The number of people on a bus decreased from 25 to 18 after one stop. Find the percentage decrease in the number of people on the bus.

14 After a price increase of 20%, the cost of entry to a museum rose to $25.80. Find the original price.



15 The total price of an item including GST (at 10%) is $120. How much GST is paid, to the nearest cent?

Hint for Q15: $120 represents 110% of the price before GST is added.

16 An investor starts with $1000.

a After a bad day the initial investment is reduced by 10%. Find the balance at the end of the day.

b The next day is better and the balance is increased by 10%. Find the balance at the end of the second day.

c The initial amount decreased by 10% on the first day and increased by 10% on the second day. Explain why the balance on the second day didn't return to $1000

Repeated increase and decrease — 17, 18

17 If the cost of a pair of shoes was increased twice, by 10% from an original price of $80 and then another 15% from this new price, the final price would be

$$\$80 \times 1.10 \times 1.15 = \$101.20$$

Use a similar technique to find the final price of these items. Round to the nearest cent.

a Skis starting at $450 and increasing by 20% and 10%

b A computer starting at $2750 and increasing by 6% and 11%

c A DVD player starting at $280 and decreasing by 10% and 25%

d A circular saw starting at $119 and decreasing by 18% and 37%

18 If an amount is increased by the same percentage each time, powers can be used. For example, 50 kg increased by 12% three times would increase to

$$50 \text{ kg} \times 1.12 \times 1.12 \times 1.12$$
$$= 50 \text{ kg} \times (1.12)^3$$
$$= 70.25 \text{ kg (to 2 d.p.)}$$



Hint for Q18: You have a power/index key on your calculator.

Use a similar technique to find the final value in these situations. Round to two decimal places.

a The mass of a rat initially at 60 grams grows at a rate of 10% every month for 3 months.

b The cost of a new car initially at $80 000 increases by 5% every year for 4 years.

c The value of an apartment initially at $380 000 decreases by 4% per year for 3 years.

d The length of a pencil initially at 16 cm decreases through being sharpened by 15% every week for 5 weeks.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

2D Profits and discounts


CONSOLIDATING

Learning intentions

- To know the terms associated with profits and discounts
- To be able to find the selling price or original cost price from a mark-up or discount
- To be able to determine the profit or loss made on a sale

Key vocabulary: profit, selling price, cost price, loss, mark-up, percentage, discount

Percentages are widely used in the world of finance. Profits, losses, commissions, discounts and taxation are often expressed and calculated using percentages.



Examples of percentages used in the media.

Lesson starter: The best discount

Two book shops are selling the same book at a discounted price. The recommended retail price for the book is the same for both shops. Each shop has a sign near the book with the given details:

- Shop A. Discounted by 25%
- Shop B. Reduced by 20% then take a further 10% off that.

Which shop offers the bigger discount?
Is the difference equal to 5% of the retail price?

Key ideas

- **Profit** is the amount of money made on a sale.
  Profit = **selling price** − **cost price**
- A **loss** is made when the selling price is less than the cost price.
  Loss = cost price − selling price
- **Mark-up** is the amount added to the cost price to produce the selling price.
  Selling price = cost price + mark-up
- The percentage profit or loss can be found by dividing the profit or loss by the cost price and multiplying by 100.

$$\%\ \text{profit/loss} = \frac{\text{profit or loss}}{\text{cost price}} \times 100$$

- **Discount** is the amount by which an item is marked down.
  Discount = % discount × original price
  New price = original price − discount amount

Exercise 2D

Understanding

1–4 | 1

1 Match the definition in the left column with the correct word in the right-hand column.

a	The amount of money made on a sale	**A**	Loss
b	The amount by which an item is marked down	**B**	Mark-up
c	The result when the selling price is less than the cost price	**C**	Profit
d	The amount added to the cost price	**D**	Discount

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

2 Write the missing numbers in the table.

Cost price ($)	7	18	460.95	3250
Selling price ($)	10	15.50	395	4430
Profit/loss ($)				



Hint for Q2: Profit/loss is the difference between cost price and selling price.

3 Copy and complete the table of mark-ups.

Cost price ($)	30.95	99.95		18 000
Mark-up ($)	10	80	700	
Selling price ($)			1499.95	26 995



Hint for Q3: Mark-up increases price.

4 Copy and complete the table of discounts.

Original price ($)	100	29.95		2215
New price ($)	72	22.70	176	
Discount ($)			23	178



Hint for Q4: Discount reduces price.

Fluency

5–8 5–8

Example 10 Calculating selling price from mark-up

An electrical store marks up all entertainment systems by 30%.
If the cost price of one entertainment system is $8000, what will be its selling price?

Solution

Mark-up = 30% of $8000
= 0.3 × 8000
= $2400

Selling price = $8000 + $2400
= $10 400

Alternatively, multiply cost price by 130% or 1.3.

Explanation

Change percentage to a decimal or fraction and multiply by the cost price.

Selling price = cost price + mark-up

130% is a 30% increase on the cost price

Now you try

A computer store marks up all notebook computers by 24%.

If the cost price of one notebook is $1200, what will be its selling price?

5 Copy and complete this table by calculating the selling price of each item.

Item	Cost price	% mark-up	Selling price
Jeans	$60	28%	
Toaster	$40	80%	
Car	$22 000	45%	
Can of drink	$1.20	140%	
Loaf of bread	$1.80	85%	
Handbag	$80	70%	
Electronic tablet	$320	35%	



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

2D

Example 11 Finding the discount amount

An electrical store advertises a 15% discount on all equipment as a holiday special. Find the sale price on a projection system that has a marked price of $18 000.

Solution

Discount = 15% of $18 000
= 0.15 × 18 000
= $2700

The new price = $18 000 − $2700
= $15 300

Alternatively, multiply the original price by 85% or 0.85.

Explanation

Change the percentage to a decimal and evaluate.

New price is original price minus discount.

A 15% discount leaves 85%.

Now you try

A department store offers a post-Christmas discount of 35% on all decorations. Find the sale price on a wreath that has a marked price of $80.

6 Copy and complete the table by writing in the missing values.

Item	Cost price	% discount	Selling price
Camera	$900	15%	
Car	$24 000	20%	
Bike	$600	25%	
Shoes	$195	30%	
Blu-ray player	$245	50%	
Electric razor	$129	20%	
Lawn mower	$880	5%	



Example 12 Determining profit

A manufacturer produces an item for $400 and sells it for $540.

a Determine the profit made.
b Express this profit as a percentage of the cost price.

Solution

a Profit = $540 − $400
= $140

b $\%\ \text{profit} = \frac{140}{400} \times 100$
$= 35\%$

Explanation

Profit = selling price − cost price

$\%\ \text{profit} = \frac{\text{profit}}{\text{cost price}} \times 100$

Now you try

A jeweller produces a necklace for $36 and sells it for $49.50.

a Determine the profit made.
b Express this profit as a percentage of the cost price.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

7 Find the missing values in these tables by finding the profit or loss and expressing this as a percentage of the cost price. Round to two decimal places where necessary.

a

Cost price ($)	Selling price ($)	Profit ($)	Profit (%)
10	15		
24	30		
100	150		
250	255		
17.50	20		

b

Cost price ($)	Selling price ($)	Loss ($)	Loss (%)
10	8		
16	12		
100	80		
34	19		
95	80.75		

Example 13 Calculating the original price

A toy store discounts a toy by 10% in a sale. If the sale price was $10.80, what was the original price?

Solution

90% of the original = $10.80

1% of the original = 10.80 ÷ 90

1% of the original = 0.12

100% of the original = $12

The original price was $12.

Alternate method: 90% of P = $10.80

$0.9 \times P = \$10.80$

$P = \$10.80 \div 0.9$

$= \$12$

Explanation

The discount factor = 100% − 10% = 90%.
Thus, $10.80 is 90% of the original price.

Use the unitary method to find 1%.

Multiply by 100 to find the original amount.

Write the answer in words.

Write an equation using the pronumeral P to represent the original price.

Solve the equation for P by dividing both sides by 0.9

Now you try

An outdoor equipment store discounts a tent by 20% in a sale.

If the sale price was $176, what was the original price?


8 Answer the following questions relating to finding the original price (cost price).

a Find the original price if a coffee mug was discounted by 20% and sold for $4.40.

b Find the cost price of a pair of shoes that sold for $250 after a mark-up of 25%.

c Find the cost price after a discount of 10% was given on a surfboard that sold for $1350.

d Find the original price on a concert ticket for a major recording artist if it was marked up by 100% and sold for $250.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

2D

Problem-solving and reasoning 9–11 11–13

9 A manufacturer produces and sells items for the prices shown.
- **i** Determine the profit made.
- **ii** Express this profit as a percentage of the cost price.

a Cost price $10, selling price $12
b Cost price $20, selling price $25
c Cost price $120, selling price $136.80
d Cost price $1400, selling price $3850

10 Lenny marks up all computers in his store by 12.5%. If a computer cost him $890, what will be the selling price of the computer?



11 A used-car dealer purchases a vehicle for $13 000 and sells it for $18 500. Determine the percentage mark-up on the vehicle to one decimal place.


Hint for Q11: What is the increase as a percentage of the original price?

12 A refrigerator is discounted by 25%. If Paula pays $460 for it, what was the original price? Round to the nearest cent.

13 An electrical store buys a computer from the wholesaler for $500. The store marks up the computer by 80%.
a What is the amount of the mark-up in dollars?
b What is the retail price of the computer after the store's mark-up?
c The store offers the computer on sale for a discount of 15%. What is the price now?
d If the 15% discount was calculated on the original $500 cost price, and then the computer was marked up 80% after that, would it make a difference to the sale price?

House value — 14

14 The graph shows the changes in the value of a particular house in southern Sydney.





a How much was the house worth in 2006?
b Find the percentage increase, to one decimal place, in the value of the house from 2006 to:
i 2010 **ii** 2012 **iii** 2014
c In 2026 the owners hope to sell for 1.25 million dollars. What percentage increase is needed in the two years from 2024 to 2026 so they could sell for that price?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

2E Income

Learning intentions

- To understand the different ways that workers can be paid
- To know how wages, overtime, salaries and commission work
- To be able to use wages and salaries to calculate hourly and yearly incomes
- To be able to calculate overtime earnings
- To be able to calculate commission on sales amounts

Key vocabulary: wage, overtime, time and a half, double time, salary, commission, retainer

There are many different ways of earning a living. You can be self-employed or work for someone else. Your income is usually related to the skills you have. It can be calculated and paid in different ways. You can earn, for example, a salary, a wage, a commission or possibly a royalty.

Lesson starter: Types of income

As a class, write down one example of a job that earns each of the following types of income.

- Salary
- Wage (i.e. hourly rate of pay)
- Overtime
- Commission

In the newspaper classifieds or online, find a job advertisement for each of the income types above.

Key ideas

- Workers who earn a **wage** (for example, a casual waiter) are paid a fixed rate per hour. Hours outside the normal working hours (public holidays etc.) are paid at a higher rate called **overtime**. This can occur in a couple of common ways:
  - **Time and a half:** pay is 1.5 times the usual hourly rate
  - **Double time:** pay is twice the usual hourly rate
- Workers who earn a **salary** (for example, an engineer) are paid a fixed amount per year, say, $125 000. This is often paid monthly or fortnightly.
  - 12 months in a year and approximately 52 weeks in a year = 26 fortnights
- **Commission** is a proportion of the overall sales amount. Salespeople may receive a commission on their sales as well as a set weekly or monthly fee called a **retainer**.
  - Commission = % commission × total sales

Exercise 2E

Understanding

1–4 | 1, 4

1 Match the pay description (**a**–**d**) with the income type (**A**–**D**).

a	Sally earns $22.70 per hour working in a shop	**A**	Overtime
b	Maths teacher Stephanie earns $72 000 per year	**B**	Commission
c	Ross earns 5% of all sales he makes	**C**	Wage
d	Matt earns time and a half pay on Saturdays	**D**	Salary

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

2E

2 Tom earns $12.70 an hour. How much does he earn for:
a 2 hours of work? **b** 8 hours of work? **c** 38 hours of work?

3 Sela's hourly rate of pay is $24. Calculate her overtime rate at:
a time and a half **b** double time



4 William earns a salary of $136 875 each year. Approximately how much is this:
a each month?
b each week (to the nearest cent)?
c each day?

Hint for Q4: To find approximate monthly salary, divide annual salary by 12.
To find approximate weekly salary, divide annual salary by 52.
To find approximate daily salary, divide annual salary by 365.

Fluency 5–7, 8(½), 9 | 5, 6, 8(½), 9

Example 14 Comparing wages and salaries

Ken earns an annual salary of $90 000 and works a 38-hour week. His wife Brooke works part time in retail and earns $61.80 per hour.
a Calculate how much Ken earns per week.
b Determine who has the higher hourly rate of pay.
c If Brooke works on average 18 hours per week, what is her yearly income?

Solution	Explanation
a Weekly rate = $90 000 ÷ 52	$90 000 pay in a year.
= $1730.77	There are approximately 52 weeks in a year.
∴ Ken earns $1730.77 per week	Divide by 52 to find the weekly wage.
b Brooke: $61.80/h	Ken works 38 hours in the week.
Ken: $1730.77 ÷ 38	Hourly rate = weekly rate ÷ number of hours. Round to the nearest cent.
= $45.55/h	
∴ Brooke is paid more per hour.	Compare hourly rates.
c In one week: $61.80 × 18	Weekly income = hourly rate × number of hours
= $1112.40	
Yearly income = $1112.40 × 52	Multiply by 52 weeks to get yearly income.
= $57 844.80	

Now you try

Mali earns an annual salary of $77 200 and works a 38-hour week. Her partner Ben works part time as a photographer and earns $75 per hour.
a Calculate how much Mali earns per week.
b Determine who has the higher hourly rate of pay.
c If Ben works on average 15 hours per week, what is his yearly income from photography?


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5 Talib earns $58 000 per year at a fast food restaurant. His sister works part time as a waitress and earns $33.20 per hour.

- **a** How much does Talib earn each week?
- **b** Each week Talib works 38 hours. Calculate if his hourly rate of pay is higher than his sister's.
- **c** If his sister averages 10 hours of work per week, what is her yearly income?

6 Paul earns $790 each week. How much does he earn each:

- **a** year?
- **b** month?
- **c** hour, if he worked 40 hours each week?



7 Copy and complete the table.

Employee	Hourly rate	Hours worked	Income
Adam	$20.40	8	
Betty	$15.50	$8\frac{1}{2}$	
Ceanna	$19.70	15	
David	$24.30	38	
Edward	$57.85	42	
Francis	$30	27	
George	$35.20	7.25	

Example 15 Calculating overtime

Georgio works some weekends and late nights, and earns overtime for that work. His hourly rate of pay is $32 per hour.

- **a** Calculate Georgio's time and a half rate of pay per hour.
- **b** Calculate Georgio's double time rate of pay per hour.
- **c** Calculate Georgio's weekly wage for a week where he works 18 hours at his normal rate, 2 hours at time and a half, and 1 hour at double time.

Solution

a Time and a half $= \$32 \times 1.5$
$= \$48$

b Double time $= \$32 \times 2$
$= \$64$

c Wage $= \$32 \times 18 + \$48 \times 2 + \$64 \times 1$
$= \$736$

Explanation

Time and a half is 1.5 times the hourly rate.

Double time is 2 times the hourly rate.

Find the sum of:

- the normal hourly rate ($32) multiplied by the number of hours worked at the normal rate (18)
- the time and a half hourly rate ($48) multiplied by the number of hours worked at that rate (2)
- the double time hourly rate ($64) multiplied by the number of hours worked at that rate (1).

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

2E

Now you try

Kane works some weekends and some public holidays and earns overtime for that work. His hourly rate of pay is $24 per hour.

a Calculate Kane's time and a half rate of pay per hour.
b Calculate Kane's double time rate of pay per hour.
c Calculate Kane's weekly wage for a week where he works 15 hours at his normal rate, 6 hours at time and a half and 3 hours at double time.

8 A job has a normal working hours pay rate of $29.20 per hour. Calculate the pay, including overtime, from the following hours worked.

a 3 hours at the normal rate and 4 hours at time and a half
b 4 hours at the normal rate and 6 hours at time and a half
c 14 hours at the normal rate and 3 hours at double time
d 20 hours at the normal rate and 5 hours at double time
e 10 hours at the normal rate and 8 hours at time and a half and 3 hours at double time
f 34 hours at the normal rate and 4 hours at time and a half and 2 hours at double time



Hint for Q8:
For time and a half: × 1.5
For double time: × 2

Example 16 Calculating commission

A part-time saleswoman is paid a retainer of $1500 per month. She also receives a commission of 5% on the value of goods she sells. If she sells goods worth $5600 during the month, calculate her earnings for that month.

Solution

Commission = 5% of $5600
= 0.05 × $5600
= $280

Earnings = $1500 + $280
= $1780

Explanation

Calculate the commission on sales. Change the percentage to a decimal and evaluate.

Earnings = retainer + commission

Now you try

A real estate agent is paid a retainer of $2200 per month. She also receives a commission of 0.4% on the value of houses she sells. If she sells houses worth $1 575 000 during the month, calculate her earnings for that month.

9 Copy and complete the table.

Person	Weekly retainer	Rate of commission	Commission earned ($)	Weekly wage ($)
Adina	$0	12% on $7000		
Byron	$160	8% on $600		
Cindy	$300	5% on $680		
Deanne	$260	5% on $40 000		
Elizabeth	$500	8% on $5600		
Faruq	$900	2% on $110 000		
Gary	$1000	1.5% on $45 000		

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Problem-solving and reasoning 10–12 11, 13–15

10 Calculate how many hours at the standard hourly rate the following working hours are the same as:

a 3 hours and 2 hours at double time

b 6 hours and 8 hours at time and a half

c 15 hours and 12 hours at time and a half



Hint for Q10: 2 hours at double time is 4 hours at the standard hourly rate.



Hint for Q11: Calculate the number of hours at the standard hourly rate.

11 Jim, a part-time gardener, earned $522 in a week. If he worked 12 hours during normal working hours and 4 hours overtime at time and a half, what was his hourly rate of pay?

12 Sally earned $658.80 in a week. She worked 10 hours during the week, 6 hours on Saturday at time and a half and 4 hours on Sunday at double time. What was her hourly rate of pay?

13 Amy works at Best Bookshop. During one week she sells books valued at $800. If she earns $450 per week plus 5% commission, how much does she earn in this week?



14 Jason works for a campervan company. If he sells $84 000 worth of campervans in a month, and he earns $1650 per month plus 4% commission on sales, how much does he earn that month?

15 Stephen earns an hourly rate of $34.60 for the first 38 hours, time and a half for the next 3 hours and double time for each extra hour above that. Calculate his earnings if he works 44 hours in a week.

Pay Slips — 16

16 Workers at a fast food restaurant are paid $21.63 per hour for working Monday to Friday up until 7 p.m. and time and a half after 7 p.m. They earn time and a half on Saturdays and double time on Sundays. They are given an unpaid 30 minute meal break for any shift over 5 hours.
Donna's shifts for the week are given below.

Monday	Tuesday	Wednesday	Thursday	Friday	Saturday	Sunday
–	4.30 p.m.–	4.30 p.m.–	5 p.m.–	5 p.m.–	10 a.m.–	10 a.m.–
–	7 p.m.	7.30 p.m.	10 p.m.	8 p.m.	6.30 p.m.	1.30 p.m.

a How much can Donna expect to earn, to the nearest cent, if she works the hours she is rostered for?

b The restaurant decides on a new workplace deal, gets rid of all overtime and creates a flat hourly rate of $24 per hour. How much worse off will Donna be for the week above?

c How many extra hours a week does Donna need to work during the week to make up the extra income? Answer in a whole number of hours.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Progress quiz

2A **1** Express the following in the required form.

a 32% as a simplified fraction
b 7% as a decimal
c 0.23 as a percentage
d 0.125 as a percentage
e $\frac{7}{20}$ as a percentage
f 4.5% as a simplified fraction

2A **2** Express the following as percentages.

a 20c out of $2
b 32 marks out of 40
c 5 kg out of 40 kg

2B **3** Find the following amounts.

a 20% of 80 km
b 35% of $120
c $33\frac{1}{3}$% of 180 g
d 4.2% of $500

2B **4** Determine the original amount if:

a 25% of the amount is $100
b 12% of the amount is $36
c 8% of the amount is $60

2C **5** Complete the following.

a Increase 82 by 10%
b Increase 220 by 8%
c Decrease 110 by 15%
d Decrease 250 by 6%

2C **6** **a** The number of students in a school increased from 450 to 540. Find the percentage increase.

b The cost of petrol dropped from 121c per litre to 103c per litre overnight. Find the percentage decrease correct to one decimal place.

2C **7** After offering a special ticket deal for students, a concert's ticket sales rose by 24% to 961 ticket sales. How many tickets had been sold before the ticket deal?

2D **8** Find the selling price of the following items.

a A $1200 tablet computer marked up by 16%
b A pair of $126 sneakers discounted by 20%

2D **9** Jane has a stall at the market selling bracelets that she makes. If each one costs $6.20 to produce and she sells them for $10, find her percentage profit correct to one decimal place.

2D **10** Find the original price if:

a a dress was discounted by 15% and sold for $187
b a computer game was marked up by 12% and sold for $72.80

2E **11** Find the weekly earnings in the following work situations.

a Seb earns an annual salary of $74 100. (Assume 52 weeks in the year.)
b Jodie works 22 hours in a week at $18.20 per hour.
c Ned works in a fast food restaurant and has a normal hourly rate of $15.10 per hour. In a week he works 4 hours at the normal rate, 3 hours at time and a half and 2 hours at double time.
d Harper is paid a weekly retainer of $200 and she also earns a 5% commission on her sales. In a particular week she makes $8400 worth of sales.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

2F Taxation

Learning intentions

- To know the difference between gross income and net income
- To understand how taxable income is determined
- To know what is meant by income tax
- To know how to calculate net income from gross income and deductions
- To be able to calculate tax as a percentage of the taxable income
- To be able to calculate income tax using the tax table from the Australian Taxation Office

Key vocabulary: gross income, net income, taxable income, income tax, deductions

All wage and salary earners have some deductions taken out of their pay. Deductions usually include income tax. Tax is paid to the government. The government uses it to pay for community welfare, education and a number of other services.



Calculating your own tax is an important life skill.

Lesson starter: Deductions from pay

What types of deductions can you and your class think of that might be taken out of someone's pay?

Discuss, as a class, what superannuation and the Medicare levy is, and who pays it.

Key ideas

- **Gross income** = the total of all income
- **Net income** = gross income minus **deductions** (expenses that reduce income)
- **Taxable income** = gross income minus tax deductions
- **Income tax** is paid to the government once a person's taxable income passes a set amount in the year.
- Income tax is calculated by the current tax table available through the Australian Taxation Office. The following rates apply as of 2024/2025.

Taxable income	Tax on this income
\$0 – \$18 200	Nil
\$18 201 – \$45 000	16c for each \$1 over \$18 200
\$45 001 – \$135 000	\$4288 plus 30c for each \$1 over \$45 000
\$135 001 – \$190 000	\$31 288 plus 37c for each \$1 over \$135 000
\$190 001+	\$51 638 plus 45c for each \$1 over \$190 000

* Does not include the Medicare levy (2% of taxable income).

Exercise 2F

Understanding

1–4 | 4

1 Copy and complete the table by inserting the net income amounts.

Gross income	Deductions	Net income
\$5600	\$450	
\$87 000	\$28 000	
\$50 000	\$6700	



Hint for Q1: Recall that net income = gross income minus deductions.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

2F

2 Find Hoang's gross weekly income if he earns:
- $1200 a week as a teacher
- $60 an hour for tutoring, and he tutors for 3 hours a week
- $25 in interest on his bank account per week

3 John has a gross income of $45 000 and a net income of $20 000. How much did his deductions come to?

4 Use the table in the Key ideas to interpret the tax paid on an income of
a $12 000 **b** $37 000 **c** $80 000

Fluency

5–6(½), 7, 8(½) | 5–6(½), 8(½)

Example 17 Calculating tax to find net income

Liam starts a new job with an annual salary of $52 800. His payslip each month shows deductions for taxation of $968.

a Calculate Liam's net income each month.
b What percentage of Liam's monthly pay is being paid to the government by his employer for taxation?
c Liam's salary is increased to $65 000 and the taxation rate for Liam's salary changes to 24% with the first $18 200 tax free. Calculate Liam's net income for the year.

Solution | **Explanation**

a Monthly pay $= \$52\,800 \div 12$
$= \$4400$

Calculate gross income per month.

$\therefore$ net monthly income $= \$4400 - \$968$
$= \$3432$

Net income = gross income − taxation

b $\%\ \text{tax} = \frac{968}{4400} \times 100$
$= 22\%$

Calculate what fraction $968 is of the monthly income $4400. Multiply by 100 to convert to a percentage.

c Salary for tax purposes $= \$65\,000 - \$18\,200$
$= \$46\,800$

First $18 200 is not taxed.

Tax amount $= 24\%$ of $\$46\,800$
$= 0.24 \times \$46\,800$
$= \$11\,232$

Calculate tax amount on $46 800. Convert percentage to a decimal and evaluate.

$\therefore$ net income $= \$65\,000 - \$11\,232$
$= \$53\,768$

Net income = gross income − tax amount.

Now you try

Poppy has a job with an annual salary of $74 400. Her payslip each month shows deductions for taxation of $1488.

a Calculate Poppy's net income each month.
b What percentage of Poppy's monthly pay is being paid to the government by her employer for taxation?
c Poppy's salary is increased to $82 800 and the taxation rate for Poppy's salary changes to 28% with the first $18 200 tax free. Calculate Poppy's net income for the year.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5 For each of the following find:
- **i** the annual net income
- **ii** the percentage of gross income paid as tax. Round to one decimal place where necessary.

a Gross annual income = $48 241, tax withdrawn = $8206
b Gross annual income = $67 487, tax withdrawn = $13 581.20
c Gross monthly income = $4041, tax withdrawn = $606.15
d Gross monthly income = $3219, tax withdrawn = $714.62



Hint for Q5: Net income = gross income − deductions

6 Calculate the amount of tax to be paid using the following annual salaries and tax rates if the first $18 200 is tax free.
a salary = $30 400, tax rate = 15%
b salary = $56 500, tax rate = 21%
c salary = $69 700, tax rate = 24.5%
d salary = $96 400, tax rate = 30.4%



Hint for Q6: Remember to subtract the $18 200 to find the salary to calculate tax on.

7 Ed earns $1400 per week and pays 27% of his annual income in tax.
a Calculate the amount of income tax that Ed pays in one year.
b Find Ed's annual net income.

Example 18 Using the tax table

Use the tax table below to find the income tax for an income of $84 000.

Taxable income	Tax on this income
$0 − $18 200	Nil
$18 201 − $45 000	16c for each $1 over $18 200
$45 001 − $135 000	$4288 plus 30c for each $1 over $45 000
$135 001 − $190 000	$31 288 plus 37c for each $1 over $135 000
$190 001+	$51 638 plus 45c for each $1 over $190 000

* Does not include the Medicare levy (2% of taxable income).

Solution

Tax = $4288 + 0.3 × ($84 000 − $45 000)
= $4288 + $11 700
= $15 988

Explanation

Find the tax bracket in which the amount ($84 000) lies ($45 001 − $135 000).

Write down the values in this bracket, remembering that 30c in the dollar is 30%, or 0.3.

Subtract 45 000 from 84 000 to find the amount of income that the 30% applies to.

Use your calculator to find the answer.

Now you try

Use the tax table above to find the income tax for an income of $75 000.

8 Use the tax table in Example 18 to find the income tax payable on each of these incomes.

a $10 000 **b** $30 000 **c** $80 000
d $156 000 **e** $200 000 **f** $1 000 000



Hint for Q8: Find the right category (tax bracket) first.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

2F

Problem-solving and reasoning

9, 10 | 9–12



Hint for Q9: What % of gross income does $53 246 represent?

9 Mel has a net annual income of $53 246 after 21% of her income is withdrawn for tax purposes. What was her gross income?

10 Fred earns $50 000 each year as a shop clerk and an extra $1200 each year for doing gardening on some weekends.

- **a** Calculate his annual gross salary.
- **b** Use his gross salary to find his income tax, using the tax table in Example 18.
- **c** What is his annual net income, after tax?
- **d** What is his approximate fortnightly net income, to the nearest cent, after tax?



11 William earns $1600 a week. Using the tax table in Example 18, how much tax is deducted from his pay each week? Assume 52 weeks in a year.

12 John pays $5000 in tax a year.

- **a** Using the tax table in Example 18, which tax bracket does John fall into?
- **b** Work backwards from this amount to work out how much John earns per year. Answer to the nearest dollar.

Other types of deductions and taxable income

— | 13–15

The lower a person's income, the less tax they must pay. People therefore try to lower their taxable income by claiming allowable tax deductions.

Work-related expenses such as uniforms, stationery and job-related travel expenses are all examples of allowable tax deductions. The income tax is therefore calculated on what we call a person's taxable income.

Taxable income = gross income − allowable tax deductions

Use the tax table from Example 18 to compare the following jobs.

13 An young accountant earns $3120 a fortnight, with no allowable tax deductions.

- **a** What is the accountant's taxable income?
- **b** How much tax does the accountant owe in a year?
- **c** What is the accountant's annual net pay?
- **d** What is the accountant's fortnightly net pay?

14 A young lawyer earns $6520 a fortnight with allowable tax deductions totalling $2560 a year.

- **a** What is the lawyer's taxable income?
- **b** How much tax does the lawyer owe in a year?
- **c** What is the lawyer's annual net pay?
- **d** What is the lawyer's fortnightly net pay?



15 How much more money does the lawyer in question **14** have each week compared to the accountant in question **13**?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

2G Simple interest

Learning intentions

- To understand how simple interest is calculated
- To be able to calculate simple interest and the final amount
- To be able to calculate simple interest using different time periods

Key vocabulary: simple interest, principal, per annum

Interest is charged when a person or institution borrows money. The interest is an extra amount that must be paid back, on top of the borrowed amount.

Interest is also earned, when a person or institution invests money.

Simple or flat rate interest is usually charged or earned each year. It is calculated on the full amount borrowed or invested at the beginning of the loan.



You can calculate the interest you would earn on savings or pay on a loan.

Lesson starter: Developing the rule

$\$5000$ is invested in a bank and 5% simple interest is paid to the investor every year. In the table below, the amount of interest paid is shown for Year 1, and the amount of total interest is shown for Years 1 and 2.

- Complete the table.
- How much interest would the investor earn in 10 years?

Year	Interest paid that year	Total interest
0	$\$0$	$\$0$
1	$\frac{5}{100} \times \$5000 = \$250$	$1 \times \$250 = \$250$
2		$2 \times \$250 = \$500$
3		
4		
5		

Key ideas

- **Simple interest** is interest calculated each time period on the initial amount.
- To compute simple interest, we apply the formula:

$$I = \frac{Prt}{100} \text{ or } I = P \times \frac{r}{100} \times t$$

I is the amount of simple interest (in \$)
P is the **principal** amount; the money borrowed or invested (in \$)
r is the annual interest rate expressed as a percentage
t is the number of years

- The total amount ($\$A$) equals the principal plus interest

$$A = P + I$$

- p.a. means '**per annum**' or 'per year'.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Exercise 2G

Understanding

1–3 | 1, 2

1 $12 000 is invested at 6% p.a. for 42 months.
- **a** What is the principal amount?
- **b** What is the interest rate?
- **c** What is the time period in years?

2 Jann earns $560 p.a. in simple interest on an investment. How much would he earn on the investment in:
- **a** 2 years?
- **b** 5 years?
- **c** 10 years?



3 Use the rule $I = \frac{Prt}{100}$ to find the simple interest (I) earned in these financial situations.
- **a** $P = \$10\,000$, $r = 10$, $t = 3$
- **b** $P = \$6000$, $r = 12$, $t = 5$
- **c** $P = \$5200$, $r = 4$, $t = 2$

Hint for Q3:
I = interest
P = principal
r = interest rate
t = time (years)

Fluency

4–6, 8 | 4, 6–8

Example 19 Using the simple interest formula

Calculate the simple interest earned if the principal is $1000, the rate is 5% p.a. and the time is 3 years.

Solution	Explanation
$P = 1000, r = 5, t = 3$	List the information given.
$I = \frac{Prt}{100}$	Write the formula and substitute the given values.
$= \frac{1000 \times 5 \times 3}{100}$	Cancel the zeros, leaving $10 \times 5 \times 3$.
$= 150$	
Interest = $150	Answer the question.

Now you try

Calculate the simple interest earned if the principal is $2000, the rate is 4% p.a and the time is 5 years.

4 Find the simple interest earned on:
- **a** $5000 at 6% p.a. for 1 year
- **b** $5000 at 6% p.a. for 3 years
- **c** $8000 at 4% p.a. for 5 years
- **d** $15 000 at 3% p.a. for 7 years
- **e** $7250 at 5.5% p.a. for 3 years



5 Wally invests $15 000 at a rate of 6% p.a. for 3 years. Calculate the simple interest and the amount available at the end of 3 years.

Hint for Q5:
Amount = principal + interest

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Example 20 Using other time periods

Calculate the simple interest on \$7000 invested at $6\frac{1}{4}\%$ p.a. for 18 months.


Solution	Explanation
$P = 7000, r = 6\frac{1}{4} = 6.25$	List the information.
$t = 18$ months $= \frac{18}{12}$ or 1.5 years	Convert 18 months into years by dividing by 12.
$I = \frac{Prt}{100}$	Write the formula.
$I = \frac{7000 \times 6.25 \times 1.5}{100}$	Substitute in the values and evaluate.
$= 656.25$	
Interest = \$656.25	

Now you try

Calculate the simple interest on \$4500 invested at $5\frac{1}{2}\%$ for 30 months.

6 Calculate the simple interest earned on:

- **a** \$500 at 7% p.a. for 18 months
- **b** \$1000 at 5% p.a. for 24 months
- **c** \$2000 at 4% p.a. for 6 months
- **d** \$4700 at $4\frac{1}{2}\%$ p.a. for 15 months (Round to the nearest cent.)
- **e** \$50 000 at 3.75% p.a. for 200 days (Round to the nearest cent.)



Example 21 Calculating the final balance

Allan and Rachel plan to invest some money for their child Kaylan. They invest \$4000 for 30 months in a bank that pays 4.5% p.a. Calculate the simple interest and the amount available at the end of the 30 months.

Solution	Explanation
$P = 4000, r = 4.5, t = \frac{30}{12} = 2.5$	t is written in years since interest rate is per annum.
$I = \frac{Prt}{100}$	Write the formula, substitute and evaluate.
$= \frac{4000 \times 4.5 \times 2.5}{100}$	
$= 450$	
Interest = \$450	
Total amount = \$4000 + \$450 = \$4450	Total amount = principal + interest

Now you try

Joy wins some money which she decides to invest. She invests the \$5000 for 36 months in a bank account that pays 3.8% p.a. Calculate the simple interest and the amount available at the end of the 36 months.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

2G

7 Annie invests $22 000 at a rate of 4% p.a. for 27 months. Calculate the simple interest and the amount available at the end of 27 months.

8 Copy and complete the table.

	Principal, P ($)	Annual interest, rate, r	Time period, t	Interest, I	Final balance, $A = P + I$
a	7000	3%	4 years		
b	1500	7%	8 years		
c	40 000	2.5%	18 months		
d	70 000	$3\frac{1}{4}$%	2 years		
e	2000	4%	30 months		

Problem-solving and reasoning

9, 10 | 10, 11

9 A finance company charges 14% p.a. simple interest. If Lyn borrows $2000 to be repaid over 2 years, calculate her total repayment.

10 Markus borrows $20 000 to buy a car. He is charged simple interest at 18% p.a. for a period of 5 years.

a How much interest is Markus charged each year?

b Calculate the total interest Markus will pay on this loan.

c What is the total amount that Markus will have paid at the end of the loan period?

11 Wendy wins $5000 during a chess tournament. She wishes to invest her winnings, and has the two choices given below. Which one gives her the greater total at the end of the time?
Choice 1: 8.5% p.a. simple interest for 4 years
Choice 2: 8% p.a. simple interest for 54 months



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Simple interest tables and graphs — 12, 13

12 The table shows the amount of simple interest payable (in $, to the nearest $) on loans at a certain interest rate.

Amount of loan	1 year	2 years	3 years	5 years	10 years
50	6	13	19	32	65
100	13	26	39	65	129
500	65	129	194	323	645
1000	129	258	387	645	1290
5000	645	1290	1935	3225	6450
10 000	1290	2580	3870	6450	12 900
50 000	6450	12 900	19 350	32 250	64 500
100 000	12 900	25 800	38 700	64 500	129 000


Use the table to find the interest payable on the following loans.

a $5000 for 1 year
b $500 for 3 years
c $100 for 10 years
d $150 for 1 year
e $85 500 for 5 years
f $9550 for 10 years
g $5000 for 4 years
h $50 000 for 9 years
i $100 000 for 10 years

13 The graph on the right shows the annual interest earned on investments for interest rates of 4% p.a. and 6% p.a. Use the graph to answer the following.

a Find the annual interest earned on an investment of:
  i $300 at 4% p.a.
  ii $520 at 6% p.a.
  iii $250 at 4% p.a.

b What investments would earn annual interest of (to the nearest $5):
  i $20 at 6% p.a.?
  ii $20 at 4% p.a.?
  iii $14 at 6% p.a.?



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

2H Applications of simple interest

Learning intentions

- To be able to use the simple interest formula to calculate the investment period, interest rate or principal
- To be able to calculate repayment amounts given the terms of the deal

Key vocabulary: simple interest, principal, repayment

Financial calculations are a critical component of the thinking behind the decisions people make as to where to borrow or invest money.



Understanding how loans work is essential for making smart financial decisions.

Lesson starter: Where do I invest?

Bank A: $4000 at 5% p.a. for 8 years
Bank B: $5000 at 8% p.a. for 4 years
Bank C: $8000 at 4% p.a. for 5 years

Calculate the simple interest earned on each investment option. What do you notice? Which bank would you choose and why?

Key ideas

- The simple interest formula, $I = \frac{Prt}{100}$, can be used to calculate any of I, P, r or t depending on what other information is given.
  - To find an unknown, substitute the given information and solve for the unknown.
- To repay a loan, you must repay the amount borrowed (the principal) and the interest.
- **Repayments** are the amount of money, usually the same amount each time period, used to repay a loan.

Exercise 2H

Understanding

1–4 | 4

1 If Phil earns $100 in simple interest in one year, how long would it take him to earn:
a $200 **b** $400 **c** $5000 **d** $250



Hint for Q2: How much interest is earned on $100 at 10% p.a. for 1 year?

2 How many years does it take for $100 to earn $50 in interest if the simple interest rate is 10% p.a.?

3 Jodie repays $200 a month for 12 months to pay back her loan and interest. How much does she repay?

4 A loan of $4000 has interest of $500 added to it. Calculate the size of each of the 10 repayments needed to repay the loan.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Fluency

5, 6, 8, 9 | 5, 7, 8, 9

Example 22 Determining the investment period

Remy invests $2500 at 8% p.a. simple interest, for a period of time, to produce $50 interest. For how long did she invest the money?

Solution

$I = 50, P = 2500, r = 8$

$I = \frac{Prt}{100}$

$50 = \frac{2500 \times 8 \times t}{100}$

$50 = 200t$

$\therefore t = \frac{50}{200}$

$= 0.25$

Time $= 0.25$ years

$= 0.25 \times 12$ months

$= 3$ months

Explanation

List the information.

Write the formula.

Substitute the known information and simplify.

Solve the remaining equation for t by dividing both sides by 200.

Convert decimal time to months where appropriate.

Now you try

Jas invests $3000 at 6% p.a. simple interest, for a period of time, to produce $450 interest. For how long did he invest the money?

5 Alvi invests $5000 at 8% p.a. simple interest and wants to earn $1200 in interest. For how many years should Alvi invest his money?

6 Sam earns $288 interest on his $1600 investment. If the interest was calculated at 4% p.a., how many years did Sam invest the money for?

7 $8000 earns $600 interest at 5% p.a. over how many months?

Example 23 Determining the interest rate

Bank East advertises $450 interest a year on an investment of $7500. Calculate the simple interest rate for this investment.

Solution

$I = 450, P = 7500, t = 1, r = ?$

$I = \frac{Prt}{100}$

$450 = \frac{7500 \times r \times 1}{100}$

$450 = 75 \times r$

$r = 450 \div 75$

$r = 6$

Interest rate is 6% p.a.

Explanation

List the information.

Write the formula and substitute $I = 450$, $P = 7500$ and $t = 1$.

Simplify and solve for r.

Write r as a percentage.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

2H

Now you try

Oz Loans offers $610 interest a year on an investment of $12 200. Calculate the simple interest rate for this investment.

8 Find the annual simple interest rate needed for each of the following situations.

- **a** $4000 earns $500 in 2 years
- **b** $500 earns $120 in 12 years
- **c** $18 000 earns $3510 in 3 years
- **d** $950 earns $470.25 in 9 years
- **e** $3000 earns $945 in 18 months
- **f** $2500 earns $393.75 in 4.5 years



Hint for Q8: Set up the formula $I = \frac{Prt}{100}$ where r is the unknown.

Example 24 Calculating repayments

'Deals 4 You' offers a loan of $24 000 at 16% p.a. simple interest if the loan is repaid in equal monthly instalments over 5 years.

- **a** How much interest is charged on the loan?
- **b** What is the total amount of the loan and the interest?
- **c** Calculate the size of each repayment.

Solution

a $I = \frac{Prt}{100}$

$P = 24\,000, r = 16, t = 5$

$\therefore I = \frac{24\,000 \times 16 \times 5}{100}$

$= \$19\,200$

b Total $= \$24\,000 + \$19\,200$

$= \$43\,200$

c Repayments $= \frac{43\,200}{60}$

$= 720$

Repayments come to $720 a month.

Explanation

Write down the simple interest formula and list the information.

Substitute in the values and evaluate.

Total = principal + interest

Divide the total amount by the number of months in 5 years ($5 \times 12 = 60$).

Now you try

A loan of $35 000 is repaid in equal monthly instalments over 6 years at 8% p.a. simple interest.

- **a** How much interest is charged on the loan?
- **b** What is the total amount of the loan and the interest?
- **c** Calculate the size of each repayment.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

9 Copy and complete this table and find the monthly repayment for each loan.

Amount borrowed	Annual simple interest rate	Number of years	Interest	Total amount to be repaid	Monthly repayment
$5000	21%	5			
$14 000	15%	5			
$10 000	6%	4			
$55 000	8%	10			
$250 000	7%	30			

Problem-solving and reasoning

10, 11 | 10, 12, 13

10 Calculate the principal amount which earns $500 simple interest over 3 years at a rate of 8% p.a. Round to the nearest cent.

Hint for Q10: Substitute into $I = \frac{Prt}{100}$ and solve for P.



11 Charlotte borrows $9000 to buy a second-hand car. The loan must be repaid over 5 years at 12% p.a. simple interest. Calculate:

a the total amount to be repaid

b the monthly repayment amount if the repayments are spread equally over the 5 years



12 If $5000 grows to $11 000 in 12 years, find the simple interest rate.

13 An investor invests $\$P$ and wants to double this amount of money.

a How much interest must be earned to double this initial amount?

b What simple interest rate is required to double the initial amount in 8 years?

c If the simple interest rate is 5% p.a.:

i how many years will it take to double the investment?

ii how many years will it take to triple the investment amount?

iii how do the investment periods in parts **i** and **ii** compare?



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

2H

Compound interest — 14, 15

With simple interest, the principal and the interest earned each year remain the same for the period of the investment.

However, with compound interest, each time the interest is calculated it is added to the principal to give a new value. This means that the next time the interest is calculated, it is done so using a larger amount.

In the following questions you will be asked to do repeated applications of simple interest to find the final compounded amount.

14 $500 is invested for 4 years at 10% p.a. interest compounded annually.

a Complete the table to find the final value of the investment at the end of this time and the total.

Time (years)	Amount (A)	Interest (I)	New amount ($A + I$)
1	500	$500 \times 0.1 = 50$	$500 + 50 = 550$
2	550	$550 \times 0.1 =$	
3			
4			

b How much interest did the investment earn over the 4 years?

15 **a** Complete the following table to find the final value of an investment of $4500 compounded at 5% p.a. annually for 5 years.

Time (years)	Amount (A)	Interest (I)	New amount ($A + I$)
1	4500	225	4725
2	4725		
3			
4			
5			

b Type 4500×1.05^5 into your calculator. What do you notice about this answer?

c Can you explain how the answers to part **a** and part **b** relate?

d Use the formula $r = \frac{100I}{Pt}$ to find out what simple interest rate would be needed to create the same final amount over the 5 years. Answer correct to one decimal place.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Facebook cake-decorating business

More and more individuals are setting up a business using Facebook. For example, a successful cake-decorating business can be run from home while looking after the kids or while working normal business hours at other jobs.

As with any business, an understanding of financial mathematics is important to the success of the business. Skills such as calculating costs and profits, percentages and taxation are important for any manager.



1 Calculate the total cost of buying each of the following cake tin sets:
Round set: 6, 8, 9 and 12 inches by 3 inches deep at $64.
Round set: 6, 8, 10 and 12 inches by 4 inches deep at $99.
Square set: 6, 8 and 10 inches by 3 inches deep at $45.
Square set: 6, 8, 10 and 12 inches by 4 inches deep at $86.

2 Convert the following measurements from inches (US standard) to whole number of centimetres by using the following conversion rate: 1 inch = 2.54 cm.
- **a** 3 inches
- **b** 4 inches
- **c** 9 inches
- **d** 10 inches

3 Imagine that you spend 2 hours out of $9\frac{1}{2}$ work hours on the internet promoting your business. Write this as a percentage, rounded to the nearest whole percentage.

4 Fondant icing comes in different colours and in different-sized tubes. Managing your budget means looking for the best buy.
- **a** Which of the following represents the best buy for each colour of fondant listed below?

White fondant	Red fondant	Blue fondant
100 g at $3.25	2.5 kg at $36.95	100 g at $2.95
500 g at $5.50	1 kg at $19.90	500 g at $7.95
5 kg at $40	100 g at $3.25	750 g at $11.95
1 kg at $10.50	500 g at $7.95	1 kg at $19.90

- **b** What is the average cost per 100 grams for white fondant icing?
- **c** Under what circumstances would someone buy a size that was not the best buy?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5 A customer has the following quotes for a large 21st birthday cake from four different Facebook cake suppliers: \$195, \$290, \$225 and \$215.
For each of the following, state answers to the nearest whole number.
- **a** What is the mean or average cost for this type of cake?
- **b** What is the percentage change from the lowest quote to the highest quote?
- **c** If it costs each supplier \$50 in product to make the cake, calculate the percentage profit for each of the four quotes given above.
- **d** If it takes each supplier on average $5\frac{1}{2}$ hours to make and decorate the cake, what is each person charging per hour, excluding the \$50 product costs?

Using technology

6 Imagine that you have started a Facebook cake-making business. To analyse possible profits, set up an Excel spreadsheet as shown below and enter formulas in the shaded cells. Note that this is a simplified analysis and excludes power, gas and equipment costs.



Hint for Q6: Format all \$ cells as 'currency' with zero decimal places. Format profit % cells as 'percentage' with zero decimal places.

	A	B	C	D	E	F	G
1	CELEBRATION CAKES						
2	Cakes (30 serves, decorated)	Product costs in \$	Selling price in \$	Profit in \$	Profit %	Prep time in hours	Hourly rate
3	**Birthday** - caramel mud	\$35	\$105			3.25	
4	**Birthday** - choc fudge mud	\$38	\$112			3	
5	**Birthday** - choc Jaffa	\$35	\$108			3.5	
6	**Birthday** - child's theme	\$58	\$295			5.5	
7	**21st Birthday** - 4 tiers cupcakes	\$52	\$220			4	
8	**Valentine's Day** - red velvet choc	\$36	\$140			3	
9	**Wedding** - 2 tiers 70 serves	\$85	\$360			6	
10	**Wedding** - 3 tiers 120 serves	\$120	\$545			9	

- **a** How much profit would be made from selling 2 Valentines' Day cakes and 3 choc Jaffa birthday cakes?
- **b** List, in ascending order, the 3 cakes that bring the lowest percentage profit.
- **c** List, in descending order, the 3 cakes that pay the highest hourly rates.
- **d** Suggest a reason why the cakes in **c** cost the customer more in hourly rates.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Saving for a holiday

Sally is an accounts clerk earning a salary of $70 348. She plans to save 20% of her fortnightly after tax take home pay, for any extra expenses and to put towards a holiday. In fact, Sally is trying to save for an overseas holiday in 12 months and she estimates she will need $12 000 for her planned itinerary.

To reach her goal she decides to get a second job working at her local late night supermarket. They offer her three $2\frac{1}{2}$ hour shifts a week paying $25 an hour. She plans to work here for a total of 20 weeks, thinking that this will give her enough for her trip.

Present a report for the following tasks and ensure that you show clear mathematical workings and explanations where appropriate. Give answers to the nearest cent.

1 Preliminary task

a At her job at the supermarket, how much does Sally earn per shift?

b How much would she earn per week at the supermarket?

c Over the 20 weeks, calculate Sally's total gross income from the supermarket.

d As an accounts clerk, Sally earns a salary. What is the gross fortnightly income Sally earns from this job?

e What is her total annual taxable income from both jobs, assuming she only works for the 20 weeks at the supermarket?

2 Modelling task

Formulate

a The problem is to determine if Sally has saved enough to afford to go on her trip after tax. Write down all relevant information that will help you solve this problem.

Solve

b The firm that Sally works for as an accounts clerk take out income tax from her fortnightly pay on her behalf, they do this by using her salary and finding the tax owed by using the current personal income tax table found on the ATO website.

By using the personal income tax table below, calculate the annual tax that the firm deducts from Sally's pay.

Taxable income	Tax on this income
$0 – $18 200	Nil
$18 201 – $45 000	16c for each $1 over $18 200
$45 001 – $135 000	$4288 plus 30c for each $1 over $45 000
$135 001 – $190 000	$31 288 plus 37c for each $1 over $135 000
$190 001+	$51 638 plus 45c for each $1 over $190 000

* Does not include the Medicare levy (2% of taxable income).

c Calculate the fortnightly net pay (take home pay) that Sally receives from her firm.

d Sally aims to save 20% of her take home pay from her firm each fortnight. How much will Sally have saved at the end of 12 months?

e By considering her total taxable income from both jobs, calculate her total income tax liability for the year.

f Hence, calculate Sally's annual net income from her second job.

g Calculate her total tax liability, by including the Medicare levy of 2%.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Evaluate and verify

h When Sally takes in account all her finances and tax, will she have enough left from her savings to take her proposed trip?

Communicate

i Summarise your findings and make a recommendation about Sally's savings and proposed trip at the end of the year.

3 Extension questions

a What percentage pay rise is needed from her firm for Sally to no longer need her second job?

b Investigate what happens to Sally's tax liability from her salary from her firm if she has allowable tax deductions of:

i $2000 **ii** $4000



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Coding tax

Key technology: Spreadsheets



People will often use an online tax calculator to get an idea of how much income tax they will have to pay each year. Such calculators will use an in-built algorithm to help determine the tax rates depending on the given income. An added factor is the Medicare levy which is added on to most tax liabilities.

We will use the following income tax rates for this investigation.

Taxable income	Tax on this income
$0 – $18 200	Nil
$18 201 – $45 000	16 cents for each $1 over $18 200
$45 001 – $135 000	$4288 plus 30 cents for each $1 over $45 000
$135 001 – $190 000	$31 288 plus 37 cents for each $1 over $135 000
$190 001 and over	$51 638 plus 45 cents for each $1 over $190 000

The Medicare levy is 2% of taxable income.

1 Getting started

a Find the amount of tax paid, not including the Medicare levy, for the following taxable incomes.
i $15 000 **ii** $32 000 **iii** $95 000 **iv** $150 000 **v** $225 000

b Calculate the total tax payable on the following taxable incomes, including the Medicare levy, which is 2% of your taxable income.
i $60 000 **ii** $140 000

2 Applying an algorithm

Consider the flowchart on the following page which aims to calculate the amount of tax a person should pay.

a Complete the flowchart by filling in the empty components.
b Explain what the formula $T = T + 0.02I$ does in the flowchart.
c Test the algorithm in the flowchart on some of the taxable income amounts from part **1b** above.

3 Using technology

The following spreadsheet is a tax calculator which applies the algorithm outlined in the flowchart using nested IF statements.

	A	B
1	**Tax calculator**	
2		
3	**Taxable income**	250000
4		
5	**Income tax**	=IF(B3<=18200,0,IF(B3<=45000,0.16*(B3-18200),IF(B3<=135000,4288+0.3*(B3-45000),IF(B3<=190000,31288+0.37*(B3-135000),51638+0.45*(B3-190000)))))
6		
7	**Medicare levy**	=0.02*B3

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


Technology and computational thinking

a Try entering the formulas into a spreadsheet. Be careful to put all the brackets and commas in the correct positions.

b Test your spreadsheet calculator by entering different taxable incomes studied earlier in this investigation.

c Use your tax calculator to find the total tax payable for the following taxable incomes.

i $67 000 **ii** $149 000 **iii** $310 000



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

1 The answers to the clues are hidden in the wordfind. Can you find all 16 words?

S	A	G	R	O	S	S	W	H	K	L	O	M
A	P	E	R	C	E	N	T	A	G	E	C	O
L	A	R	C	O	M	M	I	S	S	I	O	N
A	S	E	R	V	I	N	T	E	R	E	S	T
R	T	Y	H	E	T	S	L	O	S	S	T	H
Y	E	F	O	R	T	N	I	G	H	T	I	L
W	S	M	O	T	A	X	A	T	I	O	N	Y
E	I	O	L	I	D	I	D	N	P	Y	I	E
K	M	N	Q	M	O	N	T	R	N	I	S	C
E	P	Y	U	E	D	I	S	C	O	U	N	T
M	L	R	E	P	A	Y	M	E	N	T	A	H
D	E	D	U	C	T	I	O	N	S	I	X	L

a A fixed annual income

b A percentage of the value of goods sold, which you earn as an income

c Working longer than normal working hours

d Money from your income given to the government

e Two weeks

f The total of all income

g Money taken from total pay

h Yearly

i 12 times a year

j Flat-rate interest

k Meaning 'out of 100'

l Money given to repay a loan

m Money earned on an investment

n An item offered for a sale price has had this happen

o The original price of an item

p You incur this when you sell an item for less than you paid for it

2 Find out the four classical elements of the world by answering the following simple interest problems. Match the letter beside each question to its corresponding answer in each grid.

E = \$600 at 6% p.a. for 1 year	**H** = \$796 at 5% p.a. for 4 months
R = \$12 500 at $6\frac{1}{4}$% p.a. for 2 years	**A** = \$7000 at 5% p.a. for 3 years
I = \$1000 at 1% p.a. for 100 years	**F** = \$576.50 at 19% p.a. for 18 months
W = \$36 000 at 2% p.a. for 5 years	**T** = \$550 at 10% p.a. for 6 months

\$36	\$1050	\$1562.50	\$27.50	\$13.27


\$1050	\$1000	\$1562.50

\$164.30	\$1000	\$1562.50	\$36

\$3600	\$1050	\$27.50	\$36	\$1562.50

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter summary



ISBN 978-1-009-59315-1
© Greenwood et al. 2025
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter checklist

A version of this checklist that you can print out and complete can be downloaded from your Interactive Textbook.

		✓
2A	**1 I can convert between percentages and fractions.** e.g. Write **a** $\frac{7}{40}$ as a percentage **b** 17.5% as a fraction	
2A	**2 I can convert between percentages and decimals.** e.g. Write **a** 0.3 as a percentage **b** 72% as a decimal	
2A	**3 I can write a quantity as a percentage.** e.g. Write 27 cm out of 1.8 m as a percentage.	
2B	**4 I can find a percentage of a quantity.** e.g. Find 35% of $75	
2B	**5 I can find the original amount from a percentage.** e.g. Determine the original amount if 12% of the amount is $72	
2C	**6 I can increase and decrease by a percentage.** e.g. **a** Increase $80 by 12% **b** Decrease $40 by 8%	
2C	**7 I can find a percentage change.** e.g. The price of a gym membership increased from $320 to $370. Find the percentage increase correct to one decimal place.	
2C	**8 I can find the original amount after an increase or decrease.** e.g. A decrease of 22% reduced the population of a town to 1014. What was the original population of the town?	
2D	**9 I can calculate the selling price from a mark-up or discount.** e.g. A store marks up all white goods by 20%. If the cost price of a fridge is $1100, what will be its selling price?	
2D	**10 I can determine percentage profit.** e.g. A stall holder makes candles for $8 and sells them for $13. Find the profit and express this profit as a percentage of the cost price.	
2D	**11 I can calculate the original price before discount.** e.g. A department store discounts all Christmas trees by 15%. If the sale price of a tree was $106.25, what was the original price?	
2E	**12 I can compare wages and salaries.** e.g. Tony has an annual salary of $88 000 and Jodie earns $72 per hour. Calculate **a** Tony's hourly rate of pay if he works a 38-hour week **b** Jodie's yearly income if she works on average 22 hours per week.	

Chapter checklist

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter checklist

		✔
2E	**13 I can calculate overtime.** e.g. Calculate Julian's weekly wage for a week where he works 12 hours at his normal hourly rate of $28, 4 hours at time and a half and 3 hours at double time.	
2E	**14 I can calculate commission.** e.g. A salesperson is paid a retainer of $2200 per month and receives an 8% commission on sales. If one month he makes sales worth $10 400, calculate his earnings for that month.	
2F	**15 I can calculate net income.** e.g. Danielle has an annual salary of $64 800. She has monthly tax deductions on her payslip of $1296. Calculate her net income each month.	
2F	**16 I can calculate net income using income tax rates.** e.g. Anna has an annual salary of $78 000 and a taxation rate of 26% with the first $18 200 tax free. Calculate Anna's net income for the year.	
2F	**17 I can use the tax table to calculate income tax.** e.g. Use the tax table from Example 18 to find the income tax for an income of $90 000.	
2G	**18 I can use the simple interest formula and find the final balance.** e.g. Calculate the simple interest earned if the principal is $4000, the rate is 3% p.a. and the time is 4 years. Hence, what is the amount at the end of the 4 years?	
2G	**19 I can work with simple interest using other time periods.** e.g. Calculate the simple interest on $6000 invested at $4\frac{1}{2}\%$ p.a. for 42 months.	
2H	**20 I can determine the investment period or interest rate for simple interest.** e.g. Joshua invests $3500 at 6% p.a. simple interest, for a period of time, to produce $315 interest. For how long did he invest the money?	
2H	**21 I can calculate repayments.** e.g. Syd takes out a loan to purchase a yacht. The loan is for $32 000 at 8% p.a. simple interest if the loan is repaid in equal monthly instalments over 4 years. Calculate the interest charged on the loan and hence the total amount owing and the size of the required monthly repayments.	

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Short-answer questions

2A **1** Copy and complete the table shown.

Decimal	Fraction	Percentage
0.6		
	$\frac{1}{3}$	
		$3\frac{1}{4}\%$
	$\frac{3}{4}$	
1.2		
		200%

2B **2** Find:
- **a** 25% of \$310
- **b** 110% of 1.5

2B **3** Determine the original amount if:
- **a** 20% of the amount is 30.
- **b** 72% of the amount is 18.

2C **4**
- **a** Increase 45 by 60%.
- **b** Decrease 1.8 by 35%.
- **c** Find the percentage change if \$150 is reduced by \$30.

2C **5** The mass of a cat increased by 12% to 14 kg over a 12 month period. What was its previous mass?

2D **6** Determine the discount given on a \$15 000 car if it is discounted by 12%.

2D **7** A couch at a cost price of \$3500 is to be marked up by 25%. Find the selling price.



2D **8** The cost price of an article is \$150. If it is sold for \$175:
- **a** determine the profit made
- **b** express the profit as a percentage of the cost price.

2E **9** Determine the hourly rate of pay for each of the following cases:
- **a** a person with an annual salary of \$76 076 working a 38-hour week
- **b** a person who earns \$429 working 18 hours at the hourly rate and 8 hours at time and a half.

2F **10** Jo's monthly income is \$5270 however 20% of this is paid straight to the government in taxes. What is Jo's net yearly income?

2G **11** Find the simple interest earned on \$1500 at 7% p.a. for 5 years.

2H **12** Bill invests \$6000 at 4% simple interest, for a period of time, to produce \$720. For how long did he invest the money?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter review

2F 13 Use the tax table below to find the income tax payable on an income of $140 000.

Taxable income	Tax on this income
$0 – $18 200	Nil
$18 201 – $45 000	16c for each $1 over $18 200
$45 001 – $135 000	$4288 plus 30c for each $1 over $45 000
$135 001 – $190 000	$31 288 plus 37c for each $1 over $135 000
$190 001+	$51 638 plus 45c for each $1 over $190 000

* Does not include the Medicare levy (2% of taxable income).

Multiple-choice questions

2A 1 2.8% as a decimal is:
A 2.8 B 0.28 C 0.028 D 0.0028 E 280

2A 2 What percentage of $2 is 50 cents?
A 4% B 40% C 25% D $2\frac{1}{2}\%$ E 400%

2A 3 $12\frac{1}{2}\%$ as a simple fraction is:
A $\frac{12}{100}$ B $\frac{1}{8}$ C $\frac{3}{25}$ D 0.125 E 12.5

2B 4 $33\frac{1}{3}\%$ of $660 is the same as:
A $660 ÷ 2 B $660 × 0.3 C $660 × 0.03 D $660 ÷ 3 E $660 ÷ $\frac{1}{3}$

2B 5 15% of $1600 is equal to:
A 24 B 150 C $240 D $24 E 240

2B 6 If 110% of a number is 528, then the number is:
A 475.2 B 52.8 C 480 D 580.8 E 475

2C 7 9670 increased by 12% becomes:
A 9682 B 9658 C 10830.4 D 1160.4 E 8509.6

2E 8 Jane is paid a wage of $27.80 per hour. She works 12 hours at this rate during a week, plus 4 hours on a public holiday where she gets paid at time and half. Her earnings in the week are:
A $500.40 B $444.80 C $556 D $667.20 E $278

2E 9 Simon earns a weekly retainer of $370 and 12% commission of any sales he makes. If he makes $2700 worth of sales in a particular week, he will earn:
A $595 B $652 C $694 D $738.40 E $649.60

2G/H 10 $1200 is invested with a simple interest rate of 10% for two years. The total balance at the end of the two years is:
A $252 B $1452 C $1450 D $240 E $1440

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Extended-response questions



1 Pauline buys a debutante dress at cost price from her friend Tila. Pauline paid $420 for the dress, which is normally marked up by 55%.
   a How much did she save?
   b What is the normal selling price of the dress?
   c If Tila gets a commission of 15%:
      i how much commission did she get?
      ii how much commission did Tila lose by selling the dress at cost price rather than the normal selling price?

2 Adam starts a new job and works a 38-hour week for a wage of $975.84.
   a Calculate his hourly rate of pay.
   b If overtime is calculated at time and a half, what is Adam's overtime rate?
   c How much does Adam earn for 4 hours of overtime work?
   d How many hours of overtime did Adam work in a week if his wage for that week was $1226.22?
   e If Adam usually works the amount of overtime in part **d** in the 52 weeks of the year he works, and he pays 27% of his pay in tax, what is his net annual income?
   f If Adam invests 10% of his net income in an account earning 8% p.a. simple interest for 18 months, how much extra income will he have earned?



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.



3 Expressions and equations

Essential mathematics: why skills with algebraic, expressions and formulas are important

Algebraic expressions and equations are powerful tools used in various fields to solve problems and make predictions. Understanding how to manipulate and solve these expressions can help in many real-life situations.

Computer programmers use algebra all the time when writing code. For example, sorting data with methods like quicksort or mergesort relies on algebra to calculate how fast it works. Programmers also use algebra to make sure searches are speedy and accurate. Knowing algebra helps them write better and more efficient code.

Engineers use equations to design and analyse structures, making sure they can withstand different forces. To check if materials can handle applied loads, they might use the stress formula, $\sigma = \frac{F}{A}$, where σ represents stress, F is the load applied, and A is the cross-sectional area over which the force is measured. By calculating stress, engineers can determine if the material will deform or fail under specific loads, ensuring the safety and stability of their designs.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.



In this chapter

3A Algebraic expressions (Consolidating)
3B Adding and subtracting algebraic expressions (Consolidating)
3C Multiplying and dividing algebraic expressions (Consolidating)
3D Expanding algebraic expressions
3E Solving linear equations (Consolidating)
3F Solving linear equations involving fractions
3G Solving equations with brackets
3H Solving equations with pronumerals on both sides
3I Solving word problems
3J Solving linear inequalities
3K Using formulas

Victorian Curriculum 2.0

This chapter covers the following content descriptors in the Victorian Curriculum 2.0:

NUMBER

VC2M9N01

ALGEBRA

VC2M9A02, VC2M9A03, VC2M9A06

Please refer to the curriculum support documentation in the teacher resources for a full and comprehensive mapping of this chapter to the related curriculum content descriptors.

© VCAA

Online resources

A host of additional online resources are included as part of your Interactive Textbook, including HOTmaths content, video demonstrations of all worked examples, auto-marked quizzes and much more.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025
Photocopying is restricted under law and this material must not be transferred to another party.
Cambridge University Press

Warm-up quiz

1 Write in simplified form.

a $3 \times x$
b $4 \times a \times b$
c $2 \times 5 \times x$
d $3 \times b \times 7$

2 If $x = 3$, find the value of:

a $x + 5$
b $10 - x$
c $4x$
d $\frac{18}{x}$
e $2x + 3$
f $2(x + 4)$

3 Write algebraic expressions for:

a 3 more than x
b the product of a and b
c 2 lots of y less 3
d the sum of x and 2, all divided by 3

4 Calculate:

a -6×3
b -7×4
c $-8 \times (-5)$
d $-1 \times (-3)$
e $-2 + 5$
f $-8 + 3$
g $7 - 11$
h $-2 - 6$

5 Simplify by collecting like terms.

a $2x + 5x - 4x$
b $7y + 5 - y$
c $4x - (-4x)$
d $3x + 12y - 3x + 5y$
e $3a + 4a^2 + 7a + 5a^2$
f $3xy - 4y + 2yx - 3y$

6 Simplify the following.

a $3 \times 2a$
b $7x \times (-3y)$
c $\frac{8b}{2}$
d $\frac{9mn}{6n}$

7 Expand the following.

a $2(x + 3)$
b $3(a - 5)$
c $4(2x + 1)$

8 Evaluate the following if $a = 3$ and $b = -2$.

a $2a - 5$
b $ab + 4$
c $\frac{9}{a} - b$
d $2a(b + 1)$

9 To which of the following equations is $x = 4$ a solution?

a $2x + 3 = 9$
b $\frac{x}{2} + 3 = 5$
c $\frac{2x + 1}{3} = 3$
d $5 - 2x = -1$

10 Find the value of a that makes the following true.

a $a + 4 = 13$
b $a - 3 = 7$
c $2a + 1 = 9$
d $\frac{a - 1}{4} = 5$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3A Algebraic expressions CONSOLIDATING

Learning intentions

- To review the terms associated with algebraic expressions
- To be able to identify terms, coefficients and constant terms in expressions
- To know the notation for multiplication and division involving pronumerals
- To be able to convert words and word problems into algebraic expressions
- To be able to substitute into expressions and evaluate

Key vocabulary: pronumeral, variable, expression, term, coefficient, constant term, substitution, evaluate

Algebra is central to the study of mathematics and is commonly used to solve problems in many everyday situations and in more complex problems. Algebra involves working with unknown values in maths. Pronumerals (or variables) are used to represent these unknown values.

Lesson starter: What does the letter mean?

Consider the following situations and how they could be written as algebraic expressions.

1 Fidel has 7 marbles and Jack has an unknown number of marbles, which we will call m. They then each get 4 more marbles.
   - **a** How many marbles does Fidel have now?
   - **b** What mathematical operation (+, −, ×, ÷) did you use to answer part **a**?
   - **c** Using the same operation, how could you describe how many marbles Jack has now?

2 Fidel can buy a chocolate bar for $2. Jack can buy a different chocolate bar but he will pay an unknown amount for it. We'll call the amount Jack pays p (dollars).
   - **a** If Fidel buys 3 chocolate bars, how much will he pay?
   - **b** If Fidel buys 10 chocolate bars, how much will he pay?
   - **c** What mathematical operation did you use to answer parts **a** and **b**?
   - **d** Using the same operation, how could you write how much Jack will pay for 5 chocolate bars?


Key ideas

- In algebra, letters are used to represent numbers. These letters are called **pronumerals** (sometimes written pro-numerals).
  - Pronumerals can also be called **variables** if they can take a range of possible values.
- In algebra, the multiplication and division signs are generally not shown.
  For example: $3 \times x$ is written as $3x$ and $x \div 4$ is written as $\frac{x}{4}$.
- An **expression** is a combination of numbers and pronumerals connected by the four operations +, −, × and ÷. Brackets can also be used.
  For example: $5x + 4y - 1$ and $3(x + 2) - \frac{y}{5}$.
- A **term** is a combination of numbers and pronumerals connected with only multiplication and division. Terms are separated with the operations + and −. For example: $5x + \frac{7}{y}$ is a two-term expression with $5x$ and $\frac{7}{y}$ as the two terms.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3A

- **Coefficients** are the numbers being multiplied by pronumerals.



- **Constant terms** consist of a number only.
  For example: -2 in $x^2 + 4x - 2$ (The sign must be included.)
- **Substitution** is the process of replacing a pronumeral with its numeric value.
  Expressions can be evaluated by substituting a number for a pronumeral.
  For example: if $a = 2$ then $a + 6 = 2 + 6 = 8$
- Order of operations should be followed when evaluating expressions:
  1 Brackets
  2 Powers
  3 Multiplication and division
  4 Addition and subtraction

Exercise 3A

Understanding

1–4 | 4

1 Use the words from the list below to complete the following.
expression, *coefficient*, *constant term*, *terms*, *pronumeral*

a $3x - 2y + 7$ is an example of an algebraic _____.
b In $2a + b - 3$, -3 is the _____.
c In $4m + 3$, m is called a _____.
d In $3 - 2y + 4z$, 4 is the _____ of z.
e In the expression $3s + 7t$, $3s$ and $7t$ are the _____ .

2 Express in simplified form without the $\times$ or $\div$ symbols.

a $7 \times y$
b $-2 \times x$
c $a \times b$
d $y \div 2$
e $x \div y$
f $2 \div a$

3 Substitute $x = 2$ into the following and evaluate.

a $x + 3$
b $10 - x$
c $7x$
d $\frac{8}{x}$



Hint for Q3: Substitute means to replace the pronumeral with the given numeral.

4 State, yes or no, if 2 is the coefficient of x in the following.

a $2x + 7$
b $y - 2x$
c $3xy - 4y + 2x$
d $\frac{x}{2} - 3$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Fluency

5, 6, 7–8(½) | 5, 6, 7–8(½)

Example 1 Identifying parts of algebraic expressions

Consider the expression $5a + 2ab + 7 - 4b$.

a How many terms are in the expression?

b State the coefficient of:

i a ii b

c What is the constant term?

Solution

a There are 4 terms.

b i 5 is the coefficient of a.

ii -4 is the coefficient of b.

c 7 is the constant term.

Explanation

The terms in the expression are separated by + or −.

$5a$ is $5 \times a$; 5 is the number being multiplied by a.

$-4b$ is $-4 \times b$; the negative sign is included.

7 is the number with no pronumeral part.

Now you try

Consider the expression $3ab + 5 - 2a + 6b$.

a How many terms are in the expression?

b State the coefficient of:

i a ii b

c What is the constant term?

5 Complete each of the parts **i–iii** for the algebraic expressions below.

i How many terms are in the expression?

ii What is the coefficient of y?

iii What is the constant term?

a $5x + 2y + 3$ b $2x - 3y$

c $3xy + 7y - 4$ d $2x^2 - 1 + 4x + \frac{y}{2}$



Hint for Q5: Remember: the coefficient of a pronumeral includes the sign before it; in $3 - 2x$, -2 is the coefficient of x.

Example 2 Writing algebraic expressions for word problems

Write an algebraic expression for the following.

a The number of tickets needed for 3 boys and r girls

b The cost in dollars of P pies at \$3 each

c The amount for each person if a \$300 prize is shared equally among m friends

Solution

a $3 + r$

b $\$3P$

c $\frac{\$300}{m}$

Explanation

3 tickets plus the number of girls, r.

2 pies would cost $\$3 \times 2 = \$6$. The cost is 3 × the number of pies, so P pies costs $\$3 \times P = \$3P$

\$300 shared between 3 people is \$300 divided into 3 parts, so \$300 divided into m parts is $\$300 \div m = \frac{\$300}{m}$.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3A

Now you try

Write an algebraic expression for the following.

a The number of tickets required for 5 children and g adults
b The cost of 4 tickets at $\$x$ each
c The amount for each child if 600 mL of juice is shared equally among y children

6 Write an algebraic expression for the following.

a The number of tickets required for:
  i 4 boys and r girls
  ii t boys and 2 girls
  iii x boys, y girls and z adults

b The cost in dollars of:
  i P pies at \$6 each
  ii 10 pies at $\$n$ each
  ii D drinks at \$2 each
  iv P pies at \$5 and D drinks at \$2

c The number of grams of lollies for one child if 500 g of lollies is shared equally among C children.



Hint for Q6:
For part bi:
1 pie costs $\$6 \times 1 = \$6$
2 pies cost $\$6 \times 2 = \$12$
5 pies cost $\$6 \times 5 = \$30$
P pies cost …

Example 3 Converting words to expressions

Write an algebraic expression for the following.

a Five less than x
b Three more than twice x
c The sum of a and b is divided by 4
d The square of the sum of x and y

Solution	Explanation
a $x - 5$	5 below x is 5 subtracted from x.
b $2x + 3$	Twice x is $2 \times x$, then add 3.
c $\frac{a+b}{4}$	The sum of a and b is done first $(a + b)$ and the result divided by 4.
d $(x+y)^2$	The sum of x and y is done first and then the result is squared (recall that the square of a is $a^2 = a \times a$).

Now you try

Write an algebraic expression for the following.

a Seven more than y
b Six less than twice x
c The sum of m and n is divided by 3
d The square root of the sum of a and b

7 Write an algebraic expression for each of the following.

a The sum of 2 and x
b The sum of ab and y
c 5 less than x
d 7 subtracted from $2y$
e The product of x and 3
f Three times the value of p
g Four more than twice x
h The sum of x and y is divided by 5
i 10 less than the product of 4 and x
j 3 lots of x subtracted from 1
k The sum of 3 and y is divided by 2
l Half of 1 more than x
m The square of the sum of m and n
n The sum of the squares of m and n



Hint for Q7: Sum is +
Product is ×
Twice is × 2
Square of x is x^2

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Example 4 Substituting values into expressions

Evaluate these expressions if $a = 5$, $b = 6$ and $c = -3$.

a $3a + b$ **b** $2a - (b + c)$ **c** $a^2 - bc$

Solution

a $3a + b = 3 \times 5 + 6$
$= 15 + 6$
$= 21$

b $2a - (b + c) = 2 \times 5 - (6 + (-3))$
$= 10 - 3$
$= 7$

c $a^2 - bc = (5)^2 - 6 \times (-3)$
$= 25 - (-18)$
$= 25 + 18$
$= 43$

Explanation

Substitute the value 5 for a and the value 6 for b. Recall $3a = 3 \times a$.

Substitute the values for a, b and c. Then apply order of operations, with brackets evaluated first, before moving onto multiplication and division, then addition and subtraction. $6 + (-3) = 6 - 3 = 3$

Evaluate powers before the other operations: $5^2 = 5 \times 5$.
A positive (6) times a negative (-3) gives a negative (-18).
Subtracting a negative means to add.

Now you try

Evaluate these expressions if $a = 4$, $b = -2$ and $c = 5$.

a $a + 4c$ **b** $2b - (a + c)$ **c** $c^2 - ab$

8 Evaluate these expressions if $a = 4$, $b = -3$ and $c = 8$.

a $3a + c$ **b** $ac - 7$ **c** $2c - (a + b)$
d $a^2 - 2c$ **e** $\frac{a + c}{2}$ **f** $b + 2(c - a)$
g $2c - ab$ **h** abc **i** $b - c + a^2$



Hint for Q8: When adding or subtracting a negative number:
$8 + (-3) = 8 - 3 = 5$
$8 - (-3) = 8 + 3 = 11$

Problem-solving and reasoning 9–11 10–13

9 A rectangular garden bed is 12 m long and 5 m wide.

a Find the area of the garden bed.
b The length is increased by x m and the width is decreased by y m. Find the new length and width of the garden.
c One length and one width of the rectangular garden will be lined with paving. Write an expression for the total distance around the garden that will be paved.



Hint for Q9: Draw a diagram to help. Area of a rectangle $= l \times w$

10 Jelena earns $(10 + 8t)$ dollars for each shift that she works, where t is the number of hours worked in the shift.


a How much does Jelena earn if $t = 2$?
b How much does Jelena earn from a 5-hour shift?
c Will she earn \$100 from a 10-hour shift?
d How many whole hours must she work to earn more than \$100?



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3A

11 The expression for the area of a trapezium is $\frac{1}{2}(a+b)h$, where a, b and h represent the lengths shown.



- **a** Find the area of the trapezium with $a = 5$, $b = 7$ and $h = 3$.
- **b** A trapezium has $h = 4$ and area 12. If a and b are whole numbers, what possible values can the pronumeral a have?

12 Christina earns $\$A$ from selling 20 glasses of lemonade. Write an expression for:


- **a** the cost of one glass of lemonade
- **b** the cost of three glasses of lemonade
- **c** the cost of n glasses of lemonade




13 For each of these shapes, write an expression for:

i the perimeter **ii** the area


a



b



c





Hint for Q13: Perimeter is the distance around the outside of the shape. Area of a rectangle is length × width. Area of a triangle is $\frac{1}{2}$ × base × height.

Should there be brackets? — 14

14 Brackets are used to ensure that the mathematics inside the brackets occurs first.

- **a** By substituting values for x, decide whether the following are the same.
  - **i** $2(x+1)$ and $2x+1$
  - **ii** $3+\frac{x}{2}$ and $\frac{3+x}{2}$
- **b** Write down two different possible expressions that could describe each of the following.
  - **i** 3 lots of x plus 1
  - **ii** The sum of 5 and x divided by 3
  - **iii** Half of x plus y
- **c** Write down a clear statement that describes the following expressions.
  - **i** $4(x+2)$
  - **ii** $3+\frac{x}{2}$
  - **iii** $\frac{1}{3}(m+n)$
  - **iv** $5x+7$
  - **v** $\frac{x+y}{2}$
  - **vi** $\frac{1}{2}a+b$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3B Adding and subtracting algebraic expressions

CONSOLIDATING

Learning intentions

- To understand that only like terms can be combined under addition or subtraction
- To be able to identify like terms
- To be able to collect like terms under addition or subtraction

Key vocabulary: like terms, pronumeral, coefficient

Just as $2+2+2+2=4\times 2$, so $x+x+x+x=4\times x$ or $4x$. We say that the expression $x+x+x+x$ is simplified to $4x$; i.e. 4 lots of x. Similarly, $3x+2x=x+x+x+x+x=5x$ and $4x-3x=x+x+x+x-(x+x+x)=x$.

All these expressions have like terms and can be simplified to an expression with a smaller number of terms by adding or subtracting their coefficients.



$x+x+x+x=4x$

Lesson starter: Two groups

Place the terms in the following four sets into groups of similar terms on either side of the dividing line. The first set has been started for you.

Set 1	Set 2	Set 3	Set 4
~~$3x$~~ $7y$ $2x$ $8x$	$4ab$ $2a$ $5ab$	2 $5y$ 7 1	$3ab$ $4a^2b$ $2ab$ $10ab$
y $5x$ ~~$2y$~~	$7a$ ab $6ab$	$3y$ $2y$ 8 25	$5a^2b$ $12ab$ $9a^2b$
$3x \mid 2y$			

- Describe how you classified each set into two groups.
- What do the terms in each group have in common?
- Simplify the sum of all the terms in each of the eight groups.

Key ideas

- $4x$ means $4\times x$ or $x+x+x+x$.
  - x means $1\times x$ but we leave off the 1 and express it simply as x.
- **Like terms** have the same pronumerals and powers.
  For example: $5\boldsymbol{x}$ and $7\boldsymbol{x}$ are like terms
  $3\boldsymbol{ab}$ and $-8\boldsymbol{ab}$ are like terms
  - Since $a\times b=b\times a$ then ab and ba are also like terms
- The pronumeral part of a term is often written in alphabetical order; e.g. xy rather than yx.
- Like terms can be collected to form a single term by adding or subtracting the coefficients.
  For example: $4x+7x=11x$
  $10ab-2ab=8ab$
- Unlike terms do not have the same pronumeral factors.
  For example: $5x$, x^2, xy, $2xy^2$ and 3 are all unlike terms.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Exercise 3B

Understanding

1–3 | 3

1 Fill in the missing word or number.

a $4a$ and $7a$ are an example of _____ terms.

b To add like terms, we add the _____.

c Like terms have the same _____ and powers.

d The coefficient of y in $2x + y$ is _____.



Hint for Q2: Like terms must have exactly the same pronumerals and powers.

2 Decide whether the following pairs of terms are like terms (Yes or No).

a $4ab$ and $3ab$ **b** $2x$ and $7xy$ **c** 5 and $4m$

d $7yz$ and $-yz$ **e** $2mn$ and $9nm$ **f** $3x^2y$ and $7xy^2$

3 Which of the following represent the same expression as $w - x + y - z$?

a $w - z + y - x$

b $w - y + x - z$

c $y + w - z - x$

d $-x + w - z + y$

e $y - w - x + z$

f $y - z - x + w$



Hint for Q3: Make sure the correct pronumerals are being added or subtracted.

Fluency

4, 5–6(½) | 4–6(½)

Example 5 Identifying like terms

Choose the pair(s) of like terms from the following sets.

a $3x,\ 3,\ 4xy,\ 5x,\ y$ **b** $4b,\ 4bc,\ b,\ 2abc,\ -7bc,\ 2bc^2$

Solution

a $3x$ and $5x$ are like terms.
The rest are not.

b $4b$ and b are like terms.
$4bc$ and $-7bc$ are like terms.

Explanation

$3x$ and $5x$ have the same pronumeral; i.e. x.

$4b$ and b have the same pronumeral.
$4bc$ and $-7bc$ have the same pronumerals.
$2bc^2$ does not, due to the power of 2.
All other terms do not have the same pronumerals and powers.

Now you try

Choose the pair(s) of like terms from the following sets.

a $5m, 3mn, 7, n, 2m$ **b** $7y, 4xz, 5xy^2, -2xyz, zx, -y$

4 Choose the pair(s) of like terms from the following sets.

a $4y,\ 5,\ 3xy,\ 2y,\ 7xy$ **b** $3x,\ 3y,\ 3,\ xy,\ 7x$

c $7ab,\ 2a,\ -3ab,\ b^2,\ 5a$ **d** $2a^2,\ 4a,\ 3ab,\ a,\ -3a^2$

e $2xy,\ 3x^2y,\ -5x,\ 7yx^2$ **f** $5ab^2,\ 3ab,\ 2b^2,\ 4ab^2,\ 7ba$



Hint for Q4: st and ts are like terms since $s \times t = t \times s$.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Example 6 Collecting like terms

Simplify the following by collecting like terms.

a $4a + 7a$ **b** $3x + 4 - 2x$

Solution

a $4a + 7a = 11a$

Explanation

Since $4a$ and $7a$ are like terms they can be simplified to one term by adding their coefficients: $4 + 7 = 11$.

b $3x + 4 - 2x = 3x - 2x + 4$
$= x + 4$

Collect like terms ($3x$ and $-2x$). The sign belongs to the term that follows. Combine their coefficients: $3 - 2 = 1$. Recall that $1x$ is written as x.

Now you try

Simplify the following by collecting like terms.

a $5t + 3t$ **b** $9y + 3 - 5y$

5 Simplify the following by collecting like terms.

a $3a + 7a$ **b** $4n + 3n$ **c** $12y - 4y$
d $5x + 2x + 4x$ **e** $6ab - 2ab$ **f** $7mn + 2mn - mn$
g $4y + 3y + 8$ **h** $7x + 5 - 4x$ **i** $5m + 2 - m$
j $5ab + 3a + 7ab$ **k** $6xy + xy + 4y$ **l** $5bc - 4 - 2bc$



Hint for Q5: In order to add or subtract coefficients, the terms must be like terms.

Example 7 Combining like terms

Simplify the following by collecting like terms.

a $3x + 2y + 4x + 7y$ **b** $3xy + 4x + xy - 6x$ **c** $8ab^2 - 9ab - 5ab^2 + 3ba$

Solution

a $3x + 2y + 4x + 7y = 3x + 4x + 2y + 7y$
$= 7x + 9y$

Explanation

Collect like terms ($3x$ and $4x$ and $2y$ and $7y$) and combine their coefficients: $3 + 4 = 7$ and $2 + 7 = 9$.

b $3xy + 4x + xy - 6x = 3xy + xy + 4x - 6x$
$= 4xy - 2x$

Collect the like terms ($3xy$ and xy and $4x$ and $-6x$). Combine their coefficients (recall $xy = 1xy$): $3 + 1 = 4$ and $4 - 6 = -2$

c $8ab^2 - 9ab - 5ab^2 + 3ba$
$= 8ab^2 - 5ab^2 - 9ab + 3ab$
$= 3ab^2 - 6ab$

Collect like terms in ab^2 and ab. Remember that $ba = ab$ and the + or − sign belongs to the term that follows; i.e. $-5ab^2$.
Combine coefficients: $8 - 5 = 3$ and $-9 + 3 = -6$.

Now you try

Simplify the following by collecting like terms.

a $4a + 3b + 2a + 10b$ **b** $5st + 3s - 2st + s$ **c** $2x^2y - 5xy + 3x^2y - yx$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3B

6 Simplify the following by collecting like terms.

a $2a + 4b + 3a + 5b$
b $4x + 3y + 2x + 2y$
c $xy + 8x + 4xy - 4x$
d $6t + 5 - 2t + 1$
e $5x + 1 + 6x + 3$
f $3mn + 4 + 4nm - 5$
g $4ab + 2a + ab - 3a$
h $3st - 8ts + 2st + 3ts$
i $8ac - 6c + 2ac - 3c$
j $5xy^2 - 2xy^2$
k $8m^2n - 6nm^2 + m^2n$
l $2x^2y - 4xy + 5yx^2$
m $x^2 - 7x + 3x^2 + 4x$
n $a^2b - 4ab + 3a^2b + ba$
o $10pq^2 - 2qp - 3pq^2 - 6pq$

Problem-solving and reasoning

7, 8 | 8(½), 9, 10


7 A farmer has x pigs and y chickens.




a Write an expression for the total number of heads.
b Write an expression for the total number of legs.

8 Write simplified expressions for the perimeter of the following shapes.


a




b




c




d



e




f





Hint for Q8: Sides with matching dash marks are equal in length.



9 A rectangular pool's length is three times its width, x metres. Write an expression for its perimeter.


10 Fill in the missing term to make each of the following correct.

a $11t + 3 - \square = 7t + 3$
b $\square + 3y - 2x = 7x + 3y$
c $3x + 4y - \square + 5y = x + 9y$
d $4a + 7b - \square - 2b = 5b - 3a$
e $3mn - \square + nm + 4n = 4mn - n$
f $2pq + 2p - \square - 5pq = 2p - 7pq$
g $3x^2y - 11xy + 3yx^2 - \square = 6x^2y - 13xy$
h $4b^2 - 3b + 2b^2 + \square = 6b^2$

Around the edge

— | 11

11 Consider the shape shown.



a If the unknown sides (?) have length $2b$, what is the perimeter in simplified form?
b If $a = 3$ and $b = 2$ and the perimeter is 23 units, what is the length of the unknown sides?
c If the perimeter is $(5a + 3b)$, what is the length of each unknown side?
d If the perimeter is $(7a + b)$, what is the length of each unknown side?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3C Multiplying and dividing algebraic expressions

CONSOLIDATING

Learning intentions

- To understand that order does not matter when multiplying
- To be able to multiply algebraic terms by multiplying coefficients and pronumerals
- To be able to divide algebraic terms by cancelling the highest common factor

Key vocabulary: pronumeral, coefficient, factor, highest common factor

We have seen that $3x + 3x$ can be simplified to $6x$ and that $3x + 3x$ can also be written as $2 \times 3x$. Thus, $2 \times 3x = 6x$. Also, $4 \times 5a = 4 \times 5 \times a = 20a$. Any two or more terms can be simplified using multiplication; we do not need the terms to be like terms.

A single term such as $2 \times 5 \times x \div 10$ can also be simplified using multiplication and division, so $2 \times 5 \times x \div 10 = \frac{10x}{10} = x$, since $\frac{10}{10}$ simplifies to 1 by cancelling the common factor.

Lesson starter: Are they equivalent?

These expressions can be separated into two groups. Group them so that the expressions in each group are equivalent.

$4x$ $\quad 2x - y$ $\quad 6x - x - x$ $\quad 10x - y - 8x$

$\frac{24x}{6}$ $\quad y + 2x - y + 2x$ $\quad 2 \times x - 1 \times y$ $\quad -y + 2x$

$\frac{1}{2} \times 8x$ $\quad 2 \times 2x$ $\quad \frac{8x^2}{2x}$ $\quad -1 \times y + \frac{2x^2}{x}$

Key ideas

- The symbols for multiplication ($\times$) and division ($\div$) are usually not shown in simplified algebraic terms.
  For example: $5 \times a \times b = 5ab$ and $-7 \times x \div y^2 = -\frac{7x}{y^2}$
- Order does not matter in multiplication; i.e. $2 \times 3 = 3 \times 2$, $a \times 2 \times b = 2 \times a \times b$.
- Algebraic terms are multiplied by multiplying coefficients with the pronumeral parts.
  For example: $4a \times 3b = 4 \times 3 \times a \times b = 12ab$.
- The factors of a number are the set of numbers that divide evenly into it. For example, the **factors** of 6 are 1, 2, 3 and 6.
- When dividing algebraic expressions, common factors can be cancelled. Cancel the highest common factor of the numerals and then the pronumerals.
  For example: $\frac{{}^{1}\cancel{7}x}{{}^{2}\cancel{14}} = \frac{x}{2}$, $\frac{{}^{2}\cancel{8}xy^{1}}{{}^{3}\cancel{12}y^{1}} = \frac{2x}{3}$, $\frac{a^2b}{a} = \frac{{}^{1}\cancel{a} \times a \times b}{{}^{1}\cancel{a}} = ab$

Essential Mathematics Core for the Victorian Curriculum Year 9 ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Exercise 3C

Understanding

1, 2 | 2

1 Complete the simplified form of the following.

a $7 \times x \times y = \square$ **b** $3 \times 2 \times m = \square m$ **c** $4 \times x \times 5 = \square$ **d** $a \times a = a^{\square}$

2 Answer true (T) or false (F) to the following statements.

a $2 \times x \times 3 = 2 \times 3 \times x$

b $7a \times 3b = 7 \times 3 \times a \times b$

c $4x \times 5xy = 4 \times 5 \times x \times y$

d $\frac{8x}{16} = 2x$

e $5a^2 \div (10a) = \frac{10a}{5a^2}$

f $\frac{3xy^2}{y} = 3xy$



Hint for Q2: Simplify by cancelling the highest common factor of the numerator and denominator:
$\frac{6}{10} = \frac{2_1 \times 3}{2_1 \times 5} = \frac{3}{5}$

Fluency

3–5(½) | 3–5(½)

Example 8 Multiplying algebraic terms

Simplify the following.

a $3 \times 2b$ **b** $-2a \times 3b$

Solution

a $3 \times 2b = 3 \times 2 \times b$
$= 6b$

b $-2a \times 3b = -2 \times 3 \times a \times b$
$= -6ab$

Explanation

Multiply the coefficients.

Recall that multiplication can occur in any order; i.e. $2 \times 3 \times 4 = 2 \times 4 \times 3$ etc.
Multiply the coefficients and simplify.

Now you try

Simplify the following.

a $4 \times 5x$ **b** $-3m \times 6p$

3 Simplify the following.

a $5 \times 2m$ **b** $2 \times 6b$
c $3 \times 5p$ **d** $3x \times 2$
e $3p \times 6r$ **f** $4m \times 4n$
g $-2x \times 7y$ **h** $5m \times (-3n)$
i $-4c \times 3d$ **j** $2a \times 3b \times 5$
k $-4r \times 3 \times 2s$ **l** $5j \times (-4) \times 2k$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Example 9 Dividing algebraic terms

Simplify the following.

a $\frac{4x}{8}$

b $10ab \div (15b)$

Solution

a $\frac{{}^{1}\cancel{4}x}{{}_{2}\cancel{8}} = \frac{x}{2}$

b $10ab \div (15b) = \frac{10ab}{15b}$

$= \frac{{}^{2}\cancel{10} \times a \times \cancel{b}_{1}}{{}_{3}\cancel{15} \times \cancel{b}_{1}}$

$= \frac{2a}{3}$

Explanation

Deal with numerals and pronumerals separately, cancelling any common factors.
The highest common factor of 4 and 8 is 4.

Write division as a fraction first. Write numerator and denominator as a product and cancel the common factors 5 and b.

Now you try

Simplify the following.

a $\frac{9x}{27}$

b $18pq \div (4p)$

4 Simplify the following by cancelling.

a $\frac{8b}{2}$ b $\frac{2a}{6}$ c $\frac{4ab}{6}$ d $\frac{3mn}{6n}$

e $\frac{5xy}{20y}$ f $\frac{10st}{6t}$ g $\frac{3xy}{xy}$ h $\frac{27pq}{6p}$

i $2x \div 4$ j $12ab \div (2a)$ k $7mn \div (3n)$ l $8y \div (20xy)$



Hint for Q4: Make sure they are in fraction form before cancelling; e.g. $2x \div 4 = \frac{2x}{4}$.

Example 10 Multiplying and dividing with squared pronumerals

Simplify the following.

a $6x \times 5xy$

b $\frac{12a^2b}{3ab}$

Solution

a $6x \times 5xy = 6 \times 5 \times x \times x \times y$

$= 30x^2y$

b $\frac{12a^2b}{3ab} = \frac{{}^{4}\cancel{12} \times {}^{1}\cancel{a} \times a \times \cancel{b}^{1}}{{}^{1}\cancel{3} \times \cancel{a}^{1} \times \cancel{b}^{1}}$

$= 4a$

Explanation

Multiply the coefficients and the pronumerals.
Recall that $x \times x$ is written as x^2 (x squared).

Write numerator and denominator as a product, with $a^2 = a \times a$. Cancel common factor (3) for numerals, then cancel common factors for pronumerals.

Now you try

Simplify the following.

a $7ab \times 6a$

b $\frac{20xy^2}{4xy}$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3C

5 Simplify the following.

a $4n \times 6n$ **b** $-3q \times q$ **c** $5s \times 2s$

d $7a \times 3ab$ **e** $5mn \times (-3n)$ **f** $-3gh \times (-6h)$

g $\frac{24ab^2}{8ab}$ **h** $\frac{25x^2y}{5xy}$ **i** $\frac{9m^2n}{18mn}$

j $\frac{2xy}{8xy^2}$ **k** $\frac{6a^2b}{10a}$ **l** $\frac{45p^2q^2}{15pq}$



Hint for Q5:
Deal with numerals and pronumerals separately.
$\frac{24^3}{8_1}$ and $\frac{ab^2}{ab} = \frac{a_1 \times b \times b_1}{a^1 \times b_1} = b$

Problem-solving and reasoning

6(½), 7 | 6(½), 7–9

6 Simplify the following by first writing in fraction form.

a $4 \times x \div y$ **b** $3 \times m \div 9$

c $4 \times a \div (2b)$ **d** $5x \times 4 \div (2y)$

e $6 \times 4mn \div (3m)$ **f** $8a \times 5b \div (8a)$

g $10m \times 4n \div (8mn)$ **h** $4x \times 3xy \div (2x)$


i $3pq \times p \div (6q)$ **j** $5ab^2 \times 4 \div (10b)$




Hint for Q6: Apply one operation at a time from left to right in these; e.g.
$4 \times x \div y = 4x \div y$
$= \frac{4x}{y}$

7 Write a simplified expression for the area of the following shapes.


a




b



c



d





Hint for Q7: Area of a triangle = $\frac{1}{2} \times$ base $\times$ height

8 A bag of lollies contains x jelly beans. Jean buys 6 bags of lollies to share equally among her three grandchildren. Write a simplified expression for how many jelly beans each grandchild receives.

9 Fill in the missing term to make each of the following correct.

a $3m \times \square = 12mn$ **b** $4x \times \square = 28xy$ **c** $-2xy \times \square = -18xy^2$


d $6a \times \square = 30a^2b$ **e** $\frac{\square}{7} = 2x$ **f** $\frac{15}{\square} = \frac{3}{2y}$

g $\frac{\square}{18ab} = \frac{2}{3b}$ **h** $\frac{\square}{4x} = 2xy$ **i** $\frac{\square}{2x} = 5x$

Rectangular paddocks

— | 10, 11

10 The length of a rectangular paddock is three times its width. Its width is x metres. It is bordered on one side by a stream.



a Write a simplified expression for the:

i area of the paddock

ii length of fencing needed for the three sides of the paddock

b If $x = 50$, use your answers to part **a** to give the:

i area of the paddock **ii** length of fencing material required.

11 Another rectangular paddock has its area given by the expression $4ab^2$. Find the width of the paddock if the length is:

a ab **b** $2b$ **c** $4ab$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3D Expanding algebraic expressions

Learning intentions

- To know how the distributive law is used to expand and remove brackets
- To be able to apply the distributive law to expand brackets
- To be able to simplify expressions involving brackets and like terms

Key vocabulary: distributive law, expand, like terms

We know from previous sections that, when brackets are used in expressions, we need to do the mathematics inside the brackets first.


For example, the expression $2(x + 3)$ means 'first add three to x, and then double the result'. If $x = 10$, we get $2 \times (10 + 3) = 2 \times 13 = 26$.

Similarly, if we multiply each term inside the brackets separately by 2, the result is also 26: $(2 \times 10) + (2 \times 3) = 20 + 6 = 26$.

We can apply this to algebraic expressions, so that $2(x + 3)$ can also be written as $2 \times x + 2 \times 3 = 2x + 6$. This uses the distributive law for addition.


Lesson starter: Rectangular distributions

This rectangle has the dimensions as shown. Given that the area of a rectangle is length × width:



- describe two different ways to find the area of the rectangle
- fill in the boxes below to show the two methods used
  $\square(\square + \square) = \square \times \square + \square \times \square$

This diagram also shows two joined rectangles with the given dimensions, with one unknown, x.



- Use your methods from above to find two different ways to write expressions for the combined area of the two rectangles.
- Complete the following: $4(_____) = _____ + _____$

Key ideas

- The **distributive law** is used to **expand** and remove brackets.
  - It states that adding numbers in brackets, *then* multiplying the total, gives the same answer as multiplying each number in the brackets separately first, *then* adding the products.
- To expand brackets:
  - The term on the outside of the brackets is multiplied by each term inside the brackets.

    $a(b + c) = ab + ac$ or $a(b - c) = ab - ac$

    For example:

    $2(x + 4) = 2 \times x + 2 \times 4 = 2x + 8$
  - If the number in front of the brackets is negative, the sign of each of the terms inside the brackets will change when expanded.
    For example: $-2(x - 3) = -2x + 6$ since $-2 \times x = -2x$ and $-2 \times (-3) = 6$


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Exercise 3D

Understanding

1–4 | 4

1 This diagram shows two joined rectangles with the given dimensions.



a Write an expression for the area of:
- **i** the larger rectangle (x by 5)
- **ii** the smaller rectangle (2 by 5)

b Use your answers from part **a** to find the combined area of both rectangles.

c Write an expression for the total side length of the side involving x.

d Use your answer from part **c** to find the combined area of both rectangles. You will need brackets.

e Complete this statement: $5(x+2) =$ _____ + _____

2 Complete the following:
To expand brackets the _____ on the outside of the brackets is _____ by each term _____ the brackets.

3 Using the distributive law, $x(y-z)$ is equivalent to

A $xy - z$ **B** $xy + z$ **C** $xy - xz$ **D** $-xyz$

4 Fill in the boxes to complete the following.

a $3(x+5) = \square \times x + \square \times 5$
$= 3x + \square$

b $-4(x+2) = \square \times x + \square \times 2$
$= \square - \square$

Fluency

5–7(½) | 5–8(½)

Example 11 Expanding simple expressions with brackets

Expand the following.

a $3(x+4)$ **b** $5(x-11)$

Solution

a $3(x+4) = 3x + 12$

b $5(x-11) = 5x - 55$

Explanation

The 3 on the outside is multiplied by each term inside the brackets; i.e.

$3(x+4) = 3 \times x + 3 \times 4$

$5 \times x = 5x$ and $5 \times (-11) = -55$

Now you try

Expand the following.

a $4(x+2)$ **b** $6(x-10)$

5 Expand the following.

a $2(x+3)$ **b** $5(x+12)$ **c** $2(x+7)$
d $7(x+9)$ **e** $3(x-7)$ **f** $6(x-11)$
g $5(x-9)$ **h** $10(x-1)$ **i** $3(2+x)$
j $7(3+x)$ **k** $4(7-x)$ **l** $5(8-x)$



Hint for Q5: Remember that the number in front of the brackets is multiplied by each term inside the brackets when expanding: $2(x+3) = \ldots$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Example 12 Expanding brackets with a negative number in front

Expand the following.

a $-2(x+5)$ b $-3(x-4)$

Solution

a $-2(x+5) = -2x - 10$

b $-3(x-4) = -3x + 12$

Explanation

-2 is multiplied by each term inside the brackets.
$-2 \times x = -2x$ and $-2 \times 5 = -10$

Multiplying by a negative changes the sign of each term in the brackets.
$-3 \times x = -3x$ and $-3 \times (-4) = +12$

Now you try

Expand the following.

a $-3(x+7)$ b $-5(x-9)$

6 Expand the following.

a $-3(x+2)$ b $-2(x+11)$ c $-5(x-3)$ d $-6(x-6)$
e $-4(2-x)$ f $-13(3+x)$ g $-8(9+x)$ h $-300(1-x)$

Example 13 Expanding brackets and simplifying

Expand the following.

a $4(x+3y)$ b $2x(4x-3)$

Solution

a $4(x+3y) = 4 \times x + 4 \times 3y$
$= 4x + 12y$

b $2x(4x-3) = 2x \times 4x + 2x \times (-3)$
$= 8x^2 - 6x$

Explanation

Multiply each term inside the brackets by 4.
$4 \times x = 4x$ and $4 \times 3 \times y = 12y$.

Each term inside the brackets is multiplied by $2x$.
$2x \times 4x = 2 \times 4 \times x \times x = 8x^2$ and
$2x \times (-3) = 2 \times (-3) \times x = -6x$.

Now you try

Expand the following.

a $5(a+4b)$ b $3y(5y-4)$

7 Expand the following.

a $2(a+b)$ b $5(a+2b)$
c $3(2m+y)$ d $8(2x-5)$
e $-3(4x+5)$ f $4x(x-2y)$
g $t(2t-3)$ h $a(3a+4)$
i $d(2d-5)$ j $2b(3b-5)$
k $2x(4x+1)$ l $5y(1-3y)$



Hint for Q7: $x \times x$ is written as x^2.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3D

Example 14 Simplifying by removing brackets

Expand the following and collect like terms.

a $4(x+5)-10$ b $2-3(x-4)$

Solution

a $4(x+5)-10 = 4x+20-10$
$= 4x+10$

b $2-3(x-4) = 2-(3x-12)$
$= 2-3x+12$
$= 14-3x$

Explanation

Expand brackets first: $4(x+5) = 4\times x+4\times 5$.
Then collect like terms and simplify.

Expand brackets.
$3(x-4) = 3x-12$.
$-(3x-12) = -1(3x-12)$, so multiplying by negative 1 changes the sign of each term inside the brackets;
i.e. $-1\times 3x = -3x$ and $-1\times(-12) = +12$.

Now you try

Expand the following and collect like terms.

a $7(x+3)-15$ b $8-2(y-5)$

8 Expand the following and collect like terms.


a $2(x+4)+3$ b $6(x+3)+4$ c $5(x+2)-4$
d $3(x+4)-2$ e $3+4(x-2)$ f $7+2(x-3)$
g $2-3(x-2)$ h $1-5(x-4)$ i $5-2(x+3)$
j $12-3(x+4)$ k $7-(x+4)$ l $4-(3x-2)$



Hint for Q8: Expand the brackets first then collect like terms.

Problem-solving and reasoning 9–11 10–13

9 The diagram shows the route taken by a salesperson who travels from A to D via B and C.





a If the salesperson then returns directly to A, write an expression (in simplest form) for the total distance travelled.
b If $y = x+1$, write an expression for the total distance the salesperson travels in terms of x only. Simplify your expression.
c When $y = x+1$, how much would the distance have been reduced by (in terms of x) if the salesperson had travelled directly from A to D and straight back to A?





Hint for Q9: Subtract the distance from A to D and back again from the total distance.

10 Find the area of these basic shapes in expanded form. All angles are 90°.

a 

b 

c 

d 



Hint for Q10: Remember to include brackets when multiplying length by width.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

11 Murray has \$60 in savings. One Saturday, he does some odd jobs for a neighbour and earns \$$x$, which he adds to his savings. Murray's parents think he's doing a good job of saving money, so they decide to double his savings. Write an expanded expression for the amount of savings Murray has now.



12 Identify the errors in these expressions then write out the correct expansion.

a $2(x+6) = 2x+6$

b $x(x-4) = 2x-4x$

c $-3(x+4) = -3x+12$

d $-7(x-7) = -7x-49$

e $5(x+2)+4 = 5x+6$

f $5-2(x-7) = 5-2x-14$

$= -9-2x$



Hint for Q12: Watch out for the negative numbers.

13 Jill pays tax at 20c in the dollar for every dollar earned over \$10 000. Jill earns \$$x$ and $x > 10\,000$.

a Write an expression for the amount that Jill pays tax on.

b Write an expression for the amount Jill pays in tax. Expand your answer.



Hint for Q13: 20c in the dollar is 0.2 of each dollar. In \$10, this is $0.2 \times 10 = \$2$.

Pairs of brackets — 14, 15

14 Expand each pair of brackets first, then collect like terms.

a $2(x+3)+3(x+2)$

b $2(x+4)+2(x-1)$

c $3(2x+4)+5(x-1)$

d $4(3x+2)+5(x-3)$

e $-2(x+2)+3(x-1)$


f $2(4x-3)-2(3x-1)$

g $x(x+2)+2x(x+4)$


h $3(x-1)+x(x+2)$

15 Find the area of these shapes in expanded form. All angles are 90°.


a



b



c





Hint for Q15: Split these shapes into rectangles and add or subtract areas.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3E Solving linear equations CONSOLIDATING

Learning intentions

- To know what represents an equation
- To understand that equivalent equations can be created by applying the same operation to both sides of the equation
- To review the steps for solving simple linear equations involving backtracking
- To be able to solve simple linear equations

Key vocabulary: equation, equivalent equation, solution, substitution, solve

A mathematical statement containing an equals sign, a left-hand side and a right-hand side is called an equation.

$5 = 10 \div 2$, $3x = 9$, $x^2 + 1 = 10$ and $\frac{1}{x} = 4$ are examples of equations. Linear equations can be written in the form $ax + b = c$, where the power of x is 1.

$4x - 1 = 6$, $3 = 2(x + 1)$ and $5x = 2x + 1$ are all linear equations.

Equations are solved by finding the value of the pronumeral that makes the equation true. This can be done by inspection for very simple linear equations (for example, if $3x = 15$ then $x = 5$, since $3 \times 5 = 15$). More complex linear equations can be solved through a series of steps where each step produces an equivalent equation.

Lesson starter: Is $x = 4$ a solution?

State which of the following equations has the solution $x = 4$.

$5x = 20$ $\quad 2x - 1 = 5$ $\quad 10 - x = 14$

$x + 9 = 13$ $\quad 3x + 2 = 14$ $\quad -4x = 16$

- How did you find the equations for which $x = 4$ is a solution?
- The equations you have listed should be equivalent equations to $x = 4$. Starting from $x = 4$, describe what you would do to produce each equivalent equation; for example, 'add 1 to both sides'.

Key ideas

- An **equation** is a mathematical statement containing an equals sign.
  For example: $3x + 1 = 7$.
- The **solution** of an equation is the value of the pronumeral that makes the equation true.
- **Equivalent equations** are created by:
  - adding the same number to or subtracting the same number from both sides of the equation
  - multiplying or dividing both sides of the equation by the same number (not including 0).
- Linear equations can be **solved** by creating equivalent equations and using opposite operations (backtracking).
- The solution to an equation can be checked by substituting the solution into the original equation and checking that both sides are equal.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Exercise 3E

Understanding

1–4 | 3, 4

1 Use substitution to state which of these equations has a solution of $x = 3$.

a $x + 5 = 9$
b $1 - x = -4$
c $3x - 2 = 7$
d $4 - 3x = -5$



Hint for Q1: If $x = 3$ is a solution, the left hand side must equal the value that is on the right hand side of the = sign.

2 State the value of x that makes these equations true.

a $x + 4 = 10$
b $x - 3 = 5$
c $5x = 20$



Hint for Q2: What value plus 4 equals 10?

3 Obtain an equivalent equation by completing the step given in brackets.

a $2x + 5 = 11$ (subtract 5 from both sides)
b $3x - 4 = 17$ (add 4 to both sides)
c $4x = 36$ (divide both sides by 4)

Hint for Q3:

$2x + 5 = 11$
-5 ( ) -5
$2x = \ldots$



4 State the single operation that has been performed to obtain these pairs of equivalent equations; for example, 'add 3 to both sides'.

a $x + 9 = 11$
$x = 2$

b $2x - 1 = 5$
$2x = 6$

c $3x = 27$
$x = 9$

Fluency

5(½), 6 | 5–7(½)

Example 15 Solving simple linear equations

Solve each of the following equations.

a $2x + 5 = 9$ **b** $3x - 7 = 11$

Solution

a $2x + 5 = 9$
$2x = 4$
$x = 2$

Check:
LHS $= 2x + 5$ RHS $= 9$
$= 2 \times (2) + 5$
$= 4 + 5$
$= 9$

Explanation

Subtract 5 from both sides to undo the +5.
Divide both sides by 2, since $2x = 2 \times x$ and $\frac{{}^{1}\cancel{2} \times x}{\cancel{2}^{1}} = x$.

Check the answer by substituting $x = 2$ into the original equation to check that the LHS = RHS.

Continued on next page

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3E

b $3x - 7 = 11$
$3x = 18$
$x = 6$

Add 7 to both sides to undo the -7, since $-7 + 7 = 0$.
$3x = 3 \times x$, so divide both sides by 3.

Check:

Check your solution by substituting $x = 6$.

LHS $= 3x - 7$ RHS $= 11$
$= 3 \times (6) - 7$
$= 18 - 7$
$= 11$

Now you try

Solve each of the following equations.

a $3x + 8 = 23$ **b** $4x - 2 = 10$

5 Solve each of the following equations. Check your answers using substitution.

a $2x + 7 = 15$ **b** $5a + 6 = 11$ **c** $3m + 4 = 16$
d $2x + 4 = 12$ **e** $2n + 13 = 17$ **f** $3x + 5 = -7$
g $5b - 3 = 12$ **h** $3y - 2 = 19$ **i** $7a - 8 = 20$
j $4b - 7 = 25$ **k** $2x - 4 = -6$ **l** $10y - 35 = -15$



Hint for Q5: Think carefully about each step to make sure you do produce an equivalent equation.

Example 16 Solving equations with negative coefficients

Solve the following equations.

a $9 - 2x = 15$ **b** $7 - 4x = 10$

Solution **Explanation**

a $9 - 2x = 15$
$-2x = 6$
$x = -3$

Subtract 9 from both sides, leaving $-2x$ on the LHS.
Divide both sides by -2; recall that a positive number divided by a negative number is negative.
Check your answer.

Check:

LHS $= 9 - 2x$ RHS $= 15$
$= 9 - 2 \times (-3)$
$= 9 - (-6)$
$= 15$

b $7 - 4x = 10$
$-4x = 3$
$x = -\frac{3}{4}$

Subtract 7 from both sides.
Divide both sides by -4. Leave your answer in fraction form; i.e. $3 \div (-4) = -\frac{3}{4}$.

Check:

LHS $= 7 - 4x$ RHS $= 10$
$= 7 - 4 \times \left(-\frac{3}{4}\right)$
$= 7 - (-3)$
$= 10$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Now you try

Solve the following equations.

a $10 - 3x = 16$ **b** $11 - 5x = 15$

6 Solve each of the following equations.

a $12 - 2x = 18$ **b** $2 - 7x = 9$
c $15 - 5x = 25$ **d** $3 - 2x = 19$
e $2 - 5x = -8$ **f** $24 - 7x = 10$
g $14 - 8x = -2$ **h** $3 - 4x = -21$

7 Solve these equations. Leave your answer in fraction form.

a $2x + 5 = 8$
b $5x - 1 = 2$
c $11x - 3 = 7$
d $6 - 2x = 5$
e $4 - 3x = 5$
f $8 - 5x = 4$



Hint for Q7: If $5x = 3$, $x = \frac{3}{5}$ (divide both sides by 5). The answer can be left as a fraction.

Problem-solving and reasoning

8, 9 | 8(½), 9, 10

8 For each of the following, write an equation and solve it to find the unknown value. Use x as the unknown value.

a If 8 is added to a certain number, the result is 34.
b Seven less than a certain number is 21.
c I think of a number, double it and add 4. The result is 10.
d I think of a number, triple it and subtract 4. The result is 11.
e Four less than three times a number is 20.
f I start with the number 25 and subtract twice a certain number. The result is 7.



Hint for Q8: Form the algebraic expression (e.g. $x + 8$) and set it equal to the result to make the equation.

9 Describe the error made in each of these incorrect solutions.

a $3x + 2 = 8$
$3x = 10$
$x = \frac{10}{3}$

b $2x - 1 = 5$
$2x = 6$
$x = 12$

c $5 - x = 12$
$x = 7$

d $2x + 3 = 8$
$x + 3 = 4$
$x = 1$

10 Solve these equations.

a $-3 + 4x = 17$ **b** $-5 - 2x = 13$ **c** $-10 - 3x = -4$

Is the car speeding?

— | 11

11 A car is travelling in zones where speed signs are in metres per second (m/s).

a When it first sets out, its speed (s m/s) is given by $s = 2t$, where t is time in seconds. How long does it take for the car to reach a speed of 20 m/s (this is 72 km/h)?
b The car starts to accelerate again, and its speed is now given by $s = 16 + 1.5t$. For how many seconds can it continue this acceleration before it reaches 28 m/s (about 100 km/h)?



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3F Solving linear equations involving fractions

Learning intentions

- To know how to write pronumerals with fractional coefficients in fraction form
- To know that multiplication and division are inverse operations
- To understand the order in which steps need to be applied to solve equations involving fractions
- To be able to solve linear equations involving fractions

Key vocabulary: coefficient, equation, equivalent equation, inverse

We saw in the previous section that to solve an equation such as $2x = 8$, we divide each side of the equation by 2.

What happens if the equation involves a fraction? For the equation $\frac{1}{2}x = 2$, we could similarly divide both sides by $\frac{1}{2}$ to solve it. But we can also write the equation in a different way, to make it easier to solve.

A half of x is equivalent to dividing x by 2, so $\frac{1}{2}x = 5$ can be written as $\frac{x}{2} = 5$.

To solve this new equation, we can multiply both sides by 2, since $2 \times \frac{x}{2} = x$. This gives us $x = 10$.

From this process, we can see that we could solve this equation by either dividing both sides by $\frac{1}{2}$ *or* multiplying both sides by 2.

For more complex equations we'll also need to consider the order of operations.

Lesson starter: Fractions of x

Consider these problems involving fractions.

1. If half of x is 8, what is x?
2. If a third of x is 5, what is x?
3. If three-quarters of x is 30, what is x?
   - Discuss the methods you used to find the value of x.
   - Write an equation to represent each case.
   - Discuss how your method for finding the answer relates to the fraction in the equation.

Key ideas

- Pronumerals with a fraction coefficient can be expressed in fraction form; i.e. $\frac{1}{3}x = \frac{x}{3}$ and $\frac{3}{5}x = \frac{3x}{5}$.
- Multiplication and division are **inverse** operations; i.e. $2 \times 3 = 6$ and $\frac{6}{2} = 3$.
- Equations involving fractions are also solved by creating a series of simpler equivalent equations.

  For example: $\frac{2x}{3} = 8$

  $2x = 24$

  $x = 12$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

- At some stage both sides of the equation can be multiplied by the number in the denominator of the fraction. The order of when this happens is important.

For example: $\frac{x-1}{2}=4$ is different from $\frac{x}{2}-1=4$

$x-1=8$ $\frac{x}{2}=5$

$x=9$ $x=10$

Exercise 3F

Understanding

1–4 | 4

1 Fill in the boxes to produce an equivalent expression.

a $\frac{1}{5}x=\frac{x}{\square}$

b $\frac{3x}{4}=\square x$

c $-\frac{2}{3}x=-\frac{\square}{3}$

2 Write down the value of x that is the solution to these equations. No written working is required.

a $\frac{x}{2}=10$

b $\frac{x}{4}=6$

c $\frac{x+1}{3}=4$

d $\frac{x}{5}-2=1$



Hint for Q2: In **a**, ask yourself which number divided by 2 gives 10.

3 Which one of the following is an equivalent equation to $\frac{x}{4}+1=8$?

A $x+1=32$
B $x=36$
C $\frac{x}{4}=7$
D $\frac{x+1}{4}=2$

4 Match each equation in the first column with its equivalent equation in the second column and state the single operation that generates the equivalent equation (for example, multiply both sides by 5).

a	$\frac{x}{3}-1=2$	**i**	$x+2=3$
b	$\frac{x}{4}=2$	**ii**	$x-1=6$
c	$\frac{x+2}{3}=1$	**iii**	$x=8$
d	$\frac{x-1}{2}=3$	**iv**	$\frac{x}{3}=3$



Hint for Q4: Think about what step you would apply first to produce a simple equivalent equation.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3F

Fluency

5–6(½) | 5–7(½)

Example 17 Solving linear equations with fractional coefficients

Solve each of the following equations.

a $\frac{2x}{3} = 8$ b $\frac{x}{4} - 3 = 7$

Solution | **Explanation**

a $\frac{2x}{3} = 8$

$2x = 24$

$x = 12$

Multiply both sides by 3 to remove the fraction on the left-hand side. Then divide both sides by 2.

Check that $x = 12$ makes LHS = RHS.

Check:

LHS $= \frac{2x}{3}$ RHS $= 8$

$= \frac{2 \times (12)}{3}$

$= \frac{24}{3}$

$= 8$

b $\frac{x}{4} - 3 = 7$

$\frac{x}{4} = 10$

$x = 40$

Add 3 to both sides first to undo the −3.

Multiply both sides by 4.

Check the answer by substituting $x = 40$ into $\frac{x}{4} - 3$.

Check:

LHS $= \frac{x}{4} - 3$ RHS $= 7$

$= \frac{40}{4} - 3$

$= 10 - 3$

$= 7$

Since this equals 7, $x = 40$ is the solution.

Now you try

Solve each of the following equations.

a $\frac{3x}{4} = 6$ b $\frac{x}{2} - 5 = 10$

5 Solve each of the following equations.

a $\frac{x}{5} = 3$ b $\frac{y}{2} = 7$ c $\frac{2y}{5} = 4$

d $\frac{3x}{2} = 6$ e $\frac{x}{4} + 3 = 5$ f $\frac{x}{2} + 4 = 5$

g $\frac{b}{3} + 5 = 9$ h $\frac{t}{2} + 5 = 2$ i $\frac{a}{3} + 4 = 2$

j $\frac{y}{5} - 4 = 2$ k $\frac{x}{3} - 7 = -2$ l $\frac{s}{2} - 3 = -1$

m $\frac{x}{4} - 5 = -2$ n $\frac{m}{4} - 2 = -3$ o $\frac{y}{5} + 4 = -3$



Hint for Q5: In part e, the first step is to subtract 3 from both sides.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Example 18 Solving simple fractional equations

Solve the equation $\frac{x+4}{6} = 2$.

Solution

$\frac{x+4}{6} = 2$

$x + 4 = 12$

$x = 8$

Check:

LHS $= \frac{x+4}{6}$ RHS $= 2$

$= \frac{(8)+4}{6}$

$= \frac{12}{6}$

$= 2$

Explanation

Multiply both sides by 6 first, since all of $(x+4)$ is divided by 6.

Subtract 4 from both sides and check the answer.

Now you try

Solve the equation $\frac{x-5}{3} = 4$.

6 Solve each of the following equations. Check your answers.

a $\frac{x+1}{3} = 4$ **b** $\frac{x+4}{2} = 5$ **c** $\frac{y+4}{3} = 2$

d $\frac{b+6}{2} = 3$ **e** $\frac{y-2}{3} = 6$ **f** $\frac{t-4}{5} = 7$

g $\frac{k-1}{7} = 8$ **h** $\frac{x-7}{9} = 7$ **i** $\frac{x+2}{4} = -2$

j $\frac{b+3}{7} = -3$ **k** $\frac{y-3}{5} = -5$ **l** $\frac{y-8}{7} = -1$



Hint for Q6: Think carefully about the order of the steps here.

7 Solve this mixed selection of equations.

a $2t + 1 = 7$ **b** $\frac{m}{4} - 2 = 8$ **c** $5 - 2y = 9$

d $\frac{x-4}{5} = 3$ **e** $3 + \frac{y}{6} = 2$ **f** $4t - 5 = -8$

Problem-solving and reasoning

8, 9 | 8(½), 9, 10, 11(½)

8 For each of the following, write an equation and solve it to find the unknown value. Use x as the unknown value.

a If a certain number is divided by 3, the result is 12.

b A certain number is doubled, then divided by 5. The result is 4.

c I think of a number, halve it and subtract 4. The result is 10.

d I think of a number, add 3 and divide this by 4. The result is 6.

e A number is multiplied by 7 and the product is divided by 3. The final result is 8.

f 5 is added to a third of a certain number. The result is 16.


Hint for Q8: x halved is $\frac{x}{2}$. A third of x is $\frac{x}{3}$.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3F

9 A bag of chocolate eggs containing x eggs is shared equally between India and her two brothers. After India eats 2, she has 5 left. How many eggs were in the bag?





Hint for Q9: Start with an expression for the number of eggs India first receives.

10 Describe the error made in each of these incorrect solutions. Then write out the correct solution.

a $\frac{x+2}{3} = 7$

$\frac{x}{3} = 5$

$x = 15$

b $\frac{x}{3} - 4 = 2$

$x - 4 = 6$

$x = 10$

11 Solve these equations involving more than two steps.

a $\frac{2x}{3} - 1 = 7$ **b** $\frac{3x}{4} - 2 = 7$ **c** $\frac{x}{2} + 3 = 1$

d $\frac{3b-8}{2} = 5$ **e** $\frac{2x+2}{3} = 4$ **f** $\frac{7m-8}{3} = 9$

g $\frac{5-y}{3} = 2$ **h** $\frac{4-2t}{6} = 3$ **i** $1 - \frac{4x}{3} = 5$



Hint for Q11: Keep operating on both sides of the equation until you find the solution.

Fractional coefficients — 12, 13

12 Solve these equations involving fractional coefficients.

a $\frac{2}{3}x = 6$ **b** $\frac{5}{6}x = 30$ **c** $-\frac{2}{7}x = 2$

d $\frac{3}{4}x - 1 = 5$ **e** $\frac{7}{5}x - 3 = 4$ **f** $1 - \frac{3}{5}x = 7$

13 If $\frac{7}{5}x = 14$, does it matter whether you multiply by 5 first or divide by 7 first? Which method would be best here?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Progress quiz

3A **1** Consider the expression $4x - 5xy - 3y + 8$.
- **a** How many terms are in the expression?
- **b** What is the coefficient of y?
- **c** What is the constant term?

3A **2** Write algebraic expressions for the following.
- **a** The cost of 3 drinks at $\$d$ each
- **b** $\$500$ winnings shared equally between x people
- **c** Four less than three lots of y
- **d** The sum of m and n doubled

3A **3** Evaluate these expressions if $x = 3$, $y = -4$ and $z = 7$.
- **a** $2y + z$
- **b** $xy - y$
- **c** $y^2 + x - z$

3B **4** Simplify the following by collecting like terms.
- **a** $5x + 7x$
- **b** $8b - b$
- **c** $12a + 5 - 3a$
- **d** $6xy - 4x + 2xy$
- **e** $4a + 7b - a + 3b$
- **f** $5mn + 2n + mn - 5n$
- **g** $4xy^2 - 5xy - 2xy^2 + 3yx$
- **h** $3a^2b + 2ab^2 - a^2b + 5ab^2$
- **i** $-m^2n + 5mn^2 - m^2n - 7mn^2$

3C **5** Simplify the following.
- **a** $6 \times 4b$
- **b** $3a \times 8c$
- **c** $-5x \times 4y$
- **d** $7ab \times 4b$
- **e** $\frac{3x}{12}$
- **f** $\frac{6xy}{8x}$
- **g** $\frac{9x^2y}{6x}$
- **h** $6x \times 4y \div (3y)$
- **i** $4ab \times 5b \div (8b)$

3D **6** Expand the following.
- **a** $3(x + 5)$
- **b** $4(y - 1)$
- **c** $-2(x + 5)$
- **d** $-5(a - 2)$
- **e** $6(2x - 5y)$
- **f** $3m(4m + n)$

3D **7** Expand the following and collect like terms.
- **a** $3(2x - 5) + 7$
- **b** $8 - 4(x + 3)$
- **c** $1 - (2x - 1)$

3E **8** Solve the following equations.
- **a** $3x + 2 = 11$
- **b** $5x - 3 = 17$
- **c** $8 - 2x = 26$
- **d** $12 - 5x = 15$

3F **9** Solve the following equations involving fractions.
- **a** $\frac{x}{7} = 4$
- **b** $\frac{x}{3} + 6 = 22$
- **c** $\frac{x - 7}{2} = 10$
- **d** $\frac{x + 1}{3} = -7$

3E/3F **10** Write an equation to find the unknown number x in the following and solve it.
- **a** Five more than three times the number is 32.
- **b** Three less than the number is divided by 4 to give 5.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3G Solving equations with brackets

Learning intentions

- To know that equations involving brackets can be solved by first expanding the brackets
- To understand that equations with a common factor can be simplified by first dividing both sides by the factor
- To be able to solve equations involving brackets

Key vocabulary: expand, equation

Just as we have seen expressions involving brackets, linear equations may also involve brackets. Brackets can be removed by expanding and then the remaining equation can be solved. Collection of like terms may also be required.

Lesson starter: To expand or not to expand?

The steps to solve two problems involving brackets are listed here in the incorrect order.

1 $3(x+4) = 14$

$x = \frac{2}{3}$

$3x + 12 = 14$

$3x = 2$

2 $3(x+2) = 12$

$x = 2$

$x + 2 = 4$

- Arrange the steps in the correct order, working from top to bottom.
- By considering all the steps in the correct order, explain what has happened in each step.
- Describe the two different approaches to solving the problems.
- Can you see why these methods have been used in each case?
- Apply method 1 to the second equation.

Key ideas

- Equations with brackets can be solved by first expanding the brackets and then solving the remaining equation.
  For example: $3(x+1) = 2$ becomes $3x + 3 = 2$.
- An alternate method is to deal with the common factor first.
  For example: $5(x-1) = 15$ becomes $x - 1 = 3$ by first dividing both sides by 5.
- For equations like $2(x+4) + 3x = 28$ the brackets are always expanded first, then collect any like terms on the LHS and continue to solve as usual.

Exercise 3G

Understanding

1–3 | 3

1 Complete the missing parts to produce an equation without brackets.

a

$3(x+7) = 25$

$3 \times \square + \square \times 7 = 25$

$3x + \square = 25$

b

$4(2x-1) = 15$

$\square \times 2x + 4 \times \square = 15$

$\square - 4 = 15$

2 Give the equivalent equation that results from dividing both sides of the equation by the number in front of the brackets.

a $3(x+2) = 9$

b $4(x-3) = 20$

c $-2(x+6) = 8$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3 Expand these expressions and simplify.

a $3(x-4)+x$ **b** $5(x-3)-3x$ **c** $4(x+2)-6x$

Fluency

4–5(½) 4–6(½)

Example 19 Solving equations with brackets

Solve $2(3x-4)=11$.

Solution

$2(3x-4)=11$

$6x-8=11$

$6x=19$

$x=\frac{19}{6}$ or $3\frac{1}{6}$

Explanation

Expand the brackets:

$2(3x-4)=2\times 3x+2\times(-4)$

Solve the remaining equation by adding 8 to both sides, then dividing both sides by 6.

Leave your answer in fraction form.

Now you try

Solve $3(2x-5)=7$.

4 Solve each of the following equations by first expanding the brackets.

a $2(x+3)=11$ **b** $5(a+3)=18$

c $3(m+4)=31$ **d** $5(y-7)=12$

e $4(p-5)=15$ **f** $2(k-5)=9$

g $4(5-b)=21$ **h** $2(1-m)=13$

i $5(3-x)=19$ **j** $7(2a+1)=8$

k $4(3x-2)=30$ **l** $3(3n-2)=0$

m $5(3-2x)=16$ **n** $6(1-2y)=8$



Hint for Q4: Once you have expanded the brackets, the equation can be solved in the usual way.

Example 20 Solving equations with brackets and collecting like terms

Solve $6(x+3)-4x=32$.

Solution

$6(x+3)-4x=32$

$6x+18-4x=32$

$2x+18=32$

$2x=14$

$x=7$

Explanation

Expand the brackets first, then collect any like terms: $6x-4x=2x$. Solve the remaining equation by subtracting 18 from both sides, and then divide both sides by 2.

Now you try

Solve $5(x+4)-2x=35$.

5 Expand and simplify, then solve each of the following equations.

a $2(x+4)+2x=12$ **b** $2(x-3)+3x=4$

c $6(x+3)-2x=26$ **d** $5(x+2)-2x=46$

e $3(2x-3)+x=12$ **f** $4(3x+1)-3x=19$

g $2(3x+5)-8x=20$ **h** $4(1-x)-x=9$



Hint for Q5: Expand the brackets and then combine like terms before solving.

6 Solve these equations.

a $4(x-1)+x-1=0$ **b** $3(x+2)-2+4x=18$ **c** $3(2x+3)-1-4x=10$

d $2(4x-1)-5x+6=31$ **e** $2(4x+2)-10x-6=6$ **f** $5(2x+3)-15x-7=3$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3G

Problem-solving and reasoning

7, 8 | 7(½), 9, 10

7 Using x for the unknown number, write down an equation then solve it to find the number. The first one is done for you.

a Three times 1 more than a number is 4. $(3(x+1)=4;\ x=\frac{1}{3})$

b Twice 2 less than a number is 19.

c The product of 2 and 3 more than a number is 7.

d The product of 3 and 4 less than a number is 8.

e When 2 less than 3 lots of a number is doubled, the result is 5.

f When 5 more than 2 lots of a number is tripled, the result is 10.



Hint for Q7: You will need to use brackets when setting up the expressions in these equations.

8 Since Tara started her job, her original hourly wage ($\$x$) has been tripled, then decreased by $\$6$. Her pay is now to be doubled so that she earns $\$18$ an hour. What was her original hourly wage?





Hint for Q8: Tripling Tara's pay of $\$x$ is $3 \times x = 3x$.

9 Consider the equation $3(x-2)=9$.

a Solve the equation by first dividing both sides by 3.

b Solve the equation by first expanding the brackets.

c Which of the above two methods is preferable and why?



Hint for Q9: You may recall this idea from the 'Lesson starter' activity at the beginning of the section.

10 Consider the equation $3(x-2)=7$.

a Solve the equation by first dividing both sides by 3.

b Solve the equation by first expanding the brackets.

c Which of the above two methods is preferable and why?

Brackets or no brackets?

— | 11

11 Decide if brackets need to be inserted into the following equations to make $x=4$ the solution. If so, insert the brackets where required.

a $3x-2=10$ **b** $6x-12=12$ **c** $4x+1=20$

d $2x-4+3x+1=13$ **e** $3x+2+2x-5=12$ **f** $2x-3-3x-1=-4$

Using a calculator 3G: Solving equations

This activity can be found in the More Resources section of the Interactive Textbook in the form of a printable PDF.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3H Solving equations with pronumerals on both sides

Learning intentions

- To be able to collect pronumerals on one side of the equation using addition or subtraction
- To be able to solve equations with pronumerals on both sides of the equation

Key vocabulary: pronumeral, expand, equation, coefficient

More complex linear equations may have pronumerals on both sides of the equation and/or brackets. Examples are $3x = 5x - 1$ or $4(x + 2) = 5x$.

We've seen that brackets can be removed by expanding. Equations with pronumerals on both sides can be solved by collecting like terms using addition and subtraction.



It's important to keep track of your steps while solving equations with pronumerals on both sides.

Lesson starter: What is it worth?

Consider these two cases, where red buttons and blue buttons are shown on either side of a dividing line. In each case, the buttons on either side of the line have an equivalent value.

1 

2 

- In each case, find how many red buttons one blue button is worth.
- Describe the steps that helped you arrive at your answer.
- Carry out similar steps in the following cases to find what x is worth if each side of the dividing line is equivalent.

1

$4x$	$2x$
$+10$	

2

$5x$	x
$+4$	$+12$

- Use the same thought processes to write out the steps for solving:
  $4x = 2x + 10$
  $5x + 4 = x + 12$

Key ideas

- If an equation has pronumerals on both sides, collect them on one side by adding or subtracting one of the terms.
  For example: $3x + 4 = 2x - 3$ becomes $x + 4 = -3$ by subtracting $2x$ from both sides.
- The resulting equation will be simplest if the pronumeral is collected on the side with the higher coefficient. For example, in $5x + 2 = 4x + 7$, collect the pronumeral on the left hand side, since 5 is greater than 4.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Exercise 3H

Understanding

1–3 | 3

1 State which side (left or right) you would collect the x terms on to keep the x coefficient positive.

a $5x = 2x + 3$ **b** $6x + 1 = 2x + 5$

c $3x - 5 = 7x + 3$ **d** $1 - 2x = 3x - 9$



Hint for Q1: The x coefficient is the numeral multiplied by the x.



Hint for Q2: For $3 - x$, adding x makes 3.

2 State the step you would carry out to collect the x terms on the left hand side; e.g. 'subtract $2x$ from both sides'.

a $7x = 3x + 4$ **b** $9x + 2 = 5x + 10$

c $3x - 1 = 3 - x$ **d** $4x + 2 = 8 - 2x$

3 Show the next step only for the given equations and instructions.

a $3x = x + 6$ (subtract x from both sides)

b $4x - 3 = 2x + 1$ (subtract $2x$ from both sides)

c $2x = 1 - 3x$ (add $3x$ to both sides)

d $4 - x = 8 + 5x$ (add x to both sides)

Fluency

4–6(½) | 4–6(½)

Example 21 Solving equations with pronumerals on both sides

Solve these equations with pronumerals on each side.

a $4x = 2x + 10$ **b** $3x = 16 - 5x$

Solution

a $4x = 2x + 10$

$2x = 10$

$x = 5$

Check:

LHS $= 4x$ RHS $= 2x + 10$

$= 4 \times (5)$ $= 2 \times (5) + 10$

$= 20$ $= 20$

b $3x = 16 - 5x$

$8x = 16$

$x = 2$

Check:

LHS $= 3x$ RHS $= 16 - 5x$

$= 3 \times (2)$ $= 16 - 5 \times (2)$

$= 6$ $= 16 - 10$

$= 6$

Explanation

Collect x terms on LHS (since the x coefficient is larger on that side) by subtracting $2x$ from both sides: $4x - 2x = 2x$.

Solve by dividing both sides by 2 and check that $x = 5$ gives the same value for $4x$ and $2x + 10$.

Collect x terms by adding $5x$ to both sides (collect on LHS since $3x$ coefficient is positive). Divide both sides by 8 and check the solution.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Now you try

Solve these equations with pronumerals on each side.

a $8x = 5x + 12$ **b** $2x = 18 - 4x$

4 Solve these equations by first collecting the pronumeral on one side.

a $7x = 5x + 8$ **b** $8x = 3x + 15$
c $9x = 5x + 12$ **d** $10x = 9x + 7$
e $4x = x - 9$ **f** $6x = 2x - 8$
g $3x = 10 - 2x$ **h** $x = 12 - 2x$
i $x = 12 - x$ **j** $4x = 14 - 3x$



Hint for Q4: Remember: It is easier to collect on the side that will keep the x coefficient positive.

Example 22 Solving equations with pronumerals and numerals on both sides

Solve each of the following equations.

a $5x + 2 = 3x + 6$
b $3 - 2x = 5x - 4$

Solution

a $5x + 2 = 3x + 6$
$2x + 2 = 6$
$2x = 4$
$x = 2$

Check:

LHS $= 5 \times (2) + 2$ RHS $= 3 \times (2) + 6$
$= 12$ $= 12$

b $3 - 2x = 5x - 4$
$3 = 7x - 4$
$7 = 7x$
$1 = x$
$\therefore x = 1$

Check:

LHS $= 3 - 2 \times (1)$ RHS $= 5 \times (1) - 4$
$= 1$ $= 1$

Explanation

Collect x terms on one side by subtracting $3x$ from both sides: $5x - 3x = 2x$. Collect on the side with the larger x term to keep it positive. Solve the remaining equation.
Check by substituting $x = 2$ into $5x + 2$ and $3x + 6$ to see that it gives the same result for both.

Add $2x$ to both sides to collect x terms on the side with the positive x coefficient. Solve the remaining equation by adding 4 to both sides and dividing both sides by 7.
Rewrite with x on the left.

Now you try

Solve each of the following equations.

a $7x + 3 = 4x + 15$
b $8 - x = 4x - 2$

5 Solve each of the following equations.

a $5x - 3 = 4x + 5$ **b** $9a + 3 = 8a + 6$
c $3m - 8 = 2m$ **d** $12x - 3 = 10x + 5$
e $8x + 4 = 12x - 16$ **f** $3x + 7 = 8x - 8$
g $2x + 6 = 9 - x$ **h** $3y + 6 = 14 - y$
i $5m - 18 = 15 - 6m$ **j** $4 - 7x = 2x - 23$
k $8 - 2b = 4b + 14$ **l** $3 - 4m = 3m + 24$



Hint for Q5: Collect the pronumerals on one side first.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3H

Example 23 Solving equations with brackets and pronumerals on both sides

Solve these equations involving brackets.

a $7(x+2) = 3x+2$
b $3(2x+4) = 8(x+1)$

Solution

a $7(x+2) = 3x+2$
$7x+14 = 3x+2$
$4x+14 = 2$
$4x = -12$
$x = -3$

b $3(2x+4) = 8(x+1)$
$6x+12 = 8x+8$
$12 = 2x+8$
$4 = 2x$
$2 = x$
$\therefore x = 2$

Explanation

Expand the brackets first, then subtract $3x$ from both sides to collect x terms.
Subtract 14 from both sides ($2 - 14 = -12$) and divide by 4.
Recall that a negative number divided by a positive number is negative ($-12 \div 4 = -3$).

Expand the brackets on each side.
Subtract $6x$ from both sides. (Subtracting $6x$ keeps the x-coefficient positive, as alternatively subtracting $8x$ would end up with $-2x + 12 = 8$.)
Solve the equation and make x the subject.


Now you try

Solve these equations involving brackets.

a $5(x+3) = 3x+7$
b $4(2x-1) = 6(x+2)$

6 Solve each of the following equations.

a $5(x-2) = 2x+11$
b $3(a+1) = a+13$
c $3(y+4) = y+16$
d $2(x+5) = x-4$
e $5b-4 = 6(b+2)$
f $2(4m-5) = 4m+2$
g $3(2a-3) = 5(a+2)$
h $4(x-3) = 3(3x+1)$
i $3(x-2) = 5(x+4)$
j $3(n-2) = 4(n-5)$
k $2(a+5) = 2(2a+3)$
l $4(x+2) = 3(2x+1)$



Problem-solving and reasoning


7, 8 | 7(½), 8, 9

7 Using x for the unknown number, write down an equation then solve it to find the number. The first one has been done for you as an example.

a Twice a number is equal to 4 less than 3 times a number. ($2x = 3x - 4$; $x = 4$)
b 4 more than 2 lots of a number is equal to 5 times the number.
c 10 more than 3 lots of a number is equivalent to 5 lots of the number.
d 2 more than 3 times the number is equivalent to 6 less than 5 times the number.
e 1 less than a doubled number is equivalent to 5 more than 3 lots of the number.
f 4 more than 2 lots of a number is equivalent to the number subtracted from 13.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

8 Mardy is x years old. He correctly tells his mother that she is 6 years older than 3 times his age. His mother replies that she is also 10 years younger than 5 times his age. How old is Mardy?






9 Consider the equation $3x + 1 = 5x - 7$.

- **a** Solve the equation by first subtracting $3x$ from both sides.
- **b** Solve the equation by first subtracting $5x$ from both sides.
- **c** Which method above do you prefer and why? Describe the differences.


Matching sides — 10

10 Use the properties of these basic shapes to find their perimeters.


a




b



c



d





ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3I Solving word problems ★

Learning intentions

- To be able to define a variable to represent the unknown in a problem
- To be able to write an equation to represent a word problem
- To be able to use algebra to solve a word problem

Key vocabulary: pronumeral, variable, equation

Many types of problems can be solved by writing and solving linear equations. Often problems are expressed only in words. Reading and understanding the problem, defining a pronumeral and writing an equation become important steps in solving the problem.

Lesson starter: Too much television?



Three friends, Rick, Kate and Sue, compare how much television they watch in a week at home. Kate watches 3 times the amount of television that Rick watches. Sue watches 4 hours less television than Kate. In total they watch 45 hours of television. How many hours of television did Rick watch?

- Underline each piece of important information in the question.
- Circle the item in the question that you are being asked to find.

Let x hours be the number of hours of television watched by Rick.

- Write expressions for the number of hours of television watched by:
  a Kate **b** Sue
- Add up the number of hours watched by Rick, Kate and Sue using your expressions.
- Write an equation using all the information in the question.
- Solve the equation.
- Answer the question in the original problem.

Key ideas

- To solve a word problem using algebra:
  - Read the problem and find out what the question is asking for.
  - Define a pronumeral and write a statement such as: 'Let x be the number of …'
    The pronumeral will often represent what you have been asked to find.
  - Highlight the key pieces of information in the question.
  - Write an equation that links the facts in the question using your defined pronumeral.
  - Solve the equation using algebra.
  - Answer the question in words.

Exercise 3I

Understanding 1–3 3

1 Match the written scenario with its possible equation.

a 7 is 3 more than a number
b 2 less than a number is 10
c 16 is 1 more than 3 lots of a number
d Half of a number is 3

i $3x + 1 = 16$
ii $\frac{a}{2} = 3$
iii $n + 3 = 7$
iv $y - 2 = 10$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

2 Arrange the following steps in order when solving a word problem using algebra.
- **i** Solve the equation.
- **ii** Read the problem.
- **iii** Answer the question in words.
- **iv** Set up an equation connecting the facts in the question using the defined variable.
- **v** Define a pronumeral to represent the unknown.

3 The sum of the ages of Sam and his brother Bernard is 34. If Sam is 4 years older than Bernard, fill in the following to find their ages.

Let x be the ☐ of Bernard.
The age of Sam is ☐.
The sum of their ages is 34.
$\therefore\ x + \square = 34$
$\square + 4 = 34$
$2x = \square$
$x = \square$
$\therefore$ Bernard is _____ years old and Sam is _____ years old.



Hint for Q3: Start by underlining the key information.

Fluency

4–8 5–9

Example 24 Turning a word problem into an equation

Five less than three times a certain number is 13. Write an equation and solve it to find the number.

Solution	Explanation
Let x be the number. $3x - 5 = 13$ $3x = 18$ $x = 6$	Define the unknown using a pronumeral. Interpret the wording bit by bit to construct the equation. 3 times the number is $3x$, 5 less than this is $3x - 5$ and this equals 13.
The number is 6. Check: $3 \times (6) - 5 = 13$	Solve the equation by adding 5 to both sides, then dividing both sides by 3. Write the answer in words and check your solution.

Now you try

Four more than five times a certain number is 29. Write an equation and solve it to find the number.

4 Write an equation and solve it to find the unknown number in the following.
- **a** 4 less than 2 times a number is 10.
- **b** When a certain number is doubled, it results in a number that is 5 more than the original number.
- **c** I think of a number, divide it by 3 and add 5. The result is 12.
- **d** I think of a number, take away 2 and multiply the result by 4. This gives 24.
- **e** 3 less than a certain number is 9 less than 4 times the number.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

31

Example 25 Applying algebra to a word problem

A bicycle shop hires out bikes. It charges an initial fee of $10 for hiring a bike and then $8 per hour. Leah returns her bike and is charged $42. For how many hours did she hire the bike?

Solution

Let h be the number of hours of hire.

$10 + 8h = 42$

$8h = 32$

$h = 4$

$\therefore$ Leah had the bike for 4 hours.

Explanation

Define the unknown value using a pronumeral.
Write an equation from the information: the cost is $10 plus $8 per hour ($8 \times h$), which equals $42.
Solve the equation for h.

Answer the question in words.


Now you try

A babysitter, Jess, charges an initial fee of $20 and then $15 per hour. If Jess earns $95, for how many hours did she babysit?

5 Toby rented a car for a total cost of $290. The rental company charged $40 per day, plus a hiring fee of $50.

- **a** Define a pronumeral for the number of days Toby rented the car.
- **b** Write an equation in terms of your pronumeral in part **a** to represent the problem.
- **c** Solve the equation in part **b** to find the unknown value.
- **d** For how many days did Toby rent the car?

6 A jeweller earns a weekly amount of $200 plus $10 per item she sells. If in one week she earned $680, how many items did she sell?



Example 26 Solving word problems with more complex equations

David and Usman made 254 runs between them in a cricket match. If Usman made 68 more runs than David, how many runs did each of them make?

Solution

Let the number of runs for David be r.
Number of runs Usman made is $r + 68$.

$r + (r + 68) = 254$

$2r + 68 = 254$

$2r = 186$

$r = 93$

David made 93 runs and Usman made $93 + 68 = 161$ runs.

Explanation

Define the unknown value as a pronumeral.
Write all other unknown values in terms of r.
Usman made 68 more runs than David: $r + 68$.
Write an equation: number of runs for David + number of runs for Usman = 254.
Collect like terms then solve the equation.

Answer the question in words.

Now you try

Max and Tim kicked 11 goals between them in a football match. If Tim kicked 3 more goals than Max, how many goals did each of them kick?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

7 Leonie and Emma scored 28 goals between them in a netball match. Leonie scored 8 more goals than Emma.

- **a** Define a pronumeral for the number of goals scored by Emma.
- **b** Write the number of goals scored by Leonie in terms of the pronumeral in part **a**.
- **c** Write an equation in terms of your pronumeral to represent the problem.
- **d** Solve the equation in part **c** to find the unknown value.
- **e** How many goals did each of them score?

8 A rectangle is four times as long as it is wide and its perimeter is 560 cm.

- **a** Define a pronumeral for the unknown width.
- **b** Write an expression for the length in terms of your pronumeral in part **a**.
- **c** Write an equation involving your pronumeral and the perimeter to represent the problem.
- **d** Solve the equation in part **c**.
- **e** What is the length and width of the rectangle?



Hint for Q8: Draw and label a rectangle to help.

9 A prize of $1000 is divided between Adele and Benita so that Adele receives $280 more than Benita. How much did each person receive?

Problem-solving and reasoning

10–12 | 11–14

10 Andrew, Brenda and Cammi all work part-time at a supermarket. Cammi earns $20 more than Andrew's wage. Brenda earns $30 less than twice Andrew's wage. If their total combined wage is $400, find how much each of these workers earns.



11 Ed walked a certain distance, and then ran twice as far as he walked. He then caught a bus for the last 2 km. If he travelled a total of 32 km, find how far Ed walked and ran.



Hint for Q11: Remember to follow these steps


1. Define a pronumeral for the unknown.
2. Set up an equation using the information from the question.
3. Solve the equation.
4. Answer the question in words.


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

31

12 Kate is three times as old as her son. If Kate is 30 years older than her son, what are their ages?

13 Two paddocks in the shapes shown below are to be fenced with wire. If the same total amount of wire is used for each paddock, what are the side lengths of each paddock, in metres?







Hint for Q13: Solve for w first.

14 Consecutive integers can be represented algebraically as x, $x+1$, $x+2$ etc.

a Find three consecutive numbers that add to 84.

b **i** Write three consecutive even numbers starting with x.

ii Find three consecutive even numbers that add to 18.



Hint for Q14: Make sure that you are still setting up an equation.

Profit and revenue — 15

15 Tedco produces a teddy bear which sells for \$24. Each teddy bear costs the company \$8 to manufacture and there is an initial start-up cost of \$7200.

a Write a rule for the total cost, \$$T$, of producing x teddy bears.

b If the cost of a particular production run was \$9600, how many teddy bears were manufactured in that run?

c If x teddy bears are sold, write a rule for the revenue, \$$R$, received by the company.

d How many teddy bears were sold if the revenue was \$8400?

e If they want to make an annual profit of \$54 000, how many teddy bears do they need to sell?



Hint for Q15: Revenue is the total amount of money collected. Profit is the amount left over from the revenue after the costs have been taken out. Profit = revenue − costs



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3J Solving linear inequalities

Learning intentions

- To be able to work with the symbols $>$, $\geqslant$, $<$, $\leqslant$
- To understand that an inequality represents an infinite set of values
- To be able to illustrate an inequality on a number line using known conventions
- To understand when an inequality symbol needs to be reversed
- To be able to solve a linear inequality

Key vocabulary: inequality, solution set

An inequality (or inequation) is a mathematical statement which uses a $<$, $\leqslant$, $>$ or $\geqslant$ symbol. Here are some examples of inequalities:

$$2 < 6, 5 \geqslant -1,\ 3x + 1 \leqslant 7 \text{ and } 2x + \frac{1}{3} > \frac{x}{4}$$

Inequalities can represent an infinite set of numbers. For example, the inequality $2x < 6$ means that $x < 3$ and this is the infinite set of all real numbers less than 3.



The altitude, h, that passenger planes can successfully and economically fly at can be represented by an inequality: $0 < h \leqslant 40\,000$ feet or $0 < h \leqslant 12$ km.

Lesson starter: Infinite solutions

Greg, Kevin and Greta think that they all have a correct solution to this inequality:

$$4x - 1 \geqslant x + 6$$

Greg says $x = 4$ is a solution.

Kevin says $x = 10$ is a solution.


Greta says $x = 100$ is a solution.

- Use substitution to show that they are all correct.
- Can you find the smallest whole number that is a solution to the inequality?
- Can you find the smallest number (including fractions) that satisfies the inequality? What method leads you to your answer?


Key ideas

- **Inequalities** can be illustrated using a number line because a line represents an infinite number of points.
  - Use an open circle when showing $>$ (greater than) or $<$ (less than).

For example: $x > 1$




$x < 1$




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3J


- Use a **closed circle** when showing $\geqslant$ (greater than or equal to) or $\leqslant$ (less than or equal to).
  For example: $x \geqslant 1$



$x \leqslant 1$



- A **set** of numbers may have both an upper and lower bound.
  For example: $-2 < x \leqslant 3$



- Linear inequalities can be solved in a similar way to linear equations.
  - All the numbers that satisfy an inequality are called a **solution set**.
  - If we multiply or divide both sides of an inequality by a negative number, the inequality symbol is reversed.
    For example: $5 < 8$ but $-5 > -8$, so if $-x > 1$ then $x < -1$.
  - If we swap the sides of an inequality, then the inequality symbol is reversed.
    For example: $3 < 7$ but $7 > 3$ so if $2 > x$ then $x < 2$.

Exercise 3J

Understanding

1, 2 | 1, 2

1 State the missing symbol < or > to make each statement true.

a $3 \square 2$ **b** $-1 \square 4$ **c** $-7 \square -3$ **d** $5 \square -50$


2 Match the inequalities **a**, **b** and **c** with the number lines **A**, **B** and **C**.

a $x < 2$


b $x \geqslant -1$

c $-3 \leqslant x < 0$


A



B



C



Fluency

3–7(½) | 3–8(½)

Example 27 Representing inequalities on a number line

Show each of the following examples on a number line.


a x is less than 6, $(x < 6)$.

b x is greater than or equal to 2, $(x \geqslant 2)$.


c x is greater than -2 but less than or equal to 3, $(-2 < x \leqslant 3)$.

Solution

a



b




Explanation

An open circle is used to indicate that 6 is not included.

A closed circle is used to indicate that 2 is included.

Essential Mathematics Core for the Victorian Curriculum Year 9 ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

c



An open circle is used to indicate that -2 is not included and a closed circle is used to indicate that 3 is included.

Now you try

Show each of the following examples on a number line.

a x is less than 4, $(x < 4)$.

b x is greater than or equal to -3, $(x \geqslant -3)$.

c x is greater than -3 but less than or equal to 1, $(-3 < x \leqslant 1)$.

3 Show each of the following inequalities on a number line.

a $x > 2$ **b** $x \leqslant 4$ **c** $x \geqslant -1$

d $-1 < x < 1$ **e** $0 \leqslant x \leqslant 3$ **f** $-3 < x \leqslant 4$

Example 28 Solving linear inequalities

Find the solution set for each of the following inequalities.

a $x - 3 < 7$ **b** $5 - 2x > 3$

Solution	Explanation
a $x - 3 < 7$	Add 3 to both sides.
$x < 10$	
b $5 - 2x > 3$	Subtract 5 from both sides.
$-2x > -2$	Divide both sides by -2, and reverse the inequality symbol.
$x < 1$	

Now you try

Find the solution set for each of the following inequalities.

a $x - 2 < 9$ **b** $4 - 3x > 7$

4 Find the solution set for each of the following inequalities.

a $x + 5 < 8$ **b** $b - 2 > 3$

c $y - 8 > -2$ **d** $5x \geqslant 15$

e $4t > -20$ **f** $y + 10 \geqslant 0$

g $3m - 7 < 11$ **h** $4a + 6 \geqslant 12$

i $7x - 5 < 2$ **j** $3x - 1 < 0$

5 Find the solution set for each of the following inequalities.

a $4 - 3x > -8$

b $2 - 4n \geqslant 6$

c $4 - 5x \leqslant 1$

d $7 - a \leqslant 3$

e $5 - x \leqslant 11$

f $7 - x \leqslant -3$

g $-2x - 3 > 9$

h $-4t + 2 \geqslant 10$

i $-6m - 14 < 15$



Hint for Q5: Remember that if you multiply or divide by a negative number, you flip the inequality symbol.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3J

Example 29 Solving more complex linear inequalities

Find the solution set for each of the following inequalities.

a $\frac{d}{4} - 3 \geqslant -11$ **b** $2a + 7 \leqslant 6a + 3$

Solution	Explanation
a $\frac{d}{4} - 3 \geqslant -11$	Add 3 to both sides.
$\frac{d}{4} \geqslant -8$	Multiply both sides by 4; the inequality symbol does not change.
$d \geqslant -32$	
b $2a + 7 \leqslant 6a + 3$	Gather pronumerals on one side by subtracting $2a$ from both sides.
$7 \leqslant 4a + 3$	
$4 \leqslant 4a$	Subtract 3 from both sides and then divide both sides by 4.
$1 \leqslant a$	Place the variable a on the left and reverse the inequality symbol.
$a \geqslant 1$	

Now you try

Find the solution set for each of the following inequalities.

a $\frac{a}{3} + 1 \geqslant -2$ **b** $3a - 1 \leqslant 5a + 6$

6 Find the solution set for each of the following inequalities.

a $\frac{x}{2} - 5 \leqslant 3$ **b** $3 - \frac{x}{9} \geqslant 4$ **c** $\frac{2x}{5} \leqslant 8$

d $\frac{2x + 6}{7} < 4$ **e** $\frac{3x - 4}{2} > -6$ **f** $\frac{1 - 7x}{5} \leqslant 3$

7 Solve these inequalities involving brackets.

a $4(x + 2) < 12$ **b** $-3(a + 5) > 9$

c $5(3 - x) \geqslant 25$ **d** $2(3 - x) > 1$

e $5(y + 2) < -6$ **f** $-7(1 - x) < -11$



Hint for Q7: You may wish to expand brackets first.



8 Find the solution set for each of the following inequalities.

a $2x + 9 \leqslant 6x - 1$ **b** $6t + 2 > t - 1$

c $7y + 4 \leqslant 7 - y$ **d** $3a - 2 < 4 - 2a$


e $1 - 3m \geqslant 7 - 4m$ **f** $7 - 5b > -4 - 3b$

Hint for Q8: Put all the pronumerals on one side and collect like terms.

Problem-solving and reasoning 9–11 10–12

9 Wendy is x years old and Jay is 6 years younger. The sum of their ages is less than 30. Write an inequality involving x and solve it. What can you say about Wendy's age?

10 The perimeter of a particular rectangle needs to be less than 50 cm. If the length of the rectangle is 12 cm and the width is w cm, write an inequality involving w and solve it. What width does the rectangle need to be?



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

11 Two car rental companies have the following payment plans:
Carz: \$90 per week and 15c per kilometre
Renta: \$110 per week and 10c per kilometre
What is the maximum whole number of kilometres that can be travelled in one week with Carz if it is to cost less than it would with Renta?



12 Consider the equation $9 - 2x > 3$.
- **a** Solve the equation by first adding $2x$ to both sides, then solve for x.
- **b** Solve the equation by first subtracting 9 from both sides.
- **c** What did you have to remember to do in part **b** to ensure that the answer is the same as in part **a**?

Many step inequalities — 13(½)

13 Combine all your knowledge from this chapter so far to solve these inequalities for x. For parts **g–i** a, b and c are positive numbers.

- **a** $\frac{2(x+1)}{3} > x + 5$
- **b** $2x + 3 \geqslant \frac{x-6}{3}$
- **c** $\frac{2-3x}{2} < 2x - 1$
- **d** $\frac{4(2x-1)}{3} \leqslant x + 3$
- **e** $1 - x > \frac{7(2-3x)}{4}$
- **f** $2(3 - 2x) \leqslant 4x$
- **g** $ax - b > -c$
- **h** $\frac{bx}{c} \leqslant a$
- **i** $a(x + b) < c$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3K Using formulas ★

Learning intentions

- To know which variable is the subject of a formula
- To be able to work with a formula by substituting known values
- To be able to find the unknown value in a formula by solving an equation
- To be able to transpose a formula to make a different variable the subject

Key vocabulary: subject, formula, variable, substitute, transpose, equation

A formula (or rule) is an equation that relates two or more variables. You can find the value of one of the variables if you are given the value of all other unknowns. The following are some examples of formulas.

- $A = \pi r^2$ is the formula for finding the area, A, of a circle given its radius, r.
- $V = lwh$ is the formula for finding the volume of a rectangular prism given its length, l, width, w, and height, h.
- $F = \frac{9}{5}C + 32$ is the formula for converting degrees Celsius, C, to degrees Fahrenheit, F.

A, V and F are said to be the subjects of the formulas given above.



Degrees Celsius and degrees Fahrenheit can be converted to one another using the formula that relates them.

Lesson starter: Common formulas

As a class group, try to list at least 10 formulas that you know.

- Write down the formulas.
- Try to describe what each variable in your formulas represents.
- Which variable is the subject of each formula?

Key ideas

- A **formula** is a rule for finding the value of one quantity given the value of the others.
- The **subject** of a formula is a variable that usually sits on its own on the left side.

  For example, the A in $A = lw$ is the subject of the formula.
- A variable in a formula can be evaluated by substituting numbers for all other variables.
- If the unknown is not the subject, a simple equation will need solving.
- **Transposing** is when a formula is rearranged to make another variable the subject. Steps are similar to those applied when solving an equation.

  $A = lw$ can be transposed (i.e. rearranged) to give $l = \frac{A}{w}$ with l now the subject.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Exercise 3K

Understanding

1–3 | 3

1 Write down 3 different formulas that you know for the area of shapes.

2 State the letter which is the subject of these formulas.

a $A = \frac{1}{2}bh$ **b** $D = b^2 - 4ac$

c $M = \frac{a+b}{2}$ **d** $A = \pi r^2$



Hint for Q2: In $V = lwh$, V is the subject.

3 Complete the step given in brackets to transpose the following formulas to make x the subject.

a $ax = b$ (divide both sides by a)

b $\frac{x}{c} = d$ (multiply both sides by c)

c $x - ab = d$ (add ab to both sides)

d $\sqrt{x} = c$ (square both sides)



Hint for Q3: Transpose means to make a different variable the subject.

Fluency

4, 5, 6(½) | 4–6(½)

Example 30 Substituting values into formulas

Substitute the given values into the formula to find the value of the subject.

a $A = \frac{1}{2}(a+b)h$, when $a = 3$, $b = 7$ and $h = 5$

b $E = \frac{1}{2}mv^2$, when $m = 4$ and $v = 5$

Solution

a $A = \frac{1}{2}(a+b)h$

$A = \frac{1}{2} \times (3+7) \times 5$

$= \frac{1}{2} \times 10 \times 5$

$= 25$

b $E = \frac{1}{2}mv^2$

$E = \frac{1}{2} \times 4 \times 5^2$

$= \frac{1}{2} \times 4 \times 25$

$= 50$

Explanation

Substitute $a = 3$, $b = 7$ and $h = 5$. Work out the sum in the brackets first, then multiply out to find the value of A.

Substitute $m = 4$ and $v = 5$.
$5^2 = 5 \times 5 = 25$, then work out E by multiplying.

Now you try

Substitute the given values into the formula to find the value of the subject.

a $M = \frac{a+b}{2}$, when $a = 5$ and $b = 9$

b $F = \frac{mn}{r^2}$, when $m = 12$, $n = 6$ and $r = 3$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3K

4 Substitute the given values into each of the following formulas to work out the subject. Round to two decimal places where appropriate.

a $A = bh$, when $b = 3$ and $h = 7$

b $F = ma$, when $m = 4$ and $a = 6$

c $m = \frac{a+b}{4}$, when $a = 14$ and $b = -6$

d $t = \frac{d}{v}$, when $d = 18$ and $v = 3$

e $A = \pi r^2$, when $r = 12$

f $V = \frac{4}{3}\pi r^3$, when $r = 2$

g $c = \sqrt{a^2 + b^2}$, when $a = 12$ and $b = 22$

h $Q = \sqrt{2gh}$, when $g = 9.8$ and $h = 11.4$

i $I = \frac{MR^2}{2}$, when $M = 12.2$ and $R = 6.4$

j $x = ut + \frac{1}{2}at^2$ when $u = 0$, $t = 4$ and $a = 10$

Example 31 Finding the unknown value in a formula

Substitute the given values into each formula, then solve to find the unknown value. Round to one decimal place in part **b**.

a $S = \frac{d}{t}$ given $S = 15$ and $d = 60$

b $C = 2\pi r$ given $C = 30$

Solution

a
$$S = \frac{d}{t}$$
$$15 = \frac{60}{t}$$
$$15t = 60$$
$$t = 4$$

b
$$C = 2\pi r$$
$$30 = 2\pi r$$
$$\frac{30}{2\pi} = r$$
$$r = 4.8 \text{ (to 1 d.p.)}$$

Explanation

Write the formula and substitute the given values of S and d.

Solve for t by first multiplying both sides by t.

Divide both sides by 15.

Substitute $C = 30$. Solve for r by dividing both sides by 2π in one step. $\frac{30}{2\pi} = 4.77464\ldots$

Note: On a calculator you will need to insert brackets, $30 \div (2\pi)$, or set up using the fraction template, $\frac{\square}{\square}$.

Now you try

Substitute the given values into each formula, then solve to find the unknown value. Round to one decimal place in part **b**.

a $R = \frac{pl}{A}$ given $R = 16$, $p = 6$ and $l = 8$

b $S = 2\pi rh$ given $S = 100$ and $h = 5$

5 Substitute the given values into each of the following formulas. Solve the equations to find the unknown value. Round to two decimal places where appropriate.

a $m = \frac{F}{a}$, when $m = 12$ and $a = 3$


b $A = lw$, when $A = 30$ and $l = 6$

c $v = u + at$, when $v = 20$, $u = 5$ and $a = 2$

d $p = \frac{m}{v}$, when $p = 2$ and $m = 50$

e $A = \pi r^2$, when $A = 48$ and $(r > 0)$

f $A = \frac{1}{2}(a + b)h$, when $A = 64$, $b = 12$ and $h = 4$



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Example 32 Transposing formulas

Transpose each of the following to make the variable in red the subject.

a $ax + by = c$

b $A = \pi r^2 \ (r > 0)$

Solution

a $ax + by = c$

$by = c - ax$

$y = \dfrac{c - ax}{b}$

Explanation

We need to obtain y on one side by itself. Follow steps as if you are solving the equation for y.

Subtract ax from both sides.

by is $b \times y$. Just as for $2y$ you would divide both sides by 2, divide both sides by b.

b $A = \pi r^2$

$\dfrac{A}{\pi} = r^2$

$r^2 = \dfrac{A}{\pi}$

$r = \sqrt{\dfrac{A}{\pi}}$

Divide both sides by π.

Make r^2 the subject.

Take the square root of both sides, since $\sqrt{r^2} = r$ if $r > 0$.

Now you try

Transpose each of the following to make the variable in red the subject.

a $v = u + at$

b $E = \dfrac{mv^2}{2}, \ v > 0$

6 Transpose each of the following formulas to make the pronumeral in red the subject.

a $C = \pi d$

b $a + bx = d$

c $p = m(x + n)$

d $I = \dfrac{Prt}{100}$

e $P = \dfrac{v^2}{R} \quad (v > 0)$

f $A = 2\pi rh$

g $V = \pi r^2 h \quad (r > 0)$

h $\sqrt{A} + B = 4C$

i $c^2 = a^2 + b^2 \quad (a > 0)$

j $\sqrt{b + c} = a$



Hint for Q6: Follow similar steps as if you were solving to find the letter in red, such as $d = \ldots$

Problem-solving and reasoning 7, 8 8–10

7 The formula $s = \dfrac{d}{t}$ gives the speed s km/h of a car which has travelled a distance of d km in t hours.

a Find the speed of a car which has travelled 400 km in 4.5 hours. Round to two decimal places.

b **i** Transpose the formula $s = \dfrac{d}{t}$ to make d the subject.

ii Find the distance covered if a car travels at 75 km/h for 3.8 hours.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3K

8 The velocity, v m/s, of an object is described by the rule $v = u + at$, where u is the initial velocity in m/s, a is the acceleration in m/s^2 and t is the time in seconds.

a Find the velocity after 3 seconds if the initial velocity is 5 m/s and the acceleration is 10 m/s^2.

b Find the time taken for a body to reach a velocity of 20 m/s if its acceleration is 4 m/s^2 and its initial velocity is 12 m/s.



Hint for Q8: Write down the value of the known variables first; i.e. $t = 3$, $u = 5$ etc.



Hint for Q8: Velocity is another word for speed.

9 The volume of water (V litres) in a tank is given by $V = 4000 - 0.1t$, where t is the time in seconds after a tap is turned on.

a Over time, does the water volume increase or decrease according to the formula?

b Find the volume after 2 minutes. (Note: t is in seconds.)

c Find the time it takes for the volume to reach 1500 litres. Round to the nearest minute.

10 The formula $F = \frac{9}{5}C + 32$ converts degrees Celsius, C, to degrees Fahrenheit, F.

a Find what each of the following temperatures is in degrees Fahrenheit.

i 100°C **ii** 38°C

b Calculate what each of the following temperatures is in degrees Celsius. Round to one decimal place where necessary.

i 14°F **ii** 98°F



Hint for Q10: $\frac{9}{5}C$ can also be written as $\frac{9C}{5}$.

Basketball formulas — 11

11 The formula $T = 3x + 2y + f$ can be used to calculate the total number of points made in a basketball game where:

x = number of three-point goals
y = number of two-point goals
f = number of free throws made
T = total number of points

a Find the total number of points for a game where 12 three-point goals, 15 two-point goals and 7 free throws were made.

b Find the number of three-point goals made if the total number of points was 36, with 5 two-point goals made and 5 free throws made.

The formula $V = \dfrac{\left(p + \frac{1.5r}{2} + 2a + \frac{1.5s}{2} + 2b\right) - (1.5t + 2f + m - o)}{g}$ can be used to calculate the value, V, of a basketball player where:

p = points earned
a = number of assists
b = number of blocks
f = number of personal fouls
o = number of offensive rebounds
r = number of rebounds
s = number of steals
t = number of turnovers
m = number of missed shots
g = number of games played

c Calculate the value of a player with 350 points earned, 2 rebounds, 14 assists, 25 steals, 32 blocks, 28 turnovers, 14 personal fouls, 24 missed shots, 32 offensive rebounds and 10 games.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Plumber

Plumbers use many skills in their day-to-day jobs. The training course is up to 5 years of on-the-job experience via an apprenticeship and TAFE courses.

Mathematics is an important skill for a plumber. They measure, cut, bend and connect pipes according to project specifications. Equations and formulas play an important role in the life of a plumber, especially if working on big construction sites as calculations need to be accurate for buildings to satisfy Australian regulations.



Maths@Work

1 Water expands when heated. Plumbers need to allow for this additional volume when selecting the most suitable water storage unit for a project. The volume E (litres) of water expansion is calculated with this formula:

$E = L \times (T - t) \times 0.000375$ where

L = initial volume of water in litres

T = highest temperate in °C

t = lowest temperature in °C

0.000375 is the coefficient of expansion for water.

Calculate the volume of water expansion in litres (to 2 d.p.) that would occur when:

a 600 L of water is heated from 22° to 50°

b 500 L of water is heated from 18° to 60°

c 1000 L of water is heated from 20° to 70°

d 800 L of water increased in temperature by 50°

2 Plumbers need to calculate water pressure, P, measured in kPa (kilopascals) for any given height, h (m), above an outlet. The following formula is used:

$P = h \times 9.81$ where 9.81 is the gravitational pressure exerted by 1 m height of water.


a Calculate the water pressure for a pressure head of:

i 10 m **ii** 20 m **iii** 30 m **iv** 40 m

b Rewrite the above formula with h as the subject and calculate how many metres (to 2 d.p.) of water are in a non-pressurised tank if the pressure gauge reads:

i 24.6 kPa **ii** 24.1 kPa **iii** 13.5 kPa **iv** 27.9 kPa

3 A plumber needs to install the pipes that connect a house to the underground mains water supply and to the council sewage disposal system. Water flows under pressure up to the house and the sewerage pipe is sloped downwards to allow waste to flow away under the force of gravity. The difference in level between each end of a sloping pipe is called the fall measurement.



Fall (mm) $= \dfrac{\text{pipe length in mm} \times x}{100}$, where the gradient $x\%$ = the fall in mm per 100 mm of pipe.


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Calculate the total fall in mm required to install the following pipes. Round to the nearest mm.

a A water pipe is 6 m long and installed at a 2.5% gradient.
b A water pipe is 17 m long and installed at a 2.1% gradient.
c A sewerage pipe is 9 m in length installed at a 1.67% gradient.
d A sewerage pipe is 18 m in length installed at a 1.85% gradient.

Using technology

4 The number of downpipes required for a house depends on the roof area and the size of the downpipes. Each size of downpipe can drain a maximum area of roof.
The number of downpipes needed can be estimated using the rain catchment area of a roof, which is the equivalent flat area of the roof, not the actual sloping area.





$$\text{Number of downpipes} = \frac{\text{total roof catchment area}}{\text{maximum area per downpipe}}$$

Set up an Excel spreadsheet as shown below. Use the given roof dimensions applied to the roof plan above and determine the number of downpipes required for each house.



Hint for Q4: The whole number of downpipes is manually determined by rounding up the decimal number of downpipes.

	A	B	C	D	E	F	G	H	I
1	**The number of downpipes for a house**								
2	House	Length m	Width m	x m	y m	Total roof catchment area in m^2	Maximum area in m^2 per downpipe	Decimal number of downpipes	Whole number of downpipes
3	**i**	14	8.5	4	3		47		
4	**ii**	15	9	3	4		45		
5	**iii**	17	10	3	3.5		46		
6	**iv**	22	11	3	5		47		

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Theme Parks

Theme parks and amusement parks all over the world offer a vast array of exhilarating and thrilling rides, such as Buzzsaw and The Giant Drop.

Hugo and Alice live near two theme parks called Extreme Thrill and Adventure World. They plan to run tourist bus trips to the theme parks and want to advertise the best deal at each theme park depending on the number of rides.

The theme parks charge different entry prices and cost for each ride.

Theme park	Entry cost	Fee per ride
Extreme Thrill	\$42	\$3
Adventure World	\$30	\$5

Present a report for the following tasks and ensure that you show clear mathematical workings, explanations and diagrams where appropriate.

1 Preliminary task

Tickets are bought at the entry for a certain number of rides.

a Find the cost of a ticket at Extreme Thrill for
- **i** 4 rides
- **ii** 10 rides

b Find the cost of a ticket at Adventure World for
- **i** 4 rides
- **ii** 10 rides

2 Modelling task

Formulate

The problem is to determine the cheapest theme park depending on the number of rides taken.

a Write down all the relevant information that will help solve this problem.

b Let T = total ticket price and n = number of rides.
Construct a formula for the total ticket price T for n rides at
- **i** Extreme Thrill
- **ii** Adventure World

Solve

c Hugo spends \$75 for a ticket into Extreme Thrill.
- **i** Write the formula for the ticket price T, for n rides at Extreme Thrill.
- **ii** Set up an equation by substituting $T = 75$
- **iii** Solve this equation algebraically for n.
- **iv** How many rides were included in Hugo's \$75 ticket?

d Alice spends \$75 for a ticket into Adventure World.
- **i** Write the formula for the ticket price T, for n rides at Adventure world.
- **ii** Set up an equation by substituting $T = 75$
- **iii** Solve this equation algebraically for n.
- **iv** How many rides were included in Alice's \$75 ticket?

e Compare your answers to **c** and **d** above, and state which theme park offers the best deal for a \$75 ticket. Give a reason for your answer.

f Hugo and Alice buy 2 identical tickets for Extreme Thrill costing a total of \$138.
- **i** Write the formula for 2 Extreme Thrill tickets and form an equation by substituting $T = 138$.
- **ii** Solve this equation for n and state how many rides they each paid for.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

g Hugo decides to calculate the number of rides that make the ticket prices equal at each theme park.

- **i** Set up the equation by equating the ticket price formulas.
  i.e. Extreme Thrill ticket formula = Adventure World ticket formula
- **ii** Solve this equation algebraically for n.
- **iii** For how many rides does each theme park have the same ticket price?
- **iv** Calculate this ticket price for each theme park.

Evaluate and verify

h For how many rides is Extreme Thrill the cheaper option? Justify your answer with a mathematical calculation.

i For how many rides is Adventure world the cheaper option? Justify your answer with a mathematical calculation.

Communicate

j Summarise your results using a table like the one below and describe any key findings.

Ticket prices							
Number of rides n							
Extreme Thrill							
Adventure world							

3 Extension questions

A new theme park called Monster Rides charges $22 entry plus $7 per ride.

a Construct a formula for the total ticket price T for n rides at Monster Rides.

b Set up equations by equating ticket price formulas and then solve each equation to find the number of rides n that make the Monster Rides ticket price equal to the:

- **i** Extreme Thrill ticket price
- **ii** Adventure World ticket price

c If, on average, people buy tickets for 7 or more rides, which theme park out of the three, would be expected to make the largest profit? Justify your answer with some mathematical calculations.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Protein balls

Key technology: Spreadsheets

Carrie runs a small business making and selling protein balls, which are a combination of muesli and chocolate. To make a batch of protein balls she combines chocolate and muesli so that the total weight of the batch is 10 kg. The cost of chocolate is $6 per kilogram and the cost of muesli is $4 per kilogram.



1 Getting started

When Carrie started her business, the total cost of chocolate and muesli for one batch was $48.

- Let x be the amount of chocolate purchased for one batch in kilograms.
- Let y be the amount of muesli purchased for one batch in kilograms.

a Given the total cost for one batch is $48 write down an equation including the variables x and y. We will call this Equation 1.

b Use your equation to find the amount of muesli purchased if the following amounts of chocolate are purchased.

i 2 kg
ii 3 kg
iii 5 kg

c Write down a second equation using the information given in the introduction regarding the total weight. We will call this Equation 2.

d Solve your equations 1 and 2 simultaneously to find the values of x and y and hence the amounts of chocolate and muesli purchased for each batch of protein balls.

2 Using technology

a Rearrange your Equation 1 and Equation 2 to make y the subject.

b We will now set up a spreadsheet to solve this pair of simultaneous equations numerically. Enter the shown details into a spreadsheet.

	A	B	C
1	**Making protein balls**		
2			
3	**Total cost of one batch**	48	
4			
5			
6	**x value**	**Equation 1**	**Equation 2**
7	0	=(B$3-6*A7)/4	=10-A7
8	=A7+1		

c Fill down from cells A8, B7 and C7.

d In your spreadsheet, search for a value of x for which the y values are equal for both equations. Describe what you have found.

e Change the total cost value in cell B3 to $50, then search for the solution to the simultaneous equations to find the amount of chocolate and muesli for one batch.

Essential Mathematics Core for the Victorian Curriculum Year 9 ISBN 978-1-009-59315-1 © Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Technology and computational thinking

3 Applying an algorithm

For selected values for the total cost in cell B3, you may have noticed that the values of x and y are integers. This will not be the case for all chosen values.

a Change the total cost value in cell B3 to $\$51$ then search for the value of x for which the y values are the same. What do you notice? Between what two integer x values will the y values for both equations be equal?

b Alter both the starting value of x and the increment in column A so that you can search for solutions between integer x values. Suggestion: Change the starting number in A7 to 4 and the formula in A8 to $= A7 + 0.5$. Fill down once again.

c Now alter the total cost in cell B3 to $\$47.60$ and apply this algorithm to find the solution.
- Step 1: Alter the total cost in cell B3 as chosen.
- Step 2: Look for the value of x for which the y values are closest.
- Step 3: Adjust the starting value of x to be nearer to the solution.
- Step 4: Adjust the increment in column A so that smaller increases in the value of x can be considered.
- Step 5: Repeat from Step 2 until the solution is found.

d Try the above algorithm for finding the solution for when the total cost is $\$51.50$.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

1 In a magic square, all the rows, columns and main diagonals add to the same total. The total for this magic square is 15. Find the value of x then complete the magic square.

$2x - 2$	$3x$	$x - 1$
x		

2 Simplify $2xy - 3x(y - 2x) - 8x^2y \div (2y) + 2x \times 6xy \div (4y)$.

3 Reveal the word below by solving each equation for x. The value of x gives the number of the letter in the alphabet; e.g. $x = 1$ would be A, $x = 2$ would be B etc.

$\frac{x-3}{4} = 5$	$6x = 4x + 30$	$2(x - 10) = 16$	$1 - 2x = -7$

4 Twelve years ago Eric's father was seven times as old as Eric was. If Eric's father is now 54 years old, how old is Eric now?

5 In a yacht race the second leg was half the length of the first leg. The third leg was two-thirds of the length of the second leg. The last leg was twice the length of the second leg. If the total distance was 153 km, find the length of each leg.



6 A group of office workers had some prize money to distribute among themselves. When all but one took \$9 each, the last person only received \$5. When they all took \$8 each, there was \$12 left over. How much had they won?

7 Try finding a solution to these more complex linear equations by first finding a common denominator.

a $\frac{x}{3} + \frac{x}{2} = 2$ **b** $\frac{x+1}{5} - \frac{x-1}{7} = 1$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter summary

Expressions and equations

Substitution
The process of replacing a pronumeral with a given value;
E.g. if $x = 3$, $2x + 4 = 2 \times (3) + 4$
$= 6 + 4$
$= 10$
E.g. if $x = 2$, $y = -4$,
then $xy = 2 \times (-4)$
$= -8$

Expanding brackets
E.g. $2(x + 3) = 2 \times x + 2 \times 3$
$= 2x + 6$
$-3(4 - 2x) = -3 \times 4 + (-3) \times (-2x)$
$= -12 + 6x$

Expressions
E.g. $2x - 3y + 5$
term, coefficient, constant
E.g. 2 more than 3 lots of m is $3m + 2$
3 less than x is $x - 3$


Addition/subtraction
Only like terms can be combined under addition or subtraction.
Like terms have the same pronumerals and powers; i.e. $4xy$ and $2xy$.
E.g. $5a + 2 + 3a$
$= 5a + 3a + 2$
$= 8a + 2$
E.g. $3x + 2y - x + 4y$
$= 3x - x + 2y + 4y$
$= 2x + 6y$

Solving word problems
1 Define a pronumeral for the unknown.
2 Set up an equation.
3 Solve the equation.
4 Check the answer and write in words.

Solving linear equations
The solution of a linear equation is the value that makes the equation true.
E.g. $2x + 5 = 9$
$2x = 4$ (subtract 5)
$x = 2$ (divide by 2)
E.g. $5(x - 1) = 2x + 7$
$5x - 5 = 2x + 7$ (expand)
$3x - 5 = 7$ (subtract $2x$)
$3x = 12$ (add 5)
$x = 4$ (divide by 3)

Multiplication/ Division
In simplified form the multiplication sign ($\times$) and division sign ($\div$) are not shown.
That is, $2 \times x = 2x$ and $x \div 2 = \frac{x}{2}$.
Multiplying:
$2a \times 3b = 2 \times 3 \times a \times b$
$= 6ab$
Dividing:
$4x \div 2 = \frac{4x}{2} = 2x$
$\frac{6ab^2}{9b} = \frac{2 \times 3 \times a \times b \times b}{3 \times 3 \times b} = \frac{2ab}{3}$

Inequalities
These can be represented using $>$, $<$, $\geqslant$, $\leqslant$ rather than $=$.
e.g. $x > 2$ 2 not included (open circle)



Solving inequalities uses the same steps as solving equations except when multiplying or dividing by a negative number. In this case, the inequality symbol must be reversed.
e.g. $4 - 2x < 10$ (-4)
$-2x < 6$ $(\div -2)$
$x > -3$ (reverse symbol)

Equations involving fractions
E.g. $\frac{x}{3} + 1 = 5$
$\frac{x}{3} = 4$ (subtract 1)
$x = 12$ (multiply by 3)
E.g. $\frac{x-2}{4} = 1$
$x - 2 = 4$ (multiply by 4)
$x = 6$ (add 2)
E.g. $\frac{5x}{2} = 10$
$5x = 20$ (multiply both sides by 2)
$x = 4$ (divide both sides by 5)

Formulas
Some common formulas are
$A = lw$, $C = 2\pi r$
An unknown value can be found by substituting values for the other variables.
A formula can be transposed to make a different variable the subject.
E.g. make r the subject in $A = \pi r^2$
$A = \pi r^2$
$\frac{A}{\pi} = r^2$
$\sqrt{\frac{A}{\pi}} = r$
$\therefore\ r = \sqrt{\frac{A}{\pi}}$ where $r > 0$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter checklist

A version of this checklist that you can print out and complete can be downloaded from your Interactive Textbook.

		✔
3A	**1 I can identify parts of an algebraic expression.** e.g. Consider the expression $3x - 5y + 6$. State: **a** the number of terms **b** the coefficient of y **c** the constant term	
3A	**2 I can convert words and word problems to expressions.** e.g. Write an algebraic expression for **a** the cost of n movie tickets at \$15 each **b** four less than three times x **c** the square of y is divided by 4	
3A	**3 I can substitute values into expressions and evaluate.** e.g. Evaluate these expressions if $p = -4$, $q = 6$ and $r = 5$: **a** $4q - r$ **b** $3p - (q + r)$	
3B	**4 I can identify like terms.** e.g. Choose the pair(s) of like terms from the following set: $4x, 5, 3xy, -5x, 2xy^2, yx$	
3B	**5 I can collect like terms.** e.g. Simplify the following by collecting like terms: **a** $5a - 4 + 3a$ **b** $4xy + 6y - xy + 4y$	
3C	**6 I can multiply algebraic terms.** e.g. Simplify the following: **a** $4s \times 5t$ **b** $-3y \times 8xy$	
3C	**7 I can divide algebraic terms.** e.g. Simplify the following: **a** $\frac{15x}{12}$ **b** $16a^2b \div (24ab)$	
3D	**8 I can expand expressions with brackets.** e.g. Expand the following and simplify: **a** $4(x + 5)$ **b** $-3(x - 6)$ **c** $2x(3x - y)$	
3D	**9 I can expand and collect like terms.** e.g. Expand the following and collect like terms: **a** $3(x + 2) - 4$ **b** $5 - 2(x + 1)$	
3E	**10 I can solve simple linear equations.** e.g. Solve the following equations: **a** $2x - 5 = 7$ **b** $8 - 3x = 15$	
3F	**11 I can solve linear equations with fractional coefficients.** e.g. Solve each of the following equations **a** $\frac{3x}{4} = 9$ **b** $\frac{x}{5} - 4 = 2$	

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter checklist

		✔
3F	**12 I can solve simple fractional equations.** e.g. Solve the equation $\frac{x-5}{3}=6$	
3G	**13 I can solve equations with brackets.** e.g. Solve $3(3x+5)=22$	
3G	**14 I can solve equations with brackets and a common factor.** e.g. Solve $4(2x+3)=32$ by first dividing by the common factor	
3G	**15 I can solve equations with brackets and like terms.** e.g. Solve $2(4x+7)-5x=20$	
3H	**16 I can solve equations with pronumerals on both sides.** e.g. Solve these equations: **a** $8x=3x+25$ **b** $10-3x=x+2$	
3H	**17 I can solve equations with brackets and pronumerals on both sides.** e.g. Solve $4(2x-1)=2(x-8)$	
3I	**18 I can turn a word problem into an equation.** e.g. 8 more than two times a certain number is 30. Write an equation and solve it to find the number.	
3I	**19 I can apply algebra to a word problem.** e.g. A surf shop hires out surfboards. It charges an initial fee of \$15 for hiring a board and then a charge of \$12 per hour. Jett returns his board and is charged \$51. For how many hours did he hire the board?	
3I	**20 I can solve more complex word problems.** e.g. Toni and Trey earn \$370 between them walking dogs. If Toni earned \$140 more than Trey, how much money did each of them make?	
3J	**21 I can represent an inequality on a number line.** e.g. Show x is less than or equal to 3 ($x \leqslant 3$) on a number line.	
3J	**22 I can solve a linear inequality.** e.g. Find the solution set for $8-3x<20$.	
3K	**23 I can substitute values into formulas and find the unknown.** e.g. Substitute the given values into the formula to find the value of the unknown. Round to one decimal place where necessary. **a** $A=\frac{1}{2}xy$ when $x=7$ and $y=10$ **b** $V=\pi r^2h$ when $V=120$ and $r=3$	
3K	**24 I can transpose a formula.** e.g. Transpose the formula $v^2=u^2+2as$ to make u the subject ($u>0$).	

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Short-answer questions

3A **1** Write algebraic expressions to represent the following.
- **a** The sum of x and y
- **b** The product of m and 7
- **c** The cost of 3 movie tickets at m dollars each
- **d** 3 less than n, all divided by 4

3A **2** Evaluate the following if $x = 2$, $y = -1$ and $z = 5$.

a $xz + 1$ **b** $4x + y$ **c** $x - 2yz$ **d** $x(y + z)$ **e** $x^2 - 3z$

3C **3** Simplify.

a $2 \times 4n$ **b** $3x \times 2y$ **c** $8a \div 2$ **d** $\frac{4x^2y}{12x}$

3B **4** Simplify by collecting like terms.

a $2b + 4b + b$ **b** $6x + 3 - 2x$
c $4p - 3q - p + 5q$ **d** $3mn + 2m - 6mn + n$

3D **5** Expand and simplify the following.

a $2(x + 7)$ **b** $-3(2x + 5)$ **c** $2x(3x - 4)$
d $-2a(5 - 4a)$ **e** $4(x + 2) + 5$ **f** $3(x - 2) - 1$

3E **6** Solve the following linear equations for x.

a $5x + 6 = 51$ **b** $7x - 4 = 10$ **c** $4 - x = 7$ **d** $3 - 2x = 21$

3F **7** Solve these linear equations involving fractions.

a $\frac{2x}{5} = 4$ **b** $\frac{x}{2} - 1 = 3$ **c** $\frac{x + 2}{4} = 7$ **d** $\frac{1}{3}x + 2 = 4$

3I **8** Write an equation to represent each of the following and then solve it for the pronumeral.
- **a** A number, n, is doubled and increased by 3 to give 21.
- **b** A number of lollies, l, is decreased by 5 and then shared equally among 3 friends so that they each get 7.
- **c** 5 less than the result of Toni's age, x, divided by 4 is 0.

3G **9** Solve the following linear equations by first expanding the brackets.

a $2(x + 4) = 18$ **b** $3(2x - 3) = 2$

3H **10** Solve these equations with variables on both sides.
- **a** $8x = 2x + 24$
- **b** $5x - 2 = 3x + 2$
- **c** $3 - 4x = 7x - 8$
- **d** $5(2x + 4) = 7x + 5$

3I **11** Nick makes an initial bid of $\$x$ in an auction for an old cricket bat. By the end of the auction he has paid $\$550$, which is $\$30$ more than twice his initial bid, x. Set up and solve an equation to determine Nick's initial bid.

3J **12** Represent each of the following on a number line.

a $x > 1$ **b** $x \leqslant 7$ **c** $x \geqslant -4$
d $x < -2$ **e** $2 < x \leqslant 8$ **f** $-1 < x < 3$

3J **13** Solve the following inequalities.

a $x + 8 < 20$ **b** $2m - 4 > -6$ **c** $3 - 2y \leqslant 15$
d $\frac{7 - 2x}{3} > -9$ **e** $3a + 9 < 7(a - 1)$ **f** $-4x + 2 \leqslant 5x - 6$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter review

3K **14** Find the value of the unknown in each of the following formulas.

a $E = \sqrt{PR}$ when $P = 90$ and $R = 40$

b $v = u + at$ when $v = 20$, $u = 10$, $t = 2$

c $V = \frac{1}{3}Ah$ when $V = 20$, $A = 6$

3K **15** Rearrange the following formulas to make the variable in brackets the subject.

a $v^2 = u^2 + 2ax \quad (x)$

b $P = RI^2,\ I > 0 \quad (I)$

Multiple-choice questions

3A **1** The algebraic expression that represents 2 less than 3 lots of n is:

A $3(n-2)$ **B** $2-3n$ **C** $3n-2$ **D** $3+n-2$ **E** n

3C **2** The fully simplified form of $2ab \div (8a)$ is:

A $\frac{2ab}{8a}$ **B** $\frac{4}{b}$ **C** $\frac{b}{4}$ **D** $4b$ **E** $4a$

3B **3** The simplified form of $6ab + 14a - 2ab + 3a$ is:

A $21ab$ **B** $4ab + 11a$ **C** $ab + 7a$ **D** $4ab + 17a$ **E** $4 + 17a$

3E **4** The solution to the equation $2x - 3 = 7$ is:

A $x = 2$ **B** $x = -5$ **C** $x = 5$ **D** $x = 6.5$ **E** $x = 0.5$

3F **5** The solution to $\frac{x}{3} - 1 = 4$ is:

A $x = 13$ **B** $x = 7$ **C** $x = 9$ **D** $x = 15$ **E** $x = \frac{5}{3}$

3D **6** The expanded form of $3(2y - 1)$ is:

A $6y - 3$ **B** $5y - 1$ **C** $6y - 1$ **D** $6y + 2$ **E** $3y - 3$

3H **7** An equivalent equation to $5x = 2x + 9$ is:

A $2x = 5$ **B** $3x = 9$ **C** $0 = 3x + 9$ **D** $7x = 9$ **E** $x = 9$


3G **8** The solution to the equation $2(x - 3) = 7$ is:

A $x = 5$ **B** $x = \frac{13}{2}$ **C** $x = 2$ **D** $x = \frac{1}{2}$ **E** $x = \frac{7}{3}$

3I **9** Eli is x years old. His sister is 2 years older. The sum of their ages is 22. A simplified equation to represent this is:

A $x + 2 = 22$ **B** $2x + 2 = 22$ **C** $2(x + 2) = 22$ **D** $x(x + 2) = 22$ **E** $2x = 22$

3J **10** The inequality representing the x-values on the number line below is:



A $x < -1$ **B** $x > -1$ **C** $x \leqslant -1$ **D** $x \geqslant -1$ **E** $-1 < x < 3$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3J **11** The solution to the inequality $1 - 2x > 9$ is:

A $x < -4$ **B** $x < 4$ **C** $x < -5$ **D** $x > -4$ **E** $x > 5$

3K **12** If $A = \frac{1}{2}bh$ with $A = 360$ and $h = 40$, then the value of b is:

A 90 **B** 4.5 **C** 20 **D** 18 **E** 9

Extended-response questions


1 Julie hires a jumping castle for her daughter's birthday party. It costs \$60 for the set-up, plus \$25 for each hour that it is hired.



a What is the cost for:
- **i** 1 hour of hire?
- **ii** 2 hours of hire?

b Julie is charged \$210 for the hire of the jumping castle.
- **i** Define a variable to represent the number of hours for which Julie hired the castle.
- **ii** Set up an equation using your variable.
- **iii** Solve the equation to find how many hours Julie hired the jumping castle for.

2 A new backyard deck is being designed in the shape of the trapezium shown.



The area of a trapezium is given by $A = \frac{1}{2}(a + b)h$.

Currently the dimensions are set such that $a = 12$ m and $b = 8$ m.

a Substitute the given values and rearrange the equation to make h the subject.

b Use your answer to part **a** to find the width of the deck (h m) required to have a deck area of 100 m^2.

c It is decided to extend the width (h) to 12 m to increase the area to 150 m^2. If b is fixed at 8 m, find the value of a that gives the required area.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

4

Pythagoras' theorem and trigonometry



Essential mathematics: why understanding Pythagoras' theorem and trigonometry is important

Pythagoras' theorem and trigonometry are essential tools used in various fields to solve problems involving distances, angles, and heights

Surveyors use trigonometry extensively to ensure accurate measurements and effective design in their work. Using trigonometric ratios like sine, cosine, and tangent, they calculate the steepness of slopes, which is crucial for assessing terrain. To ensure water does not pool on roads, they design the road's slope using trigonometry. By measuring the angle of the slope and the horizontal distance, they determine the vertical height difference, allowing for proper drainage system design.

Javelin **athletes** and **coaches** use trigonometry to calculate the ideal angle that balances both horizontal and vertical components of the throw to maximise distance. Athletes like Jan Železný and Thomas Röhler have used these principles to achieve record-breaking throws. Trigonometry helps them fine-tune their techniques, ensuring they hit the right angles and achieve the best results.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025
Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.



In this chapter

4A Exploring Pythagoras' theorem
4B Finding the length of the hypotenuse
4C Finding the lengths of the shorter sides
4D Applying Pythagoras' theorem ★
4E Trigonometric ratios
4F Finding side lengths
4G Solving for the denominator ★
4H Finding an angle
4I Applying trigonometry ★

Victorian Curriculum 2.0

This chapter covers the following content descriptors in the Victorian Curriculum 2.0:

NUMBER

VC2M9N01

MEASUREMENT

VC2M9M03

SPACE

VC2M9SP01, VC2M9SP03

Please refer to the curriculum support documentation in the teacher resources for a full and comprehensive mapping of this chapter to the related curriculum content descriptors.

© VCAA

Online resources


A host of additional online resources are included as part of your Interactive Textbook, including HOTmaths content, video demonstrations of all worked examples, auto-marked quizzes and much more.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


Warm-up quiz

1 What is the length of the longest side given on these triangles?


a



b



c



2 Calculate each of the following.

a 3^2 **b** 7^2 **c** $2^2 + 4^2$ **d** $9^2 - 6^2$

3 Round off each number correct to four decimal places.

a 0.45678 **b** 0.34569 **c** 0.04562 **d** 0.27997

4 Round off each number correct to two decimal places.


a 4.234 **b** 5.678 **c** 76.895 **d** 23.899

5 Use a calculator to calculate each of the following correct to two decimal places.


a $\sqrt{35}$ **b** $\sqrt{236}$ **c** $\sqrt{23.6}$ **d** $\sqrt{65.4}$

6 Find the value of x in these diagrams.


a




b




c




d



e



f



7 Solve each of the following equations to find the value of x. Multiply or divide both sides to start.

a $3x = 6$ **b** $4x = 12$ **c** $5x = 60$ **d** $8x = 48$

e $\frac{x}{3} = 4$ **f** $\frac{x}{5} = 6$ **g** $\frac{x}{7} = 4$ **h** $\frac{x}{13} = 14$

8 Solve each of the following equations to find the value of x correct to one decimal place. First multiply both sides by x, then divide to finish.

a $\frac{3}{x} = 5$ **b** $\frac{4}{x} = 7$ **c** $\frac{32}{x} = 15$ **d** $\frac{14}{x} = 27$

9 Given that x is a positive number, find its value in each of the following equations. Note, for example, that if $x^2 = 9$ then $x = \sqrt{9} = 3$ (provided that $x > 0$).

a $x^2 = 4$ **b** $x^2 = 16$ **c** $x^2 = 3^2 + 4^2$ **d** $x^2 = 12^2 + 5^2$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

4A Exploring Pythagoras' theorem


Learning intentions

- To know which side of a right-angled triangle represents the hypotenuse
- To know that Pythagoras' theorem relates to right-angled triangles
- To know the relationship between the square of the sides of a right-angled triangle (Pythagoras' theorem)
- To be able to write Pythagoras' theorem for a triangle using variables or numbers

Key vocabulary: hypotenuse, Pythagoras' theorem, right-angled, square

The philosopher Pythagoras was born in Greece in the 6th century BCE. He travelled to Egypt and Persia where he developed his ideas in mathematics and philosophy. His students and followers were called the Pythagoreans. They made many advances in mathematics.


The Pythagoreans discovered the famous theorem, which is named after Pythagoras, and the existence of irrational numbers such as $\sqrt{2}$, which cannot be written down as a fraction or terminating decimal. The Pythagoreans called these numbers 'unutterable' numbers because any member who mentioned these numbers in public might be put to death.



A woodcut engraving of Pythagoras


Lesson starter: Discovering Pythagoras' theorem

Use a ruler to measure the sides of these right-angled triangles to the nearest mm. Then complete the table.



Triangle	a	b	c	a^2	b^2	c^2
1						
2						
3						

- Can you see any relationship between the numbers in the columns for a^2 and b^2 and the number in the column for c^2?
- Can you write down this relationship as an equation?
- Explain how you might use this relationship to calculate the value of c if it was unknown.
- Research how you can cut the two smaller squares (a^2 and b^2) to fit the pieces into the largest square (c^2).




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

4A

Key ideas

- The **hypotenuse**:
  - is the longest side of a **right-angled** (90°) triangle.
  - is opposite the right angle.
- **Pythagoras' theorem** states:
  - the square of the hypotenuse is equal to the sum of the squares of the other two shorter sides.
    $a^2 + b^2 = c^2$ or $c^2 = a^2 + b^2$



Exercise 4A


Understanding

1–3 | 3


1 Write the missing words in this sentence.
The _____ is the longest side of a right-angled _____.

2 Which letter marks the length of the hypotenuse in these triangles?


a



b



c





Hint for Q1 and Q2: The hypotenuse is the longest side of a right-angled triangle.



3 Decide if these equations are true or false.

a $2^2 + 3^2 = 4^2$ **b** $6^2 + 8^2 = 10^2$

c $5^2 - 3^2 = 4^2$ **d** $10^2 - 5^2 = 5^2$


Hint for Q3: Does LHS = RHS?

Fluency

4, 5 | 4, 5

Example 1 Writing Pythagoras' theorem

Write down Pythagoras' theorem for this triangle.



Solution


$p^2 + q^2 = r^2$

Explanation

r is the length of the hypotenuse and p and q are the lengths of the two shorter sides.
Alternatively, $r^2 = p^2 + q^2$.

Now you try


Write down Pythagoras' theorem for this triangle.




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

4 Write down Pythagoras' theorem for these triangles using the given pronumerals.


a



b




c






Hint for Q4: Replace the letters a, b and c in $a^2 + b^2 = c^2$ with the given letters.


d



e




f



Example 2 Substituting with Pythagoras' theorem

Substitute the numbers on this triangle into Pythagoras' theorem $a^2 + b^2 = c^2$.



Solution


$6^2 + 8^2 = 10^2$

Explanation

10 is the length of the hypotenuse and 6 and 8 are the lengths of the two shorter sides.


Now you try

Substitute the numbers on this triangle into Pythagoras' theorem $a^2 + b^2 = c^2$.




5 Substitute the numbers on these triangles into $a^2 + b^2 = c^2$.


a




b




c




d



e



f




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

4A


Problem-solving and reasoning 6–8 7–10

6 Write down Pythagoras' theorem using the pronumerals given in these diagrams.


a



b



c



7 Complete this table and answer the questions below.

a	b	c	a^2	b^2	$a^2 + b^2$	c^2
3	4	5				
6	8	10				
8	15	17				

a Which two columns give equal results?

b What would be the value of c^2 if $a^2 + b^2 = c^2$ and:

i $a^2 = 4$ and $b^2 = 9$? **ii** $a^2 = 7$ and $b^2 = 13$?

c What would be the value of $a^2 + b^2$ if $a^2 + b^2 = c^2$ and:

i $c^2 = 25$? **ii** $c^2 = 110$?


8 A cable connects the top of a 30 m mast to a point on the ground. The cable is 40 m long and connects to a point 20 m from the base of the mast.

a Using $c = 40$, decide if $a^2 + b^2 = c^2$.


b Do you think the triangle formed by the mast and the cable is right angled? Give a reason.

9 If $a^2 + b^2 = c^2$, we know that the triangle must have a right angle. Which of these triangles must have a right angle?


a




b




c




d



e



f





Hint for Q9: Substitute into $a^2 + b^2 = c^2$ to see if it's true.

10 (3, 4, 5) and (5, 12, 13) are Pythagorean triples, since $3^2 + 4^2 = 5^2$ and $5^2 + 12^2 = 13^2$. Find 10 more Pythagorean triples using whole numbers less than 100.



Hint for Q10: If (3, 4, 5) is a Pythagorean triple, then (6, 8, 10) is also a triple.

Pythagorean triples — 11

11 Find the total number of Pythagorean triples with whole numbers of less than 100. Include any found from Question **10** above.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


4B Finding the length of the hypotenuse

Learning intentions

- To be able to use Pythagoras' theorem to find the length of the hypotenuse given the other two sides
- To be able to find hypotenuse lengths both as rounded decimals or exact values including surds

Key vocabulary: hypotenuse, Pythagoras' theorem, right-angled, substitute, surds, square root


From our knowledge of formulas and equations we know that the value of one variable can be found if the values of all the other variables are known. With the help of Pythagoras' theorem, this means that an unknown side length of a right-angled triangle can be found if the other two sides are known. In this section we will find the length of the hypotenuse given the two shorter sides.



A geometric visualisation of Pythagoras' theorem.

Lesson starter: Correct substitution

Here are three attempts at using Pythagoras' theorem for the given triangle.



$c^2 = a^2 + b^2$
$= 3^2 + 4^2$
$= 25$
$\therefore c = \sqrt{25}$
$= 5$

$c^2 = a^2 + b^2$
$= 4^2 + 3^2$
$= 25$
$\therefore c = \sqrt{25}$
$= 5$


$c^2 = a^2 + b^2$
$= (3 + 4)^2$
$= 49$
$\therefore c = 7$

- Which set of working is incorrect? Give reasons.
- Why are the other two sets of working correct? Are they identical?

Key ideas

- To use Pythagoras' theorem to find the length of the hypotenuse:
  - Use the shorter lengths to substitute for a and b. This can be done in any order.
  - Find the value of $a^2 + b^2$.
  - Take the square root to find the value of c. Round if required.

- Lengths can be expressed with exact values using **surds**. $\sqrt{2}$, $\sqrt{28}$ and $2\sqrt{3}$ are examples of surds.
  - When expressed as a decimal, a surd is an infinite non-recurring decimal with no pattern; for example, $\sqrt{2} = 1.4142135623...$



$c^2 = a^2 + b^2$
$= 4^2 + 6^2$
$= 16 + 36$
$= 52$
$\therefore c^2 = \sqrt{52}$
$= 7.21$ (to 2 d.p.)

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


Exercise 4B

Understanding


1–4 | 3, 4

1 State the length of the hypotenuse (c) in these right-angled triangles.


a



b




c




2 Write down Pythagoras' theorem using the given pronumerals and values for these right-angled triangles.

a



b





Hint for Q2: Write a statement such as $c^2 = 2^2 + 3^2$.

3 Identify the surds from the following list.

a $\sqrt{3}$ **b** 7 **c** 6.4 **d** $\sqrt{25}$ **e** $0.\dot{3}$ **f** $\sqrt{7}$

4 Find the value of c ($c > 0$) in the following using the square root. Round to one decimal place in parts **a** and **b** and answer as a surd in parts **c** and **d**.


a $c^2 = 45$ **b** $c^2 = 10 + 30$ **c** $c^2 = 22$ **d** $c^2 = 3^2 + 5^2$

Fluency

5(½), 6 | 5–7(½)

Example 3 Finding the length of the hypotenuse

Find the length of the hypotenuse in this right-angled triangle.



Solution

$c^2 = a^2 + b^2$

$= 5^2 + 12^2$

$= 169$

$\therefore c = \sqrt{169}$

$= 13$


Explanation

Write the rule and substitute the lengths of the two shorter sides. Any order is okay.

If $c^2 = 169$ then $c = \sqrt{169} = 13$.

Now you try


Find the length of the hypotenuse in this right-angled triangle.




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5 Find the length of the hypotenuse in each of the following right-angled triangles.


a




b




c




d




e




f




g



h



i






Hint for Q5: Substitute the two shorter sides for a and b in the equation $c^2 = a^2 + b^2$.

Example 4 Using rounding with Pythagoras' theorem

Find the length of the hypotenuse in this right-angled triangle, rounding to two decimal places.



Solution

$c^2 = a^2 + b^2$

$= 7^2 + 9.5^2$

$= 139.25$

$\therefore c = \sqrt{139.25}$

$= 11.80$ (to 2 d.p.)


Explanation

The order for a and b does not matter since $7^2 + 9.5^2 = 9.5^2 + 7^2$.

$\sqrt{139.25} = 11.8004\ldots$ and the third decimal place is zero, so round down.


Now you try

Find the length of the hypotenuse in this right-angled triangle, rounding to two decimal places.




6 Find the length of the hypotenuse in each of these right-angled triangles, correct to two decimal places.


a




b




c




d



e



f






Hint for Q6: Use a calculator to find the value of c and round your answer to two decimal places.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

4B

Example 5 Finding the length of the hypotenuse using exact values

Find the length of the hypotenuse in this right-angled triangle, leaving your answer as an exact value.



Solution

$c^2 = a^2 + b^2$

$= 5^2 + 2^2$

$= 29$

$\therefore\ c = \sqrt{29}$


Explanation

Apply Pythagoras' theorem to find the value of c.

Express the answer exactly using a surd.


Now you try

Find the length of the hypotenuse in this right-angled triangle, leaving your answer as an exact value.




7 Find the length of the hypotenuse in these triangles, leaving your answer as an exact value.


a




b




c




d



e



f






Hint for Q7: Leave your answer as a surd; e.g. $\sqrt{3}$ or $\sqrt{26}$.


Problem-solving and reasoning

8–10, 11(½) | 10, 11(½), 12–14


8 Find the length of the diagonal steel brace required to support a wall of length 3.5 m and height 2.6 m. Give your answer correct to one decimal place.



9 A miniature rocket blasts off at an angle of 45° and, after a few seconds, reaches a height of 350 m above the ground. At this point it has also covered a horizontal distance of 350 m. How far has the rocket travelled to the nearest metre?




10 Find the length of the longest rod, as shown, that will fit inside a cylinder of height 2.1 m and with circular end surface of 1.2 m diameter. Give your answer correct to one decimal place.




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

11 For each of these triangles, first calculate the length of the hypotenuse then find the perimeter, correct to two decimal places.


a




b




c




d



e



f






12 A helicopter hovers at a height of 150 m above the ground and is a horizontal distance of 200 m from a beacon on the ground. Find the direct distance of the helicopter from the beacon.

Hint for Q12: First draw a right-angled triangle.



13 Find the perimeter of this triangle.






Hint for Q13: You will need to find AB and BC first by considering two different triangles.

14 One way to check whether a four-sided figure is a rectangle is to make sure that both its diagonals are the same length. What should the length of the diagonals be if a rectangle has side lengths 3 m and 5 m? Answer to two decimal places.

The dog kennel — 15

15 A dog kennel has the dimensions shown in the diagram on the right. Give your answers to each of the following, correct to two decimal places.

- **a** What is the width of the kennel?
- **b** What is the total height, h m, of the kennel?
- **c** If the sloping height of the roof was to be reduced from 55 cm to 50 cm, what difference would this make to the total height of the kennel? (Assume that the width is the same as in part **a**.)



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

4C Finding the lengths of the shorter sides


Learning intentions

- To be able to set up Pythagoras' theorem where the unknown is a shorter side
- To be able to solve the resulting equation from Pythagoras' theorem to find the length of a shorter side

Key vocabulary: Pythagoras' theorem, hypotenuse, equation, square root, surd

We know that the sum $7 = 3 + 4$ can be written as a difference: $3 = 7 - 4$ or $4 = 7 - 3$. Likewise, if $c^2 = a^2 + b^2$ then $a^2 = c^2 - b^2$ or $b^2 = c^2 - a^2$.


Applying this idea to a right-angled triangle means that we can now find the length of one of the shorter sides if the other two sides are known.



The height of a roof can be found using Pythagoras' theorem.

Lesson starter: True or false

Below are some mathematical statements relating to a right-angled triangle with hypotenuse 10 and the two shorter sides 6 and b.



Some of these mathematical statements are true and some are false. Can you sort them into true and false groups?


$6^2 + b^2 = 10^2$ $\quad 6 = \sqrt{10^2 - b^2}$ $\quad 10^2 - 6^2 = b^2$ $\quad 6^2 - 10^2 = b^2$

$10 = \sqrt{6^2 + b^2}$ $\quad b = \sqrt{6^2 - 10^2}$ $\quad 10 = \sqrt{6^2 - b^2}$ $\quad 10^2 - b^2 = 6^2$

Key ideas

- When finding the length of a shorter side:
  - Substitute known values into Pythagoras' theorem.
  - Solve this equation to find the unknown value.

  For example:



$c^2 = a^2 + b^2$	Write out Pythagoras' theorem.
$25^2 = a^2 + 15^2$	Substitute in the known values.
$625 = a^2 + 225$	Subtract 225 from both sides.
$400 = a^2$	To find a, take the square root of both sides.
$\sqrt{400} = a$	
$20 = a$	
or $a = 20$	

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Exercise 4C

Understanding

1–3 | 3

1 Write the missing word or number.

a To solve for a in $a^2 + 9 = 25$, the first step would be to _____ 9 from both sides.

b To solve for b in $16 + b^2 = 49$, the first step would be to subtract _____ from both sides.

c If $a^2 + 25 = 36$, then $a^2 =$ ____.

2 If $a^2 + 64 = 100$, decide if the following are true or false.

a $a^2 = 100 - 64$ **b** $64 = 100 + a^2$ **c** $a = 6$ **d** $a = 10$

3 Find the value of b in these equations. (b is a positive number.)

a $b^2 + 9 = 25$ **b** $b^2 + 49 = 625$ **c** $36 + b^2 = 100$




Hint for Q3: Solve for b^2 first.

Fluency

4–5(½) | 4-5(½)

Example 6 Finding the length of a shorter side

Find the value of the pronumeral.



Solution

$a^2 + 15^2 = 17^2$

$a^2 + 225 = 289$

$a^2 = 64$

$\therefore a = \sqrt{64}$

$a = 8$

Explanation

Write the rule and substitute the known sides.


Square 15 and 17.

Subtract 225 from both sides.

Take the square root of both sides.


Now you try

Find the value of the pronumeral.




4 Find the value of the pronumeral.


a




b




c




d



e



f






Hint for Q4: Substitute for a, b and c, then solve for the remaining pronumeral.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

4C

Example 7 Using rounding to approximate the length of a shorter side


Find the value of x, rounding to two decimal places.



Solution	Explanation
$x^2 + 7.6^2 = 10^2$	Write the rule and substitute $c = 10$ and $b = 7.6$
$x^2 + 57.76 = 100$	
$x^2 = 42.24$	Subtract 57.76 from both sides.
$\therefore x = \sqrt{42.24}$	Take the square root of both sides.
$x = 6.50$ (to 2 d.p.)	Round to two decimal places.


Now you try

Find the value of x, rounding to two decimal places.




5 Find the value of the pronumeral. Express your answers correct to two decimal places.


a




b




c




d



e



f






Hint for Q5: Use your calculator to get your answer, then round to two decimal places.


Problem-solving and reasoning

6–8 | 7–10

6 The base of a ladder leaning against a wall is 1.5 m from the base of the wall. The ladder is 5.5 m long. Find how high the top of the ladder is above the ground, correct to one decimal place.



7 A 32 m communication tower is supported by 35 m cables stretching from the top of the tower to a position at ground level. Find the distance from the base of the tower to the point where the cable reaches the ground, correct to one decimal place.




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

8 If a television has a screen size of 63 cm, it means that the diagonal length of the screen is 63 cm. If the vertical height of a 63 cm screen is 39 cm, find how wide the screen is to the nearest centimetre.




9 A 1.3 m vertical fence post is supported by a 2.27 m bar, as shown in the diagram on the right. Find the distance (d metres) from the base of the post to where the support enters the ground. Give your answer correct to two decimal places.



Example 8 Using Pythagoras' theorem and exact values

Find the value of x, giving your answer as a surd (exact value).



Solution

$x^2 + x^2 = 3^2$

$2x^2 = 9$

$x^2 = \frac{9}{2}$

$\therefore x = \sqrt{\frac{9}{2}}$ or $\frac{3}{\sqrt{2}}$

Explanation

Two sides are of length x.


Add like terms.

Divide both sides by 2.

Take the square root of both sides. To express as an exact answer, do not calculate the square root.


Now you try

Find the value of x, giving your answer as a surd (exact value).




10 Find the value of x as an exact answer. Note that the triangles are isosceles.


a



b




c




First find a missing length — 11

11 Find the value of x. Give an exact answer each time.


a



b



c



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

4D Applying Pythagoras' theorem ★

Learning intentions

- To be able to identify right-angled triangles in diagrams
- To be able to calculate unknown side lengths using given information and Pythagoras' theorem
- To be able to use Pythagoras' theorem to solve a problem

Key vocabulary: Pythagoras' theorem, right-angled


To apply Pythagoras' theorem to solve a real problem, it can help to first draw a right-angled triangle. This makes it easier to identify the unknown side. As long as two sides of the right-angled triangle are known, the length of the third side can be found.



The length of each cable on the Anzac Bridge, Sydney, can be calculated using Pythagoras' theorem.

Lesson starter: But where is the right-angled triangle?


A shed with two walls of length 3 m and 7 m has a sloping 10 m roof. We need to determine the distance between the two walls.



- Can you identify a right-angled triangle?
- What two side lengths on the right-angled triangle do you know?
- Show how Pythagoras' theorem can be used to find the unknown side.
- How would you put the answer in words?

Key ideas

- When applying Pythagoras' theorem to real-world problems:
  - Identify and draw right-angled triangles that may help to solve the problem (shown in red on diagram).
  - Label the sides with their lengths or with a letter (pronumeral) if the length is unknown.
  - Use Pythagoras' theorem to solve for the unknown.
  - Solve the problem by making any further calculations and answering in words.
  - Check that the answer that you have appears to be reasonable in the given situation.




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Exercise 4D


Understanding

1, 2 | 2


1 Two vertical posts are connected at the top by a wire as shown (right), Which of the following triangles could be used to help find the cable length?




A



B



C




2 Match each problem (**a**, **b** or **c**) with both a diagram (**A**, **B** or **C**) and its solution (**I**, **II**, **III**).

a Two trees stand 20 m apart. They are 32 m and 47 m tall. What is the distance between the tops of the two trees?


b A kite is flying with a kite string of length 35 m. Its horizontal distance from its anchor point is 17 m. How high is the kite flying?

c A man walks due north for 2 km then north-east for 3 km. How far north is he from his starting point?


A



B



C



I The kite is flying at a height of 30.59 m.

II The man has walked a total of $2+2.12 = 4.12$ km north from his starting point.

III The distance between the top of the two trees is 25 m.

Fluency


3, 4, 6 | 3, 5, 6

Example 9 Identifying right-angled triangles in a diagram

An area of lawn is in the shape of a trapezium, as shown.


a Draw a right-angled triangle to help find the unknown side length.

b Find the unknown side length correct to two decimal places.




Solution

a



Explanation

The top side of the triangle is 20 m − 11 m = 9 m.



Continued on next page

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

4D


b $a^2 + b^2 = c^2$
$a^2 + 9^2 = 13^2$
$a^2 + 81 = 169$
$a^2 = 88$
$\therefore a = \sqrt{88}$
$= 9.38$ (to 2 d.p.)
The unknown length is 9.38 m.

Use Pythagoras' theorem with the hypotenuse as 13 m.
Solve for a and round as required.


Now you try

A paved area is in the shape of a trapezium, as shown.


a Draw a right-angled triangle to help find the unknown side length.
b Find the unknown side length correct to two decimal places.



3 The side of a sculpture is in the shape of a trapezium, as shown.
a Draw a right-angled triangle to help find the unknown side length.
b Find the unknown length correct to two decimal places.




4 A scale drawing of the front of a dog kennel is shown. Find the unknown side length correct to two decimal places.




5 Find the direct distance between the points A and B in each of the following, correct to one decimal place.

a




b






Hint for Q5: Draw in the right-angled triangle that will help to find the distance AB. The first one is shown for you.

c




d



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


Example 10 Applying Pythagoras' theorem

Two skyscrapers are located 25 m apart and a cable links the tops of the two buildings. Find the length of the cable if the buildings are 50 m and 80 m in height. Give your answer correct to two decimal places.



Solution

Let x m be the length of the cable.




$$c^2 = a^2 + b^2$$
$$x^2 = 25^2 + 30^2$$
$$x^2 = 625 + 900$$
$$= 1525$$
$$\therefore x = \sqrt{1525}$$
$$= 39.05 \text{ (to 2 d.p.)}$$

$\therefore$ The cable is 39.05 m long.

Explanation

Draw a right-angled triangle and label the measurements and pronumerals.




Set up an equation using Pythagoras' theorem and solve for x.


Answer the question in words.

Now you try

Two buildings are 20 m apart. A cable links the top of the two buildings. Find the length of the cable if the buildings are 40 m and 65 m in height. Give your answer correct to two decimal places.



6 Two poles are located 2 m apart. A wire links the tops of the two poles. Find the length of the wire if the poles are 2.8 m and 5 m in height. Give your answer correct to one decimal place.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

4D

Problem-solving and reasoning


7, 8 | 7, 9, 10

7 Two skyscrapers are located 25 m apart and a cable of length 62.3 m links the tops of the two buildings. If the taller building is 200 metres tall, what is the height of the shorter building? Give your answer correct to one decimal place.



Hint for Q7: First draw a right-angled triangle just like in Example 10. Then find the difference in their heights.

8 A garage is to be built with a skillion roof (a roof with a single slope). The measurements are given in the diagram. Calculate the pitch line length, to the nearest millimetre. Allow 500 mm for each of the eaves.



9 Two bushwalkers are standing on different mountain sides. According to their maps, one of them is at a height of 2120 m and the other is at a height of 1650 m. If the horizontal distance between them is 950 m, find the direct distance between the two bushwalkers. Give your answer correct to the nearest metre.



Hint for Q9: You will need to find the hypotenuse length of a right-angled triangle. Use a diagram like Example 10

10 A 100 m radio mast is supported by six cables in two sets of three cables. They are anchored to the ground at an equal distance from the mast. The top set of three cables is attached at a point 20 m below the top of the mast. Each cable in the lower set of three cables is 60 m long and is attached at a height of 30 m above the ground. If all the cables have to be replaced, find the total length of cable required. Give your answer correct to two decimal places.



Hint for Q10: First draw a diagram and add all the given information.


The suspension bridge

— | 11

11 A suspension bridge is built with two vertical pylons and two straight beams of equal length. They extend from the top of the pylons to meet at a point, C, above the centre of the bridge, as shown in the diagram on the right.

a Calculate the vertical height of the point C above the tops of the pylons.

b Calculate the distance between the pylons; that is, the length of the span of the bridge correct to one decimal place.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

4E Trigonometric ratios

Learning intentions

- To know how to label the sides of a right-angled triangle as opposite, adjacent and hypotenuse
- To understand that the opposite and adjacent sides of a triangle are relative to the angle involved
- To know the trigonometric ratios for sine, cosine and tangent
- To understand that the trigonometric ratios are always the same if the angles in a right-angled triangle are the same
- To be able to write the correct trigonometric ratio for a right-angled triangle based on the given information

Key vocabulary: trigonometric ratio, opposite, adjacent, hypotenuse, sine, cosine, tangent, right-angled


The branch of mathematics called trigonometry deals with the relationship between the side lengths and angles in triangles. Trigonometry dates back to the ancient Egyptian and Babylonian civilisations where a basic form of trigonometry was used in the building of pyramids and in the study of astronomy. In the first century CE, Claudius Ptolemy advanced the study of trigonometry by writing 13 books called the *Almagest*. Ptolemy also developed tables of values linking the sides and angles of a triangle.




A basic form of trigonometry was used in the study of ancient astronomy.

Lesson starter: Constancy of sine, cosine and tangent

In geometry we would say that similar triangles have the same shape but are of different size. Here are three similar right-angled triangles. The angle θ (theta) is the same for all three triangles.



We will now calculate three special ratios for the angle θ in the above triangles: sine, cosine and tangent. We use the sides labelled Hypotenuse (H), Opposite (O) and Adjacent (A) as shown at right.



- Complete this table, simplifying all fractions.
- What do you notice about the value of:
  - **a** $\sin\theta$ for all three triangles?
  - **b** $\cos\theta$ for all three triangles?
  - **c** $\tan\theta$ for all three triangles?
- Why are the three ratios ($\sin\theta$, $\cos\theta$ and $\tan\theta$) the same for all three triangles? Discuss.


Triangle	$\frac{O}{H}$ ($\sin\theta$)	$\frac{A}{H}$ ($\cos\theta$)	$\frac{O}{A}$ ($\tan\theta$)
A	$\frac{5}{13}$		
B		$\frac{24}{26} = \frac{12}{13}$	
C			$\frac{15}{36} = \frac{5}{12}$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

4E


Key ideas

- If a right-angled triangle has one angle θ, then:
  - the longest side is called the hypotenuse (H)
  - the side opposite θ is called the **opposite** (O)
  - the remaining side is called the **adjacent** (A) (next to angle θ)



- For a right-angled triangle with a given angle θ, the three **trigonometric ratios: sine** (sin), **cosine** (cos) and **tangent** (tan) are given by:
  - sine of angle θ (or $\sin\theta$) $= \dfrac{\text{length of the opposite side}}{\text{length of the hypotenuse}}$
  - cosine of angle θ (or $\cos\theta$) $= \dfrac{\text{length of the adjacent side}}{\text{length of the hypotenuse}}$
  - tangent of angle θ (or $\tan\theta$) $= \dfrac{\text{length of the opposite side}}{\text{length of the adjacent side}}$

- For any right-angled triangle with the same angles, these ratios are always the same.
- The mnemonic SOHCAHTOA is useful when trying to remember the three ratios.



SOH CAH TOA


$\sin\theta = \dfrac{\text{Opposite}}{\text{Hypotenuse}}$ $\cos\theta = \dfrac{\text{Adjacent}}{\text{Hypotenuse}}$ $\tan\theta = \dfrac{\text{Opposite}}{\text{Adjacent}}$

Exercise 4E

Understanding


1, 2 | 2

1 For this triangle:



- **a** What is the length of the hypotenuse?
- **b** What is the length of the side opposite θ?
- **c** What is the length of the side adjacent to θ?

2 Write the missing word in these sentences.



- **a** H stands for the word _____.
- **b** O stands for the word _____.
- **c** A stands for the word _____.
- **d** $\sin\theta =$ _____ ÷ Hypotenuse.
- **e** $\cos\theta$ = Adjacent ÷ _____.
- **f** $\tan\theta$ = Opposite ÷ _____.


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Fluency


3–5 | 3–4(½), 5, 6

Example 11 Labelling the sides of triangles

Copy this triangle and label the sides as opposite to θ (O), adjacent to θ (A) and hypotenuse (H).



Solution




Explanation

Draw the triangle and label the side opposite the right angle as hypotenuse (H), the side opposite the angle θ as opposite (O) and the remaining side next to the angle θ as adjacent (A).


Now you try

Copy this triangle and label the sides as opposite to θ (O), adjacent to θ (A) and hypotenuse (H).




3 Copy each of these triangles and label the sides as opposite to θ (O), adjacent to θ (A) and hypotenuse (H).

a




b






Hint for Q3: Alternatively, pencil the letters O, A and H onto this page.


c




d




e




f



g



h




Essential Mathematics Core for the Victorian Curriculum Year 9 ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

4E


Example 12 Writing trigonometric ratios

Write a trigonometric ratio (in fraction form) for each of the following triangles.


a



b




c



Solution

a $\cos\theta = \frac{A}{H}$

$\cos\theta = \frac{5}{7}$




Explanation

Side length 7 is opposite the right angle so it is the hypotenuse (H).
Side length 5 is adjacent to angle θ so it is the adjacent (A).
Use $\cos\theta$ since we have A and H.

b $\sin\theta = \frac{O}{H}$


$\sin\theta = \frac{4}{9}$



Side length 9 is opposite the right angle so it is the hypotenuse (H).
Side length 4 is opposite angle θ so it is the opposite (O).
Use $\sin\theta$ since we have O and H.

c $\tan\theta = \frac{O}{A}$

$\tan\theta = \frac{3}{5}$




Side length 3 is opposite angle θ so it is the opposite (O).
Side length 5 is the adjacent side to angle θ so it is the adjacent (A).
Use $\tan\theta$ since we have O and A.


Now you try

Write a trigonometric ratio (in fraction form) for each of the following triangles.


a



b




c




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

4 Write a trigonometric ratio (in fraction form) for each of the following triangles and simplify where possible.


a



b




c






Hint for Q4: Use sin θ if you have O and H, use cos θ if you have A and H, and use tan θ if you have O and A.


d




e




f




g



h




i




5 For each of these triangles, write a ratio (in simplified fraction form) for sin θ, cos θ and tan θ.


a




b



c



6 Here are two similar triangles, A and B.





Hint for Q6: Simplify your fractions from the larger triangle to help see the connection.

a i Write the ratio sin θ (as a fraction) for triangle A.

ii Write the ratio sin θ (as a fraction) for triangle B.

iii What do you notice about your two answers from parts **a i** and **a ii** above?

b i Write the ratio cos θ (as a fraction) for triangle A.

ii Write the ratio cos θ (as a fraction) for triangle B.

iii What do you notice about your two answers from parts **b i** and **b ii** above?

c i Write the ratio tan θ (as a fraction) for triangle A.

ii Write the ratio tan θ (as a fraction) for triangle B.

iii What do you notice about your two answers from parts **c i** and **c ii** above?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

4E

Problem-solving and reasoning

7–9 | 8–11

7 The facade of a Roman temple has the given measurements. Write down the ratio for:

a $\sin\theta$


b $\cos\theta$

c $\tan\theta$



The Pantheon, a Roman temple that was built in 126 CE.

8 A vertical flag pole casts a shadow 20 m long. If the pole is 15 m high, find the ratio for $\tan\theta$.





Hint for Q8: Decide which side is the opposite to θ and which side is adjacent to θ.

9 For the triangle shown, state the length of the side that corresponds to:


a the hypotenuse

b the side opposite angle θ

c the side opposite angle α

d the side adjacent to angle θ


e the side adjacent to angle α.



10 For the triangle shown on the right, write a ratio (in fraction form) for:

a $\sin\theta$ **b** $\sin\alpha$ **c** $\cos\theta$

d $\tan\alpha$ **e** $\cos\alpha$ **f** $\tan\theta$



11 **a** Draw a right-angled triangle and mark one of the angles as θ. Mark in the length of the opposite side as 15 units and the length of the hypotenuse as 17 units.

b Using Pythagoras' theorem, find the length of the adjacent side.

c Determine the ratios for $\sin\theta$, $\cos\theta$ and $\tan\theta$.




Hint for Q11: Use $a^2 + b^2 = c^2$ for part **b** with $c = 17$ and $b = 15$.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

The complementary connection — 12, 13

12 This triangle has angles $90°$, $60°$ and $30°$ and side lengths 1, 2 and $\sqrt{3}$.



a Write a ratio for:


i $\sin 30°$ **ii** $\cos 30°$ **iii** $\tan 30°$

iv $\sin 60°$ **v** $\cos 60°$ **vi** $\tan 60°$

b What do you notice about the following pairs of ratios?

i $\cos 30°$ and $\sin 60°$ **ii** $\sin 30°$ and $\cos 60°$

13 **a** Measure all the side lengths of this triangle to the nearest millimetre.



b Use your measurements from part **a** to find an approximate ratio for:

i $\cos 40°$ **ii** $\sin 40°$ **iii** $\tan 40°$

iv $\sin 50°$ **v** $\tan 50°$ **vi** $\cos 50°$

c Do you notice anything about the trigonometric ratios for $40°$ and $50°$?



Surveyors measure distances and angles and apply trigonometry to calculate other angles and distances.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

4F Finding side lengths

Learning intentions

- To know how to use a calculator to evaluate sin, cos or tan in degrees
- To be able to set up a trigonometric ratio to find a missing side length
- To know that at least one angle and one side length must be known to find another side length using trigonometry
- To be able to use a trigonometric ratio to find an unknown side length

Key vocabulary: trigonometric ratio, opposite, adjacent, hypotenuse, sine, cosine, tangent, right-angled, numerator

Since ancient times, mathematicians have attempted to tabulate the three trigonometric ratios for varying angles. Here are the ratios for some angles in a right-angled triangle, correct to three decimal places.

Angle (θ)	$\sin\theta$	$\cos\theta$	$\tan\theta$
0°	0	1	0
15°	0.259	0.966	0.268
30°	0.5	0.866	0.577
45°	0.707	0.707	1
60°	0.866	0.5	1.732
75°	0.966	0.259	3.732
90°	1	0	undefined

In modern times these values can be evaluated using calculators to a high degree of accuracy and can be used to help solve problems involving triangles with unknown side lengths.



Architects use trigonometry to design buildings.


Lesson starter: Calculator start up

All scientific or CAS calculators can produce accurate values of $\sin\theta$, $\cos\theta$ and $\tan\theta$.

- Ensure that your calculator is in degree mode.
- Find the value of the following, correct to three decimal places.
  - $\sin 50°$
  - $\cos 81°$
  - $\tan 36°$
- Use trial and error to find (to the nearest degree) an angle, θ, which satisfies these conditions.
  - $\sin\theta = 0.454$
  - $\cos\theta = 0.588$
  - $\tan\theta = 9.514$

Key ideas


- If θ is in degrees, the ratios for $\sin\theta$, $\cos\theta$ and $\tan\theta$ can accurately be found using a calculator in degree mode.
- If the angles and one side length of a right-angled triangle are known, then the other side lengths can be found using the $\sin\theta$, $\cos\theta$ or $\tan\theta$ ratios.



$\sin 30° = \frac{x}{3}$

$x = 3 \times \sin 30°$


$= 1.5$



$\cos 42° = \frac{x}{7.2}$

$x = 7.2 \times \cos 42°$

≈ 5.35



$\tan 71° = \frac{x}{4}$

$x = 4 \times \tan 71°$

≈ 11.62

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


Exercise 4F

Understanding


1–3 | 2, 3

1 For the marked angle θ, decide if x represents the length of the opposite (O), adjacent (A) or hypotenuse (H) side.


a



b




c




2 Decide whether you would use $\sin\theta = \frac{\text{O}}{\text{H}}$, $\cos\theta = \frac{\text{A}}{\text{H}}$ or $\tan\theta = \frac{\text{O}}{A}$ to help find the value of x in these triangles. Do not find the value of x; just state which ratio would be used.


a



b



c



3 Which first step would be used to solve for x in the equation $\sin 70° = \frac{x}{4}$?

A Add 4 to both sides
B Multiply both sides by 4
C Subtract 4 from both sides
D Divide both side by 4

Fluency

4–6(½) | 4–6(½)

Example 13 Using a calculator

Use a calculator to evaluate the following, correct to two decimal places.

a $\sin 50°$ **b** $\cos 16°$ **c** $\tan 77°$

Solution

a $\sin 50° = 0.77$ (to 2 d.p.)

b $\cos 16° = 0.96$ (to 2 d.p.)

c $\tan 77° = 4.33$ (to 2 d.p.)

Explanation

$\sin 50° = 0.766044\ldots$ the third decimal place is greater than 4, so round up.

$\cos 16° = 0.961261\ldots$ the third decimal place is less than 5, so round down.

$\tan 77° = 4.331475\ldots$ the third decimal place is less than 5, so round down.

Now you try

Use a calculator to evaluate the following, correct to two decimal places.

a $\sin 20°$ **b** $\cos 38°$ **c** $\tan 67°$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

4F

4 Use a calculator to evaluate the following correct to two decimal places.

a $\sin 20°$ **b** $\cos 37°$ **c** $\tan 64°$ **d** $\sin 47°$
e $\cos 84°$ **f** $\tan 14.1°$ **g** $\sin 27.4°$ **h** $\cos 76.2°$

Example 14 Solving for x in the numerator of a trigonometric ratio

Find the value of x in the equation $\cos 20° = \frac{x}{3}$, correct to two decimal places.

Solution

$\cos 20° = \frac{x}{3}$

$x = 3 \times \cos 20°$
$= 2.82$ (to 2 d.p.)

Explanation

Multiply both sides of the equation by 3 and round as required.

Swap sides to put x on the left.

Now you try

Find the value of x in the equation $\cos 40° = \frac{x}{5}$, correct to two decimal places.

5 Find the value of x correct to two decimal places.

a $\sin 50° = \frac{x}{4}$ **b** $\tan 81° = \frac{x}{3}$ **c** $\cos 33° = \frac{x}{6}$

d $\cos 75° = \frac{x}{3.5}$ **e** $\sin 24° = \frac{x}{4.2}$ **f** $\tan 42° = \frac{x}{10}$

g $\frac{x}{7.1} = \tan 18.4°$ **h** $\frac{x}{5.3} = \sin 64.7°$ **i** $\frac{x}{12.6} = \cos 52.9°$




Hint for Q5: For part a, for example, multiply by 4 and use a calculator to evaluate $4 \times \sin 50°$.


Example 15 Finding side lengths

For each triangle, find the value of x correct to two decimal places.


a



b



c



Solution


a $\sin 38° = \frac{O}{H}$

$\sin 38° = \frac{x}{7}$

$x = 7 \sin 38°$
$= 4.31$ (to 2 d.p.)

Explanation

Since the opposite side (O) and the hypotenuse (H) are involved, the $\sin\theta$ ratio must be used. Use $\theta = 38°$.



Multiply both sides by 7 and evaluate using a calculator.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

b $\tan 42° = \frac{O}{A}$


$\tan 42° = \frac{x}{4}$

$x = 4 \tan 42°$

$= 3.60$ (to 2 d.p.)

Since the opposite side (O) and the adjacent side (A) are involved, the $\tan \theta$ ratio must be used. Use $\theta = 42°$.

Multiply both sides by 4 and evaluate.



c $\cos 24° = \frac{A}{H}$


$\cos 24° = \frac{x}{10}$

$x = 10 \cos 24°$

$= 9.14$ (to 2 d.p.)

Since the adjacent side (A) and the hypotenuse (H) are involved, the $\cos \theta$ ratio must be used.


Multiply both sides by 10.




Now you try

For each triangle, find the value of x correct to two decimal places.


a



b




c




6 Find the value of x correct to two decimal places.


a



b




c






Hint for Q6: First decide which sides you have, including the unknown (O, A or H), then choose the correct ratio ($\sin \theta$, $\cos \theta$ or $\tan \theta$).


d




e




f




g




h




i




j




k




l



m



n




ISBN 978-1-009-59315-1
© Greenwood et al. 2025
Photocopying is restricted under law and this material must not be transferred to another party.


4F

Problem-solving and reasoning 7–9 8–11

7 A pole is supported by a 12 m cable that makes a 60° angle with the ground. How tall is the pole, correct to two decimal places?




8 Amy walks 5.4 m up a ramp that is inclined at 12° to the horizontal. How high (correct to two decimal places) is she above her starting point?




9 Kane wanted to measure the width of a river at a certain point. He placed two markers, A and B, 72 m apart along the bank. C is a point directly opposite marker B. Kane measured angle CAB to be 32°. Find the width of the river correct to two decimal places.



Hint for Q9: The 'width' is opposite (O) the 32° angle and the 72 m length is adjacent (A) to the 32° angle.



10 One end of a 12.2 m rope is tied to a boat. The other end is tied to an anchor, which is holding the boat steady in the water. If the anchor is making an angle of 34° with the vertical, how deep is the water? Give your answer correct to two decimal places.






Hint for Q10: The depth length is adjacent (A) to the 34° angle.

11 For this right-angled triangle:


- **a** Find the value of angle $\angle C$.
- **b** Calculate the value of x correct to three decimal places using the sine ratio.
- **c** Calculate the value of x correct to three decimal places but instead use the cosine ratio.




Combining multiple trigonometric calculations — 12

12 Find the length AB in these diagrams. A combination of calculations is required. Round to two decimal places where necessary.

a



b




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


Progress quiz

4A **1** Write down Pythagoras' theorem for these triangles using the pronumerals and numbers given.


a



b




c




4B **2** Find the length of the hypotenuse in these right-angled triangles. Give exact answers for parts **a** and **b** and round to one decimal place in part **c**.


a



b




c




4C **3** Find the value of the pronumeral in these right-angled triangles. Round to two decimal places where necessary.


a




b




4C **4** A 3 m wooden plank is supporting an advertising billboard that is 2 m high. What is the distance of the plank from the base of the billboard? Round to one decimal place.



4D **5** A backyard deck is in the shape of a trapezium as shown. By drawing an appropriate right-angled triangle, find the width of the deck, w metres, correct to two decimal places.



4D **6** Two telegraph poles 3.2 m and 5.4 m high are 2.4 m apart. A wire connects the top of the two poles. Find the length of the wire correct to one decimal place.




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


Progress quiz

4E **7** Write a trigonometric ratio (in fraction form) for each of the following triangles.


a



b




c




4F **8** Find the value of x, correct to two decimal places, in each of the following triangles.


a



b



4F **9** A ladder leaning against a wall makes a 60° angle with the ground as shown. If the ladder is 3 m long, how far up the wall (x metres) does it reach? Round to one decimal place.





ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

4G Solving for the denominator ★

Learning intentions

- To be able to use a trigonometric ratio to find an unknown in the denominator

Key vocabulary: trigonometric ratio, opposite, adjacent, hypotenuse, sine, cosine, tangent, right-angled, denominator

So far we have constructed trigonometric ratios using a pronumeral that has always appeared in the numerator. For example: $\sin 40° = \frac{x}{5}$. This makes it easy to solve for x where both sides of the equation can be multiplied by 5. If, however, the pronumeral appears in the denominator, there are a number of algebraic steps that can be taken to find the solution.



The Pont de Normandie bridge in France is supported by cables of varying lengths but fixed angle. The length is determined by the distance of the anchor to the vertical support.

Lesson starter: Solution steps

Three students attempt to solve $\sin 40° = \frac{5}{x}$ for x.

Nick says $x = 5 \times \sin 40°$


Sharee says $x = \frac{5}{\sin 40°}$

Dori says $x = \frac{1}{5} \times \sin 40°$

- Which student has the correct solution?
- Can you show the algebraic steps that support the correct answer?

Key ideas

- If the unknown value of a trigonometric ratio is in the denominator, you need to rearrange the equation to make the pronumeral the subject.



For example, for the triangle shown, multiplying both sides by x removes the fraction.

$$\cos 30° = \frac{5}{x}$$

$$x \times \cos 30° = 5$$

Dividing both sides by $\cos 30°$ solves for x.

$$x = \frac{5}{\cos 30°}$$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


Exercise 4G

Understanding


1–3 | 3

1 Write the trigonometric ratio that would be used to find the value of x in these triangles.


a



b



c



2 Which of the following equations is produced by multiplying both sides of $\sin 50° = \frac{3}{x}$ by x?

A $x = 3 \times \sin 50°$ **B** $x = \frac{\sin 50°}{3}$ **C** $x \times \sin 50° = 3$

3 Complete the following solutions by filling in the boxes.

a $x \times \cos 25° = 2$

$$x = \frac{2}{\square}$$

b $x \times \tan 65° = 10$

$$x = \frac{10}{\square}$$

c $x \times \sin 72° = 6$

$$x = \frac{\square}{\square}$$

Fluency

4–5(½) | 4–5(½)

Example 16 Solving for x in the denominator

Solve for x in the equation $\cos 35° = \frac{2}{x}$, correct to two decimal places.

Solution	Explanation
$\cos 35° = \frac{2}{x}$	
$x \cos 35° = 2$	Multiply both sides of the equation by x.
$x = \frac{2}{\cos 35°}$	Divide both sides of the equation by $\cos 35°$.
$= 2.44$ (to 2 d.p.)	Evaluate and round off to two decimal places.

Now you try

Solve for x in the equation $\tan 40° = \frac{3}{x}$, correct to two decimal places.

4 Find the value of x correct to two decimal places.

a $\cos 43° = \frac{3}{x}$ **b** $\sin 36° = \frac{4}{x}$ **c** $\tan 9° = \frac{6}{x}$

d $\tan 64° = \frac{2}{x}$ **e** $\cos 67° = \frac{5}{x}$ **f** $\sin 12° = \frac{3}{x}$

g $\sin 38.3° = \frac{5.9}{x}$ **h** $\frac{45}{x} = \tan 21.4°$ **i** $\frac{18.7}{x} = \cos 32°$




Hint for Q4: First multiply both sides by x.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Example 17 Finding side lengths

Find the value of x correct to two decimal places.



Solution

$\sin 35° = \frac{O}{H}$

$\sin 35° = \frac{5}{x}$


$x \sin 35° = 5$

$x = \frac{5}{\sin 35°}$

$= 8.72$ (to 2 d.p.)

Explanation

Since the opposite side (O) is given and we require the hypotenuse (H), use $\sin\theta$.




Multiply both sides of the equation by x then divide both sides of the equation by $\sin 35°$.

Evaluate on a calculator and round off to two decimal places.


Now you try

Find the value of x correct to two decimal places.




5 Find the value of x correct to two decimal places using the sine, cosine or tangent ratios.


a



b




c




Hint for Q5:
Are you going to use $\sin\theta = \frac{O}{H}$, $\cos\theta = \frac{A}{H}$ or $\tan\theta = \frac{O}{A}$?




d




e




f




g




h




i




j



k



l




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

4G

Problem-solving and reasoning

6–8 | 7–10


6 A kite is flying at a height of 27 m above the anchor point. If the string is inclined at 42° to the horizontal, find the length of the string correct to the nearest metre.






Hint for Q6 and Q7: First decide if you are going to use sin θ (SOH), cos θ (CAH) or tan θ (TOA).

7 A hang glider flying at a height of 800 m descends at an angle of 12° to the horizontal. How far (to the nearest metre) has it travelled in descending to the ground?



Example 18 Finding two side lengths on one triangle

Find the value of x and y correct to two decimal places.



Solution

$\tan 28° = \frac{O}{A}$

$\tan 28° = \frac{19}{x}$

$x \tan 28° = 19$

$x = \frac{19}{\tan 28°}$

$= 35.73$ (to 2 d.p.)

$\sin 28° = \frac{19}{y}$


$y \sin 28° = 19$

$y = \frac{19}{\sin 28°}$

$= 40.47$ (to 2 d.p.)

Explanation


Since the opposite side (O) is given and the adjacent (A) is required, use tan θ.



Multiply both sides of the equation by x.

Divide both sides of the equation by $\tan 28°$ and round answer to two decimal places.


Use the opposite (19) as it is an exact length, so use $\sin \theta = \frac{O}{H}$.



Alternatively, y can be found by Pythagoras' theorem once x is known.

Now you try


Find the value of x and y correct to two decimal places.




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

8 Find the value of each pronumeral correct to one decimal place.


a




b




c




d




e



f




9 A mine shaft is dug at an angle of 15° to the horizontal. The horizontal length of the mine is 100 m. Answer the questions to the nearest metre.




a How far below ground level is the end of the shaft?

b How long is the mine shaft?



10 In calculating the value of x for this triangle, correct to two decimal places, two students come up with these answers.

A $x = \frac{5}{\sin 31°} = \frac{5}{0.52} = 9.62$ B $x = \frac{5}{\sin 31°} = 9.71$




a Which of the above two answers is more correct and why?

b What advice would you give to the student whose answer is not accurate?


Trigonometry and perimeters — 11

11 Find the perimeter of these triangles correct to one decimal place. You will first need to find the lengths of all the sides.

a



b



Using a calculator 4G: Trigonometry

This activity can be found in the More Resources section of the Interactive Textbook in the form of a printable PDF.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

4H Finding an angle

Learning intentions


- To know that inverse sine, inverse cosine and inverse tangent can be used to find angles in right-angled triangles
- To know that two side lengths must be known to find an unknown angle in a right-angled triangle
- To be able to evaluate the inverse trigonometric ratios on a calculator
- To be able to use the inverse trigonometric functions to find an unknown angle

Key vocabulary: trigonometric ratio, inverse sine, inverse cosine, inverse tangent, opposite, adjacent, hypotenuse, sine, cosine, tangent

Given two sides of a right-angled triangle, we can work out the unknown angles.

By using one of the three trigonometric ratios, we can use inverse trigonometric functions to find a value of θ.

For example, if $\cos\theta = 0.5$, then $\theta = \cos^{-1}(0.5) = 60°$






Surveyors use trigonometry to measure the height and angles of land.

Lesson starter: Trial and error can be slow


We know that for this triangle, $\sin\theta = \frac{1}{3}$



- Guess the angle θ.
- For your guess, use a calculator to see if $\sin\theta = \frac{1}{3} (= 0.333\ldots)$ for your chosen value of θ.
- Update your guess and use your calculator to check once again.
- Repeat this trial-and-error process until you think you have the angle θ correct to three decimal places.
- Now evaluate $\sin^{-1}\left(\frac{1}{3}\right)$ and check your guess.

Key ideas

- **Inverse sine** ($\sin^{-1}$), **inverse cosine** ($\cos^{-1}$) and **inverse tangent** ($\tan^{-1}$) can be used to find angles in right-angled triangles.
  - $\sin\theta = \frac{3}{5}$ means $\theta = \sin^{-1}\left(\frac{3}{5}\right)$
  - $\cos\theta = \frac{4}{5}$ means $\theta = \cos^{-1}\left(\frac{4}{5}\right)$
  - $\tan\theta = \frac{3}{4}$ means $\theta = \tan^{-1}\left(\frac{3}{4}\right)$



- Note that $\sin^{-1} x$ does *not* mean $\frac{1}{\sin x}$.
- The $\sin^{-1}$, $\cos^{-1}$, and $\tan^{-1}$ functions can be found on most calculators.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


Exercise 4H

Understanding


1–3 | 3

1 Write the trigonometric ratio for these triangles, e.g. $\sin\theta = \frac{8}{9}$ in part **a**.


a




b



c



d





Hint for Q1: Choose sin θ if using O and H, cos θ if using A and H, and tan θ if using O and A.

2 Use a calculator to evaluate the following, rounding to two decimal places.

a $\sin^{-1}(0.2)$ **b** $\cos^{-1}(0.43)$ **c** $\tan^{-1}(2.5)$

3 Complete the following by filling in the boxes.

a If $\cos\theta = \frac{1}{2}$ then $\theta = \cos^{-1}(\square)$

b If $\sin\theta = 0.6$ then $\theta = \sin^{-1}(\square)$

c If $\tan\theta = \frac{5}{4}$ then $\theta = \square$

Fluency

4–6(½) | 4–6(½)

Example 19 Using inverse trigonometric ratios

Find the value of θ to the nearest degree if $\sin\theta = 0.3907$.

Solution	Explanation
$\sin\theta = 0.3907$	
$\theta = \sin^{-1}(0.3907)$	Use the $\sin^{-1}$ key on your calculator
$= 23°$ (to nearest degree)	Round off to the nearest whole number.

Now you try

Find the value of θ to the nearest degree if $\sin\theta = 0.464$.

4 Find the value of θ to the nearest degree.

a $\sin\theta = 0.5$ **b** $\cos\theta = 0.5$ **c** $\tan\theta = 1$

d $\cos\theta = 0.8660$ **e** $\sin\theta = 0.7071$ **f** $\tan\theta = 0.5774$

g $\sin\theta = 1$ **h** $\tan\theta = 1.192$ **i** $\cos\theta = 0$

j $\cos\theta = 0.5736$ **k** $\cos\theta = 1$ **l** $\sin\theta = 0.9397$



Hint for Q4: Use the $\sin^{-1}$, $\cos^{-1}$ or $\tan^{-1}$ functions.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

4H

Example 20 Using inverse trigonometric ratios with fractions

Find the value of θ correct to two decimal places if $\tan\theta = \frac{1}{2}$.

Solution

$\tan\theta = \frac{1}{2}$

$\theta = \tan^{-1}\left(\frac{1}{2}\right)$

$= 26.57°$ (to 2 d.p.)

Explanation

Use the $\tan^{-1}$ key on your calculator and round the answer to two decimal places.

Now you try


Find the value of θ correct to two decimal places if $\cos\theta = \frac{2}{3}$.

5 Find the angle θ correct to two decimal places.

a $\sin\theta = \frac{4}{7}$ **b** $\sin\theta = \frac{1}{3}$ **c** $\sin\theta = \frac{9}{10}$


d $\cos\theta = \frac{1}{4}$ **e** $\cos\theta = \frac{4}{5}$ **f** $\cos\theta = \frac{7}{9}$

g $\tan\theta = \frac{3}{5}$ **h** $\tan\theta = \frac{8}{5}$ **i** $\tan\theta = 12$



Example 21 Finding an angle

Find the value of θ to the nearest degree.



Solution

$\sin\theta = \frac{O}{H}$


$= \frac{6}{10}$

$\theta = \sin^{-1}\left(\frac{6}{10}\right)$

$= 37°$ (to nearest degree)

Explanation


Since the opposite side (O) and the hypotenuse (H) are given, use $\sin\theta$.



Use the $\sin^{-1}$ key on your calculator and round as required.

Now you try


Find the value of θ to the nearest degree.




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

6 Find the value of θ to the nearest degree.


a




b




c




d




e




f




g



h



i







Hint for Q6: First set up an equation (like those in Question 5), then use a calculator to find θ.


Problem-solving and reasoning 7–9 9–12

7 A road rises at a grade of 3 in 10. Find the angle (to the nearest degree) the road makes with the horizontal.





8 When a 2.8 m long seesaw is at its maximum height, it is 1.1 m off the ground. What angle (correct to two decimal places) does the seesaw make with the ground?





Hint for Q8: The 1.1 m length is opposite the angle and the 2.8 m length forms the hypotenuse.

9 Write the missing number.

a If $\sin 30° = \frac{1}{2}$ then $30° = \sin^{-1}(_____)$.

b If $\cos 50° = 0.64$ then $_____ = \cos^{-1}(0.64)$.

c If $\tan 45° = 1$ then $_____ = \tan^{-1}(_____)$.


d If $\sin 45° = 0.707$ then $45° = \sin^{-1}(_____)$.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


4H

10 Adam holds one end of a plank of wood 1.8 m above the ground. The other end of the plank rests on the ground. The plank of wood is 4.2 m long. Find the angle that the plank makes with the ground (θ), correct to one decimal place.






11 A ladder has a length of 5.8 m. The ladder is resting against the top of a platform that is 2.6 m high. Find the angle the ladder makes with the ground, correct to one decimal place.



12 For what value of θ is $\sin\theta = \cos\theta$? Choose a value of θ between $0°$ and $90°$.

Viewing angle — 13

13 Joe has trouble with his eyesight but every Sunday he goes to view his favourite painting at the gallery. His eye level is at the same level as the base of the painting and the painting is 1 metre tall.



Answer the following to the nearest degree for angles and to two decimal places for lengths.

a If $x = 3$, find the viewing angle θ.
b If $x = 2$, find the viewing angle θ.
c If Joe can stand no closer than 1 metre to the painting, what is Joe's largest viewing angle?
d When the viewing angle is $10°$, Joe has trouble seeing the painting. How far is he from the painting at this viewing angle?
e What would be the largest viewing angle if Joe could go as close as he would like to the painting?



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

4I Applying trigonometry ★

Learning intentions

- To be able to solve word problems using trigonometry by drawing a diagram and identifying appropriate right-angled triangles
- To know that angles of elevation and depression are measured from the horizontal
- To know that the angle of elevation and angle of depression between the same two points is equal
- To be able to work with angles of elevation and depression in diagrams and word problems

Key vocabulary: angle of elevation, angle of depression and horizontal

In many practical maths problems, trigonometry can be used to find unknown lengths and angles. Two special angles are called the angle of elevation and the angle of depression. These angles are measured from the horizontal.



A cat and a bird in a trigonometric stand-off.

Lesson starter: The cat and the bird

For the situation below, draw a detailed diagram showing these features:


- an angle of elevation (rising from the horizontal)
- an angle of depression (falling from the horizontal)
- any given lengths
- a right-angled triangle that will help to solve the problem

A cat and a bird eye each other from their respective positions. The bird is 20 m up a tree and the cat is on the ground 30 m from the base of the tree. Find the angle that their line of sight makes with the horizontal.


Compare your diagram with others in your class. Is there more than one triangle that could be drawn and used to solve the problem?


Key ideas

- To solve application problems involving trigonometry:
  - Draw a diagram and label the key information.
  - Identify and draw the appropriate right angled triangles separately.
  - Solve using trigonometry to find the missing measurements.
  - Express your answer in words.




- **Angles of elevation** and **depression** are always measured from the **horizontal**.
  - angle of elevation is looking up at an object
  - angle of depression is looking down at an object





- Between the same two points the angle of elevation is equal to the angle of depression.




ISBN 978-1-009-59315-1
© Greenwood et al. 2025
Photocopying is restricted under law and this material must not be transferred to another party.

Exercise 4I

Understanding

1–4 | 3, 4

1 Draw this diagram and complete these tasks.






Hint for Q1: The angles are measured from the corresponding horizontal

a Mark in the following.
- **i** The angle of elevation (θ) of B from A
- **ii** The angle of depression (α) of A from B


b Is $\theta = \alpha$ in your diagram? Why?

2 Choose the diagram (**A**, **B** or **C**) that matches this situation. A boy views a kite at an angle of elevation of 28°.


A



B




C



3 For this diagram:

a What is the angle of elevation of B from A?


b What is the angle of depression of A from B?



Hint for Q3: Why do you think the two angles are equal in this diagram?



4 Find the values of the pronumerals in this diagram.






ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Fluency

5, 7, 9, 10 | 5, 6, 8–10

Example 22 Using angles of elevation

The angle of elevation of the top of a tower from a point on the ground 30 m away from the base of the tower is 28°. Find the height of the tower to the nearest metre.



Solution

Let the height of the tower be h m.

$$\tan 28° = \frac{\text{O}}{\text{A}}$$

$$= \frac{h}{30}$$


$$h = 30 \tan 28°$$

$$= 16 \text{ (to nearest metre)}$$

The height is 16 m.

Explanation


Since the opposite side (O) is required and the adjacent (A) is given, use $\tan\theta$.




Multiply both sides by 30 and evaluate, rounding to the nearest metre.
Write the answer in words.

Now you try

The angle of elevation of the top of a tower from a point on the ground 40 m from the base of the tower is 36°. Find the height of the tower to the nearest metre.



5 The angle of elevation of the top of a castle wall from a point on the ground 25 m from the base of the castle wall is 32°. Find the height of the castle wall to the nearest metre.




6 From a point on the ground, Emma measures the angle of elevation of an 80 m tower to be 27°. Find how far Emma is from the base of the tower, correct to the nearest metre.



Hint for Q6: First draw a simple triangle and label your side lengths and given angle.

7 From a pedestrian overpass, Chris spots a landmark at an angle of depression of 32°. How far away (to the nearest metre) is the landmark from the base of the 24 m high overpass?






ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

41


8 From a lookout tower, David spots a bushfire at an angle of depression of 25°. If the lookout tower is 42 m high, how far away (to the nearest metre) is the bushfire from the base of the tower?



Example 23 Finding an angle of depression

From the top of a vertical cliff, Andrea spots a boat out at sea. If the top of the cliff is 42 m above sea level and the boat is 90 m away from the base of the cliff, find Andrea's angle of depression to the boat to the nearest degree.

Solution



$\tan\theta = \frac{O}{A}$

$= \frac{42}{90}$

$\theta = \tan^{-1}\left(\frac{42}{90}\right)$

$\theta = 25°$ (to nearest degree)

The angle of depression is 25°.

Explanation

Draw a diagram and label all the given measurements. The angle of depression is below the horizontal.
Use alternate angles in parallel lines to mark θ inside the triangle.

Since the opposite (O) and adjacent sides (A) are given, use $\tan\theta$.

Use the $\tan^{-1}$ key on your calculator and round off to the nearest degree.


Express the answer in words.

Now you try


From the top of a lighthouse, a yacht is spotted out at sea. If the top of the lighthouse is 34 m above sea level and the yacht is 120 m away from the base of the lighthouse (at sea level), find the angle of depression from the top of the lighthouse to the yacht to the nearest degree.

9 From the top of a vertical cliff, Josh spots a swimmer out at sea. If the top of the cliff is 38 m above sea level and the swimmer is 50 m away from the base of the cliff, find the angle of depression from Josh to the swimmer, to the nearest degree.





10 From a ship, a person is spotted floating in the sea 200 m away. If the viewing position on the ship is 20 m above sea level, find the angle of depression from the ship to the person in the sea. Give your answer to the nearest degree.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Problem-solving and reasoning 11, 12 11–14


11 A power line is stretched from a pole to the top of a house. The house is 4.1 m high and the power pole is 6.2 m high. The horizontal distance between the house and the power pole is 12 m. Find the angle of elevation of the top of the power pole from the top of the house, to the nearest degree.

12 A road has a steady gradient of 1 in 10. It rises 1 m for every 10 m across.


a What angle does the road make with the horizontal? Give your answer to the nearest degree.

b A car starts from the bottom of the inclined road and drives 2 km along the road. How high, vertically, has the car climbed? Use your rounded answer from part **a** and give your answer correct to the nearest metre.






13 A house is to be built using the design shown on the right. The eaves are 600 mm and the house is 7200 mm wide, excluding the eaves. Calculate the length (to the nearest mm) of a sloping edge of the roof, which is pitched at 25° to the horizontal.



14 A plane flying at 1850 m starts to climb at an angle of 18° to the horizontal when the pilot sees a mountain peak 2450 m high, 2600 m away from him in a horizontal direction. Will the pilot clear the mountain?





Plane trigonometry — 15

15 An aeroplane takes off and climbs at an angle of 20° to the horizontal, at 190 km/h along its flight path for 15 minutes.

a Find:

i the distance the aeroplane travels in 15 minutes

ii the height the aeroplane reaches after 15 minutes correct to two decimal places.

b If the angle at which the plane climbs is twice the original angle but its speed is halved, will it reach a greater height after 15 minutes?

c If the plane's speed is doubled and its climbing angle is halved, will the plane reach a greater height after 15 minutes?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Maths@Work

Carpenter

Carpenters work with wood, building house frames, verandahs or decks, kitchen cabinets, built-in wardrobes and so on. They often work outdoors and have to be proficient with tools such as augers, drills, saws, chisels, rulers, squares, compasses and levels.


As with most trades, mathematical skills are used regularly. Carpenters must be able to measure, convert between units, order material and calculate lengths using measuring devices, as well as apply Pythagoras' theorem and trigonometry rules.



1 Carpenters work in millimetres as their base unit for measurement of length. Convert each of the following into millimetres.


a 45 cm **b** 2.4 m **c** 9.01 m **d** 270 cm **e** 46.0 m

2 A carpenter needs to cut mitred joints to join 4 pieces of wood of equal width to construct a picture frame. A mitre joint is formed when cuts at exactly 45° are made so that two pieces of timber fit together perfectly. Copy the diagram and show how the mitre cuts can be marked out by using isosceles triangles instead of a set square. Shade the final shape of the wood after the cuts are made. The top piece is done for you.




3 For each box below, find the minimum dimensions of a single rectangular sheet of plywood required to construct that closed box. Draw and label the least wasteful arrangement of the sides and ignore any overlap for joining. State the answers in millimetres. Diagrams are not to scale. You will need Pythagoras' theorem in part **b**.

a




b



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


4 A carpenter is constructing a set of shelves near the kitchen doorway. To avoid sharp corners and allow more walking space, a $60°$ triangle is cut from each shelf as shown in this diagram. The carpenter carefully measures AB and BC to achieve the same $60°$ angle on each shelf.



Calculate the unknown length (BC or AB) to the nearest mm, given these measurements.

a $AB = 125$ mm **b** $AB = 184$ mm **c** $BC = 108$ mm **d** $BC = 210$ mm

Many tradespeople work together constructing a house. Use the measurements on this diagram of a basic house for questions **5**, **6** *and* **7**. *It is not drawn to scale.*



5 The house roof is tiled and the verandah has a steel roof. If the builder uses 10 tiles/m^2 calculate the number of tiles needed for this house, to the nearest 10 tiles. You will need to calculate the value of w.

6 The rainfall catchment area of any roof is equal to the ground area that the roof covers. Calculate the total possible rainfall catchment area of this house, to the nearest 10 m^2.

7 The litres of rainfall that a house roof can collect per year is given by the formula:

volume of rain in litres = annual rainfall in mm × roof catchment area in m^2 × percentage of run-off

Copy and complete the following Excel spreadsheet to calculate the average annual volume of water available from this house situated in various locations around Australia. Use your answer to Question **6** for the catchment area.

Recall that 1 mm of rain over an area of 1 m^2 equals 1 litre of rain.


	A	B	C	D	E
1	**Average annual volume of water saved from house roof catchments**				
2	Location	Annual average rainfall in mm	Rainfall catchment area in m^2	percentage run-off	Volume of rain in litres
3	Strahan, TAS	1521		90	
4	Melbourne, VIC	620		90	
5	Port Lincoln, SA	508		90	
6	Perth, WA	728		90	
7	Newcastle, NSW	1200		90	
8	Tully, QLD	4105		90	
9	Alice Springs, NT	243		90	

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Modelling

Viewing angle

Lucas is keen to maximise his experience at the outdoor cinema and usually sits at a distance of 20 metres horizontally from the screen. The base of the screen is 3 metres above the ground and the screen is 5 metres in height. Lucas' eye level is 1 metre above ground level as shown.




Present a report for the following tasks and ensure that you show clear mathematical workings and explanations where appropriate. Use two decimal places for all rounding.


1 Preliminary task

a What is the vertical distance between Lucas' eye level and the base of the screen?

b Find the angle of elevation (α) from Lucas' eye level to the base of the screen.



c Find the angle of elevation (β) from Lucas' eye level to the top of the screen.



d Find Lucas' viewing angle (θ).

2 Modelling task

Formulate

The problem is to determine how close Lucas should sit to maximise his viewing angle.

a Write down all the relevant information that will help solve this problem, and illustrate using diagrams.

Solve

b Determine Lucas' viewing angle if the 20 metre horizontal distance is changed to:

i 14 metres **ii** 8 metres **iii** 2 metres.

c Choose integer values for the horizontal distance between Lucas and the screen and determine which integer horizontal distance delivers the largest viewing angle and state the corresponding viewing angle.

Evaluate and verify

d Find the viewing angle for the following horizontal distances:

i your answer from part **c** plus 0.1 metres **ii** your answer from part **c** minus 0.1 metres.

e Do your results from part **d** confirm that you have found a distance which maximises the viewing angle? Explain why.

f Refine your choices for the horizontal distance and determine a distance correct to one decimal place that maximises Lucas' viewing angle.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

g Summarise your results into a spreadsheet, like the one below.

	A	B	C	D	E	F
1	distance	h1	α	h2	β	θ
2	20	2		7		
3						
4						

h Draw a scale diagram illustrating where Lucas should sit to maximise his viewing angle. Decide if this position for Lucas is reasonable.

3 Extension question

a If the screen is lowered to 2 metres above the ground, rather than the current height of 3 metres, investigate how this changes the position that Lucas should sit to maximise his viewing angle.



Essential Mathematics Core for the Victorian Curriculum Year 9 ISBN 978-1-009-59315-1 © Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Technology and computational thinking

Generating Pythagorean triples

Key technology: Spreadsheets and programming

Pythagorean triples are sets of three positive integers which satisfy Pythagoras' theorem. Perhaps the most well-known example is (3, 4, 5), which is the triple with the smallest numbers such that $3^2 + 4^2 = 5^2$. The triple (3, 4, 5) is also a primitive (or base) triple from which a family of triples can be generated. The family is generated by multiplying all numbers in the triple by a positive integer. So by multiplying by 2 and 3, for example, we generate (6, 8, 10) and (9, 12, 15). Similarly (5, 12, 13) is a primitive triple for a different family of Pythagorean triples. It is useful to know some simple Pythagorean triples and we know that a triangle which uses these numbers as their side lengths will contain a right angle.



1 Getting started

a Decide if the following sets of three numbers form a Pythagorean triple. That is, test if the three numbers satisfy the rule $a^2 + b^2 = c^2$.

i (4, 5, 6) **ii** (6, 8, 10) **iii** (9, 15, 17)
iv (7, 12, 13) **v** (7, 24, 25)

b Use the following primitive triples to write down five other triples.

i (3, 4, 5) **ii** (5, 12, 13) **iii** (8, 15, 17)

2 Applying an algorithm

Pythagorean triples (a, b, c) can be generated using the following rules where m and n are positive integers and $n < m$.


$$a = m^2 - n^2$$
$$b = 2mn$$
$$c = \sqrt{a^2 + b^2}$$

a Using the following values of m and n and the above formulas generate the triples (a, b, c).

i $m = 2$ and $n = 1$
ii $m = 3$ and $n = 1$
iii $m = 3$ and $n = 2$

b The flowchart shown sets out the algorithm for generating Pythagorean triples for m and n less than 10. Run through the algorithm step by step writing the output (a, b, c) into this table.

a	b	c
3	4	5
8	6	



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3 Using technology

This spreadsheet uses the previous formulas to generate Pythagorean triples.

	A	B	C	D
1	**Pythagorean triples**			
2				
3	***m* value**	5		
4				
5	***n***	***a***	***b***	***c***
6	1	=B$3^2-A6^2	=2*B$3*A6	=SQRT(B6^2+C6^2)
7	=A6+1			

a Try entering the formulas into a spreadsheet. Note that the m value can be entered into cell B3.

b Fill down at cells A7, B6, C6 and D6 to generate the triples. Fill down to $n = 4$.


c Alter your m value and fill down once again so that the highest n value is one less than the chosen m value.

d Now alter your m values and try to generate all Pythagorean triples such that all a, b and c are less than 100. How many can you find? Check the internet to confirm your answer.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


Puzzles and games

1 How many right-angled triangles are there in this diagram?




2 Look at this right-angled triangle and the squares drawn on each side. Each square is divided into smaller sections.

a Can you see how the parts of the two smaller squares would fit into the larger square?
b What is the area of each square if the side lengths of the right-angled triangle are a, b and c as marked?
c What do the answers to the above two questions suggest about the relationship between a, b and c?




3 Imagine trying to cut the largest square from a circle of a certain size and calculating the side length of the square.

a If the circle has a diameter of 2 cm, can you find a good position to draw the diameter that also helps to form a right-angled triangle?
b Can you determine the side length of the largest square if the circle has a diameter of 2 cm?
c What percentage of the area of a circle does the largest square occupy?



4 Which is a better fit: a square peg in a round hole or a round peg in a square hole? Use area calculations and percentages to investigate.

5 Two circles of radius 10 cm and 15 cm respectively are placed inside a square. Find the perimeter of the square to the nearest centimetre.





Hint for Q5: First find the diagonal length of the square using the diagram shown.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter summary

Pythagoras' theorem and trigonometry

Finding the hypotenuse

$c^2 = 2^2 + 1^2$
$= 5$
$c = \sqrt{5}$
$= 2.24$ (to 2 d.p.)

2, 1, c

Shorter sides

$a^2 + 5^2 = 7^2$
$a^2 = 7^2 - 5^2$
$= 24$
$a = \sqrt{24}$ (exact)

7, a, 5

Applications ⭐

a, c, b; a, c, b

Pythagoras' theorem

a, c, b

$c^2 = a^2 + b^2$

Finding side lengths (x in the numerator)

x cm, 22°, 7 cm

$\cos 22° = \frac{x}{7}$
$x = 7 \times \cos 22°$
$= 6.49$ (to 2 d.p.)

SOHCAHTOA

5 (H), θ, 3 (A), 4 (O)

$\sin\theta = \frac{O}{H} = \frac{4}{5}$

$\cos\theta = \frac{A}{H} = \frac{3}{5}$

$\tan\theta = \frac{O}{A} = \frac{4}{3}$

Finding side lengths ⭐ (x in the denominator)

36°, x m, 5 m

$\tan 36° = \frac{5}{x}$
$x \times \tan 36° = 5$
$x = \frac{5}{\tan 36°}$
$= 6.88$ (to 2 d.p.)

Applying trigonometry ⭐

30 m, θ, 40 m

$\tan\theta = \frac{30}{40}$

$\theta = \tan^{-1}\left(\frac{30}{40}\right)$

$= 36.87°$ (to 2 d.p.)

Elevation and depression ⭐

Depression

Elevation

Finding angles

6, θ, 7.5

$\sin\theta = \frac{6}{7.5}$

$\theta = \sin^{-1}\left(\frac{6}{7.5}\right)$











$= 53.13°$
(to 2 d.p.)

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.








Chapter checklist

Chapter checklist

A version of this checklist that you can print out and complete can be downloaded from your Interactive Textbook.

		✔
4A	**1 I can write Pythagoras' theorem for a right-angled triangle.** e.g. Write down Pythagoras' theorem for these triangles **a**  **b** 	
4B	**2 I can find the length of the hypotenuse.** e.g. Find the length of the hypotenuse in these right-angled triangles, rounding to two decimal places where necessary. **a**  **b** 	
4B	**3 I can find the length of the hypotenuse as a surd.** e.g. Find the length of the hypotenuse in this right-angled triangle, leaving your answer as an exact value. 	
4C	**4 I can find the length of a shorter side.** e.g. Find the value of the pronumeral, rounding to two decimal places. 	
4C	**5 I can find the exact length of a shorter side in a right isosceles triangle.** e.g. Find the value of x, giving your answer as a surd. 	
4D	**6 I can identify right-angled triangles in a diagram to find unknown lengths.** e.g. An apartment courtyard is in the shape of a trapezium, as shown. Find the unknown side length correct to two decimal places by first drawing a suitable right-angled triangle. 	
4D	**7 I can apply Pythagoras' theorem.** e.g. Two poles are 10 m apart. The heights of the poles are 8 m and 12 m. If a bird flies from the top of one pole in a direct line to the top of the other pole, how far does the bird fly correct to two decimal places? 	
4E	**8 I can label the sides of a triangle.** e.g. Copy this triangle and label the sides as opposite to θ (O), adjacent to θ (A) and hypotenuse (H). 	

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

		✔
4E	**9 I can write trigonometric ratios for right-angled triangles.** e.g. Write a trigonometric ratio (in fraction form) for each of the following triangles. **a**  **b** 	
4F	**10 I can find trigonometric values on a calculator.** e.g. Use a calculator to evaluate tan $35°$ correct to two decimal places.	
4F	**11 I can find an unknown side length using trigonometry.** e.g. Find the value of x in the triangle correct to two decimal places. 	
4G	**12 I can find an unknown side length in the denominator of a trigonometric ratio.** e.g. Find the value of x correct to two decimal places. 	
4G	**13 I can find two side lengths using trigonometry.** e.g. Find the value of x and y correct to two decimal places. 	
4H	**14 I can use inverse trigonometric ratios.** e.g. Find the value of θ to the nearest degree if: **a** $\cos\theta = 0.6261$ **b** $\sin\theta = \frac{3}{7}$.	
4H	**15 I can find an angle using trigonometry.** e.g. Find the value of θ to the nearest degree. 	
4I	**16 I can use angles of elevation.** e.g. The angle of elevation of the top of a light tower from a point on the ground 24 m away from the base of the tower is $16°$. Find the height of the light tower to the nearest metre. 	
4I	**17 I can work with angles of depression.** e.g. From the surface of the water, a snorkeller spots a ray below the water. If the surface of the water is 3 m above the ray and the ray is 5 m horizontally from the snorkeller, find the angle of depression from the snorkeller to the ray to the nearest degree.	


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter review


Short-answer questions

4A **1** Write down Pythagoras' theorem for these triangles, using the given pronumerals.


a



b




c




4B **2** Find the length of the hypotenuse in these triangles. Round to two decimal places where appropriate.


a




b




c




d



e




f




4C **3** Find the length of the unknown side in these right-angled triangles. Round to two decimal places.


a




b



c




4D **4** A steel support beam of length 6.5 m is connected to a wall at a height of 4.7 m from the ground. Find the distance (to two decimal places) between the base of the building and the point where the beam is joined to the ground.



4D **5** For this double triangle, find:

a the length AC

b the length CD (correct to two decimal places)



4D **6** Two different cafés on opposite sides of an atrium in a shopping centre are respectively 10 m and 15 m above the ground floor. If the cafés are linked by a 20 m escalator, find the horizontal distance (to the nearest metre) across the atrium, between the two cafés.




4E **7** Find the value of each of the following, correct to two decimal places.

a $\sin 40°$ **b** $\tan 66°$ **c** $\cos 44°$


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

4E **8** Which ratio $\left(\sin\theta = \frac{O}{H},\ \cos\theta = \frac{A}{H} \text{ or } \tan\theta = \frac{O}{A}\right)$ would be used to find the value of the unknown in these triangles?


a



b




c




4F/G/H **9** Find the value of each pronumeral, correct to two decimal places where necessary.
Use $\sin\theta = \frac{O}{H}$, $\cos\theta = \frac{A}{H}$ or $\tan\theta = \frac{O}{A}$.


a




b




★**c**




★**d**




e




f



4I ★ **10** The angle of elevation of the top of a lighthouse from a point on the ground 40 m from its base is 35°. Find the height of the lighthouse, to two decimal places.




4I ★ **11** A train travels up a slope, making an angle of 7° with the horizontal. When the train is at a height of 3 m above its starting point, find the distance it has travelled up the slope, to the nearest metre.




4I ★ **12** From a point on the ground, Ahbed measures the angle of elevation of a 120 m tower to be 34°. How far from the base of the tower is Ahbed, correct to two decimal places?

4I ★ **13** Aisha is standing on the roof of a skyscraper when she spots a car at an angle of depression of 51° below her. If the skyscraper is 78 m high, how far away is the car from the base of the skyscraper, correct to one decimal place?




Multiple-choice questions

4A **1** For the right-angled triangle shown, which is a correct statement?



A $c^2 = 5^2 + 12^2$ **B** $c^2 = 5^2 - 12^2$ **C** $c^2 = 12^2 - 5^2$
D $c^2 = 5^2 \times 12^2$ **E** $(5 + 12)^2$

4A **2** Which of the following is true for this triangle?




A $a = 5 - 4$ **B** $a^2 = 5^2 - 4^2$ **C** $a^2 = 5^2 + 4^2$
D $a = \sqrt{10}$ **E** $a^2 - 4^2 = 5^2$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter review


4C **3** For the right-angled triangle shown, which is a correct statement.



A $2x^2 = 49$ **B** $7x^2 = 2$ **C** $2x^2 = 7$

D $x^2 + 7^2 = x^2$ **E** $x^2 = \frac{2}{7}$

4E **4** For the triangle shown, which is a correct statement.



A $\sin\theta = \frac{a}{b}$ **B** $\sin\theta = \frac{c}{a}$ **C** $\sin\theta = \frac{a}{c}$


D $\sin\theta = \frac{b}{c}$ **E** $\sin\theta = \frac{c}{b}$

4E **5** The value of cos 46°, correct to four decimal places, is:

A 0.7193 **B** 0.6947 **C** 0.594

D 0.6532 **E** 1.0355


4F **6** In the diagram, the value of x, correct to two decimal places, is:



A 40 **B** 13.61 **C** 4.70

D 9.89 **E** 6.47


4G **7** The value of x in the triangle is given by:



A $8\sin 46°$ **B** $8\cos 46°$ **C** $\frac{8}{\cos 46°}$

D $\frac{8}{\sin 46°}$ **E** $\frac{\cos 46°}{8}$

4H **8** The value of θ in this triangle is given by:



A $\tan\left(\frac{4}{5}\right)$


B $\tan\left(\frac{5}{4}\right)$

C $\tan^{-1}\left(\frac{5}{4}\right)$

D $\sin^{-1}\left(\frac{4}{5}\right)$

E $\tan^{-1}\left(\frac{4}{5}\right)$

4I **9** A ladder is inclined at an angle of 28° to the horizontal. If the ladder reaches 8.9 m up the wall, the length of the ladder correct to the nearest metre is:



A 19 m


B 4 m

C 2 m

D 10 m

E 24 m

4H **10** The value of θ in the diagram, correct to two decimal places, is:



A 0.73°

B 48.19°

C 41.81°

D 33.69°

E 4.181°

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Extended-response questions


1 From the top of a 100 m cliff, Skevi sees a boat out at sea at an angle of depression of 12°.

- **a** Draw a simple diagram for this situation.
- **b** Find how far out to sea the boat is to the nearest metre.
- **c** A swimmer is 2 km away from the base of the cliff and in line with the boat. What is the angle of depression to the swimmer to the nearest degree?
- **d** How far away is the boat from the swimmer, to the nearest metre?



2 An extension ladder is initially placed so that it reaches 2 metres up a wall. The foot of the ladder is 80 centimetres from the base of the wall.

- **a** Find the length of the ladder, to two decimal places, in its original position.
- **b** Without moving the foot of the ladder, it is extended so that it reaches 1 m further up the wall. How far (to two decimal places) has the ladder been extended?
- **c** The ladder is placed so that its foot is now 20 cm closer to the base of the wall. How far up the wall can the ladder length found in part **b** reach? Round to two decimal places.





ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5 Linear relationships



Essential mathematics: why skills with linear relationships are important

Linear relationships are fundamental in understanding how different variables interact with each other in a straight-line pattern.

Farmers use linear equations to manage crop yields efficiently. To predict the amount of fertiliser needed based on the field's size, they might use the equation $F = k \times A$, where F is the fertiliser amount, k is the fertiliser rate per unit area, and A is the field size. By applying these linear relationships, farmers can optimise resource use, improve yields, and maintain sustainable agricultural practices.

Event planners use linear equations to ensure they order the right amount of equipment for varying numbers of attendees. They can calculate E, the total amount of equipment needed, using the equation $E = e \times A$, where e is the equipment required per attendee, and A is the number of attendees. By calculating E, planners can ensure they have enough chairs, tables, and safety equipment, like fire extinguishers and first aid kits. Accurate calculations prevent shortages or excesses, ensuring smooth event operations.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.



In this chapter

5A Introduction to linear relationships
5B Finding x- and y-intercepts
5C Graphing straight lines using intercepts
5D Lines with one intercept
5E Gradient
5F Gradient and direct proportion
5G Gradient-intercept form
5H Finding the equation of a line ★
5I Midpoint and length of a line segment
5J Linear modelling ★

Victorian Curriculum 2.0

This chapter covers the following content descriptors in the Victorian Curriculum 2.0:

ALGEBRA

VC2M9A03, VC2M9A04, VC2M9A06, VC2M9A07

MEASUREMENT

VC2M9M05

Please refer to the curriculum support documentation in the teacher resources for a full and comprehensive mapping of this chapter to the related curriculum content descriptors.

© VCAA

Online resources


A host of additional online resources are included as part of your Interactive Textbook, including HOTmaths content, video demonstrations of all worked examples, auto-marked quizzes and much more.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025
Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Warm-up quiz

1 Plot and label the following points on a Cartesian plane (x- and y- axes).

a $(3, 1)$ **b** $(-2, 4)$ **c** $(0, 0)$ **d** $(5, -3)$



2 Given $y = x - 2$, find the value of y. when:

a $x = 3$ **b** $x = 1$ **c** $x = -1$ **d** $x = -2$

3 Solve the following equations for x.

a $3x = 15$ **b** $-2x = 4$ **c** $-x = 3$

d $0 = x - 4$ **e** $0 = 3x + 6$ **f** $0 = 2x - 8$

4 Find the value of y in each of the following when $x = 0$.

a $y = 2x + 3$ **b** $y = 3x - 4$ **c** $x + 2y = 8$ **d** $3x - 4y = 12$

5 Complete the table of values for these rules.

a $y = x + 3$ **b** $y = 2x - 1$

	x	−2	−1	0	1	2
a	y					
b	y					

6 If $x = 2$ and $y = 1$, decide whether the following equations are true.

a $y = 2x + 1$ **b** $y = 3x - 5$ **c** $y = -2x + 5$ **d** $5x - 2y = 6$

7 Simplify:

a $\frac{4 + 10}{2}$ **b** $\frac{2 + 11}{2}$ **c** $\frac{-1 + 7}{2}$ **d** $\frac{-3 + (-7)}{2}$

8 Find the vertical distance between the following pairs of points.

a $(3, 2)$ and $(3, 7)$ **b** $(-1, 1)$ and $(-1, -3)$

c $(2, 3)$ and $(2, -2)$ **d** $(1, -4)$ and $(1, -1)$

9 Find the horizontal distance between the following pairs of points.

a $(1, 3)$ and $(5, 3)$ **b** $(-1, 2)$ and $(4, 2)$

c $(-2, -3)$ and $(5, -3)$ **d** $(-4, 1)$ and $(-1, 1)$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5A Introduction to linear relationships

Learning intentions

- To review the features of the Cartesian plane
- To know that a linear relation is a set of coordinates that form a straight line when graphed
- To be able to complete a table of values and plot points to form a linear graph
- To know the general forms of a linear relation
- To understand that for a point to be on a line its coordinates must satisfy the rule for the linear relation
- To be able to decide if a point is on a line using its rule

Key vocabulary: Cartesian plane, x-axis, y-axis, linear relation, coordinates (or coordinate pair), origin, x-coordinate, y-coordinate

If two variables are related in some way, we can use mathematical rules to more precisely describe this relationship. The most simple type of relationship is one that can be illustrated with a straight line graph. These are called linear relations.

The volume of petrol in your car at a service bowser, for example, might initially be 10 L, then increase by 1.2 L per second after that. This is an example of a linear relationship between *volume* and *time* because the volume is increasing at a constant rate of 1.2 L/s.



If the fuel bowser delivers petrol at a constant rate, there is a linear relationship between the volume of petrol in the car's tank and time.

Lesson starter: Is it linear?

Here are three rules linking x and y.


1 $y_1 = \frac{2}{x} + 1$

2 $y_2 = x^2 - 1$

3 $y_3 = 3x - 4$

First complete this simple table and plot the points on the graph.

x	1	2	3
y_1			
y_2			
y_3			




- Which of the three rules do you think is linear?
- How do the table and graph help you decide it's linear?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5A

Key ideas

- Coordinate geometry provides a link between geometry and algebra.
- The **Cartesian plane** (or number plane) consists of two axes that divide the number plane into four quadrants.
  - The **x-axis** is the horizontal axis.
  - The **y-axis** is the vertical axis.
  - The x-axis and y-axis intersect at the **origin** (O) at right angles.
  - A point is precisely positioned on a Cartesian plane using the **coordinate pair** (x, y).
    x (the **x-coordinate**) describes the horizontal position of the point from the origin.
    y (the **y-coordinate**) describes the vertical position of the point from the origin.



- In the coordinate pair $(2, -4)$, 2 is the x-coordinate and -4 is the y-coordinate. The point is 2 units to the right of the origin (horizontal direction) and 4 units below the origin (vertical direction).
- A **linear relation** is a set of ordered pairs (x, y) that, when graphed, give a straight line.
- Linear relations have rules that may be of the form:
  - $y = mx + c$ (or $y = mx + b$), for example, $y = 2x + 1$
  - $ax + by = d$ or $ax + by + c = 0$, for example, $2x - 3y = 4$ or $2x - 3y - 4 = 0$
- Each point that is on the line fits the rule for the linear relation. For example, on the line with rule $y = 2x$, each y-coordinate will be two times the x-coordinate. So, $(3, 6)$ will be on the line but $(4, 10)$ will not.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


Exercise 5A

Understanding

1–4 | 2, 4

1 **a** For the Cartesian plane shown, label the following parts.

i x-axis
ii origin
iii y-axis
iv coordinate pair



b For the point $(3, -1)$:


i 3 is the _____.
ii -1 is the _____.
iii From the origin, the point is 3 units to the _____ and 1 unit _____.

2 Refer to the Cartesian plane shown.

a Draw this set of axes (without A–F) and plot the following points.

i $(3, 1)$
ii $(-4, -2)$
iii $(0, 2)$
iv $(-1, 2)$

b Give the coordinates of all the points A–F.





Hint for Q2b: List the x-coordinate followed by the y-coordinate; e.g. $(2, 4)$

3 Which one of the following rules is linear?

A $y = x^2 + 4$ **B** $y = \frac{2}{x}$ **C** $y = 2x - 1$ **D** $y = 3^x$



Hint for Q3: Linear rules are often in the form $y = mx + c$ or $ax + by = d$.

4 In the rule $y = 3x$, the y-coordinate of each point on the line is 3 times the x-coordinate. Which point(s) lie on this line?

A $(1, 3)$ **B** $(3, 6)$ **C** $(-3, -9)$ **D** $(-1, 3)$

Fluency

5, 6 | 5, 6

Example 1 Filling in tables from rules

Complete the tables for the given rules.

a $y = 2x - 1$

x	-2	-1	0	1	2	3
y						

b $y = -3x + 2$

x	-2	-1	0	1	2
y					

Continued on next page

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5A

Solution

a $y = 2x - 1$

x	-2	-1	0	1	2	3
y	-5	-3	-1	1	3	5

Explanation

Substitute each x value in the table into the rule $y = 2x - 1$ to find its corresponding y value.

For $x = -2$, $y = 2 \times (-2) - 1$
$= -4 - 1$
$= -5$

For $x = 1$, $y = 2 \times 1 - 1$
$= 2 - 1$
$= 1$

b $y = -3x + 2$

x	-2	-1	0	1	2
y	8	5	2	-1	-4

Substitute each x value into the rule $y = -3x + 2$.
Recall that negative × negative = positive.

For $x = -2$, $y = -3 \times (-2) + 2$
$= 6 + 2$
$= 8$

For $x = 0$, $y = -3 \times 0 + 2$
$= 0 + 2$
$= 2$

Now you try

Complete the table for the given rules.

a $y = 3x - 2$
b $y = -2x + 4$

x	-2	-1	0	1	2
y					

5 Complete the tables for the given rules.

a $y = x + 2$

x	-2	-1	0	1	2	3
y						

b $y = 2x + 3$

x	-2	-1	0	1	2
y					

c $y = -2x$

x	-2	-1	0	1	2
y					

d $y = -3x - 1$

x	-2	-1	0	1	2	3
y						



Hint for Q5: A negative × positive is negative. A negative × negative is positive.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Example 2 Plotting points to graph straight lines


Using $-3 \leq x \leq 3$, construct a table of values and plot a graph for these linear relations.

a $y = x + 2$ **b** $y = -2x + 2$

Solution

a

x	-3	-2	-1	0	1	2	3
y	-1	0	1	2	3	4	5



Explanation

Use $-3 \leq x \leq 3$ to construct a table of x values as instructed. Substitute each value of x into the rule $y = x + 2$.

The coordinates of the points are read from the table; i.e. $(-3, -1)$, $(-2, 0)$ etc.


Draw a set of axes with an appropriate scale to show the points from the table.

Plot each point and join to form a straight line.

Extend the line to show it continues in either direction.

b

x	-3	-2	-1	0	1	2	3
y	8	6	4	2	0	-2	-4



Use $-3 \leq x \leq 3$ as instructed. Substitute each value of x into the rule $y = -2x + 2$.

Plot each point and join to form a straight line.

Extend the line beyond the plotted points.

Now you try

Using $-3 \leq x \leq 3$, construct a table of values and plot a graph for these linear relations.

a $y = x - 2$ **b** $y = -3x + 4$

6 Using $-3 \leq x \leq 3$, construct a table of values and plot a graph for these linear relations.

a $y = x - 1$ **b** $y = x + 3$ **c** $y = 2x - 3$

d $y = -3x$ **e** $y = -2x - 1$ **f** $y = -x + 4$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5A

Problem-solving and reasoning 7–9 8–11

Example 3 Deciding whether a point is on a line

Decide whether the point $(3, 7)$ is on the line with the given rule.

a $y = 3x - 4$ **b** $y = 2x + 1$

Solution

a $y = 3x - 4$
Substitute $x = 3$
$y = 3 \times 3 - 4$
$= 5$ (not 7)
$\therefore$ the point $(3, 7)$ is not on the line

b $y = 2x + 1$
Substitute $x = 3$
$y = 2 \times 3 + 1$
$= 7$
$\therefore$ the point $(3, 7)$ is on the line

Explanation

Find the value of y on the graph of the rule for $x = 3$.

To fit the rule, the y value needs to be 5, not 7, so $(3, 7)$ is not on the line.

By substituting $x = 3$ into the rule for the line, we find $y = 7$.

Since $(3, 7)$ fits the rule, it is on the line.

Now you try

Decide whether the point $(2, 10)$ is on the line with the given rule.

a $y = 2x + 6$ **b** $y = 5x - 4$

7 Decide whether the point $(2, 8)$ is on the line with the given rule.
a $y = 2x + 6$ **b** $y = 3x + 2$ **c** $y = -x + 6$.



Hint for Q7: When $x = 2$, does $y = 8$?

8 Decide whether the point $(-1, 4)$ is on the line with the given rule.
a $y = x + 5$ **b** $y = -2x + 2$ **c** $y = 4x$

9 Find a rule in the form $y = mx + c$ (e.g. $y = 2x - 1$) that matches these tables of values.

a

x	0	1	2	3	4
y	2	3	4	5	6

b

x	0	1	2	3
y	0	2	4	6

c

x	-2	-1	0	1	2	3
y	-3	-1	1	3	5	7

d

x	-2	-1	0	1	2	3
y	-7	-4	-1	2	5	8

e

x	-1	0	1	2	3	4
y	-3	-2	-1	0	1	2



Hint for Q9: Think about how you can find the y value from the x value. Is it double the x value each time or 3 more than the x value each time? etc.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10 The table below shows the recorded height, y cm, of a seedling x days after it sprouts. Find a linear rule in the form $y = \ldots$ that matches the data in the table.




x (day)	1	2	3	4
y (height)	3	6	9	12

11 By considering the relationship between the coordinates, match these rules **i**, **ii**, **iii** and **iv** to the graphs **a**, **b**, **c** and **d**.


i $y = 2x + 1$ **ii** $y = -x - 1$

iii $y = -2x + 3$ **iv** $x + y = 2$


a




b



c



d





Hint for Q11: Each rule shows the relationship between the x- and y-coordinates for each point on the line. In i, each y-coordinate has to be 1 more than twice its x-coordinate.

Tougher rule finding — 12

12 Find the linear rule linking x and y in these tables.

a

x	−1	0	1	2	3
y	5	7	9	11	13

b

x	−2	−1	0	1	2
y	22	21	20	19	18

c

x	1	3	5	7	9
y	1	5	9	13	17

d

x	6	7	8	9	10
y	4	4.5	5	5.5	6

e

x	0	2	4	6	8
y	4	−2	−8	−14	−20

f

x	−5	−4	−3	−2	−1
y	−21	−16	−11	−6	−1

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5B Finding x- and y-intercepts


Learning intentions

- To know which points are represented by the x-intercept and y-intercept
- To be able to identify the x-intercept and y-intercept from a table or graph
- To know how to find the x-intercept and the y-intercept of a linear relation

Key vocabulary: x-intercept, y-intercept, coordinates, x-axis, y-axis

Each point on a line satisfies a rule for a linear relation. When the linear relation is a model for a real situation, each point gives a piece of information about that situation. Some points are very significant.

For example, the graph below for the linear relation $y = -0.2x + 40$ models how much petrol a car uses. The y-axis shows the amount of petrol in the tank and the x-axis shows the number of kilometres travelled. The coordinate pair $(50, 30)$ shows that after travelling 50 km, 30 litres of petrol are left in the tank.





A car's dashboard display shows the number of kilometres until the car runs out of petrol, the x-intercept.

One important point shows us how much petrol was in the tank before the car has started the journey. Using the linear relation (substituting $x = 0$), we can see that this point is $(0, 40)$. This is where the graph cuts the y-axis and is called the y-intercept.


We also want to know when the car will run out of petrol. Again using the linear relation (substituting $y = 0$), we can see that this point is $(200, 0)$. This is where the graph cuts the x-axis and is called the x-intercept.

Lesson starter: Are we there yet?

This table shows the distance (y metres) a student is from his home (at x minutes) when he rides his bike home from school.

x	0	2	4	6	8	10
y	2000	1600	1200	800	400	0

- Copy the axes shown and plot these points to form a line.
- How far is school from home?
- How long does it take to get home?
- Circle the above two points on the graph.
- Which point is the x-intercept? Which point is the y-intercept?
- If the rule for the table is $y = -200x + 2000$, how could you find the x- and y-intercepts without sketching the graph?



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


Key ideas

- The y-intercept is the point where the line cuts (intersects) the y-axis. It always has an x-coordinate of 0.
- The x-intercept is the point where the line cuts (intersects) the x-axis. It always has a y-coordinate of 0.

x-intercept

x	-2	-1	0	1	2	3
y	8	6	4	2	0	-2

y-intercept



- The rule of the linear relation can be used to find the x- and y-intercepts.
  - To find the y-intercept, substitute $x = 0$ into the rule, since it lies on the y-axis.
  - To find the x-intercept, substitute $y = 0$ into the rule, since it lies on the x-axis.

Exercise 5B

Understanding

1–3 | 1, 3


1 Fill in the missing part of these sentences. Choose from x-intercept, x-coordinate or x-axis.



a The x-intercept is the point where the graph cuts the _____.

b The _____ of the y-intercept is always 0.


c The point on a line with coordinates $(4, 0)$ is the _____.



2 For the Cartesian plane shown:

a give the coordinates of each point that is on the x-axis

b give the coordinates of each point that is on the y-axis



3 Copy and complete the following

a The x-intercept is found in a rule by substituting _____.

b The y-intercept is found in a rule by substituting _____.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5B

Fluency

4, 5, 6(½) | 4–6(½)


Example 4 Reading off the x-intercept and y-intercept

Read off the coordinates of the x-intercept and y-intercept from this table and graph.

a

x	−2	−1	0	1	2	3
y	12	9	6	3	0	−3

b



Solution

a The x-intercept is at $(2, 0)$.
The y-intercept is at $(0, 6)$.

b The x-intercept is $(-4, 0)$.
The y-intercept is at $(0, 5)$.

Explanation

The x-intercept is at the point where $y = 0$ (on the x-axis).
The y-intercept is at the point where $x = 0$ (on the y-axis).

The x-intercept is at the point on the x axis $(y = 0)$.
The y-intercept is at the point on the y axis $(x = 0)$.


Now you try

Read off the coordinates of the x-intercept and y-intercept from this table and graph.

a

x	−2	−1	0	1	2	3
y	−2	0	2	4	6	8

b



4 Read off the coordinates of the x- and y-intercepts from these tables and graphs.

a

x	−3	−2	−1	0	1	2	3
y	4	3	2	1	0	1	2

b

x	−3	−2	−1	0	1	2	3
y	−1	0	1	2	3	4	5

c

x	−1	0	1	2	3	4
y	10	8	6	4	2	0


d

x	−5	−4	−3	−2	−1	0
y	0	2	4	6	8	10




Hint for Q4: The x-intercept is where $y = 0$. The y-intercept is where $x = 0$.

e




f




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

g



h



Example 5 Finding the y-intercept

Find the coordinates of the y-intercept for these linear relations.

a $y = 2x - 1$ **b** $2x + 3y = 6$

Solution

a $y = 2x - 1$
y-intercept (let $x = 0$):
$y = 2 \times (0) - 1$
$= -1$
$\therefore$ the y-intercept is at $(0, -1)$

Explanation

Substitute $x = 0$ into the rule to find the y value of the y-intercept.

Recall that anything multiplied by 0 is 0.

Using coordinates the y-intercept is at $(0, -1)$.

b $2x + 3y = 6$
y-intercept (let $x = 0$):
$2 \times (0) + 3y = 6$
$3y = 6$
$y = 2$
$\therefore$ the y-intercept is at $(0, 2)$

Substitute $x = 0$ to find the y-intercept.

Solve the equation $3y = 6$ by dividing both sides by 3.

Using coordinates the y-intercept is at $(0, 2)$.

Now you try


Find the coordinates of the y-intercept for these linear relations.

a $y = 3x + 5$ **b** $3x + 4y = 12$

5 Find the coordinates of the y-intercept for these linear relations.

a $y = 2x + 5$ **b** $y = 3x + 1$
c $y = x - 7$ **d** $y = 2x - 3$
e $y = -4x + 2$ **f** $y = -5x - 4$
g $y = -x - 2$ **h** $y = -2x + 10$
i $2x + y = 11$ **j** $2x + 3y = 9$
k $x + 4y = 8$ **l** $-3x + 2y = 12$
m $4x - 2y = 6$ **n** $3x - 4y = 4$
o $-2x - y = 3$ **p** $-2x - 3y = -9$





ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5B

Example 6 Finding the x-intercept

Find the coordinates of the x-intercept for these linear relations.

a $y = 2x + 4$ **b** $3x - 2y = 12$

Solution

a $y = 2x + 4$

x-intercept (let $y = 0$):

$0 = 2x + 4$

$-4 = 2x$

$-2 = x$

$x = -2$

$\therefore$ the x-intercept is at $(-2, 0)$

Explanation

Substitute $y = 0$ into the rule to find the x-coordinate of the x-intercept.

Solve the equation by subtracting 4 from both sides. Then divide both sides by 2.

Using coordinates the x-intercept is at $(-2, 0)$.

b $3x - 2y = 12$

x-intercept (let $y = 0$):

$3x - 2 \times (0) = 12$

$3x = 12$

$x = 4$

$\therefore$ the x-intercept is at $(4, 0)$

Substitute $y = 0$ to find the x-intercept.

$2 \times 0 = 0$ and then solve the remaining equation for x.

Using coordinates the x-intercept is at $(4, 0)$.

Now you try

Find the coordinates of the x-intercept for these linear relations.

a $y = 3x - 6$ **b** $5x + 4y = -15$

6 Find the coordinates of the x-intercept for these linear relations.

a $y = x + 6$ **b** $y = x - 4$ **c** $y = 2x - 8$

d $y = 3x + 6$ **e** $y = -2x - 8$ **f** $y = -4x + 12$

g $x + 2y = 7$ **h** $2x - y = 4$ **i** $3x + 4y = 12$

j $4x - 6y = 24$ **k** $-2x + 3y = -2$ **l** $-5x - 7y = 15$

Problem-solving and reasoning 7, 8 7–10

7 Find the coordinates of the x-intercept and y-intercept for these linear relations involving fractions. Leave coordinates in fraction form.

a $y = 2x - 1$ **b** $y = \frac{1}{2}x - 2$

c $y = -\frac{1}{3}x + 4$ **d** $3x + 2y = 7$



Hint for Q7b: Recall that another way of writing $\frac{1}{2}x$ is $\frac{x}{2}$.

8 The height, h, in metres, of a lift above ground after t seconds is given by $h = 100 - 5t$.

a What height does the lift start at (at $t = 0$)?

b How long does it take for the lift to reach the ground ($h = 0$)?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

9 Water is being drained from a fish tank. The amount of water in the tank, V litres, after t minutes is given by the rule $V = 80 - 4t$.

a How long will it be until the tank is empty ($V = 0$)?

b How much water was in the tank to begin with ($t = 0$)?



10 The line $y = 3x$ has both its x-intercept and y-intercept with coordinates $(0, 0)$. Can you explain how this is the case, and what it means for the graph of $y = 3x$?

Making money — 11

11 Ana makes badges to sell at the market. The rule for the amount of money, A dollars, she makes from the sale of b badges is given by $A = 2b - 60$.


a What is the value of A if she sells no badges ($b = 0$)?

b Can you explain your answer to part **a**?

c How many badges must she sell to cover her initial costs ($A = 0$)?

d What happens when she sells more badges than in your answer to part **c**?

e Use the information from your answers above to draw a graph for the rule on the axes shown. The A-axis is the y-axis and the b-axis is the x-axis.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5C Graphing straight lines using intercepts

Learning intentions

- To understand that only two points are required to sketch a straight line graph
- To know that the x-intercept and the y-intercept are two key points often used to sketch a straight line graph
- To be able to sketch a linear relation by finding the x- and y-intercepts

Key vocabulary: x-intercept, y-intercept, linear relation

When linear equations are graphed, all the points lie in a straight line. This means that it is possible to graph a straight line using only two points. Two critical points that help draw these graphs are the x-intercept and y-intercept. Once these two points are located, they can be joined to form a straight line illustrating all the other points that lie on the line.




Straight lines are frequently used in business to illustrate the relationship between variables.

Lesson starter: Two key points

Consider the relation $y = 2x - 2$.


- Write a set of steps for another student, explaining to them how to find the y-intercept using the rule.
- Write another set of steps, this time explaining how to find the x-intercept using the rule.
- Now that you have these two points, describe how you would use them to sketch the graph of $y = 2x - 2$ on the axes shown, without drawing up a table of values.



Key ideas

- A straight line can be drawn using only two points.
- Two key points we can use to graph a straight line are the x-intercept and y-intercept.
- To graph a linear relation using intercepts:
  - find the y-intercept by substituting $x = 0$ into the rule
  - find the x-intercept by substituting $y = 0$ into the rule
  - plot the two points on the axes
  - draw a straight line passing through the two intercepts.

For example: $3x + 5y = 15$



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Exercise 5C

Understanding

1–4 | 4

1 Insert the missing number.

a A minimum of _____ points are required to sketch a straight line.

b To find the y-intercept, substitute $x =$ _____.

c If $(2, a)$ is an x-intercept, the value of a is _____.

2 For these equations, find the y-intercept coordinates by letting $x = 0$.

a $2x + 3y = 9$ **b** $y = 2x - 4$

3 For these equations, find the x-intercept coordinates by letting $y = 0$.

a $2x - y = -4$ **b** $y = 3x - 6$

4 Plot each of the following pairs of points on a set of axes and join the points to form a straight line.

a $(3, 0)$ and $(0, -2)$

b $(0, 4)$ and $(2, 0)$

c $(-3, 0)$ and $(0, 6)$

d $(1, 0)$ and $(0, 3)$



Hint for Q4a: $(3, 0)$ is on the x-axis as it is 3 to the right of the origin and 0 up or down.

Fluency

5–6(½) | 5–6(½)

Example 7 Sketching linear relations of the form $ax + by = d$ using intercepts

Sketch the graph of $2x + 3y = 6$, showing the x- and y-intercepts.

Solution

$2x + 3y = 6$

y-intercept (let $x = 0$):

$2 \times (0) + 3y = 6$

$3y = 6$

$y = 2$

$\therefore$ the y-intercept is at $(0, 2)$


x-intercept (let $y = 0$):

$2x + 3 \times (0) = 6$

$2x = 6$

$x = 3$

$\therefore$ the x-intercept is at $(3, 0)$



Explanation

Only two points are required to generate a straight line.
For the y-intercept, substitute $x = 0$ into the rule and solve for y by dividing each side by 3.

State the y-intercept. Plot the point $(0, 2)$.

To find the x-intercept, substitute $y = 0$ into the rule and solve for x.

State the x-intercept. Plot the point $(3, 0)$.

Mark and label the intercepts on the axes and sketch the graph by joining the two intercepts. Continue the line past these points.

Now you try

Sketch the graph of $3x + 4y = 12$, showing the x- and y-intercepts.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5C

5 Sketch the graph of the following relations, by finding the x- and y-intercepts.

a $x + y = 2$ **b** $x + y = 5$ **c** $x - y = 3$
d $x - y = -2$ **e** $2x + y = 4$ **f** $3x - y = 9$
g $4x - 2y = 8$ **h** $3x + 2y = 6$ **i** $3x - 2y = 6$
j $y - 3x = 12$ **k** $-5y + 2x = -10$ **l** $-x + 7y = 21$



Hint for Q5: For the x-intercept, let $y = 0$. For the y-intercept, let $x = 0$.

Example 8 Sketching linear relations of the form $y = mx + c$ using intercepts

Sketch the graph of $y = 2x - 6$, showing the x- and y-intercepts.

Solution

$y = 2x - 6$

y-intercept (let $x = 0$):

$y = 2 \times (0) - 6$

$y = -6$

$\therefore$ the y-intercept is at $(0, -6)$


x-intercept (let $y = 0$):

$0 = 2x - 6$

$6 = 2x$

$x = 3$

$\therefore$ the x-intercept is at $(3, 0)$



Explanation

Substitute $x = 0$ for the y-intercept.
Simplify to find the y-coordinate.

Plot the point $(0, -6)$.

Substitute $y = 0$ for the x-intercept. Solve the remaining equation for x by adding 6 to both sides. Then divide both sides by 2.

Plot the point $(3, 0)$.

Mark in the two intercepts and join to sketch the graph.

Now you try

Sketch the graph of $y = 2x + 4$, showing the x- and y-intercepts.

6 Sketch the graph of the following relations, showing the x- and y-intercepts.

a $y = 3x + 3$ **b** $y = 2x + 2$
c $y = x - 5$ **d** $y = -x - 6$
e $y = -2x - 2$ **f** $y = -3x - 6$
g $y = -2x + 4$ **h** $y = 2x - 3$
i $y = -x + 1$ **j** $y = -2x + 1$



Hint for Q6a: To solve:
$3x + 3 = 0$
$3x = -3$ (subtract 3)
$x = -1$ (divide by 3)

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Problem-solving and reasoning 7, 8 7–9

7 The distance, d metres, of a remote controlled car from an observation point after t seconds is given by the rule $d = 8 - 2t$.

a Find the distance from the observation point initially (at $t = 0$).

b Find after what time, t, the distance, d, is equal to 0 (substitute $d = 0$).

c Sketch a graph of d versus t between the points from part **a** (the d-intercept) and part **b** (the t-intercept).



Hint for Q7: The axes will have d on the vertical axis and t on the horizontal axis.



8 Use your algebra and fraction skills to help sketch graphs for these relations by finding x- and y-intercepts.


a $\frac{x}{2} + \frac{y}{3} = 1$ **b** $y = \frac{8 - x}{4}$ **c** $\frac{y}{2} = \frac{2 - 4x}{8}$




Hint for Q8a: Recall: $\frac{0}{2} = 0$

9 Give reasons why the x-intercept on these graphs has the exact coordinates $(1.5, 0)$.

a



b



Points to rules — 10

10 Write down the rule for the graph with these axes intercepts. Write the rule in the form $ax + by = d$.

a $(0, 4)$ and $(4, 0)$ **b** $(0, 2)$ and $(2, 0)$

c $(0, -3)$ and $(3, 0)$ **d** $(0, 1)$ and $(-1, 0)$

Using a calculator 5C: Sketching straight lines

This activity can be found in the More Resources section of the Interactive Textbook in the form of a printable PDF.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5D Lines with one intercept

Learning intentions

- To understand that vertical and horizontal lines have the same x-coordinate or the same y-coordinate, respectively
- To know the equation form of vertical and horizontal lines
- To be able to graph vertical and horizontal lines and determine their rule from a graph
- To know that lines of the form $y = mx$ pass through the origin
- To be able to sketch lines that pass through the origin

Key vocabulary: origin, vertical, horizontal, parallel, x-intercept, y-intercept

Vertical and horizontal lines are special types of lines. You often see them in houses and construction.

When plotting a graph, a vertical line runs parallel to the y-axis. Since parallel lines never touch, the line will never cross the y-axis, and has only one intercept, on the x-axis.


Horizontal lines run parallel to the x-axis and they have only one intercept, on the y-axis.

Lines that pass through the origin $(0, 0)$ also have only one intercept, as the x- and y-intercepts are the same point.

Lesson starter: What rule satisfies all points?


Here is one vertical (red) and one horizontal (blue) line.

- For the vertical line shown, write down the coordinates of all the points shown as dots.
- What is always true for each coordinate pair?
- Can you think of a simple equation that describes every point on the line?
- For the horizontal line shown, write down the coordinates of all the points shown as dots.
- What is always true for each coordinate pair?
- Can you think of a simple equation that describes every point on the line?




Key ideas

- **Vertical** line: $x = b$
  - **Parallel** to the y-axis, the x-coordinate is the same for every point on the line.
  - The equation is of the form $x = b$, where b is a constant (fixed number).
  - The x-intercept coordinates are $(b, 0)$.
  - There is no y-intercept.




- **Horizontal** line: $y = c$
  - Parallel to the x-axis, the y-coordinate is the same for every point on the line.
  - The equation is of the form $y = c$, where c is a constant (fixed number).
  - The y-intercept coordinates are $(0, c)$.
  - There is no x-intercept.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

- Lines through the origin $(0, 0)$: $y = mx$
  - The y-intercept is at $(0, 0)$.
  - The x-intercept is at $(0, 0)$.
  - Since the x- and y-intercepts give only one point, a second point is required. Substitute any x value into the rule to give a second point. Use $x = 1$ for ease of calculation.



Exercise 5D

Understanding

1–4 | 2, 4

1 Classify the following lines as either *horizontal* or *vertical*.

a $y = 3$ **b** $y = -1$ **c** $x = 2$
d $y = 0$ **e** $x = -5$ **f** $x = 0$



Hint for Q1: Will they be parallel to the x-axis or y-axis?

2 Select the features from the list (**i**–**v**) that apply to:

a horizontal lines **b** vertical lines


i parallel to the y-axis
ii has a y-intercept
iii has an x-intercept
iv parallel to the x-axis
v always pass through the origin




Hint for Q2: Draw a horizontal and vertical line to help.

3 **a** What is the x-coordinate of each point on these graphs?

i




ii




b What is the y-coordinate of each point on these graphs?

i



ii



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5D

4 List which of the following rules will have graphs that pass through $(0, 0)$.

a $y = 4$ b $y = 2x$ c $y = 3x + 2$

d $x = 2$ e $y = -x$ f $y = \frac{1}{2}x$



Hint for Q4: Equations of the form $y = mx$ pass through the origin.

Fluency

5–7(½), 8 | 5–8(½)

Example 9 Graphing vertical and horizontal lines


Sketch the graph of the following vertical and horizontal lines.

a $y = 3$ b $x = -4$ c $y = 0$

Solution

Explanation


a



Each point on the line has a y-coordinate of 3, so the y-intercept is at $(0, 3)$.

Sketch a horizontal line through all points where $y = 3$.


b



Each point on the line has an x-coordinate of -4, so the x-intercept is at $(-4, 0)$.

Sketch a vertical line through all points where $x = -4$.

c



Each point on the line has a y-coordinate of 0.
Sketch a horizontal line through all points where $y = 0$; this line is the x-axis.

Now you try

Sketch the graph of the following vertical and horizontal lines.

a $y = -1$ b $x = 3$ c $x = 0$

5 Sketch the graph of the following vertical and horizontal lines.

a $x = 2$ b $x = 5$

c $y = 4$ d $y = 1$

e $x = -3$ f $x = -2$

g $y = -1$ h $y = -3$



Hint for Q5: Do they have an x-intercept or a y-intercept?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Example 10 Sketching lines that pass through the origin


Sketch the graph of $y = 3x$.

Solution

$y = 3x$
The x- and y-intercept are both at $(0, 0)$.
Another point (let $x = 1$):

$$y = 3 \times (1)$$
$$y = 3$$

Another point is at $(1, 3)$.



Explanation

The equation is of the form $y = mx$; i.e. when you substitute $x = 0$, $y = 3 \times 0 = 0$, giving $(0, 0)$ as both the x- and y-intercepts.

Two points are required to generate the straight line. Find another point by substituting $x = 1$.

Any other x value could be used but the calculation is simplest for $x = 1$.

Plot and label both points and sketch the graph by joining the points in a straight line.

Now you try

Sketch the graph of $y = 4x$.

6 Sketch the graph of the following linear relations, which pass through the origin.

a $y = 2x$ **b** $y = 5x$ **c** $y = 6x$
d $y = x$ **e** $y = -4x$ **f** $y = -3x$
g $y = -2x$ **h** $y = -x$ **i** $y = \frac{1}{2}x$



Hint for Q6: Substitute $x = 1$ to obtain a second point.

7 Sketch the graphs of these special lines all on the same set of axes and label with their equations.

a $x = -2$ **b** $y = -3$ **c** $y = 2$
d $x = 4$ **e** $y = 4x$ **f** $y = -\frac{1}{2}x$
g $y = -1.5x$ **h** $x = 0.5$ **i** $x = 0$
j $y = 0$ **k** $y = 2x$ **l** $y = 1.5x$




Hint for Q7: There are vertical lines, horizontal lines and lines through the origin.


Example 11 Finding the equation of horizontal and vertical lines

Give the equation of the following vertical and horizontal lines.

a



b



Continued on next page


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


5D

Solution	Explanation
a $x = 2$	Line is vertical, with x-coordinate always 2.
b $y = -1$	Line is horizontal, with y-coordinate always -1.


Now you try


Give the equation of the following vertical and horizontal lines.


a 

b 

8 Give the equation of the following vertical and horizontal lines.


a 


b 


c 



Hint for Q8: A horizontal line has the form $y = c$. A vertical line has the form $x = b$.

d 

e 

f 

Problem-solving and reasoning

9, 10 | 9–11

9 Find the equation of the straight line that is:

a parallel to the x-axis and passes through the point $(1, 3)$
b parallel to the y-axis and passes through the point $(5, 4)$
c parallel to the y-axis and passes through the point $(-2, 4)$
d parallel to the x-axis and passes through the point $(0, 0)$



Hint for Q9: First decide whether the line is horizontal or vertical, then use the point.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10 If, in a picture, the surface of the sea is represented by the x-axis, state the equation of the following paths.


a A plane flies horizontally at 250 m above sea level. One unit is 1 metre.

b A submarine travels horizontally 45 m below sea level. One unit is 1 metre.




11 The rules of the following graphs are of the form $y = mx$. Use the points marked with a dot to find m and hence state the equation.

a




b






Hint for Q11:
In a rule such as $y = 3x$, when $x = 1$, $y = 3 \times 1 = 3$.

c




d



Rectangular areas — 12, 13

The four lines $x = 1$, $x = 3$, $y = 2$ and $y = 5$ drawn on the one set of axes form a rectangle, as shown (shaded). The area of this rectangle is length × width = $3 \times 2 = 6$ square units.



12 Find the area of the rectangle contained within the following four lines.

a $x = 1, x = -2, y = -3, y = 2$ **b** $x = 0, x = 17, y = -5, y = -1$

13 The lines $x = -1$, $x = 3$ and $y = -2$ form three sides of a rectangle. Find the possible equation of the fourth line if:

a the area of the rectangle is:

i 12 square units **ii** 8 square units **iii** 22 square units

b the perimeter of the rectangle is:

i 14 units **ii** 26 units **iii** 31 units

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


5E Gradient

Learning intentions

- To understand what is meant by the gradient of a line
- To know that the gradient of a straight line is constant
- To know that the gradient of a line can be positive, negative, zero or undefined
- To be able to find the gradient of a line from a graph or between two given points

Key vocabulary: gradient, rise, run

The gradient of a line is a measure of its slope. It is a number that describes the steepness of a line. It is calculated by considering how far a line rises or falls between two points within a given horizontal distance. The horizontal distance between two points is called the *run*. The vertical distance is called the *rise*.






The gradient or slope of a mountain can be a challenge for mountain climbers.

Lesson starter: Which line is the steepest?

The three lines here connect the points A, B, C, D, E, F, G and H.

- Calculate the rise and run (working from left to right) and also the fraction $\frac{\text{rise}}{\text{run}}$ for these segments.
  - **i** AB **ii** BC **iii** BD
  - **iv** EF **v** GH
- What do you notice about the fractions $\left(\frac{\text{rise}}{\text{run}}\right)$ for parts **i**, **ii** and **iii**?
- How does the $\frac{\text{rise}}{\text{run}}$ for EF compare with the $\frac{\text{rise}}{\text{run}}$ for parts **i**, **ii** and **iii**? Which of the two lines is the steepest?
- Your $\frac{\text{rise}}{\text{run}}$ for GH should be negative. Why is this the case?
- Discuss whether or not GH is steeper than AD.




Use computer software (dynamic geometry) to produce a set of axes and grid.

- Construct a line segment with endpoints on the grid. Show the coordinates of the endpoints.
- Calculate the rise (vertical distance between the endpoints) and the run (horizontal distance between the endpoints).
- Calculate the gradient as the *rise* divided by the *run*.
- Now drag the endpoints and explore the effect on the gradient.
- Can you drag the endpoints but retain the same gradient value? Explain why this is possible.
- Can you drag the endpoints so that the gradient is zero or undefined? Describe how this can be achieved.


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Key ideas

- The **gradient** of a line is a measure of its steepness or slope.
- The gradient of a line can be found using gradient $(m) = \frac{\text{rise}}{\text{run}}$.
  - We work from left to right, so the run is always positive.



- The gradient can be positive, negative, zero or undefined.
  - The gradient is positive if the graph increases from left to right.
  - The gradient is negative if the graph decreases from left to right.
  - The gradient is zero if the line is horizontal.
  - The gradient is undefined if the line is vertical.




Exercise 5E

Understanding


1, 2 | 2

1 For these lines, calculate the value for: **i** run **ii** rise


a




b




c




d



e



f





Hint for Q1: Remember that the rise is negative if the line slopes downward from left to right.

2 Use the words 'positive', 'negative', 'zero' or 'undefined' to complete each sentence.

a The gradient of a horizontal line is _____.

b The gradient of the line joining $(0, 3)$ with $(5, 0)$ is _____.

c The gradient of the line joining $(-6, 0)$ with $(1, 1)$ is _____.

d The gradient of a vertical line is _____.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5E


Fluency

3(½), 5(½) | 3(½), 4, 5(½)


Example 12 Finding the gradient of a line

For each graph, state whether the gradient is positive, negative, zero or undefined, then find the gradient, where possible.


a




b



c




d



Solution

a The gradient is positive.




$$\text{Gradient} = \frac{\text{rise}}{\text{run}}$$
$$= \frac{4}{2}$$
$$= 2$$

Explanation

By inspection, the gradient will be positive since the graph increases from left to right.
Select any two points and create a right-angled triangle between them to determine the rise and run.
Substitute rise = 4 and run = 2, and simplify.

b The gradient is negative.




$$\text{Gradient} = \frac{\text{rise}}{\text{run}}$$
$$= \frac{-3}{2}$$
$$= -\frac{3}{2}$$

By inspection, the gradient will be negative since y values decrease from left to right.
Create a right-angled triangle between two points.
Rise = −3 since it 'falls' 3 units, and run = 2.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

c The gradient is positive.



$$\text{Gradient} = \frac{\text{rise}}{\text{run}} = \frac{2}{2} = 1$$

A rising graph indicates a positive gradient.
For this triangle, rise = 2. The run is from left to right, from −2 to 0, so the run is +2.


d The gradient is undefined.

The line is vertical.


Now you try

For each graph, state whether the gradient is positive, negative, zero or undefined, then find the gradient, where possible.


a




b



c




d




3 For each graph state whether the gradient is positive, negative, zero or undefined, then find the gradient where possible.

a



b





Hint for Q3:
Create a right-angled triangle to find the rise and the run.




$$\text{Gradient} = \frac{\text{rise}}{\text{run}}$$


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5E


c




d




e




f




g




h




i




j



k



l






Hint for Q3k:
In part k, from −1 to 3 is a rise of 4.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

4 Find the gradient of each line $A-F$ on this graph and grid.





Hint for Q4: Pick two known points on each line to find the gradient between them.


Example 13 Finding the gradient between two points

Find the gradient (m) of the line joining the given points.

a $A(3, 4)$ and $B(5, 6)$ **b** $A(-3, 6)$ and $B(1, -3)$

Solution

a




$$m = \frac{\text{rise}}{\text{run}}$$
$$= \frac{2}{2}$$
$$= 1$$

Explanation

Plot the two points on a set of axes and construct a right-angled triangle between them.
Observe from the graph that the gradient will be positive.
Calculate the gradient by finding the rise and run.

b



$$m = \frac{\text{rise}}{\text{run}}$$
$$= \frac{-9}{4} \text{ or } -\frac{9}{4} \text{ or } -2.25$$

Plot points A and B and join in a line. Observe that the gradient will be negative.
From left to right, -3 to 1, run $= 4$.
Rise from 6 to -3 is a 'fall' of 9 units, so rise $= -9$.

Now you try

Find the gradient (m) of the line joining the given points.

a $A(2, 3)$ and $B(4, 7)$ **b** $A(-2, 2)$ and $B(3, -6)$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5E

5 Find the gradient of the lines joining the following pairs of points.

a $A(2, 3)$ and $B(3, 5)$
b $C(0, 8)$ and $D(2, 6)$
c $E(0, 4)$ and $F(2, 0)$
d $G(2, 1)$ and $H(5, 4)$
e $A(1, 5)$ and $B(2, 7)$
f $C(-2, 4)$ and $D(1, -2)$
g $E(-3, 4)$ and $F(2, -1)$
h $G(-1, 5)$ and $H(1, 6)$
i $A(-4, -2)$ and $B(-2, 1)$
j $C(1, 1)$ and $D(3, -4)$
k $E(3, 2)$ and $F(0, 1)$
l $G(-1, 1)$ and $H(-3, -4)$



Hint for Q5: Plot the points to help visualise the rise and run.

Problem-solving and reasoning

6, 7 | 6, 8, 9

6 Find the gradient, using $\frac{\text{rise}}{\text{run}}$, corresponding to the following slopes.

a A road falls 10 m for every 200 horizontal metres.
b A cliff rises 35 metres for every 2 metres horizontally.
c A plane descends 2 km for every 10 horizontal kilometres.
d A submarine ascends 150 m for every 20 horizontal metres.



Hint for Q6: The horizontal distance is the run.

7 A firecracker ascends with a gradient of 2. How far, horizontally, has the cracker travelled after rising 80 m?

8 Sally runs up a sand dune that has a gradient of $\frac{1}{3}$. How far, horizontally, has Sally moved after rising 30 m?



9 Find the missing number.

a The gradient joining the points $(0, 2)$ and $(1, ?)$ is 4.
b The gradient joining the points $(?, 5)$ and $(3, 9)$ is 2.
c The gradient joining the points $(-3, 4)$ and $(?, 1)$ is -1.
d The gradient joining the points $(-4, ?)$ and $(-2, -12)$ is -4.




Hint for Q9a: A gradient of $4 = \frac{4}{1}$: a rise of 4 units for every 1 unit to the right.

Where does it hit?

— | 10

10 The line here has gradient $\frac{-2}{3}$, which means that it falls 2 units for every 3 units across. The y-intercept is $(0, 3)$.

a Use the gradient to find the y-coordinate on the line where:
i $x = 6$ **ii** $x = 9$
b What will be the coordinates of the x-intercept?
c What would be the x-intercept if the gradient was changed to:
i $\frac{-1}{2}$? **ii** -2? **iii** $\frac{-5}{4}$?



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5F Gradient and direct proportion

Learning intentions

- To understand that gradient represents the rate of change of one variable with respect to another
- To understand what it means for two variables to be directly proportional
- To be able to identify the form of the linear rule for two variables in direct proportion
- To be able to form and work with rules in direct proportion

Key vocabulary: rate, variable, gradient, directly proportional

We have seen that the gradient of a line is the increase or decrease (rise) of the y values compared to the change (run) in the x values.

Gradient can also be considered as a rate. It is the rate of change of y with respect to x. Speeds such as 60 km/h or 10 m/s are a common form of rates. They give the change in distance with respect to time; e.g. 60 km each hour or 10 m each second.


The connection between gradient, rate problems and direct proportion can be shown through the use of linear rules and graphs. If two variables are directly related, then the rate of change of one variable with respect to the other is constant. This means that the rule linking the two variables is linear and can be represented as a straight line graph passing through the origin.

The amount of water squirting from a hose, for example, is directly proportional to the time since it was turned on. The gradient of the graph of *water volume* versus *time* will determine the rate at which water is squirting from the hose.

Lesson starter: Average speed


Over 5 hours, Sandy travels 400 km in a car.

- What is Sandy's average speed for the trip?
- Draw a graph of distance versus time for the journey, assuming a constant speed.
- Where does your graph intersect the axes and why?
- Find the gradient of your graph. What do you notice?
- Find a rule linking distance (d) and time (t).



Key ideas

- A **rate** is the change in one variable compared with another. For example, 60 km/h (60 km per hour) is a rate. It states a 60 km change in distance for each hour in time that passes.
  - A rate in simplest form is written as a change in one variable per one unit of another variable. For example, we simplify 40 km in 2 hours to 20 km per hour or 20 km/h.
- If two variables are **directly proportional**:
  - the rate of change of one variable with respect to the other is constant
  - the graph is a straight line passing through the origin
  - the rule is of the form $y = mx$
  - the gradient (m) of the graph equals the rate of change of the y variable with respect to the x variable.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Exercise 5F

Understanding

1–3 | 3

1 Write 'Yes' or 'No' to state which of the following are rates.

a \$120 **b** 12 cm/s **c** 150 mL/min **d** 80 km




2 Write these rates in simplest form.

a 120 km in 2 hours
b 180 m in 20 seconds
c 400 L in 80 min
d \$30 for 20 litres
e \$900 in 30 hours
f 15°C in 5 min

Hint for Q2: Simplest form is per 1 unit; e.g. 50 km in 1 hour, 50 km/h.

3 This graph shows how far Caroline travels on her bike over 4 hours.



a State how far Caroline has travelled after:
- **i** 1 hour
- **ii** 2 hours
- **iii** 3 hours

b Write down the speed of the bike in km/h (rate of change of distance over time).

c Find the gradient of the graph.

d What do you notice about your answers from parts **b** and **c**?

Fluency

4–6 | 4–7

Example 14 Forming rules in direct proportion

Write a rule linking the variables A dollars and t hours if \$60 is earned in 5 hours of work at a constant rate.

Solution

$60 \div 5 = 12$
Rate is \$12/h.

$A = 12 \times t$
$\therefore A = 12t$

Explanation

$\div 5$ (\$60 in 5 hours / \$12 in 1 hour) $\div 5$

The amount earned, A, is \$12 for each hour worked.
t hours of work earns $12 \times t$ dollars.

Now you try

Write a rule linking the variables V litres and t seconds if a bucket contains 8 L of water from a hose after 4 seconds.

4 Write down a rule linking the given variables, using a constant rate.

a I travel 720 km in 12 hours. Use d for distance and t for time.
b A calf grows 12 cm in 6 months. Use g for growth height and t for time.
c The cost of petrol is \$120 for 80 litres. Use C for cost and n for the number of litres.
d The profit is \$10 000 for 500 tonnes. Use P for profit and t for the number of tonnes.



Hint for Q4: First write the rate in simplest form.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Example 15 Exploring direct proportion

Water is poured into an empty tank at a constant rate. It takes 3 hours to fill the tank with 6000 litres.

a What is the rate at which water is poured into the tank?

b Draw a graph of volume (V litres) vs time (t hours) using $0 \leq t \leq 3$.

c Find:

i the gradient of your graph

ii the rule for V

d Use your rule to find:


i the volume after 1.5 hours

ii the time to fill 5000 litres

Solution

a 6000 L in 3 hours = 2000 L/hour

b



c i gradient $= \frac{6000}{3} = 2000$

ii $V = 2000t$

d i $V = 2000t$

$= 2000 \times 1.5$

$= 3000$

$\therefore$ 3000 L after 1.5 hours

ii $V = 2000t$

$5000 = 2000t$

$2.5 = t$

$\therefore$ it takes $2\frac{1}{2}$ hours to fill 5000 L

Explanation

$\div 3$ (6000 L per 3 hours / 2000 L per 1 hour) $\div 3$

Draw axes with volume on the vertical axis and time on the horizontal axis, since volume depends on time.

Plot the two end points: $(0, 0)$ since the tank starts empty and $(3, 6000)$. Join these points with a straight line.

Using the end points, rise = 6000, run = 3.

Note that the gradient is the same as the rate.

2000 L are filled for each hour.

Substitute $t = 1.5$ into your rule.

Substitute $V = 5000$ into the rule and solve for t by dividing each side by 2000.

Now you try

Water is poured into an empty inflatable pool at a constant rate. It takes 50 minutes to fill the pool with 2500 litres.

a What is the rate at which water is poured into the tank?

b Draw a graph of volume (V litres) vs time (t minutes) using $0 \leq t \leq 50$.

c Find:

i the gradient of your graph

ii the rule for V

d Use your rule to find

i the volume after 20 minutes

ii the time to fill 2000 litres

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5F

5 A 300 litre fish tank takes 3 hours to fill using a hose at a constant rate.

a What is the rate at which water is poured into the tank?

b Draw a graph of volume (V litres) vs time (t hours) using $0 \leq t \leq 3$.

c Find:

i the gradient of your graph

ii the rule for V

d Use your rule to find:

i the volume after 1.5 hours

ii the time to fill 200 litres



Hint for Q5a: 300 litres in 3 hours; how many litres in 1 hour?



Hint for Q5b: Place time on the horizontal axis.



6 A solar powered car travels 100 km in 4 hours.

a What is the rate of change of distance over time (i.e. speed)?

b Draw a graph of distance (d km) vs time (t hours) using $0 \leq t \leq 4$.

c Find:

i the gradient of your graph

ii the rule for d


d Use your rule to find:

i the distance after 2.5 hours


ii the time to travel 40 km

7 Use the gradient to find the rate of change of distance over time (speed) for these graphs. Use the units given on each graph.


a




b



c



d



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Problem-solving and reasoning

8, 9 | 9–11

8 Who is travelling the fastest?
- Mick runs 720 m in 2 minutes.
- Sally rides 550 m in 1 minute.
- Udhav jogs 2000 m in 5 minutes.

9 Which animal is travelling the slowest?
- A leopard runs 400 m in 30 seconds.
- A jaguar runs 2700 m in 3 minutes.
- A panther runs 60 km in $1\frac{1}{3}$ hours.



Hint for Q9: To compare rates, they will need to be in the same units.



10 A car's trip computer says that the fuel economy for a trip is 8.5 L per 100 km.
- **a** How many litres would be used for 120 km?
- **b** How many litres would be used for 850 km?
- **c** How many kilometres could be travelled if the car's petrol tank capacity was 68 L?



Hint for Q10: Consider how many litres are used in 1 kilometre.

11 The area of a particular rectangle is given by $A = lw$. Its length is fixed at 12 cm but its width, w cm, can vary.
- **a** Write a rule for the area of this rectangle.
- **b** Draw a graph of A against w for $0 \leq w \leq 4$.
- **c** Decide if the area of this rectangle is directly proportional to its width. Explain.



Hint for Q11: Equations of the form $y = mx$ show that y is directly proportional to x.

Rate challenge

— | 12, 13

12 Hose A can fill a bucket in 2 minutes and hose B can fill the same bucket in 4 minutes.
- **a** What fraction of a bucket does hose A fill in 1 minute?
- **b** What fraction of a bucket does hose B fill in 1 minute?
- **c** If both hoses were used at the same time, what fraction of a bucket could they fill in 1 minute?
- **d** If both hoses were used at the same time, how long would it take to fill the bucket?

13 Vonda can vacuum an office building in 2 hours and her husband Chris can vacuum the same office building in 3 hours. How long would it take to vacuum the office building if they both vacuumed at the same time?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5G Gradient–intercept form


Learning intentions

- To know that $y = mx + c$ is the gradient–intercept form of a straight line with m the gradient and $(0, c)$ the y-intercept
- To know that any straight line equation can be expressed in gradient–intercept form
- To be able to rearrange a linear equation into gradient–intercept form
- To be able to determine the gradient and y-intercept of a straight line from gradient–intercept form
- To be able to sketch a graph using the y-intercept and the gradient

Key vocabulary: gradient–intercept form, y-intercept, gradient

Shown here is the graph of the rule $y = 2x - 1$. It shows a gradient of 2 and a y-intercept at $(0, -1)$. The fact that these numbers match numbers in the rule is no coincidence. This is why rules written in this form are called gradient–intercept form. Other examples of rules in this form include:

$$y = -5x + 2,\ y = \frac{1}{2}x - 0.5 \text{ and } y = \frac{x}{5} + 20.$$





Look at the slope of this mountain. Is the gradient larger or smaller than 1?

Lesson starter: What's in common?

Sketch the following linear relations on the same set of axes. Plot points or use the x- and y-intercepts.

a $y = 2x$ **b** $y = 2x + 2$ **c** $y = 2x - 1$

- What do these graphs have in common?
- Calculate the gradient of each line. What do you notice?
- How do the gradients relate to the rules?

Now, sketch the following linear relations on the same set of axes. Plot points or use the x- and y-intercepts.

a $y = x + 1$ **b** $y = 2x + 1$ **c** $y = -\frac{1}{2}x + 1$


- What do these graphs have in common?
- How does the common feature relate to the rules?
- Can you form a rule for a linear relation with a gradient of 3 and a y-intercept at $(0, 2)$?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Key ideas

m = gradient $(0, c)$ = y-intercept

- $y = mx + c$ (or $y = mx + b$ depending on preference) is the **gradient–intercept form** of a straight line equation.
- If the y-intercept is at $(0, 0)$, the equation becomes $y = mx$ and these graphs will pass through the origin.
- Any linear relation can be rearranged to be written in gradient–intercept form by making y the subject.
- To sketch a graph using the gradient–intercept method, find the y-intercept and use the gradient to find a second point.
  - For example, if $m = \frac{2}{5}$, move 5 right and 2 up from the y-intercept.
  - For the gradient of $-2 = \frac{-2}{1}$ move 1 right and 2 down from the y-intercept.



Exercise 5G

Understanding

1–4 | 4

1 Choose from the words *y-intercept*, *gradient* and *subject* to complete the following. In the gradient–intercept form, $y = mx + c$:

a m is the_____.

b $(0, c)$ is the _____.

c y is the _____ of the equation.

2 Match the given gradient and y-intercept in the first column with the rule in the left column.

a Gradient = 2, y-intercept = $(0, 5)$ **i** $y = 5x + 2$

b Gradient = 5, y-intercept = $(0, 2)$ **ii** $y = -2x + 3$

c Gradient = -2, y-intercept = $(0, 3)$ **iii** $y = 2x + 5$

d Gradient = -1, y-intercept = $(0, -2)$ **iv** $y = -x - 2$

3 **a** From the point $(0, 2)$, give the coordinates of the point that is:

i 2 right and 4 up **ii** 3 right and 1 down

b From the point $(0, -1)$, give the coordinates of the point that is:

i 1 right and 2 up **ii** 3 right and 4 down



Hint for Q3: Plot the point to help you visualise, if required.

4 Fill in the missing numbers.

a The gradient $\frac{5}{3}$ describes a run of _____ for a rise of _____.

b The gradient 4 describes a run of _____ for a rise of _____.

c The gradient $\frac{-1}{2}$ describes a run of _____ for a rise of _____.



Hint for Q4:
Gradient = $\frac{\text{rise}}{\text{run}}$
Run is always positive.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5G

Fluency

5–7(½) | 5–8(½)

Example 16 Stating the gradient and y-intercept

State the gradient and the y-intercept coordinates for the graphs of the following relations.

a $y = 2x - 1$ b $y = -3x$

Solution

a $y = 2x - 1$
The gradient = 2
y-intercept = $(0, -1)$

Explanation

The rule is given in gradient-intercept form, $y = mx + c$.
The gradient is the coefficient of x (the numeral multiplied by x).
The constant term is the y-coordinate of the y-intercept.

b $y = -3x$
The gradient = -3
y-intercept = $(0, 0)$

The gradient is the coefficient of x including the negative sign.
The constant term is not present so the y-intercept is at $(0, 0)$.

Now you try

State the gradient and the y-intercept coordinates for the graphs of the following relations.

a $y = 3x - 2$ b $y = -x$

5 State the gradient and y-intercept coordinates for the graphs of the following relations.

a $y = 3x - 4$ b $y = -5x - 2$ c $y = -2x + 3$
d $y = \frac{1}{3}x + 4$ e $y = -4x$ f $y = 2x$
g $y = 2.3x$ h $y = -0.7x$ i $y = x$



Hint for Q5: $y = mx + c$
gradient y-intercept y-coordinate

Example 17 Rearranging linear equations into the form $y = mx + c$

Rearrange these linear equations into the form $y = mx + c$.

a $2x + y = 7$ b $4x + 2y = 10$

Solution

a $2x + y = 7$
$y = 7 - 2x$
$y = -2x + 7$

Explanation

To have y by itself, subtract $2x$ from both sides.
$7 - 2x$ is the same as $-2x + 7$, which is in the form $mx + c$.

b $4x + 2y = 10$
$2y = -4x + 10$
$y = -2x + 5$

Solve for y by first subtracting $4x$ from both sides.
Divide both sides of the equation by 2:
$\frac{-4x + 10}{2} = \frac{-4}{2}x + \frac{10}{2} = -2x + 5$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Now you try

Rearrange these linear equations into the form $y = mx + c$.

a $3x + y = 5$ **b** $6x + 3y = 9$

6 Rearrange these linear equations into the form $y = mx + c$.

a $3x + y = 4$ **b** $2x + y = 7$ **c** $y - 3x = 2$
d $y - 2x = 7$ **e** $5x - y = 3$ **f** $4x + 2y = 8$
g $3x + 2y = 6$ **h** $4x - 2y = 5$ **i** $5x - 3y = -6$



Hint for Q6: Follow steps as if you were solving an equation for y.


Example 18 Sketching linear graphs using the gradient and y-intercept

Find the value of the gradient and y-intercept coordinates for these relations and sketch their graphs.

a $y = 2x - 1$ **b** $y = \frac{-3}{4}x + 2$

Solution

a $y = 2x - 1$
y-intercept $= (0, -1)$
Gradient $= 2 = \frac{2}{1}$




Explanation

The rule is in gradient–intercept form so we can read off the gradient (coefficient of x) and the y-intercept (constant term).
Gradient $= \frac{2 \text{ (rise)}}{1 \text{ (run)}}$; for every 1 right, move 2 up.

Label the y-intercept at $(0, -1)$.
From $(0, -1)$ move 1 to the right and 2 up to give a second point at $(1, 1)$.
Mark and join the points to form a line.

b $y = \frac{-3}{4}x + 2$
y-intercept $= (0, 2)$
Gradient $= \frac{-3}{4}$



Read off the y-intercept and gradient from the gradient–intercept form.
Rise $= -3$, run $= 4$.

Label the y-intercept and from here move 4 across to the right (run) and 3 down (due to -3) to give a second point at $(4, -1)$.
Join points to form a line.

Now you try

Find the value of the gradient and y-intercept coordinates for these relations and sketch their graphs.

a $y = 3x - 2$ **b** $y = \frac{-5}{4}x + 1$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5G

7 Find the gradient and y-intercept coordinates for these relations and sketch their graphs.

a $y = x - 2$ **b** $y = 2x + 1$ **c** $y = 3x + 2$

d $y = \frac{1}{2}x + 2$ **e** $y = -3x + 3$ **f** $y = \frac{3}{2}x + 1$

g $y = 2x$ **h** $y = \frac{-4}{3}x$ **i** $y = \frac{-2}{3}x$



Hint for Q7: Plot the y-intercept first, then use the rise and run of the gradient.

Example 19 Rearranging equations to sketch using the gradient and y-intercept

Rearrange the equation $x + 2y = 6$ to find the gradient and y-intercept coordinates, and sketch its graph.

Solution

$x + 2y = 6$


$2y = 6 - x$

$y = 3 - \frac{1}{2}x$

$= -\frac{1}{2}x + 3$

y-intercept is at $(0, 3)$

Gradient $= -\frac{1}{2} = \frac{-1}{2}$



Explanation

Make y the subject by subtracting x from both sides and then dividing both sides by 2.

$\frac{6 - x}{2} = \frac{6}{2} - \frac{x}{2} = 3 - \frac{1}{2}x$

Rewrite in the form $y = mx + c$ to read off the gradient and y-intercept.

Link the negative sign to the rise (-1) so the run is positive $(+2)$.

Mark the y-intercept and then from this point move 2 right and 1 down to give a second point at $(2, 2)$.

Note that the x-intercept will be 6. If the gradient is $\frac{-1}{2}$ then a fall of 3 needs a run of 6.

Now you try

Rearrange the equation $2x + 3y = 12$ to find the gradient and y-intercept coordinates, and sketch its graph.

8 Rearrange these equations to find the gradient and y-intercept coordinates, and sketch their graphs.

a $x + y = 4$ **b** $y - x = 6$

c $4x + 2y = 6$ **d** $x + 2y = 8$

e $2x + 3y = 6$ **f** $4x + 3y = 12$

g $x - 2y = 4$ **h** $2x - 3y = 6$

i $3x - 4y = 12$ **j** $x + 4y = 0$

k $x - 5y = 0$ **l** $8x - 2y = 0$



Hint for Q8: Solve for y to make y the subject with equations in the form $y = mx + c$.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Problem-solving and reasoning 9, 10 10–12

9 During a heavy rainstorm, a rain gauge is filling with water. The rule for the volume of water in the gauge (V mL), t hours after the start of the storm, is given by $V = 4t + 2$.

a State the gradient and V-intercept coordinates of this relation and sketch its graph.

b In this scenario, what do the gradient and the V-intercept represent?



Hint for Q9: Here, the V-intercept is like the y-intercept.



10 Vera is trying to convince her friend Li that $\frac{4x+6}{2} = 2x + 6$. She offers to pay Li \$2 if she is wrong. Does Vera lose her money? Explain.

11 Which of these linear relations have a gradient of 2 and y-intercept at $(0, -3)$?

a $y = 3 - 2x$

b $y = \frac{2x - 6}{2}$

c $y = \frac{4x - 6}{2}$

d $y = \frac{3 - 2x}{-1}$

e $2y = 4x - 3$

f $-2y = 6 - 4x$



Hint for Q11: Rewrite each in the form $y = mx + c$ and look for $y = 2x - 3$.

12 Jeremy says that the graph of the rule $y = 2(x + 1)$ has gradient 2 and y-intercept at $(0, 1)$.

a Explain his error.


b What can be done to the rule to help show the y-intercept?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5G

The missing y-intercept — 13

13 This graph shows two points $(-1, 0)$ and $(1, 4)$, with a gradient of 2. By considering the gradient (2 up for every 1 right), the y-intercept can be calculated to be at $(0, 2)$, so $y = 2x + 2$.



Use this approach to find the rule of the line passing through these points.

a $(-1, 1)$ and $(1, 5)$ **b** $(-2, 4)$ and $(2, 0)$
c $(-2, -1)$ and $(3, 9)$ **d** $(-4, 1)$ and $(2, 4)$
e $(-1, 3)$ and $(1, 4)$



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Progress quiz

5A **1** For the rules listed below:

i Complete a table of values.

x	-3	-2	-1	0	1	2	3
y							

ii Plot a graph.

a $y = 3x + 2$ **b** $y = -2x + 4$

5A **2** Decide whether the point $(2, -3)$ is on the line with the given rules.

a $y = 2x - 5$


b $y = -2x + 1$

5B **3** State the coordinates of the x- and y-intercepts from the following table and graph.

a

x	-2	-1	0	1	2
y	-6	-4	-2	0	2

b



5B/C **4** For the linear relations listed below:

i Find the coordinates of the y-intercept.

ii Find the coordinates of the x-intercept.

iii Sketch the graph showing the x- and y-intercepts.

a $y = 3x + 6$ **b** $y = -2x - 8$ **c** $2x - 5y = 10$

5C **5** A cyclist rides from school to home, with his distance from home given by the rule, $d = -\frac{1}{2}t + 10$ where d is in kilometres and t in minutes.

a Find how far school is from home $(t = 0)$.

b Find how long it takes to cycle home $(d = 0)$.

c Sketch a graph of the cycle from school to home, labelling intercepts.

5D **6** Sketch the graph of the following horizontal and vertical lines.

a $x = 5$ **b** $y = -2$ **c** $x = 0$

5D **7** On the one set of axes, sketch the graph of the following linear relations, which pass through the origin.


a $y = 6x$ **b** $y = -x$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


Progress quiz

5E **8** Find the gradient (m) of the following lines (where possible).


a




b



c



d



e the line joining the points:
- **i** $A(2, 5)$ and $B(5, 11)$
- **ii** $C(-2, -4)$ and $D(1, 3)$
- **iii** $E(-1, 3)$ and $F(1, -7)$
- **iv** $G(2, 6)$ and $H(-1, 6)$

5F **9** Write a rule linking the given variables using a constant rate.
- **a** I travel 150 km in 3 hours. Use d for distance and t for time.
- **b** I earn \$80 in 5 hours. Use E for earnings and t for time.
- **c** A cricket team makes 160 runs off 20 overs. Use r for runs and o for overs.

5F **10** A birdbath is being filled with water. It takes 5 minutes to fill the bath with 300 litres.
- **a** What is the constant rate at which the birdbath is filled with water?
- **b** Draw a graph of volume (V litres) vs time (t minutes) using $0 \leq t \leq 5$.
- **c** Find:
  - **i** the gradient of your graph
  - **ii** the rule for V
- **d** Use your rule to find:
  - **i** the volume after 2 minutes
  - **ii** the time to fill 210 litres

5G **11** State the gradient and y-intercept coordinates for the graphs of the following relations. For parts **d** and **e** you will need to first rearrange the equations into the form $y = mx + c$.

a $y = 2x - 3$ **b** $y = -4x + 1$ **c** $y = \frac{1}{2}x$ **d** $y - 2x = 3$ **e** $3y - 6x = 12$

5G **12** Find the value of the gradient and the y-intercept coordinates for these relations and sketch their graphs.

a $y = 3x - 2$ **b** $y = -\frac{1}{2}x + 3$ **c** $3x + 2y = 8$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5H Finding the equation of a line ★

Learning intentions

- To understand that all straight line graphs can be expressed in gradient–intercept form, $y = mx + c$
- To know that the gradient and one other point are required to find the equation of a line
- To know that all points on a line satisfy the equation of the line
- To be able to find the equation of a line given the y-intercept and another point or the gradient and a point

Key vocabulary: equation, gradient–intercept form, gradient, y-intercept

When data points from an experiment are plotted, they may form a straight line. This shows a linear relationship between the two variables, such as time and growth, involved in the experiment.


By finding the equation of this line, we can develop a rule to relate the two variables. This rule can be used to make further predictions about the data, and could be applied to a larger scale experiment.

Using gradient–intercept form, the rule (or equation) of a line can be found by calculating the value of the gradient and the y-intercept.

Lesson starter: But we don't know the y-intercept!

A line with the rule $y = mx + c$ passes through two points $(-1, 4)$ and $(1, -2)$.

- Using the information given, is it possible to find the value of m? If so, calculate its value.
- The y-intercept is not given on the graph. Discuss what information could be used to find the value of the constant, c, in the rule. Is there more than one way you can find the y-intercept?
- Write the rule for the line.



Key ideas

- To find the equation of a line in gradient–intercept form, $y = mx + c$, you need to find:
  - the value of the gradient (m) using $m = \frac{\text{rise}}{\text{run}}$
  - the value of the constant (c), by observing the y-intercept or by substituting another point.
- All the points (x, y) on a line satisfy the equation of the line.
  For example, $(2, 5)$ is on the line with equation $y = 2x + 1$ since substituting $x = 2$ and $y = 5$ gives a true equation: $5 = 2 \times (2) + 1$.

Exercise 5H

Understanding

1–4 | 4

1 Fill in the missing words.
To find the equation of a line in the form $y = mx + c$, the _____ (m) is required as well as the _____ or another _____.

2 Substitute the given values of m and c into $y = mx + c$ to write the rule.

a $m = 2, c = 5$

b $m = 4, c = -1$

c $m = -2, c = 5$

d $m = -1, c = -\frac{1}{2}$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5H

3 Which one of the following lines does the point $(3, 1)$ lie on?

A $y = 3x + 1$ **B** $y = x - 3$
C $y = -2x + 7$ **D** $y = 2x + 1$
E $y = -x - 2$



Hint for Q3: Which rule gives $y = 1$ when $x = 3$?

4 Substitute the point into the given rule and solve to find the value of c.
For example, using $(3, 4)$, substitute $x = 3$ and $y = 4$ into the rule.

a $(3, 4)$, $y = x + c$
b $(1, 5)$, $y = 2x + c$
c $(3, -1)$, $y = -2x + c$




Hint for Q4a: In part a, $4 = 3 + c$.
$\therefore c = 1$

Fluency

5(½), 7(½) | 5–7(½)

Example 20 Finding the equation of a line given the y-intercept and another point

Determine the equation of the straight line shown here.



Solution

The equation will have the form $y = mx + c$.

gradient, $m = \frac{\text{rise}}{\text{run}}$


$= \frac{-4}{2}$

$= -2$

y-intercept $= (0, -4)$
$\therefore y = -2x - 4$

Explanation


For the gradient–intercept form you need to find the gradient (m) and the y-coordinate of the y-intercept (c).



The y-intercept $(0, -4)$ can be read from the graph.
In $y = mx + c$, substitute $m = -2$ and $c = -4$.

Now you try


Determine the equation of the straight line shown here.




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5 Determine the equation of the following straight lines.


a




b




c




d




e



f



g






Hint for Q5: Write in the form $y = mx + c$. You will need the gradient and the y-intercept (or another point if the y-intercept is not given).

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


5H

6 Find the equation of these straight lines that have fractional gradients. Simplify any fractions.


a




b




c




d



e



f



Example 21 Finding the equation of a line given the gradient and a point

Find the equation of the line that has a gradient, m, of 3 and passes through the point $(2, -1)$.

Solution	Explanation
$y = mx + c$	Substitute $m = 3$ into $y = mx + c$.
$y = 3x + c$	Since $(2, -1)$ is on the line, it must satisfy the equation $y = 3x + c$, hence substitute the point $(2, -1)$ where $x = 2$ and $y = -1$ to find c.
$-1 = 3 \times (2) + c$	Simplify and solve for c by subtracting 6 from both sides.
$-1 = 6 + c$	
$-7 = c$	
$\therefore\ y = 3x - 7$	Write the equation in the form $y = mx + c$.

Now you try

Find the equation of the line that has a gradient, m, of 2 and passes through the point $(2, -1)$.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

7 Find the equation of the line that:
- **a** has a gradient of 3 and passes through the point $(1, 8)$
- **b** has a gradient of -2 and passes through the point $(2, -5)$
- **c** has a gradient of -3 and passes through the point $(2, 2)$
- **d** has a gradient of 1 and passes through the point $(1, -2)$
- **e** has a gradient of -3 and passes through the point $(-1, 6)$
- **f** has a gradient of 5 and passes through the point $(2, 9)$
- **g** has a gradient of -1 and passes through the point $(4, 4)$
- **h** has a gradient of -3 and passes through the point $(3, -3)$
- **i** has a gradient of -2 and passes through the point $(-1, 4)$
- **j** has a gradient of -4 and passes through the point $(-2, -1)$



Hint for Q7: In $y = mx + c$, insert the gradient value for m, then substitute the point for x and y to find c.

Problem-solving and reasoning 8, 9 8–11

8 The coordinates $(0, 1)$ mark the take-off point 1 metre above the ground for a rocket constructed as part of a science class. The positive x direction is considered to be east. Find the equation of the rocket's path if it rises at a rate of 5 m vertically for every 1 m in an easterly direction.

9 A line has gradient -2 and y-intercept at $(0, 5)$. Find its x-intercept coordinates.



Hint for Q9: First write the rule of the line.

10 Brad starts the year with \$80 in his bank account. He adds money into his account each week and does not take money out. After 4 weeks he has \$220 in his account. If the relationship is linear:
- **a** find a rule for the amount of money in the account after x weeks
- **b** use your rule to find when he will have \$500.



Hint for Q10: Plot the linear information on a graph to help.

11 For the line connecting the following pairs of points:
- **i** find the gradient
- **ii** find the equation using $y = mx + c$

a $(2, 6)$ and $(4, 10)$ **b** $(1, 7)$ and $(3, -1)$
c $(-3, 6)$ and $(5, -2)$ **d** $(-4, -8)$ and $(1, -3)$



Hint for Q11: Compare with Example 21.

Two pieces of information make a linear rule — 12, 13

Assume that the relationships between the variables in these questions are linear.

12 Water is leaking from a tank. The volume of water in the tank after 1 hour is 100 L and after 5 hours the volume is 20 L.
- **a** Find the equation of the line joining the points to find a rule. Use y for volume and x for time.
- **b** Use the rule to state the amount of water that was in the tank to start with.

13 At 1 minute before midnight, the temperature inside a house in winter was -5°C. The heater was switched on at this time. Three minutes after midnight, the temperature had reached 3°C.
- **a** Find the rule to represent this. (Plot the points on axes and find the equation of the line joining them. Use y for temperature and x for time.)
- **b** Find the x-intercept and the y-intercept coordinates.
- **c** Describe what the points in part **b** represent in a practical sense.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5I Midpoint and length of a line segment

Learning intentions

- To know that a line segment is defined between two endpoints
- To be able to find the midpoint of a line segment given the endpoints
- To understand how Pythagoras' theorem can be used to find the length of a line segment
- To be able to find the length of a line segment

Key vocabulary: midpoint, line segment, coordinates, endpoints, average, Pythagoras' theorem, hypotenuse, horizontal, vertical, length


A line continues on forever, but a line segment (or line interval) has endpoints; this means it has a defined length and therefore must have a midpoint. Both the midpoint and length can be found by using the coordinates of the endpoints.

A seesaw consists of a long beam with a pivot point (or fulcrum). We can consider the long beam to be a line segment with its midpoint at the pivot point of the seesaw.



Lesson starter: Choosing a method

This graph shows a line segment between the points at $(-2, -1)$ and $(2, 3)$.



- What is the horizontal distance between the two points?
- What is the vertical distance between the two points?
- Discuss and explain a method for finding the length of a line segment, using the right-angled triangle formed.
- What is the x-coordinate of the point halfway along the line segment?
- What is the y-coordinate of the point halfway along the line segment?
- Discuss and explain a method for finding the midpoint of a line segment.


Using graphing software or dynamic geometry software, produce a line segment like the one shown above. Label the coordinates of the endpoints and the midpoint. Also find the length of the line segment. Now drag one or both of the endpoints to a new position.

- Describe how the coordinates of the midpoint relate to the coordinates of the endpoints. Is this true for all positions of the endpoints that you choose?
- Now use your software to calculate the vertical distance and the horizontal distance between the two endpoints. Then square these lengths. Describe these squared lengths compared to the square of the length of the line segment. Is this true for all positions of the endpoints that you choose?


Essential Mathematics Core for the Victorian Curriculum Year 9 ISBN 978-1-009-59315-1 © Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Key ideas

- A **line segment** is a section of a straight line between two points (**endpoints**).
- The **midpoint** (M) of a line segment is the halfway point between the two endpoints.
  - The x-coordinate is the **average** (mean) of the x-coordinates of the two endpoints.
  - The y-coordinate is the average (mean) of the y-coordinates of the two end points.
  - $M = \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right)$



- The **length** of a line segment (or line interval) is found using Pythagoras' theorem.
  - The line segment is the hypotenuse (longest side) of a right-angled triangle.
  - The coordinates can be used to find the horizontal distance between them, and the vertical distance.



Horizontal distance from -3 to -1 is 2.
Vertical distance from 2 to 5 is 3.
Pythagoras' theorem:

$c^2 = 2^2 + 3^2$

$= 13$

$\therefore c = \sqrt{13}$

Exercise 5I

Understanding

1–3 | 3

1 Find the number that is halfway between these pairs of numbers.

a 5, 11 **b** -2, 4


c -6, 0 **d** 4, 7




Hint for Q1: If in doubt, add them and divide by 2.

2 Use Pythagoras' theorem to find the length of the hypotenuse in these right-angled triangles. Round to one decimal place where required.


a




b



c




d





Hint for Q2: Pythagoras' theorem:



3 **a** Find the horizontal distance between the points:

i (1, 4) and (4, 6) **ii** $(-2, 3)$ and $(2, -1)$

b Find the vertical distance between the points:

i $(-2, 3)$ and (0, 5) **ii** (2, 5) and $(6, -2)$



Hint for Q3: For horizontal distance, consider the x-coordinates. For vertical distance, consider the y-coordinates.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

51

Fluency

4–5(½) 4–5(½)

Example 22 Finding the midpoint of a line segment

Find the midpoint $M(x, y)$ of the line segment joining these pairs of points.

a $(1, 0)$ and $(5, 4)$
b $(-3, -2)$ and $(5, 3)$

Solution

a $x = \frac{1+5}{2} = 3$

$y = \frac{0+4}{2} = 2$

$\therefore M = (3, 2)$

b $x = \frac{-3+5}{2} = 1$


$y = \frac{-2+3}{2} = 0.5$

$\therefore M = (1, 0.5)$

Explanation

Find the average (mean) of the x-coordinates and y-coordinates for both points.

By plotting the points to form the line segment, you can check that the coordinates you find for the midpoint appear to be the halfway point of the line segment.



Now you try

Find the midpoint $M(x, y)$ of the line segment joining these pairs of points.

a $(2, 1)$ and $(6, 9)$
b $(-4, -1)$ and $(0, 6)$

4 Find the midpoint $M(x, y)$ of the line segment joining these pairs of points.

a $(0, 0)$ and $(6, 6)$
b $(0, 0)$ and $(4, 4)$
c $(0, 2)$ and $(2, 8)$
d $(3, 0)$ and $(5, 2)$
e $(-2, 0)$ and $(0, 6)$
f $(-4, -2)$ and $(2, 0)$
g $(1, 3)$ and $(2, 0)$
h $(-1, 5)$ and $(6, -1)$
i $(-3, 7)$ and $(4, -1)$
j $(-2, -4)$ and $(-1, -1)$
k $(-7, -16)$ and $(1, -1)$
l $(-4, -3)$ and $(5, -2)$






Hint for Q4: Check that your coordinates for the midpoint appear to lie halfway along the line segment.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Example 23 Finding the length of a line segment

Find the length of the segment joining $(-1, 2)$ and $(4, -2)$, correct to two decimal places.

Solution



Explanation

Plot the points and form the line segment to visualise the problem.

Form the right-angled triangle.

Use the coordinates to find the lengths of the horizontal and vertical sides.

Horizontal length = 5
Vertical length = 4

Horizontal length: from -1 to 4 is 5 units.
Vertical length: from 2 to -2 is 4 units.

$\therefore c^2 = 5^2 + 4^2$

Apply Pythagoras' theorem $c^2 = a^2 + b^2$.

$= 25 + 16$

$= 41$

From $c^2 = 41$, take the square root of both sides.

$\therefore c = \sqrt{41}$

$\therefore$ length = 6.40 (to 2 d.p.)

Round to two decimal places on a calculator as required.

Now you try

Find the length of the segment joining $(-2, 3)$ and $(1, -3)$, correct to two decimal places.

5 Find the length of the segment joining these pairs of points, correct to two decimal places.

a $(1, 1)$ and $(2, 6)$
b $(1, 2)$ and $(3, 4)$
c $(0, 2)$ and $(5, 0)$
d $(-2, 0)$ and $(0, -4)$
e $(-1, 3)$ and $(2, 1)$
f $(-2, -2)$ and $(0, 0)$
g $(-1, 7)$ and $(3, -1)$
h $(-4, -1)$ and $(2, 3)$
i $(-3, -4)$ and $(3, -1)$

Problem-solving and reasoning

6–8 | 7–10

6 A fence line on the plans for a property shows fence posts at points with coordinates $(-4, 8)$, $(2, 3)$ and $(8, -2)$. To reinforce the fence, two posts are to be placed halfway between the current posts. At what coordinates will these posts be placed?



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

51

7 Find the missing coordinates in this table if M is the midpoint of points A and B.

A	B	M
$(4, 2)$		$(6, 1)$
	$(0, -1)$	$(-3, 2)$
	$(4, 4)$	$(-1, 6.5)$



Hint for Q7: in the first row, 6 is halfway between 4 and ?, while 1 is halfway between 2 and ?.



8 A circle has its centre at $(2, 1)$. Find the coordinates of the endpoint of a diameter if the other endpoint has these coordinates.

a $(7, 1)$
b $(3, 6)$
c $(-4, -1)$

Hint for Q8: A diameter of a circle is shown: It passes through the centre.




9 Find the perimeter of these shapes correct to one decimal place.

a A triangle with vertices $(-2, 0)$, $(-2, 5)$ and $(1, 3)$.
b A trapezium with vertices $(-6, -2)$, $(1, -2)$, $(0, 4)$ and $(-5, 4)$.

Hint for Q9: Start by plotting the points to form the line segments that make up the shape.

10 For the point $(3, 2)$ shown:


a list four points that are 3 units from this point and add them to the axes
b describe the shape that would be formed by all the points that are 3 units from $(3, 2)$.



Division by ratio — 11

11 Looking from left to right, this line segment shows the point $P(-1, 0)$ which divides the segment in the ratio $1 : 2$.

a What fraction of the horizontal distance between the endpoints is P from A?
b What fraction of the vertical distance between the endpoints is P from A?
c Find the coordinates of point P on the segment AB if it divides the segment in these ratios.
i $2 : 1$ **ii** $1 : 5$ **iii** $5 : 1$



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


5J Linear modelling ★

Learning intentions

- To understand that many situations can be modelled by a linear rule or graph
- To be able to find a rule linking two variables
- To know how the dependent and independent variable are positioned on a set of axes when graphing the rule
- To be able to use a rule or graph to estimate the value of one variable given the other

Key vocabulary: dependent variable, independent variable, linear, gradient, rate

If a relationship between two variables is linear, the graph will be a straight line. The equation linking the two variables can therefore be written in gradient–intercept form. The area of mathematics which uses line graphs and rules to explore the relationship between two variables is called linear modelling. A test car, for example, increasing its speed from 100 km/h to 200 km/h in 8 seconds with constant acceleration, could be modelled by the rule $s = 12.5t + 100$. This rule could then be used to calculate the speed at different times in the test run.



Lesson starter: The test car

The above graph describes the speed of a racing car over an 8-second period.

- Explain how the rule $s = 12.5t + 100$ for the graph is found by describing what the 100 means and what the 12.5 means.
- Why might negative values of t not be considered for the graph?
- How could you accurately calculate the speed at the 6.5 second mark?
- If the car continued to accelerate at the same rate, how could you accurately predict the car's speed at the 13.2 second mark?



The speed of a racing car can be modelled by a rule linking the two variables of speed and time.

Key ideas

- Many situations can often be modelled by using a linear rule or graph.
  The key elements of linear modelling include:
  - Finding the rule linking the two variables
  - Sketching a graph
  - Using the graph or rule to predict or estimate the value of one variable given the other
  - Finding the rate of change of one variable with respect to the other variable. This is the same as finding the gradient.
- On a set of axes the **dependent variable** goes on the vertical axis and the **independent variable** goes on the horizontal axis.
  - The dependent variable changes as a result of changes that occur in the independent variable.
  - If the variables are speed and time, then speed is on the vertical axis, since the speed reached depends on the time passed.
  - Time is usually on the horizontal axis.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Exercise 5J

Understanding

1–3 | 3

1 A person gets paid \$50 plus \$20 per hour. Decide which rule describes the relationship between the pay, $\$P$, and number of hours, n.

A $P = 50 + n$ **B** $P = 50 + 20n$ **C** $P = 50n + 20$ **D** $P = 20 + 50$

2 The amount of money in a bank account is \$1000 and it is increasing by \$100 per month.

a Find the amount of money in the account after:

i 2 months **ii** 5 months **iii** 12 months

b Write a rule for the amount of money, A dollars, after n months.

3 **a** Insert the word *independent* or *dependent* in the following.

i The _____ variable goes on the vertical axis.

ii The _____ variable goes on the horizontal axis.

b Choose the dependent variable in the following cases.

i The volume of water, V, in a tank t minutes after turning on the hose.

ii The height, h, of a tree t days after it was planted.

iii The amount of money earned from the sale of m items.



Hint for Q3: Which variable changes in response to changes in the other variable?

Fluency

4–6 | 4–7

Example 24 Applying linear relations

Netshare is an internet provider. It charges \$20 up front plus \$5 per week of use.

a Write a rule for the total cost, $\$C$, of using Netshare for t weeks.

b Sketch the graph of C versus t using $0 \leq t \leq 10$.

c What is the total cost when Netshare is used for 4 weeks?

d If the total cost was \$50, for how many weeks was Netshare used?


Solution

a $C = 20 + 5t$

b C-intercept is 20

At $t = 10$, $C = 20 + 5 \times (10) = 70$

$\therefore$ end point is $(10, 70)$



Explanation

A fixed amount of \$20 plus \$5 for each week. This is $5 \times t$.

The cost, C, is the dependent variable and so it goes on the vertical (y) axis.

Let $t = 0$ to find the C-intercept (this is the point on the vertical axis).

Letting $t = 10$ gives $C = 70$ and this gives the other endpoint.

Sketch the graph using the points $(0, 20)$ and $(10, 70)$.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

c $C = 20 + 5t$
$= 20 + 5 \times (4)$
$= 40$

Substitute $t = 4$ into the rule.

The cost is \$40

Answer the question using the correct units.

d $C = 20 + 5t$
$50 = 20 + 5t$
$30 = 5t$
$\therefore t = 6$

Write the rule and substitute $C = 50$. Solve the resulting equation for t by subtracting 20 from both sides then dividing both sides by 5.

Netshare was used for 6 weeks.

Answer the question in words.

Now you try

A gardener charges \$60 upfront plus \$30 per hour of work.

a Write a rule for the total cost, \$$C$, of hiring the gardener for n hours.
b Sketch the graph of C versus n using $0 \leq n \leq 6$.
c What is the total cost for a job that takes 3 hours?
d If the total cost was \$210, how many hours was the gardener hired for?

4 A sales representative earns \$400 a week plus \$20 for each sale she makes.

a Write a rule that gives the total weekly wage, \$$W$, if the sales representative makes x sales.
b Draw a graph of W versus x using $0 \leq x \leq 40$.
c How much does the sales representative earn if, in a particular week, she makes 12 sales?
d If, in a particular week, the sales representative earns \$1000, how many sales did she make?



Hint for Q4: To sketch over $0 \leq x \leq 40$, find the endpoints; that is, the point where $x = 0$ and where $x = 40$.

5 A plumber charges a \$40 fee upfront per job, and \$50 for each hour he works.

a Find a linear equation for the total charge, \$$C$, for n hours of work.
b What will a 4-hour job cost?
c If the plumber works on a job for two days and averages 6 hours per day, what will be the total cost?

6 A catering company charges \$500 for the hire of a marquee (a giant tent), plus \$25 per guest.

a Write a rule for the cost, \$$C$, of hiring a marquee and catering for n guests.
b Draw a graph of C versus n for $0 \leq n \leq 100$.
c How much would a party catering for 40 guests cost?
d If a party cost \$2250, how many guests were catered for?



7 The cost, \$$C$, of producing a music single is \$300, plus \$120 per hour of studio time.

a Write a rule for the cost, \$$C$, of producing a single requiring t hours of studio time.
b Draw a graph of C versus t for $0 \leq t \leq 10$.
c How much does a single using 6 hours of studio time cost?
d If a single cost \$660 to make, for how long was the studio used?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5J

Problem-solving and reasoning 8, 9 9–11

8 A petrol tank holds 66 litres of petrol. If it starts with 12 litres of petrol and the petrol pump fills it at 3 litres every 10 seconds, find:

a a linear equation for the amount of fuel (F litres) in the tank after t minutes

b how long it will take to fill the tank.



Hint for Q8: Read units carefully. 3 L in 10 seconds equals 18 L in 1 minute.

9 A tank starts with 4000 L of water. Water from it is used at a rate of 20 L per minute.

a Write a rule for the volume, V litres, of water after t minutes.

b Calculate the volume after 1.5 hours.

c How long will it take for the tank to be emptied?

d How long will it take for the tank to have only 500 L?



Hint for Q9: If the water is being used, think about what should be happening to the volume of water in the tank.

10 The rule for distance travelled, d km, over a given time, t hours, for a moving vehicle is given by $d = 50 + 80t$.

a What is the speed of the vehicle?

b If the speed was actually 70 km per hour, how would this change the rule? Write the new rule.

11 The altitude, h metres, of a helicopter t seconds after it begins its descent is given by $h = 350 - 20t$.

a Use the rule to find or state:

i the rate at which the helicopter's altitude is decreasing

ii the helicopter's initial height ($t = 0$)

iii how long it will take for the helicopter to reach the ground.

b If the rule was $h = 350 + 20t$, describe what the helicopter would be doing.



Hint for Q11: Altitude is the height above ground at which the helicopter is flying.

Where will they meet? — 12

12 Car A heads out of town on the open highway travelling at an average speed of 65 km/h.

a Write a rule for the distance, d km, of the car from the town after t hours.

b Sketch a graph of distance (d) vs time (t) for $0 \leq t \leq 5$.

At the same time as Car A leaves town, Car B sets out on the highway from the opposite direction, travelling towards the town. The rule for the distance, d km, of Car B from the town after t hours is given by $d = 350 - 75t$.

c How many kilometres from town does Car B begin?

d How long will it take for Car B to reach the town?

e Sketch the graph of the rule for Car B on the same axes that you used in part **b**.

f Using your graph, find:

i how many hours into the journey the two cars will pass each other.

ii how many kilometres out of town they will be when they pass each other.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Trading in foreign currencies

Many professions use linear relations to model aspects of their jobs. FOREX traders or currency traders use their knowledge of trends and future growth to buy and sell currencies from around the world.

It is the biggest market in the world. People often trade in shares, futures and overseas currencies from the privacy of their home.



1 The currency of one country is directly proportional to the currency of another country. Each currency exchange can be written as an equation in the form $y = mx$.
The top line of this table states that 1 euro (EUR) = 1.5687 Australian dollars (AUD).
Writing this as an equation gives: $y = 1.5687 \times x$ where x = EUR, y = AUD.

	From (x: input currency)	To (y: output currency)	Exchange rate
i	The EURO (EUR)	Australian dollars (AUD)	1.5687
ii	British pounds (GBP)	AUD	2.0306
iii	USD	Chinese Yuan (CNY)	6.5181
iv	EUR	GBP	0.7725
v	AUD	EUR	0.6375
vi	AUD	United states dollar (USD)	0.7158
vii	AUD	Singapore dollar (SGD)	1.0038
viii	AUD	New Zealand dollar (NZD)	1.0753

Use the exchange rate table above to complete a table with these column headings. Work with four decimal places for the linear equation and three decimal places for y when $x = 10$ in the 'Two points' column.

x: input currency	y: output currency	Linear equation	Two points $(0, _)$ and $(10, _)$

2 Use the information in your answer to Question **1** for the following tasks.
- **a** For exchange rates **i** to **iv**, sketch each linear relation on a separate number plane. Label the currency on each axis and label each point where $x = 10$.
- **b** What feature of the straight line does the exchange rate represent in each graph?
- **c** For exchange rates **v** to **viii**, sketch each linear relation on the same number plane. Label each point where $x = 10$.
- **d** What do you notice about the slope of the line as the exchange rate increases?

Essential Mathematics Core for the Victorian Curriculum Year 9 ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Using technology

3 Currency rates fluctuate over time. Follow the Excel spreadsheet instructions below to graph a trendline and determine the trendline equation for the exchange rate of Australian dollars to Japanese yen.

May	AUD to JPY
1	80.48545
2	80.98463
3	81.15554
4	82.53038
5	82.53038
6	83.17927
7	84.55183
8	84.55183

a i Enter the data from this table into an Excel spreadsheet.

ii Select both columns.

iii Choose Insert and Scatter graph and click on the icon of unconnected points.

iv Click on one of the graphed points, right click, and select 'Add trend line'.

v From the trendline options that appear, select 'linear' and at the bottom of the callout tick 'display equation on chart'.

vi Label the graph.

b Copy the trendline equation. Define the x and y for the trendline.

c What does the gradient of this trendline tell us?

d Use the trendline equation to predict the exchange rate for May 10.

e If the exchange rate continues to follow this trend, how many Japanese yen would you expect to receive when exchanging $1500 AUD on 15 May?

f Give one reason why a currency trendline equation can't be relied on to make accurate predictions.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Running in a Triathlon

Jack, Levi and Max compete in a junior Triathlon event, first swimming 200 m, then cycling 12 km and finally running 1200 m.

Jack was the first to start the running leg and he runs at an average speed of 3 m/s. Levi runs at 3.5 m/s and Max runs at 4 m/s. However Max and Levi take longer to complete the swimming and cycling and start the run after Jack started. Can either of them overtake Jack and win the race?

For this task, d metres is the distance travelled by a runner and t seconds is the time in seconds after Jack starts running.

Present a report for the following tasks and ensure that you show clear mathematical workings, explanations and diagrams where appropriate.

1 Preliminary task

a Find the time, in seconds, taken to run 1200 m at an average speed of 2.4 m/s.

b Write the rule for distance d after t seconds when running at a speed of 2.4 m/s.

c A rule describing a different runner is given by: $d = 2.5t - 20$. Let $d = 1200$ and solve the equation to find this runner's finishing time.


2 Modelling task

Formulate

The problem is to find the times of the first, second and third place runners and their time differences.

a Write down all the relevant information that will help solve this problem.

b Using a large sheet of graph paper or digital graphing software, prepare axes like those shown below. Draw and label a horizontal axis showing time t to 400 seconds and a vertical axis showing distance d to 1200 m.



Solve

c Jack runs at an average speed of 3 m/s for the 1200 m.

- **i** How long, in seconds, does Jack take to run the 1200 m?
- **ii** Write a rule for Jack's distance (d metres) in terms of time (t seconds).
- **iii** Draw a line on the graph to represent Jack's run. Label this line: Jack.

d Levi starts running after Jack had started. Levi runs at 3.5 m/s and the rule describing Levi's distance d after t seconds is given by: $d = 3.5t - 175$.

- **i** Let $d = 0$ and find t, Levi's starting time. Plot this point and label its coordinates.
- **ii** Find Levi's distance when $t = 350$. Plot this point and label its coordinates.
- **iii** Draw a line through these points extend it to 1200 m. Label this line: Levi.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Solve

e What occurs on the graph at $t = 350$? Explain what happens between Jack and Levi at 350 s.

f Max starts the running race behind Levi and Jack. Max runs at 4 m/s and the rule describing Max's distance d after t seconds is given by: $d = 4t - 300$.

- **i** Let $d = 0$ and find t, Max's starting time. Plot this point and label its coordinates.
- **ii** Find Max's distance when $t = 250$. Plot this point and label its coordinates.
- **iii** Rule a line through these points and extend it to 1200 m. Label this line: Max.

g What occurs on the graph at $t = 250$? Explain what happens between Max and Levi at 250 seconds.

h Does Max also overtake Jack? If so, state the time and distance where this happens.

i From your graph determine who gets first, second and third place.

Evaluate and verify

j Let $d = 1200$ in each equation and solve each for t to find the runner's finishing times, to the nearest second.

k By how many seconds was the first place runner ahead of the second runner and the third runner?

Communicate

l Summarise your results and describe any key findings.

3 Extension question

A fourth runner, Elijah, started 80 seconds after Jack started and finished at the same time as Jack.

a Answer the following questions about the line on the graph that would show Elijah's run.

- **i** State the coordinates of the start and end points that would be on Elijah's line.
- **ii** Calculate the gradient of the line that joins these two points.
- **iii** What feature of Elijah's running does this gradient represent?
- **iv** The line equation is in the form: $d = mt + c$. Find the values of m and c and hence the equation of the line that would represent Elijah's run.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Hybrid linear journeys

Key technology: Graphing

When straight lines are used to describe travel situations, we use lines with different gradients and y-intercepts over different values of t (time) on the horizontal x-axis. Such relations are called hybrid or piecewise functions. For each part of the graph, we limit the x values and describe them using inequality symbols. For example, the rule $y = 50x + 100$ might be valid for $0 \leqslant x \leqslant 5$.



1 Getting started

A journey in a car involves the following three components.

- Section A: Travelling 200 km in 2 hours
- Section B: Resting for 3 hours
- Section C: Travelling 150 km in 3 hours

a Find the speed of the car for:
i Section A **ii** Section B **iii** Section C.


b What is the total distance travelled?

c Draw a travel graph for this journey using t for time in hours on the horizontal x axis and d for distance on the vertical axis in kilometres.

2 Using technology

We will now use three linear rules and a graphing program like Desmos to create the travel graph.

The image shows how to create the three rules for the different values of x. The values of a, b, c, d, f and g define the gradients and y-intercepts of the graph and are not yet correct for the given journey description in part **1** above.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

a Build a Desmos graph as shown adding sliders for the values of a, b, c, d, f and g and the points $(0,\ 0)$, $(2,\ 200)$, $(5,\ 200)$, $(8,\ 350)$. Adjust the range for each slider to suit.

b Now adjust your sliders so that the three lines pass through the correct points and that your graph matches your graph from part **1c**.

c Write down the rules for the three graphs and check that they are correct by seeing if the points satisfy the equations exactly. Do this by substituting a selected point into one of the rules which passes through that point.

3 Applying an algorithm

We will now create a new graph using technology for a four-part journey using the points $(0, 0)$, $(3, 240)$, $(5, 400)$, $(6, 400)$ and $(8, 600)$.

a Follow these steps.

- **i** Step 1: Add the five given points and adjust the graph scales to suit.
- **ii** Step 2: Add a linear relation of the form $y = ax + b$ for the time between the first two points.
- **iii** Step 3: Add sliders for the pronumerals a and b.
- **iv** Step 4: Alter the sliders so that the graph passes through the two points.
- **v** Step 5: Repeat from Step 2 moving to the next pair of points until your travel graph is complete.

b Write down the rules for the four graphs and check that they are correct by seeing if the points satisfy the equations exactly.

c Calculate the gradient of each graph and describe what the gradient means in terms of the journey.




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

1 Matches are arranged by a student such that the first three diagrams in the pattern are:



How many matches are in the 50th diagram of the pattern?

2 The first three shapes in a pattern made with matchsticks are:




How many matchsticks make up the 100th shape?

3 A tank with 520 L of water begins to leak at a rate of 2 L per day. At the same time, a second tank is being filled at a rate of 1 L per hour starting at 0 L. How long does it take for the tanks to have the same volume?

4 The points $(-1, 4)$, $(4, 6)$, $(2, 7)$ and $(-3, 5)$ are the vertices of a parallelogram. Find the midpoints of its diagonals. What do you notice?

5 Prove that the triangle with vertices at the points $A(-1, 3)$, $B(0, -1)$ and $C(3, 2)$ is isosceles.



6 Bill takes 3 days to paint a house. Rashid takes 4 days to paint a house. Lucy takes 5 days to paint a house. How long would it take to paint a house (to the nearest hour) if all three of them worked together?



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


Chapter summary

Direct proportion

$y = mx$ means y is directly proportional to x.
m = *gradient*; the rate of change of y with respect to x.
e.g. travelling at 60 km/h for t hours, distance $d = 60t$.

Finding the equation of a line

Gradient–intercept form ($y = mx + c$)
Require gradient and y-intercept or any other point to substitute.
E.g.



$m = \frac{\text{rise}}{\text{run}} = \frac{2}{1} = 2$
y-int is $(0, -2)$
$\therefore\ y = 2x - 2$

Midpoint and length of line segment

Midpoint is halfway between segment endpoints.

$M = \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right)$

E.g. midpoint of segment joining $(-1, 2)$ and $(3, 6)$


$x = \frac{-1 + 3}{2} = 1$

$y = \frac{2 + 6}{2} = 4$, i.e. $(1, 4)$

Length of segment

Use Pythagoras' theorem $c^2 = a^2 + b^2$


E.g.



$c^2 = 4^2 + 5^2$
$c = \sqrt{41}$
$c = 6.4$ (to 1 d.p.)

Gradient


This measures the slope of a line.



Gradient $m = \frac{\text{rise}}{\text{run}}$

$m = \frac{4}{2} = 2$

e.g.



Linear modelling

Define variables to represent the problem and write a rule relating the two variables.


Straight line

$y = mx + c$
gradient, y-intercept


In the form $ax + by = d$, rearrange to $y = mx + c$ form to read off gradient and y-intercept.

Lines with one intercept


Horizontal line $y = c$
e.g. $y = 3$



Vertical line $x = b$
e.g. $x = 2$




$y = mx$ passes through the origin, substitute $x = 1$ to find another point, e.g. $y = 3x$



Linear relationships

x- and y-intercepts



x-intercept has y-coordinate 0.
y-intercept has x-coordinate 0.

E.g. for $2x + 3y = 6$

y-int ($x = 0$)
$2(0) + 3y = 6$
$3y = 6$
$y = 2$
y-int is at $(0, 2)$

x-int ($y = 0$)
$2x + 3(0) = 6$
$2x = 6$
$x = 3$
x-int is $(3, 0)$

Sketching linear graphs


Sketch with two points. Often we find the x-intercept ($y = 0$) and y-intercept ($x = 0$)

E.g. $y = 2x - 4$

y-int: $x = 0$, $y = 2(0) - 4$
$y = 4$ $(0, 4)$


x-int: $y = 0$, $0 = 2x - 4$
$2x = 4$
$x = 2$ $(2, 0)$

Plot these points and join in a line.



A linear relation is made up of points (x, y) that form a straight line when plotted,
e.g. $y = 2x + 1$


x	–2	–1	0	1	2
y	–3	–1	1	3	5



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter checklist





A version of this checklist that you can print out and complete can be downloaded from your Interactive Textbook.

		✔
5A	**1 I can plot points to graph a straight line.** e.g. Complete the table of values below for the rule $y = -2x + 1$ and plot a graph. x: -2, -1, 0, 1, 2 y:	
5A	**2 I can decide whether a point is on a line.** e.g. Decide whether the point $(2, -1)$ is on the line with the given rules: **a** $y = 2x - 3$ **b** $y = 3x - 7$	
5B	**3 I can identify x- and y-intercepts.** e.g. Read off the x-intercept and y-intercept coordinates from this table and graph **a** x: -2, -1, 0, 1, 2 y: 12, 8, 4, 0, -4 **b** 	
5B	**4 I can find the y-intercept.** e.g. Find the coordinates of the y-intercept for the linear relations: **a** $y = 3x - 4$ **b** $4x - 5y = 20$	
5B	**5 I can find the x-intercept.** e.g. Find the coordinates of the x-intercept for the linear relations: **a** $y = 3x - 4$ **b** $4x - 5y = 20$	
5C	**6 I can sketch linear relations of the form $ax + by = d$ using intercepts.** e.g. Sketch the graph of $2x + 7y = 14$, showing the x- and y-intercepts	
5C	**7 I can sketch linear relations of the form $y = mx + c$ using intercepts.** e.g. Sketch the graph of $y = 3x - 9$ showing the x- and y-intercepts	
5D	**8 I can graph vertical and horizontal lines.** e.g. Sketch the graph of the following vertical and horizontal lines **a** $y = -2$ **b** $x = 5$ **c** $y = 0$	
5D	**9 I can sketch lines that pass through the origin.** e.g. Sketch the graph of $y = 5x$	


Chapter checklist

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter checklist

		✔
5D	**10 I can find the equation of horizontal and vertical lines.** e.g. Give the equation of the following vertical and horizontal lines **a**  **b** 	
5E	**11 I can find the gradient of a line.** e.g. For each graph, state whether the gradient is positive, negative, zero or undefined then find the gradient where possible. **a**  **b** 	
5E	**12 I can find the gradient between two points.** e.g. Find the gradient (m) of the line joining the points: $A(-2, 3)$ and $B(1, 7)$	
5F	**13 I can form a direct proportion rule.** e.g. Write a rule linking the variables d km and t hours if I travelled 420 km in 7 hours.	
5F	**14 I can work with direct proportion.** e.g. A 400 L aquarium is filled with water. It takes 40 minutes to fill the tank. **a** Draw a graph of volume (V litres) vs time (t minutes) using $0 \leq t \leq 40$. **b** Find the gradient of the graph and form a rule for V. **c** Use the rule for V to find the time to fill 250 L.	
5G	**15 I can state the gradient and y-intercept from $y = mx + c$.** e.g. State the gradient and the y-intercept coordinates for the graph of $y = 3x - 2$.	
5G	**16 I can rearrange linear equations into the form $y = mx + c$.** e.g. Rearrange these linear equations into the form $y = mx + c$: **a** $3x + y = 4$ **b** $6x + 2y = 12$	
5G	**17 I can sketch linear graphs using the gradient and y-intercept.** e.g. Find the value of the gradient and y-intercept coordinates for these relations and sketch their graphs: **a** $y = 4x - 3$ **b** $3x + 2y = 10$ (rearrange first)	

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

		✔
5H	**18** **I can find the equation of a line given the y-intercept and another point.** e.g. Determine the equation of the straight line shown here.  	
5H	**19** **I can find the equation of a line given the gradient and a point.** e.g. Find the equation of the line that has a gradient, m, of 2 and passes through the point $(3, -2)$.	
5I	**20** **I can find the midpoint of a line segment.** e.g. Find the midpoint $M(x, y)$ of the line segment joining $(1, -1)$ and $(5, 6)$.	
5I	**21** **I can find the length of a line segment.** e.g. Find the length of the segment joining $(-2, 1)$ and $(3, 5)$ correct to two decimal places.	
5J	**22** **I can apply linear relations.** e.g. A wedding photographer charges a $\$200$ upfront fee plus $\$80$ per hour. **a** Write a rule for the total cost, $\$C$, of hiring the photographer for t hours. **b** Sketch the graph of C vs t using $0 \leq t \leq 6$. **c** Determine the total cost if the photographer is hired for 4 hours, and how many hours the photographer is hired for if the cost is $\$400$.	

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter review

Short-answer questions


5A **1** Using $-2 \leq x \leq 3$, construct a table of values and plot a graph for $y = 2x - 3$.

5B **2** Read off the coordinates of the x- and y-intercepts from the table and graph.

a

x	-2	-1	0	1	2
y	0	2	4	6	8

b



5C **3** Sketch the following linear graphs, labelling x- and y-intercepts.

a $y = 2x - 4$ **b** $y = 3x + 9$ **c** $y = -2x + 5$ **d** $y = -x + 4$

e $2x + 4y = 8$ **f** $4x - 2y = 10$ **g** $2x - y = 7$ **h** $-3x + 6y = 12$


5D **4** Sketch the following lines with one intercept and classify their gradient as positive, negative, zero or undefined.

a $y = 3$ **b** $y = -2$ **c** $x = -4$


d $x = 5$ **e** $y = 3x$ **f** $y = -2x$

5E **5** Find the gradient of the following lines.

a



b



5E **6** Find the gradient of the line passing through these points.

a $(3, 1)$ and $(5, 5)$ **b** $(2, 5)$ and $(4, 3)$

c $(-3, -2)$ and $(1, 6)$ **d** $(-2, 6)$ and $(1, -4)$

5F **7** An empty swimming pool is being filled with water at a constant rate by a hose. It takes 4 hours to fill 8000 L.

a What is the rate at which water is poured into the pool?

b Draw a graph of volume (V litres) vs time (t hours) for $0 \leq t \leq 4$.

c By finding the gradient of your graph, give the rule for V in terms of t.

d Use your rule to find the time to fill 5000 L.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5G **8** For each of the following linear relations, state the value of the gradient and the y-intercept coordinates and then sketch using the gradient–intercept method.


a $y = 2x + 3$ **b** $y = -3x + 7$

5G **9** Rearrange the following equations into the form $y = mx + c$ and state the gradient and y-intercept coordinates.


a $2x + y = 6$ **b** $x + 2y = 4$

5H **10** Find the equations of these linear graphs in the form $y = mx + c$.


a



b



c



5H **11** Give the equation of the straight line that:

a has gradient 3 and passes through the point $(1, 4)$

b has gradient -2 and passes through the point $(2, -1)$

5I **12** For the line segment joining the following pairs of points, find:

i the midpoint **ii** the length (to two decimal places)

a $(2, 4)$ and $(6, 8)$ **b** $(5, 2)$ and $(10, 7)$


c $(-2, 1)$ and $(2, 7)$ **d** $(-5, 7)$ and $(-1, -2)$

Multiple-choice questions

5A **1** The point(s) on the Cartesian plane shown with an x-coordinate of 2 are:

A B only **B** A only **C** C, D and E

D B and D **E** E and D



5B **2** The coordinates of the x-intercept of the graph of $2x + 4y = 12$ are:

A $(4, 0)$ **B** $(0, 3)$ **C** $(6, 0)$


D $(2, 0)$ **E** $(0, 2)$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


Chapter review

5C **3** The graph of $y = 3x - 6$ is represented by:


A




B




C



D



E



5C **4** The point that is not on the straight line $y = 2x - 1$ is:


A $(1, 1)$ **B** $(0, -1)$ **C** $(3, 5)$ **D** $(-2, 7)$ **E** $(-1, -3)$

5E **5** The gradient of the line joining the points $(1, -2)$ and $(5, 6)$ is:

A 2 **B** 1 **C** 3 **D** -1 **E** $\frac{1}{3}$

5D **6** The graph shown has equation:

A $y = x - 2$
B $y = -2x$
C $y = -2$
D $x = -2$
E $x + y = -2$




5H **7** A straight line has a gradient of -2 and passes through the point $(0, 5)$. Its equation is:

A $2y = -2x + 10$ **B** $y = -2x + 5$ **C** $y = 5x - 2$
D $y - 2x = 5$ **E** $y = -2(x - 5)$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5H **8** The equation of the graph shown is:



A $y = x + 1$

B $y = \frac{1}{2}x + 4$

C $y = 2x + 1$

D $y = -2x - 1$

E $y = \frac{1}{2}x + 2$

5I **9** The length of the line segment joining the points $A(1, 2)$ and $B(4, 6)$ is:

A 5 **B** $\frac{4}{3}$ **C** 7

D $\sqrt{5}$ **E** $\sqrt{2}$

5J **10** A satellite phone call costs a 20c connection fee plus 60c per minute. A rule to represent the cost, C cents, of a call lasting n minutes is:

A $C = 20n + 60$ **B** $n = 60C + 20$ **C** $C = 60n + 20$

D $n = 20C + 60$ **E** $20C + 60n = 1$

Extended-response questions

1 Justin is hiring a clown for his son's birthday party. The clown charges a booking fee of \$100 as well as \$50 per hour at the party.

- **a** Write a rule for the cost (C dollars) of hiring the clown for t hours.
- **b** Sketch a graph of cost (C) vs time (t) for $0 \leq t \leq 5$.
- **c** How much will it cost to hire the clown for 4 hours?
- **d** At the end of the party, Justin writes out a cheque for \$225. How long was the clown at the party for?

2 Luca runs a marathon at a constant speed. The distance, d km, he has remaining in the marathon t hours after the start is given by $d = 42 - 14t$.

- **a** Find the t- and d-intercepts for the rule and sketch its graph between these points.
- **b** What is the distance of the marathon?
- **c** How long does Luca take to run the marathon?
- **d** Find the gradient of the graph.
- **e** What speed is Luca running at?



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Semester review 1

Reviewing number

Short-answer questions

1 Evaluate the following.

a $\frac{7}{4}+\frac{1}{4}$ **b** $\frac{13}{10}-\frac{3}{10}$ **c** $2\frac{1}{4}+\frac{1}{2}$ **d** $2\frac{1}{3}-1\frac{5}{9}$

2 Evaluate:

a $\frac{9}{10}\times\frac{5}{12}$ **b** $3\frac{3}{4}\div 2\frac{1}{12}$

3 Round the following to three decimal places.

a 4.1255 **b** 21.00241 **c** 0.0096

4 Divide \$800 into the following ratios.

a 1 : 4 **b** 2 : 3 **c** 7 : 1

5 Write these rates and ratios in simplest form.

a Prize money is shared between two people in the ratio 60 : 36.

b Jodie travels 165 km in 3 hours.

c 30 mL of rain falls in $1\frac{1}{4}$ hours.

6 A car averages 68 km/h on a journey. How far will it travel at this speed for 2 hours and 40 minutes? Round to one decimal place.

7 Determine the best buy if 3 kg of Royal Gala apples costs \$10.50 while 5 kg of Royal Gala apples cost \$15.

Multiple-choice questions

1 $7\times(-5)+3\times(-4)$ is equal to:

A 23 **B** −47 **C** 47 **D** −23 **E** 420

2 $45.5\div 9.5$, by first using rounding to one significant figure, gives:

A 4.9 **B** 5.35 **C** 5.4 **D** 5 **E** 4.95

3 \$450 is divided in the ratio 4 : 5. The value of the smaller portion is:

A \$210 **B** \$250 **C** \$90 **D** \$200 **E** \$220

4 Con runs 4.5 km in 20 minutes. His speed is:

A 1.5 km/h **B** 1.5 km/min **C** 13.5 km/h **D** 90 km/h **E** 22.5 km/h

5 $\frac{7}{10\,000}+\frac{13}{100}$ is equal to:

A 0.713 **B** 0.0713 **C** 0.1307 **D** 0.83 **E** 7.13

Extended-response questions

Thomas walks, on average, 6 km an hour. Phillip walks at an average speed of 8 km/h.

a How far does Thomas walk in 4 hours?

b Phillip starts at 8 a.m.

i If he stops at 9.45 a.m, how far has he walked?

ii What time will Phillip finish his walk if he walks twice as far as Thomas did in part **a**?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Financial mathematics

Short-answer questions

1 Convert each of the following to a percentage.

a 0.6 **b** $\frac{5}{16}$ **c** 2 kg out of 20 kg **d** 75c out of $3

2 Find:

a 10% of $96 **b** 85% of $900 **c** $5\frac{1}{2}$% of $200

3 **a** Increase 80 m by 5%. **b** Decrease $54 by 6%.

4 Julie spends 20% of her day (24 hours) reading a novel. How many hours and minutes did she spend reading?

5 Jamal earns a weekly retainer of $400 plus 6% of the sales he makes. If he sells $8200 worth of goods, how much will he earn for the week?

6 Clive's salary is $84 800 and the taxation rate for his salary is 26% for the amount above $18 200, which is tax free. Calculate Clive's net income for the year.

7 Sean invests $800 at 6% p.a. for 18 months.

a Calculate his simple interest.

b What is the final balance on Sean's account after the interest has been paid?

Multiple-choice questions

1 $2\frac{1}{2}$% is the same as:

A 0.25 **B** $\frac{1}{40}$ **C** $\frac{25}{100}$ **D** 2.5 **E** $\frac{5}{2}$

2 If cost price = $36 and selling price = $28.80, the percentage loss is:

A 7.2% **B** 80% **C** 92.8% **D** 20% **E** 2%

3 The simple interest earned on $660 invested at 5% p.a. for $1\frac{1}{2}$ years is:

A $49.50 **B** $40.50 **C** $594 **D** $33 **E** $709.50

4 A book that costs $27 is discounted by 15%. The new price is:

A $20.25 **B** $31.05 **C** $4.05 **D** $22.95 **E** $25.20

5 Anna is paid a normal rate of $12.10 per hour. If in a week she works 6 hours at the normal rate, 2 hours at time and a half and 3 hours at double time. How much does she earn?

A $181.50 **B** $145.20 **C** $193.60 **D** $175.45 **E** $163.35

Extended-response questions

Jill plans to trial a simple interest plan. Before investing her money she increases the amount in her account by 20% to $21 000.

a What was the original amount in her account?

b She invests the $21 000 for 4 years at a simple interest rate of 3% p.a. How much does she have in her account at the end of the four years?

c She continues with the plan in part **b** and after a certain number of years has obtained $5670 interest. For how many years has she had the money invested?

d What percentage increase does this interest in part **c** represent on her initial investment?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Semester review 1

Expressions and equations

Short-answer questions

1 Simplify the following.

a $2x + 6y - 4x + y$ **b** $-3m \times 5n$ **c** $\frac{6xy}{18x}$

2 Solve the following equations.

a $3x + 7 = 25$ **b** $\frac{x-1}{4} = 2$ **c** $4(2m + 3) = 15$

d $5a - 8 = 3a - 2$ **e** $2 - 3x < 8$ **f** $-3(x - 1) \geq 12$

3 Noah receives m dollars pocket money per week. His older brother Jake gets 1 dollar more than twice Noah's amount. If Jake receives $15:

a write an equation to represent the problem using the variable m

b solve the equation in part **a** to determine how much Noah receives each week.

4 The formula $S = \frac{n}{2}(a + l)$ gives the sum, S, of a sequence of n numbers with first term a and last term l.

a Find the sum of the sequence of 10 terms 2, 5, 8, … , 29.

b If a sequence of 8 terms has a sum of 88 and a first term equal to 4, use the formula to find the last term of this sequence.

Multiple-choice questions

1 The expression that represents 3 more than half a certain number, n, is:

A $\frac{1}{2}n + 1.5$ **B** $2n + 3$ **C** $\frac{3}{2}n$ **D** $\frac{n}{2} + 3$ **E** $\frac{n+3}{2}$

2 The simplified form of $5ab + 3a + 2ab - a$ is:

A $9ab$ **B** $10ab + 3a$ **C** $13ab - a$ **D** $7ab + 3$ **E** $7ab + 2a$

3 The expanded form of $-2(3m - 4)$ is:

A $-6m + 8$ **B** $-6m + 4$ **C** $-6m - 8$ **D** $-5m - 6$ **E** $5m + 8$

4 The solution to $\frac{d}{4} - 7 = 2$ is:

A $d = -20$ **B** $d = 15$ **C** $d = 36$ **D** $d = 1$ **E** $d = 30$

5 The formula $m = \sqrt{\frac{b-1}{a}}$, with $b = 9$ and $a = 2$, gives m equals:

A 3 **B** $\frac{3}{2}$ **C** 2 **D** 3 **E** $\frac{3}{2}$

Extended-response questions

Chris and his brother Michael play basketball. In the last game, Chris scored 12 more points than Michael. Between them they scored 38 points. Let p be the number of points scored by Michael.



a Write an expression for the number of points scored by Chris.

b Write an equation involving the unknown, p, to represent the problem.

c Solve the equation to find the number of points scored by each of Chris and Michael.


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Pythagoras' theorem and trigonometry


Short-answer questions

1 Find the value of each pronumeral, correct to one decimal place.


a




b



c




d




2 Find the value of each pronumeral correct to one decimal place.


a




b




c




d



e




f



3 A wire is to be connected from the edge of the top of a 28 m high building to the edge of the top of a 16 m high building. The buildings are 15 m apart.

a What length of wire is required? Round to two decimal places.


b What is the angle of depression from the top of the taller building to the top of the smaller building? Round to one decimal place.



4 For this diagram:


a use Pythagoras' theorem to find the value of x

b use trigonometry to find the value of y correct to two decimal places.



Multiple-choice questions


1 The length of the hypotenuse in this triangle is:



A 11 cm **B** 13 cm **C** 14.8 cm

D 169 cm **E** $\frac{1}{119}$ cm

2 The value of x in the triangle shown is:




A 3.9 **B** 113 **C** 57

D 15 **E** 2.7

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


Semester review 1

3 The value of x in the triangle shown is closest to:



A 6.7 **B** 7.4 **C** 14.9
D 13.5 **E** 9.0


4 A correct ratio for this triangle is:



A $\tan\theta = \frac{x}{y}$ **B** $\cos\theta = \frac{y}{z}$ **C** $\tan\theta = \frac{y}{x}$

D $\sin\theta = \frac{y}{x}$ **E** $\cos\theta = \frac{x}{y}$


5 A 5-metre plank of wood is leaning up against a side of a building as shown. If the wood touches the ground 3 m from the base of the building, the angle the wood makes with the building is closest to:



A 36.9° **B** 59° **C** 53.1° **D** 31° **E** 41.4°

Extended-response questions

A skateboard ramp is constructed as shown.



a Calculate the distance d metres up the ramp correct to two decimal places. Use Pythagoras' theorem.

b What is the angle of elevation (θ) between the ramp and the ground correct to one decimal place?

c If the skateboarder rides from one corner of the ramp diagonally to the other corner, what distance would be travelled? Round to one decimal place.

Linear relationships

Short-answer questions

1 Sketch the following linear graphs labelling x- and y-intercepts.

a $y = 2x - 6$ **b** $3x + 4y = 24$ **c** $y = 4x$ **d** $y = 3$

2 Find the gradient of each of the following.

a The line passing through the points $(-1, 2)$ and $(2, 4)$


b The line passing through the points $(-2, 5)$ and $(1, -4)$

c The line with equation $y = -2x + 5$

d The line with equation $-4x + 3y = 9$

3 Give the equation of the following lines in gradient–intercept form.

a The line with the given graph



b The line with gradient 3 and passing through the point $(2, 5)$


4 Find the midpoint and length of the line segment joining the following points.

a $(0, 2)$ and $(6, 10)$ **b** $(-1, 5)$ and $(2, 1)$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


Multiple-choice questions

1 The x- and y-intercepts respectively for the graph shown are at:



A $(-2, 4)$ and $(4, -2)$
B $(0, 4)$ and $(-2, 0)$
C $(-2, 0)$ and $(0, 4)$
D $(4, 0)$ and $(0, -2)$
E $(2, 0)$ and $(0, -4)$

2 The graph shown has equation:



A $y = 3x$ **B** $y = 3$
C $y = x + 3$ **D** $x = 3$
E $x + y = 3$

3 The line passing through the points $(-3, -1)$ and $(1, 1)$ has gradient:

A $\frac{1}{2}$ **B** -3 **C** 1 **D** 2 **E** -2

4 If the point $(-1, 3)$ is on the line $y = 2x + c$, the value of c is:

A 1 **B** 5 **C** -7 **D** -5 **E** -1

5 The line with equation $3x + y = 4$ has a gradient of

A -3 **B** 3 **C** 1
D 4 **E** -4

Extended-response questions

Doug works as a labourer. He is digging a trench and has 180 kg of soil to remove. He has taken 3 hours to remove 36 kg at a constant rate.

a What is the rate at which he is removing the soil?
b If he maintains this rate, write a rule for the amount of soil, S kg, remaining after t hours.
c Draw a graph of your rule.
d How long will it take to remove all of the soil?
e Doug is paid \$40 for the job plus \$25 per hour worked.
  i Write a rule for his pay P dollars for working h hours.
  ii How much will he be paid to remove all the soil?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

6 Measurement



Essential mathematics: why measurement skills are important

Measurement is fundamental in various fields to ensure accuracy and efficiency. Knowing how to measure length, perimeter, area, and volume is valuable in many real-life situations.

Tailors and fashion designers measure body dimensions for perfectly fitting garments. They optimise fabric usage by planning pattern layouts efficiently, minimising waste. By measuring the fabric roll's width and strategically placing pattern pieces, they maximise fabric use, ensuring high-quality, aesthetically pleasing, and sustainable fashion.

Plumbers accurately measure pipe lengths to cut them correctly, preventing leaks and water damage. Calculating the volume of water tanks and pipes is crucial for determining water flow rates and capacity. Plumbers also measure angles to ensure smooth water flow and prevent blockages and leaks.

Filmmakers measure focal length and distances for perfect shot composition and depth of field, creating visually compelling scenes that convey the intended mood. They consider the area within each frame, ensuring elements are well-balanced and visually appealing.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.



In this chapter

6A Errors and accuracy in measurement
6B Length and perimeter (Consolidating)
6C Circumference of a circle (Consolidating)
6D Area (Consolidating)
6E Area of a circle (Consolidating)
6F Composite shapes
6G Surface area of prisms
6H Surface area of a cylinder
6I Volume
6J Volume of a cylinder

Victorian Curriculum 2.0

This chapter covers the following content descriptors in the Victorian Curriculum 2.0:

NUMBER

VC2M9N01

MEASUREMENT

VC2M9M01, VC2M9M04, VC2M9M05

Please refer to the curriculum support documentation in the teacher resources for a full and comprehensive mapping of this chapter to the related curriculum content descriptors.

© VCAA

Online resources


A host of additional online resources are included as part of your Interactive Textbook, including HOTmaths content, video demonstrations of all worked examples, auto-marked quizzes and much more.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025
Photocopying is restricted under law and this material must not be transferred to another party.


Warm-up quiz

1 Count the number of squares in these shapes to find the area.

a



b



c




2 Find the distance (perimeter) around the shapes in Question **1**.

3 Count the number of cubes in these solids to find the volume.


a




b



c




4 **a** Name the solid shown at right.
b What is the name of the shape (shaded) on top of the solid?



5 Calculate the following by moving the decimal point right (for ×) or left (for ÷).

a 2.3×10
b 0.048×1000
c $270 \div 100$
d $52134 \div 10\,000$
e $0.0005 \times 100\,000$
f $72\,160 \div 1000$

6 Estimate the area (number of squares) in this oval.




7 Convert the following to the units shown in the brackets. Remember: 1 km = 1000 m, 1 m = 100 cm and 1 cm = 10 mm.

a 3 cm (mm)
b 20 m (cm)
c 1.6 km (m)
d 23 mm (cm)
e 3167 m (km)
f 72 cm (m)
g 20 000 m (km)
h 0.03 km (m)
i 75.6 m (km)


8 Find the area of these basic shapes using the given formulas.

a




$A = l \times w$

b




$A = \frac{1}{2}bh$

c




$A = l^2$

d




$A = \frac{1}{2}bh$

e



$A = b \times h$

f



$A = l \times w + l^2$

9 Find the circumference ($C = 2\pi r$) and area ($A = \pi r^2$) of this circle, rounding to two decimal places.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

6A Errors and accuracy in measurement

Learning intentions

- To understand that all measurements are estimates
- To be able to calculate absolute, relative and percentage errors in measurement
- To be able to find suitable upper and lower bounds for measurements and related calculations

Key vocabulary: error, absolute value, limits of accuracy, estimate, percentage error

The Hubble space telescope was launched in 1990 and has delivered millions of observations helping to produce thousands of scientific papers including stunning images of galaxies and other astronomical objects. Unfortunately, the first images produced by the telescope were fuzzy and out of focus due to an error in the shape of the main parabolic mirror. One theory is that a slight error in the calibration of the machine which polished the mirror resulted in an error of less than $\frac{1}{50}$ of the width of a human hair in the mirror's shape. This was enough to cause the received images to be out of focus. The problem was rectified by using a special lense to reverse the error in a similar way that eyeglasses are used to correct people's vision.



The Hubble Space Telescope in orbit after its lens was fixed in 1993. This repair allowed Hubble to take clear and amazing pictures of space.

Lesson starter: Underweight gobbles

A confectionery company's most popular product is called a gobble which can be purchased in bags of 50. Larry, an employee of the company, uses a digital scale which displays a whole number of grams for each measurement. Larry places one gobble on the scale which displays 7 grams. He then calculates the weight of a bag of 50 gobbles as $50 \times 7 = 350$ grams.

a Would it be possible for the actual weight of the gobble to be the following? Give a reason in each case.
i 7.2 grams **ii** 6.3 grams **iii** 6.52 grams **iv** 7.56 grams

Assume now that the weight of each gobble is actually 6.71 grams when rounded to two decimal places.

b Find the weight of a bag of gobbles.
c Find the difference in the weight of a bag of gobbles and the weight of the bag calculated by Larry.
d Express your difference in part **d** above as a percentage of the actual weight of a bag of gobbles.
e As a consumer would you be satisfied or dissatisfied by the fact that the bag label says 350 grams? Give a reason.

Key ideas

- All measurements are estimates of the actual quantity. The accuracy of each measurement may depend on:
  - the scale on the measuring instrument
  - the units used in the measurement
  - the person's ability in using the instrument and reading the scale.
- **Error** = measurement − actual value
  - The error could be positive or negative depending on the measurement.
  - The absolute value of the error gives the error as a positive number, ignoring the sign.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

6A

- Percentage error $= \frac{\text{Error}}{\text{Actual value}} \times \frac{100}{1}$
- The **limits of accuracy** are the smallest and largest possible values that a measurement could be.
  - A ruler with both cm and mm markings, for example, might produce a 36 mm recording. The lower limit would be 35.5 mm and the upper limit would be (but not including) 36.5 mm.
  - Lower limit = recorded measurement $- 0.5 \times$ smallest unit of measure
  - Upper limit = recorded measurement $+ 0.5 \times$ smallest unit of measure

Exercise 6A

Understanding


1–4 | 1–4

1 What is the smallest unit marked on these measuring devices?


a Ruler



b Thermometer



c Water container



2 Find the absolute value of the following errors. Recall that the absolute value of the error gives the error as a positive number ignoring the negative sign if necessary.

a -23 grams **b** -1.78 litres **c** -41.1 seconds

3 Danny estimates that he might spend \$100 on a trip to the city one day but he actually spends \$80.

a What is the difference between Danny's estimate and his actual expenditure?

b Express the difference calculated in part **a** as a percentage of the actual expenditure. Use the formula:

Percentage error $= \frac{\text{Error}}{\text{Actual value}} \times \frac{100}{1}$

4 Matty uses a digital scale which records the number of kilograms as whole numbers. Decide if the following actual weights would produce a digital readout of 12 kg.

a 12.3 kg **b** 12.42 kg **c** 12.61 kg **d** 11.83 kg **e** 11.49 kg

Fluency

5–6(½), 7, 8 | 5–6(½), 7, 9(½)

Example 1 Calculating absolute percentage error

Calculate the absolute percentage error for the following measurements and actual values.

a Measurement: 2.9 kg Actual value: 2.5 kg

b Measurement: 35 minutes Actual value: 40 minutes

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Solution

a Error $= 2.9 - 2.5 = 0.4$ kg

$$\%\text{Error} = \frac{0.4}{2.5} \times \frac{100}{1}$$
$$= 16\%$$

b Error $= 35 - 40 = -5$ minutes

Absolute error $= 5$ minutes

$$\%\text{Error} = \frac{5}{40} \times \frac{100}{1}$$
$$= 12.5\%$$

Explanation

Error = measurement − actual value

$$\text{Percentage error} = \frac{\text{Error}}{\text{Actual value}} \times \frac{100}{1}$$

To calculate the absolute error, ignore the negative sign after calculating the error.

Calculate the percentage error as per part **a** above.

Now you try

Calculate the absolute percentage error for the following measurements.

a Measurement: 44 seconds Actual value: 40 seconds

b Measurement: 4.7 litres Actual value: 5.0 litres

5 Calculate the absolute percentage error for the following measurements and actual values.

a Measurement: 11 kg Actual value: 10 kg
b Measurement: 19 L Actual value: 20 L
c Measurement: 26 seconds Actual value: 25 seconds
d Measurement: 2.8 kg Actual value: 3.2 kg



Hint for Q5:
Error = measurement − actual value
Percentage error $= \frac{\text{Error}}{\text{Actual value}} \times \frac{100}{1}$

6 Calculate the absolute percentage error for the following measurements and actual values.

a Measurement: 57 g Actual value: 60 g
b Measurement: 23.5 hours Actual value: 25 hours
c Measurement: 3.4 mL Actual value: 3.2 mL
d Measurement: 2.49 kL Actual value: 2.4 kL

Example 2 Finding the limits of accuracy in a measurement

A single chocolate is weighed using a digital scale which displays measurements in grams correct to one decimal place. The measurement given is 6.7 g.

a State the lower and upper limits of accuracy for this measurement.

b If the chocolates are sold in bags of 10, find the lower and upper limits of accuracy for the weight of a bag of chocolates.

Solution

a Lower limit $= 6.7 - 0.5 \times 0.1$
$= 6.65$ g

Upper limit $= 6.7 + 0.5 \times 0.1$
$= 6.75$ g

Explanation

Subtract $0.5 \times$ smallest unit (0.1 grams) to the recorded measurement.

Add $0.5 \times$ smallest unit (0.1 grams) to the recorded measurement.

Continued on next page

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

6A

b Calculated weight = $10 \times 6.7 = 67$ g

Maximum error = $10 \times 0.05 = 0.5$ g

Lower limit = $67 - 0.5$
= 66.5 g

Upper limit = $67 + 0.5$
= 67.5 g

First find the weight of the bag using the recorded measurement and calculate the maximum error possible.

Subtract or add the maximum error possible to calculate the lower and upper limits for the weight of the bag of 10 chocolates.

Now you try

A single teddy bear biscuit is weighed using a digital scale which displays measurements in grams correct to the nearest whole number. The measurement given is 58 g.

a State the lower and upper limits of accuracy for this measurement.

b If the biscuits are sold in boxes of 10, find the lower and upper limits of accuracy for the weight of a box of biscuits.

7 A single packet of chips is weighed using a digital scale which displays measurements in grams correct to one decimal place. The measurement given is 25.5 g.

a State the lower and upper limits of accuracy for this measurement.

b If the chips are sold in boxes of 20, find the lower and upper limits of accuracy for the weight of a box.



8 A single pen is weighed using a digital scale which displays in grams correct to the nearest whole number. The measurement given is 15 g.

a State the lower and upper limits of accuracy for this measurement.

b If the pens are sold in boxes of 30, find the lower and upper limits of accuracy for the weight of a box.

9 Give the limits of accuracy for these measurements.

a 36 cm	**b** 103 m	**c** 3.2 mm
d 6.8 minutes	**e** 45.24 hours	**f** 83.51 g
g 3.106 kg	**h** 0.0357 L	**i** 0.02 mL



Hint for Q9: Use $\pm 0.5 \times$ smallest unit.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Problem-solving and reasoning 10–13 11–14

10 A single bottle of drink is measured using a measuring jug which includes a scale with markings for each 10 mL. The measurement given is 1.51 L.

a State the lower and upper limits of accuracy for this measurement.

b If the drinks are sold in crates of 10, find the lower and upper limits of accuracy for the volume of drinks in a crate.



11 At a school fair you guess that there are 155 lollies in a jar when actually there were 180. Calculate your absolute percentage error. Round to one decimal place.

12 Armid measures the weight of one of 15 identical figurines. The scale gives a whole number of grams only and displays 45 grams. What is the difference between the lowest and highest possible weight of all 15 figurines?



13 Kelly measures her height and calculates a percentage error of 0.5%. If her actual height is 173.4 cm, find Kelly's possible recorded height. Round to one decimal place.

Hint for Q13: work out the error in cm first.

14 One way to reduce the error when using measuring instruments is to take multiple readings and find the mean (average) of those measurements. The distance between two towns was measured four times and the results were 22.8 km, 22.9 km, 23.2 km and 23.0 km. The actual distance is known to be 23.1 km.

a Find the mean of the four recorded measurements.

b Find the percentage error using your result from part **a**. Round to two decimal places.

World's most embarrassing mistakes — 15

15 One of the causes of some of history's most embarrassing mistakes relates to the misuse of measurement systems.

a The $125 million Mars Climate orbiter was lost due to some confusion between units. The Jet Propulsions Laboratory used the Metric system for their calculations and the Astronautics Division used Imperial units. Assuming that one inch (2.54 cm) was mistaken for a centimetre find the percentage error correct to one decimal place.

b In 1628 the Swedish Vasa Warship sank after sailing only a small distance out to sea. It was discovered that the measuring rulers used for one side of the ship were Swedish foot long rulers of length 29.69 cm including 12 inches and the rulers used for the other side of the ship were Amsterdam foot long rulers of length 28.31 cm which included 11 inches. This meant that one side of the ship contained more timber than the other side and hence the ship tilted and sank. Assuming that the Swedish 12 inch ruler was the correct ruler to use, find the percentage error made by the carpenters who used the Amsterdam 11 inch ruler if they were trying to measure a piece of wood which was supposed to be 2 feet 6 inches long. Round to one decimal place.



c Research other famous embarrassing mistakes made using measuring instruments or units. Choose one example and calculate the percentage error made.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

6B Length and perimeter CONSOLIDATING

Learning intentions

- To review the metric units of length
- To be able to convert between metric units of length
- To be able to find the perimeter of a simple closed shape

Key vocabulary: perimeter, length

Measurement is the branch of mathematics that includes the consideration of length, area and volume. This includes units of measurement, perimeter, circumference, surface area, composite shapes and capacity.

We use measurement when we design buildings, water our gardens, control satellites, fill our cars with petrol or participate in school athletics days.




A space shuttle uses mathematical formulas to track its path through space. These formulas calculate the shuttle's speed, position, and acceleration, ensuring it stays on the correct course.

Lesson starter: Not enough information?


All the angles at each vertex in this shape are 90° and the two given lengths are 10 cm and 13 cm.

- Is there enough information to find the perimeter of the shape?
- If there is enough information, find the perimeter and discuss your method.




Key ideas

- Units of length in the metric system include kilometre (km), metre (m), centimetre (cm) and millimetre (mm)
- To convert between metric units of length, multiply or divide by the appropriate power of 10.




1 km = 1000 m
1 m = 100 cm
1 cm = 10 mm

- **Perimeter** is the distance around the outside of a closed shape.
  - Sides with the same markings are of equal length



$P = 2 \times 5 + 3$
$= 13$ cm



$P = 2 \times 11 + 2 \times 4$
$= 30$ m


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Exercise 6B

Understanding

1–3 | 3

1 Fill in the gaps on this flow chart.




2 Choose from the words: *multiply*, *divide*, *add* or *subtract* to complete this sentence.


To find the perimeter of a shape you would _____ all the side lengths.

3 Write down the value of x in these diagrams.


a



b



c



Fluency

4(½), 5, 6(½) | 4(½), 6(½), 7

Example 3 Converting units of length

Convert to the units shown in the brackets.

a 5.41 cm (mm) **b** 3200 m (km)

Solution

a 5.41 cm = 5.41 × 10 mm
= 54.1 mm

Explanation

1 cm = 10 mm and you are moving to a smaller unit, so multiply.

Move the decimal point one place to the right.

b 3200 m = 3200 ÷ 1000 km
= 3.2 km

1 km = 1000 m and you are moving to a larger unit, so divide.

Move the decimal point three places to the left.

Now you try

Convert to the units shown in the brackets.

a 2.4 m (cm) **b** 420 mm (cm)

4 Convert the following length measurements into the units given in the brackets.

a 5 cm (mm) **b** 41 cm (mm)
c 2.8 m (cm) **d** 0.4 m (cm)
e 4.6 km (m) **f** 0.9 km (m)
g 521 mm (cm) **h** 36 mm (cm)
i 240 cm (m) **j** 83.7 cm (m)
k 7000 m (km) **l** 2170 m (km)



Hint for Q4: Multiply when changing to a smaller unit and divide when changing to a larger unit.


5 A steel beam is 8.25 m long and 22.5 mm wide. Write down the length and the width of the beam in centimetres.


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

6B

Example 4 Finding perimeters of simple shapes

Find the perimeter of each of the following shapes.

a 

b 

Solution

a $P = 10 + 9 + 5 + 4$
$= 28$ cm

b $P = 2 \times 12 + 2 \times 18$
$= 24 + 36$
$= 60$ m


Explanation


To find the perimeter, add all the side lengths together.

Two lengths of 12 m and two lengths of 18 m.


Now you try


Find the perimeter of each of the following shapes.


a 


b 


6 Find the perimeter of each of the following shapes.


a 


b 


c 


d 

e 


f 


g 


h 

i 

7 Find the unknown side length in these shapes with the given perimeters.

a 

b 

c 



Hint for Q7: Subtract the known lengths from the perimeter first


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Problem-solving and reasoning


8–10 | 9–12

8 Write down the values of the pronumerals in these shapes. All angles are 90°.


a



b




c



Example 5 Finding the perimeter of composite shapes

Find the perimeter of this composite shape.



Solution

Perimeter $= (2 \times 5) + 6 + 3 + 2 + 1$

$= 22$ cm

Explanation


Missing sides are:

5 cm − 3 cm = 2 cm

6 cm − 5 cm = 1 cm


Alternatively:

$2 \times 6 + 2 \times 5 = 22$ cm




Now you try

Find the perimeter of this composite shape.




9 Find the perimeter of each of the following composite shapes.


a



b




c






Hint for Q9: First label all the missing side lengths and then find the perimeter.


d



e



f



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

6B

10 A lion cage is made up of five straight fence sections. Three sections are 20 m in length and the other two sections are 15.5 m and 32.5 m. Find the perimeter of the cage.



11 Convert the following measurements into the units given in the brackets.


a 8 m (mm)
b 110 000 mm (m)
c 0.00001 km (cm)
d 0.02 m (mm)
e 28 400 cm (km)
f 62 743 000 mm (km)




Hint for Q11: In part a, first convert to cm and then to mm.

12 Find the perimeter of these shapes. All angles are right angles.


a



b




c




Picture framing — 13, 14

13 A photo 12 cm wide and 20 cm long is surrounded with a picture frame 3 cm thick. Find the outside perimeter of the framed picture.

14 A square picture of side length 20 cm is inside a frame of width x cm.





a Find the perimeter of the framed picture if:
i $x = 2$ **ii** $x = 3$ **iii** $x = 5$

b Write a rule for the perimeter, P, of the framed picture in terms of x.

c Use your rule to find the perimeter if $x = 3.7$.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

6C Circumference of a circle CONSOLIDATING

Learning intentions

- To know the formula for the circumference of a circle
- To be able to find the circumference of a circle
- To be able to find the perimeter of semicircles and quadrants


Key vocabulary: circumference, pi, radius, diameter, circle, semicircle, quadrant

For thousands of years, mathematicians have known about a special number that links a circle's diameter to its circumference.

The Egyptians, Babylonians and ancient Indians knew about this special number, and thought its value was about 3.15.

Today we call this number pi (π) and know it to be 3.14159, correct to five decimal places.

An exact value of pi cannot be written down as a decimal, as it has an infinite number of decimal places with no pattern.



The ancient Egyptians demonstrated their understanding of the concept of pi through calculations found in the Rhind Mathematical Papyrus. This text, dating back to around 1650 BCE, shows how they calculated the area of a circle.

Lesson starter: Working with string

Use a pair of compasses, a ruler and a piece of string for this activity.


The table below contains some approximate circle measurements.

Circumference (C)	Diameter (d)	$C \div d$
18.8	6.0	
12.6	4.0	
…	…	
…	…	


- Copy the table, leaving two blank rows.
- Use your compasses to draw a circle, then measure its diameter using a ruler and its circumference using your string (try to be as accurate as possible). Add this information to the table.
- Do this again, with a different-sized circle.
- Calculate $C \div d$ in the third column for each of the four circles. What do you notice?
- What does this say about how to calculate the circumference of a circle if you know the diameter?

Key ideas

- Features of a circle:
  - **Diameter** (d) is the distance across the centre.
  - **Radius** (r) is the distance from the centre to the circumference.
  - $d = 2r$
  - **Circumference** of a circle (the distance around the outside) is given by $C = 2\pi r$ or $C = \pi d$.
  - Use $\frac{22}{7}$ or 3.14 to approximate π (**pi**), or use technology for more precise calculations.



- Special circle sectors:
  - a half circle is called a **semicircle**
  - a quarter circle is called a **quadrant**




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Exercise 6C

Understanding

1–4 | 4

1 Name the features of a circle labelled on the circle shown.






Hint for Q2: diameter is twice the radius


2 **a** What is the radius of a circle if its diameter is 5.6 cm?
b What is the diameter of a circle if its radius is 48 mm?

3 Write down the rule for the circumference of a circle using:
a r, the radius **b** d, the diameter

4 Determine the **fraction** of a circle shown in these sectors. Write the fraction in simplest form.

a


b



c


Fluency

5–7(½) | 5–7(½)

Example 6 Finding the circumference of a circle

Find the circumference of this circle correct to two decimal places.



Solution

$C = 2\pi r$
$= 2 \times \pi \times 3$
$= 6\pi$
$= 18.85$ cm (to 2 d.p.)

Explanation

Use the formula $C = 2\pi r$ or $C = \pi d$ and substitute $r = 3$ (or $d = 6$).

6π would be the exact answer and 18.85 is the rounded answer.

Now you try


Find the circumference of this circle correct to two decimal places.




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5 Find the circumference of these circles correct to two decimal places. Use a calculator for the value of π.


a




b




c



d



e



f






Hint for Q5: Write the rule $C = 2\pi r$, then substitute the radius length.

Example 7 Finding circumference using the diameter


Find the circumference of this circle, correct to two decimal places.



Solution	Explanation
$C = \pi d$	Write the formula. $C = \pi d$ is preferred since d is given.
$= \pi \times 12$	Substitute $d = 12$ and multiply by π. Use a calculator and round.
$= 37.70$ m (to 2 d.p.)	Note: 37.6991 rounds to 37.70 for two decimal places.


Now you try

Find the circumference of this circle, correct to two decimal places.




6 Find the circumference of these circles correct to two decimal places.


a




b




c




d



e



f






Hint for Q6: Write the rule $C = \pi d$ and then substitute the diameter length.


7 Find the circumference of these circles without a calculator using the given approximation of π.

a




$\pi = 3.14$

b




$\pi = 3.14$

c



$\pi = \frac{22}{7}$

d



$\pi = \frac{22}{7}$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


6C

Problem-solving and reasoning

8(½), 9, 10 | 8, 10, 11

Example 8 Working with a semicircle or quadrant

Find the perimeter of this semicircle correct to two decimal places.



Solution


$$P = \frac{1}{2} \times \pi d + 10$$
$$= \frac{1}{2} \times \pi \times 10 + 10$$
$$= 25.71 \text{ cm (to 2 d.p.)}$$

Explanation

The perimeter consists of half the circumference of a circle (with diameter 10 cm) plus the 10 cm diameter across the top.


Now you try

Find the perimeter of this quadrant correct to two decimal places.




8 Find the perimeter of these sectors correct to two decimal places.


a




b




c




d



e



f





Hint for Q8: Decide what fraction of the circumference you want and don't forget to add the straight sides.

9 Find the distance around the outside of a circular pool of radius 4.5 m, correct to two decimal places.




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


10 Find the length of string required to surround the circular trunk of a tree that has a diameter of 1.3 m, correct to one decimal place.

11 Find the perimeter of these shapes, correct to two decimal places.


a



b



c






Hint for Q11: Two semicircles of the same size make a full circle. When finding the perimeter, imagine you are walking around the edge of the shape.

The rolling wheel — 12

12 A wheel of radius 30 cm is rolled in a straight line.



a Find the circumference of the wheel correct to two decimal places.

b How far, correct to two decimal places, has the wheel rolled after completing:

i 2 rotations? **ii** 10.5 rotations?

c Can you find how many rotations would be required to cover at least 1 km in length? Round to the nearest whole number.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

6D Area CONSOLIDATING


Learning intentions

- To know the formulas for the area of common shapes
- To be able to convert metric units for area
- To be able to find the area of common shapes

Key vocabulary: area, triangle, square, rectangle, rhombus, parallelogram, trapezium, kite, perpendicular

For many common shapes, such as the parallelogram and trapezium, the rules for finding their area can be found using simple rectangles and triangles.

The parallelogram, for example, can be seen as a 'pushed over' rectangle. Its area can be calculated in the same way as a rectangle.



So $A = l \times w = b \times h$




The Dockland Office Building in Hamburg, Germany, showcases a striking parallelogram shape, combining geometric precision with modern architectural design.

Lesson starter: Build the formulas


From the introduction, you can see how the rule for the area of a parallelogram is given by $A = b \times h$.

Use the given diagrams to explain the rules for the areas of each shape.


- Triangle: $A = \frac{1}{2}b \times h$



- Trapezium: $A = \frac{1}{2}(a + b)h$




- Rhombus: $A = \frac{1}{2}xy$



Key ideas


- Conversion of area units



$10^2 = 10 \times 10 = 100$

$100^2 = 100 \times 100 = 10\,000$


$1000^2 = 1000 \times 1000 = 1\,000\,000$



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


- The **area** of a two-dimensional shape is a measure of the space enclosed within its boundaries.
  - The 'height' (h) in the formulas for the area of a triangle, parallelogram and trapezium must be **perpendicular** (at 90°) to the base.

Square




Area $= l^2$

Rectangle




Area $= l \times w$

Triangle




Area $= \frac{1}{2} b \times h$

Parallelogram




Area $= b \times h$
(Can also be used for a rhombus)

Trapezium




Area $= \frac{1}{2}(a + b)h$

Rhombus



Area $= \frac{1}{2}x \times y$
(Can also be used for a square)

Kite



$A = \frac{1}{2}x \times y$

Exercise 6D

Understanding

1–3 | 3

1 Write the missing number.


a $1 \text{ cm}^2 = 10^2 \text{ mm}^2 =$ _____ mm^2

b $1 \text{ m}^2 = 100^2 \text{ cm}^2 =$ _____ cm^2


c $1 \text{ km}^2 = 1000^2 \text{ m}^2 =$ _____ m^2

2 Count the number of squares to find the area of these shapes. Each square in each shape represents one square unit.


a



b




c






Hint for Q2: For parts **d** and **e**, note that each triangle is half of a rectangle.


d



e



f



3 Name the shape that usually has the given area formula.

a $A = lw$

b $A = \frac{1}{2}xy$ (2 shapes)

c $A = \frac{1}{2}bh$

d $A = \frac{1}{2}(a + b)h$

e $A = bh$

f $A = l^2$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

6D

Fluency

4(½), 5, 6 | 4–6(½)

Example 9 Converting units of area

Convert the following area measurements into the units given in the brackets.

a 859 mm^2 (cm^2)

b 2.37 m^2 (cm^2)

Solution

a $859\ mm^2 = 859 \div 10^2\ cm^2$
$= 8.59\ cm^2$

b $2.37\ m^2 = 2.37 \times 100^2\ cm^2$
$= 23\,700\ cm^2$

Explanation

cm^2 mm^2 859.
$\div 10^2 = 100$

$\times 100^2 = 10\,000$
m^2 cm^2 2.3700

Now you try

Convert the following area measurements into the units given in the brackets.

a 32 000 cm^2 (m^2)

b 0.4 km^2 (m^2)

4 Convert the following area measurements into the units given in the brackets.


a 2 cm^2 (mm^2)
b 0.4 cm^2 (mm^2)
c 500 mm^2 (cm^2)
d 310 mm^2 (cm^2)
e 2.1 m^2 (cm^2)
f 0.2 m^2 (cm^2)
g 210 000 cm^2 (m^2)
h 3700 cm^2 (m^2)
i 0.001 km^2 (m^2)
j 4.3 km^2 (m^2)
k 3 200 000 m^2 (km^2)
l 39 400 m^2 (km^2)





Hint for Q4: Multiply or divide by:
$10^2 = 100$
$100^2 = 10\,000$
or $1000^2 = 1\,000\,000$

Example 10 Finding areas of rectangles, triangles and parallelograms

Find the area of each of the following plane figures.

a 

b 

c 

Solution

a Area $= l \times w$
$= 7 \times 3$
$= 21\ m^2$

Explanation

Use the area formula for a rectangle. Substitute $l = 7$ and $w = 3$. Include the correct units.

b Area $= \frac{1}{2} \times b \times h$
$= \frac{1}{2} \times 11 \times 5$
$= 27.5\ km^2$

Use the area formula for a triangle.

Substitute $b = 11$ and $h = 5$, where h is the height perpendicular (at 90°) to the base.

c Area $= b \times h$
$= 3 \times 1.5$
$= 4.5\ cm^2$


Use the area formula for a parallelogram.
Multiply the base length by the perpendicular height.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


Now you try

Find the area of each of the following plane figures.


a



b




c




5 Find the area of each of the following plane figures.


a




b




c




d



e



f






Hint for Q5: Choose from
$A = l \times w$,
$A = \frac{1}{2}b \times h$
or $A = b \times h$


Example 11 Finding areas of rhombuses and trapeziums

Find the area of each of the following plane figures.

a



b



Solution

a $\text{Area} = \frac{1}{2} \times x \times y$

$= \frac{1}{2} \times 10 \times 9$

$= 45 \text{ mm}^2$

b $\text{Area} = \frac{1}{2}(a + b) \times h$

$= \frac{1}{2}(4 + 6) \times 3$

$= 15 \text{ m}^2$

Explanation

Use the area formula for a rhombus.

x and y are the lengths of the diagonals.

$\frac{1}{2} \times 10 \times 9 = 5 \times 9 = 45$

Use the area formula for a trapezium.


Substitute $a = 4$, $b = 6$ and $h = 3$.

Include the correct units.


Now you try

Find the area of each of the following plane figures.

a



b




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


6D

6 Find the area of each of the following plane figures.


a



b




c






Hint for Q6: Choose from
$A = \frac{1}{2}x \times y$ or
$A = \frac{1}{2}(a+b) \times h$


d



e



f



Problem-solving and reasoning 7, 8, 9(½) 8–11


7 A piece of land has an area of one-half a square kilometre (0.5 km^2). How many square metres (m^2) is this?

8 A rectangular park covers an area of $175\,000$ m^2. Give the area of the park in km^2.




9 Find the area of each of the following mixed-plane figures.

a




b






Hint for Q9: Choose from
$A = l^2$, $A = b \times h$,
$A = \frac{1}{2}b \times h$ or
$A = \frac{1}{2}(a+b)h$


c




d



e



f



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


10 An old picture frame that was once square now leans to one side to form a rhombus. If the distances between pairs of opposite corners are 85 cm and 1.2 m, find the area inside the frame in m^2.

11 Convert the following measurements into the units given in the brackets.

a 1.5 km^2 (cm^2) **b** 0.000005 m^2 (mm^2) **c** 75 000 mm^2 (m^2)

Windows — 12

12 Six square windows of side length 2 m are placed into a 12 m wide by 8.5 m rectangular high wall, as shown. The windows are positioned so that the vertical spacing between the windows and the wall edges are equal. The horizontal spacings are also equal.



a **i** Find the horizontal distance between the windows.

ii Find the vertical distance between the windows.

b Find the area of the wall, not including the window spaces.

c If the wall included 3 rows of 4 windows (instead of 2 rows of 3), would it be possible to space all the windows so that the horizontal and vertical spacings are the same? (Horizontal doesn't have to be the same as vertical.)



Using a calculator 6D: Measurement formulas

This activity can be found in the More Resources section of the Interactive Textbook in the form of a printable PDF.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

6E Area of a circle

CONSOLIDATING

Learning intentions

- To know the formula for the area of a circle
- To be able to find the area of circles
- To be able to find the area of semicircles and quadrants


Key vocabulary: circle, sector, radius, diameter, semicircle, quadrant, pi

We know that the circumference of a circle is connected to the diameter by the special number pi (π).

Similarly, the area of a circle is also connected to pi. This time it is the product of pi and the square of the radius that gives the area; so, $A = \pi r^2$.

Lesson starter: How does a circle become a rectangle?


Consider a circle cut into small sectors, as shown, then rearranged to form a rectangular-style shape.



- Compared to the circle, what is the height of the rectangle close to?
- Compared to the circle, what is the length of the rectangle close to?
- What does this say about the area of the rectangle and hence the area of a circle?
- What could be done with the cutting up of sectors so that the rectangular arrangement is closer to a true rectangle?


Key ideas

- The area of a circle is given by $A = \pi r^2$.
  - If the diameter is given, halve it to find the radius.
  - $\pi r^2 = \pi \times r \times r$




- For the area of a **semicircle** or **quadrant**, use the appropriate fraction.

Semicircle



$A = \frac{1}{2}\pi r^2$

Quadrant



$A = \frac{1}{4}\pi r^2$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


Exercise 6E

Understanding


1–3 | 3

1 What is the radius of these circles?


a



b



c



2 Which is the correct formula for the area of a circle?


A $A = \pi d^2$ **B** $A = \frac{1}{2}\pi r^2$ **C** $A = \pi r^2$ **D** $A = (\pi r)^2$

3 What fraction of a full circle is shown by these sectors?


a



b



c




Fluency

4(½), 5 | 4(½), 5, 6

Example 12 Finding areas of circles

Find the area of this circle correct to two decimal places.



Solution

$A = \pi r^2$

$= \pi \times (5.1)^2$

$= 81.71 \text{ cm}^2$ (to 2 d.p.)


Explanation

Write the rule and substitute $r = 5.1$.

81.7128 rounds to 81.71 since the third decimal place is 2.

Now you try

Find the area of this circle correct to two decimal places.




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


6E

4 Find the area of these circles correct to two decimal places.

a




b






Hint for Q4: Substitute the radius into $A = \pi r^2$.


c




d



e




f




5 Find the area of these circles correct to two decimal places.


a



b



c





Hint for Q5: First work out the radius.

6 Find the area of these circles using the given value for pi (π). Round to two decimal places.

a $\pi = 3.14$



b $\pi = \frac{22}{7}$



c π (from calculator)




Problem-solving and reasoning

7–9 | 7, 9–11

Example 13 Finding the area of a quadrant or semicircle

Find the area of this quadrant correct to two decimal places.



Solution

$A = \frac{1}{4} \times \pi r^2$

$= \frac{1}{4} \times \pi (3)^2$

$= 7.07 \text{ m}^2$ (to 2 d.p.)


Explanation

A quadrant is one quarter of the area of a full circle.

Substitute $r = 3$ and evaluate.

Now you try


Find the area of this semicircle correct to two decimal places.




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

7 Find the area of these sectors correct to two decimal places.


a



b



c





Hint for Q7: Determine the fraction of the full circle first.

8 A pizza of radius 15 cm is divided into quarters.
Find the area of each quarter correct to the nearest cm^2.


9 A circular rug touches the edges of a square room of side length 4 m.

a What is the radius of the rug?

b Find the area of the rug correct to two decimal places.

c Find the area not covered by the rug correct to two decimal places.

d Find the percentage of the floor that is not covered by the rug, correct to one decimal place.





Hint for Q9: % non-rug area $= \frac{\text{non-rug area}}{\text{total area}} \times \frac{100}{1}$

10 A pizza shop is considering increasing the diameter of its family pizza tray from 32 cm to 34 cm.
Find the percentage increase in area, correct to two decimal places, from the 32 cm tray to the 34 cm tray.




11 You can rearrange $A = \pi r^2$ to give $r = \sqrt{\frac{A}{\pi}}$. Use this new rule to find the radius of a circle for these areas.
Round to one decimal place where necessary.

a $10\ cm^2$ **b** $117.8\ m^2$ **c** $4\pi\ km^2$

Tennis lights — 12

12 A tennis court area is lit by 4 corner lights. The area close to each light is considered to be good (G) while the remaining area is lit satisfactorily (S).
What percentage of the area is 'good'? Round to the nearest percent.





ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

6F Composite shapes

Learning intentions


- To be able to recognise the basic shapes that make up a composite shape
- To be able to find the area of simple composite shapes

Key vocabulary: composite shape, area, perimeter

Composite shapes are made up of a number of simple shapes. You can find the perimeters and areas of composite shapes by first identifying the simple shapes and then using their perimeter and area formulas.

Lesson starter: A fraction of a circle plus a square


This diagram shows a fraction of a circle plus a square.




- How could you find the area and perimeter of the shape?
- See if you can write down a full solution for finding the area and perimeter of the shape.
- See if your teacher or another student can easily follow your solution.

Key ideas

- **Composite shapes** are made up of more than one basic shape.
- Addition and/or subtraction can be used to find areas and perimeters of composite shapes.
  - Use addition



  - Use subtraction (for area)



- The layout of the relevant mathematical working needs to make sense so that the reader of your work understands each step.

Exercise 6F


Understanding 1–3 3

1 Name the two different shapes that make up these composite shapes; e.g. square and semicircle.


a



b




c



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

2 What fraction of a circle would be considered when finding the perimeter of these shapes?


a




b



c




d





Hint for Q2: Choose from $\frac{1}{4}$, $\frac{1}{2}$ or $\frac{3}{4}$.

3 This composite shape includes a square and a quadrant ($\frac{1}{4}$ circle).



a Find the area of the square.

b Find the area of the quadrant correct to two decimal places.


c Find the total area correct to two decimal places.

Fluency

4(½), 5 | 4(½), 5, 6(½)

Example 14 Finding perimeters and areas of composite shapes

Find the perimeter and area of this composite shape, rounding answers to two decimal places.



Solution

$P = 2 \times l + w + \frac{1}{2} \times \pi d$

$= 2 \times 17 + 14 + \frac{1}{2} \times \pi \times 14$

$= 69.99$ cm (to 2 d.p.)

Area (rectangle) $= l \times w$

$= 17 \times 14$

$= 238$ cm^2

A (semicircle) $= \frac{1}{2}\pi r^2$

$= \frac{1}{2} \times \pi \times 7^2$

$= 76.969\ldots$ cm^2

Total area $= 238 + 76.969\ldots$

$= 314.97$ cm^2 (to 2 d.p.)

Explanation

3 straight sides ⊐ + semicircle arc ⊂

Recall that $C = \pi d$ (or $2\pi r$).


Substitute $l = 17$, $w = 14$ and $d = 14$.

Calculate and round to two decimal places.

The total area consists of a rectangle plus a semicircle.



+



$A = l \times w$

$A = \frac{1}{2}\pi r^2$


Round to two decimal places.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

6F


Now you try

Find the perimeter and area of this composite shape.




4 Find the perimeter and the area of each of these composite shapes, rounding answers to two decimal places where necessary.


a




b




c




d



e



f






Hint for Q4: First decide which two shapes you will be dealing with in each case.

Example 15 Finding more perimeters and areas

This shape is a square with a quadrant (quarter circle) subtracted.
Find its perimeter and area correct to two decimal places.



Solution

$P = 2 \times l + \frac{1}{4} \times 2\pi r$

$= 2 \times 8 + \frac{1}{4} \times 2\pi \times 8$

$= 28.57$ cm (to 2 d.p.)

$A = l^2 - \frac{1}{4} \times \pi r^2$

$= 8^2 - \frac{1}{4} \times \pi \times 8^2$

$= 13.73$ cm^2 (to 2 d.p.)


Explanation

Includes 2 sides and a quarter of the circumference of a circle of radius 8 cm.

Includes a square minus a quarter of the area of a circle of radius 8 cm.

Now you try


This shape is a rectangle with a semicircle subtracted. Find its perimeter and area correct to two decimal places.




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5 Find the perimeter and area of these shapes correct to two decimal places.


a



b



c







Hint for Q5: Use subtraction to find the area.

6 Find the area of the shaded part of these composite shapes.

a




b






Hint for Q6: Use subtraction in each case.


c




d



e



f



Problem-solving and reasoning

7, 8 | 8, 9(½), 10

7 An area of lawn is made up of a rectangle measuring 10 m by 15 m and a semicircle of radius 5 m. Find the total area of lawn correct to two decimal places.

8 Twenty circular pieces of pastry, each of diameter 4 cm, are cut from a rectangular layer of pastry 20 cm long and 16 cm wide. What is the area, correct to two decimal places, of pastry remaining after the 20 pieces are removed?




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


6F

9 Find the area of the shaded region of each of the following shapes by subtracting the area of the clear shape from the total area. Round to two decimal places where necessary.


a




b




c




d




e




f




g



h




i




10 Find the area of each of the following composite shapes.


a



b




c




Circular challenges — 11

11 Find the perimeter and the area of each of the following composite shapes correct to two decimal places where necessary.


a




b




c




d



e



f



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

6G Surface area of prisms


Learning intentions

- To understand how the surface area of a solid can be represented using a net
- To be able to calculate the surface area of a prism


Key vocabulary: surface area, right prism, cross-section, uniform, net

Three-dimensional objects or solids have outside surfaces that together form the surface area. Nets are very helpful for determining the number and shape of the surfaces of a three-dimensional object.

For this section we will deal with right prisms. A right prism has a uniform cross-section with two identical ends and the remaining sides are rectangles at right angles to the base and top.



Right triangular prism



A triangular-based prism acts as a mirror, refracting and dispersing light into a spectrum of colours, creating a beautiful rainbow effect. This phenomenon occurs due to the prism's unique shape and the principles of light refraction and dispersion.

Lesson starter: Drawing prisms


Prisms are named by the shape of their cross-section.

- Try to draw as many different right prisms as you can.
- Describe the different types of shapes that make up the surface of your solids.
- Which solids are the most difficult to draw and why?


Key ideas

- The **surface area** of a solid is the sum of the areas of all the surfaces.
- A **net** is a two-dimensional illustration of all the surfaces of a solid.

Cube



A net of a cube







Surface area $= 6l^2$

- A **right prism** is a solid with a **uniform** (constant) **cross-section** and with remaining sides as rectangles.
  - Prisms are named by the shape of their cross-section.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

6G

- The nets for a rectangular prism (cuboid) and triangular prism are shown here.

Solid	Net	Surface area
Rectangular prism 		$A = 2 \times \text{top} + 2 \times \text{side} + 2 \times \text{end}$
Triangular prism 		$A = \text{area } ① + \text{area } ② + \text{area } ③ + 2 \times \text{area } ④$

Exercise 6G

Understanding

1–3 | 3


1 How many faces do the following solids have?


a Rectangular prism


b Cube

c Triangular prism


2 Draw a suitable net for these prisms and name each solid.

a 

b 

c 


3 Copy and complete the working to find the surface area of these solids.

a 

$A = 2 \times (8 \times 7) + 2 \times (8 \times _____) + 2 \times (_____ \times _____)$

$= _____ + _____ + _____$

$= _____ \text{ m}^2$

b 

$A = 2 \times \frac{1}{2} \times 4 \times _____ + 5 \times 7 + 4 \times _____ + _____ \times _____$

$= _____ + _____ + _____ + _____$

$= _____ \text{ cm}^2$


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Fluency

4–5(½) | 4–5(½), 6

Example 16 Finding the surface area of a rectangular prism

Find the surface area of this rectangular prism.



Solution


$A = 2 \times \text{top} + 2 \times \text{side} + 2 \times \text{end}$

$= 2 \times (5 \times 3) + 2 \times (5 \times 2) + 2 \times (2 \times 3)$

$= 30 + 20 + 12$

$= 62 \text{ cm}^2$


Explanation



Top = 5×3
Side = 5×2
End = 2×3


Now you try

Find the surface area of this rectangular prism.




4 Find the surface area of the following rectangular prisms. Draw a net of the solid to help you.

a




b






Hint for Q4: Add up all the pairs of rectangles using $A = l \times w$.


c




d



e



f




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

6G

Example 17 Finding the surface area of a right triangular prism

Find the surface area of this triangular prism.



Solution

A = area A + area B + area C + 2 × area D


$= 12 \times 4 + 12 \times 5 + 12 \times 3 + 2 \times (\frac{1}{2} \times 4 \times 3)$

$= 48 + 60 + 36 + 12$

$= 156 \text{ cm}^2$

Explanation


Draw and label the surface area net.



The surface is made up of three rectangles and two identical triangles. Use 3 cm and 4 cm for the base and height, respectively, of the triangles.


Now you try

Find the surface area of this triangular prism.




5 Find the surface area of each of the following triangular prisms.


a




b



c



d





Hint for Q5: Each prism consists of three rectangles and two identical triangles.

6 Find the surface area of a cube of side length 1 metre.


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Problem-solving and reasoning 7, 8 8–10

7 A rectangular box is to be covered in material. How much is required to cover the entire box if it has the dimensions 1.3 m, 1.5 m and 1.9 m?



8 Two wooden boxes, both with dimensions 80 cm, 1 m and 25 cm, are placed on the ground, one on top of the other, as shown. The entire outside surface, including the underside of the bottom box, is then painted. Find the area of the painted surface in cm^2.






Hint for Q8: Only include the outside surfaces.


9 The four walls and roof of a barn (shown to the right) are to be painted.

a Find the surface area of the barn, not including the floor.

b If 1 litre of paint covers 10 m^2, find how many litres are required to complete the job.



10 Cubes of side length one unit are stacked as shown.



a Complete this table.

Number of cubes (n)	1	2	3	4	5	6	7	8	9
Surface area (S)									

b Can you find the rule for the surface area (S) for n cubes stacked in this way? Write down the rule for S in terms of n.

c Use your rule to find the surface area if there are 100 cubes.




Hint for Q10: An example of a rule is $S = 2n + 5$, but this is not the rule for this question.

Pyramids — 11


11 Pyramids consist of a base and a number of triangular faces. The surface area can be calculated by adding up all the face areas, similar to that of prisms. Remember that the area of a triangle is given by $A = \frac{1}{2}bh$.

Find the surface area of each of these pyramids. Draw a net of the solid to help you.


a



b



c



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


Progress quiz

6A **1** Calculate the absolute percentage error if the recorded measurement is 1.2 m but the actual measurement is 1.5 m.


6A **2** Lengths of ribbon are measured with a tape measure with markings correct to the millimetre. A length of ribbon is measured as 25.4 cm. State the lower and upper limits of accuracy for this measurement.

6B **3** Find the perimeter of the following shapes.

a




b




6C **4** Find the circumference of these circles correct to two decimal places.

a



b




6B/D **5** Convert the following measurements into the units given in brackets.


a 824 cm (m)
b 30 cm (mm)
c 1.24 km (m)
d 3.6 m^2 (cm^2)
e 450 cm^2 (mm^2)
f 832 cm^2 (m^2)

6D **6** Find the area of the following plane figures.


a




b



c




d




6E **7** Find the area of the following circles correct to one decimal place.

a




b



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

6C/E **8** For the semicircle shown, find (correct to one decimal place):


a its perimeter **b** its area




6F **9** For each of the composite shapes, find (correct to one decimal place where necessary):

i its perimeter **ii** its area

a




b




6G **10** Find the surface area of these prisms.

a



b





ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

6H Surface area of a cylinder


Learning intentions

- To understand how the net of a cylinder can be drawn to show the surface area
- To know the formula for the total surface area of a cylinder
- To be able to calculate the total surface area of a cylinder and simple cylindrical portions

Key vocabulary: cylinder, area, prism, circumference, net, congruent, cross-section


Like a rectangular prism, a cylinder has a uniform cross-section (a circle), but because it doesn't have rectangular sides, it isn't classified as a right prism.

A cylinder's surface area can be found by considering the curved part as a rectangle with length $2\pi r$ (or πd) and height h.



Lesson starter: Curved area

- Roll a piece of paper to form the curved surface of a cylinder.
- Do not stick the ends together, so that you can allow the paper to return to a flat surface.
- What shape is the paper when lying flat on a table?
- When curved to form the cylinder, what do the sides of the rectangle represent on the cylinder? How does this help to find the surface area of a cylinder?




Key ideas

- A **cylinder** is a solid with two parallel, **congruent** circular faces connected by a curved surface.
- Surface area of a cylinder $= 2$ circles $+$ rectangle

$$= 2 \times \pi r^2 + 2\pi r \times h$$

$$A = 2\pi r^2 + 2\pi rh$$

$2\pi r^2$: 2 circular ends; $2\pi rh$: Curved area



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Exercise 6H

Understanding

1–4 | 4


1 The formula for the surface area of a cylinder is $A = 2\pi r^2 + 2\pi rh$.

a Which part of the formula works out the area of the curved surface?


b Which part of the formula works out the area of the two ends?

2 Find the circumference of the circular end of these cylinders. Round to two decimal places.


a



b



c







Hint for Q2: Use $C = \pi d$ or $C = 2\pi r$.

3 Draw a net suited to these cylinders. Label the sides using the given measurements.

a




b




4 The curved surface of these cylinders is allowed to flatten out to form a rectangle. What would be the length and width of the rectangles? Round to two decimal places where necessary.


a



b



c





Hint for Q4: Length = $2\pi r$



A rectangular piece of paper will cover the curved surface of a cylinder. The length of the piece of paper needed to fully cover the cylinder's curved surface will be the same as the height of the cylinder. The width of the paper will be the same as the circumference of the cylinder.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


6H

Fluency

5, 6 | 5, 7

Example 18 Finding the surface area of a cylinder

Find the surface area of this cylinder, rounding to two decimal places.



Solution

Area of ends $= 2 \times \pi r^2$

$= 2 \times \pi \times 5^2$

$= 50\pi$ m^2

Area of rectangle $= l \times w$

$= 2\pi r \times h$

$= 2\pi \times 5 \times 25$

$= 250\pi$ m^2


$A = 50\pi + 250\pi$

$= 942.48$ m^2 (to 2 d.p.)

Explanation


The surface area consists of two circles ($2\pi r^2$) plus a rectangle where $l \times w = 2\pi r \times h$.

When adding to find the surface area, use your exact answers in your calculator for each part, then add to get the surface area.




Now you try

Find the surface area of this cylinder, rounding to two decimal places.




5 Find the surface area of these cylinders, rounding to two decimal places.


a



b




c






Hint for Q5: Draw a net to help:




6 Find the surface area of these cylinders, rounding to one decimal place.


a



b



c






Hint for Q6: Remember that the radius of a circle is half the diameter.


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

7 Find the area of the curved surface only for these cylinders, correct to two decimal places.


a



b



c





Hint for Q7:
For the curved part only, use $A = 2\pi r \times h$.


Problem-solving and reasoning

8, 9(½) | 9(½), 10, 11

8 Find the surface area of a cylindrical plastic container of height 18 cm and with a circle of radius 3 cm at each end, correct to two decimal places.

Example 19 Finding surface areas of cylindrical portions

Find the surface area of this half cylinder, rounding to two decimal places.



Solution

Area (two semicircles) $= \pi r^2$

$= \pi \times 2^2$

$= 4\pi$ cm^2

Area (rectangles) $= 8 \times 4 + 8 \times \frac{1}{2} \times 2\pi \times 2$


$= 32 + 16\pi$ cm^2

$A = 4\pi + 32 + 16\pi$

$= 94.83$ cm^2 (to 2 d.p.)


Explanation

The two semicircles combine to form a full circle.




Now you try

Find the surface area of this quarter cylinder, rounding to two decimal places.




9 Find the surface area of these solids, rounding to two decimal places.


a




b




c




d



e



f



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

6H

10 A water trough is in the shape of a half cylinder. Its semicircular ends have diameter 40 cm and the trough length is 1 m. Find the outside surface area, in cm^2, of the curved surface plus the two semicircular ends, correct to two decimal places.






11 A solid cylinder cut in half gives half the volume but not half the surface area. Explain why.

Roller revolutions — 12

12 A cylindrical roller is used to press crushed rock in preparation for a tennis court. The entire rectangular tennis court area is 30 m long and 15 m wide. The roller has a width of 1 m and diameter 60 cm.



a Find the surface area of the curved part of the roller, in cm^2, correct to three decimal places.

b Find the area, in m^2 to two decimal places, of crushed rock that can be pressed after:

i 1 revolution **ii** 20 revolutions

c Find the minimum number of complete revolutions required to press the entire tennis court area.



Hint for Q12: $1\ m^2 = 100 \times 100\ cm^2$



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

6I Volume

Learning intentions

- To be able to convert between metric units of volume
- To understand how the volume of a solid relates to its constant cross-section and height
- To know the common units for capacity
- To know the formula for the volume of a rectangular prism
- To be able to calculate the volume of a solid with a uniform cross-section

Key vocabulary: solid, volume, cross-section, uniform, prism, perpendicular, capacity, litres, millilitres

We use volume to describe the amount of space inside a three-dimensional object. We use metric units, such as:

- cubic kilometres for the volume of water in the sea
- cubic metres for the volume of concrete poured at a building site
- cubic centimetres for the volume of space occupied by this book
- cubic millimetres for the volume of metal in a pin.


Units for capacity (millilitres, litres, kilolitres and megalitres) are used for liquids and gases.

Lesson starter: Why length × width × height?

For most people, the first thing that often comes to mind when dealing with volume is length × width × height. But this rule only applies to finding the volume of rectangular prisms.


Let's look at a rectangular prism split into two layers.

- How many cubes sit on one layer?
- What is the area of the base? What do you notice?
- What is the height and how many layers are there?
- Why is the volume rule given by $V = lwh$ in this case?




Key ideas

- **Volume** is the amount of three-dimensional space inside an object.
- Common metric units for volume include cubic kilometres (km^3), cubic metres (m^3), cubic centimetres (cm^3) and cubic millimetres (mm^3).



$1000^3 = 1\,000\,000\,000$
$100^3 = 1\,000\,000$
$10^3 = 1000$




$1\text{ cm}^3 = 10 \times 10 \times 10\text{ mm}^3$
$= 10^3\text{ mm}^3$


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

6I

- **Capacity** is the amount of liquid a container can hold.
  - For capacity, common units include:
    - megalitres (ML) $1\text{ ML} = 1000\text{ kL}$
    - kilolitres (kL) $1\text{ kL} = 1000\text{ L}$
    - litres (L) $1\text{ L} = 1000\text{ mL}$
    - millilitres (mL)
  - Also: $1\text{ cm}^3 = 1\text{ mL}$ so $1\text{ L} = 1000\text{ cm}^3$ and $1\text{ m}^3 = 1000\text{ L}$
- A **cross-section** is the plane figure formed when you slice a solid figure parallel to one of its surfaces.
  - Volume of solids with a uniform cross-section is equal to area of cross-section (A) multiplied by height (h).
    $V = A \times h$
  - The 'height' is the length of the edge that runs perpendicular to the cross-section in any solid.



- Volume of a rectangular prism:
  $V = l \times w \times h$




Exercise 6I

Understanding


1–3 | 3

1 What is the name given to the shape of the shaded cross-section of each of the following solids?


a




b




c




d



e




f




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

2 Draw the cross-sectional shape for these prisms and state the given 'height' (perpendicular to the cross-section).


a




b




c




d



e



f





Hint for Q2: 'Perpendicular' means 'at a right angle (90°)'.

3 Write the missing number.

a The number of mm in 1 cm is _____.
b The number of mm^2 in 1 cm^2 _____.
c The number of mm^3 in 1 cm^3 is _____.
d There are _____ cm^3 in 1 m^3.
e There are _____ m^3 in 1 km^3.
f There are _____ mL in 1 L.
g There are _____ L in 1 kL.
h There is _____ cm^3 in 1 mL.

Fluency

4–5(½), 6, 8 | 4–5(½), 7, 8(½), 9

Example 20 Converting units of volume

Convert the following volume measurements into the units given in the brackets.

a 2.5 m^3 (cm^3) **b** 458 mm^3 (cm^3)

Solution

a $2.5\ m^3 = 2.5 \times 100^3\ cm^3$
$= 2\,500\,000\ cm^3$

b $458\ mm^3 = 458 \div 10^3\ cm^3$
$= 0.458\ cm^3$

Explanation

$\times 100^3 = 1\,000\,000$
m^3 cm^3 2.500000

cm^3 mm^3
$\div 10^3 = 1000$ 458.

Now you try

Convert the following volume measurements into the units given in the brackets.

a 0.6 cm^3 (mm^3) **b** 4 520 000 cm^3 (m^3)

4 Convert the following volume measurements into the units given in brackets.

a 3 cm^3 (mm^3) **b** 0.3 cm^3 (mm^3) **c** 2000 mm^3 (cm^3)
d 0.001 m^3 (cm^3) **e** 8.7 m^3 (cm^3) **f** 5900 cm^3 (m^3)
g 0.00001 km^3 (m^3) **h** 21 700 m^3 (km^3) **i** 430 000 cm^3 (m^3)




Hint for Q4:
1 $km^3 = 1000^3\ m^3$
1 $m^3 = 100^3\ cm^3$
1 $cm^3 = 10^3\ mm^3$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

6I


5 Convert these units of capacity to the units given in brackets.

a 3 L (mL) **b** 0.2 kL (L) **c** 3500 mL (L)
d 0.021 L (mL) **e** 37 000 L (kL) **f** 42 900 kL (ML)
g 2 cm^3 (mL) **h** 2 L (cm^3) **i** 1 m^3 (L)



Example 21 Finding the volume of a rectangular prism

Find the volume of this rectangular prism.



Solution

Volume = $l \times w \times h$
= $1 \times 1 \times 3$
= 3 cm^3


Explanation

The solid is a rectangular prism.

Length = 1 cm, width = 1 cm and height = 3 cm


Now you try

Find the volume of this rectangular prism.




6 Find the volume of these three-dimensional rectangular prisms.


a



b




c




7 Find the volume of each of these rectangular prisms (cuboids).


a



b




c




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

8 Find the volume of each of these three-dimensional objects. The cross-sectional area has been given.


a




b




c




d



e



f






Hint for Q8: Simply use $V = A \times h$, since the area of the cross-section is given.

Example 22 Finding the volume of a triangular prism

Find the volume of this triangular prism.



Solution

Area of cross-section $= \frac{1}{2} \times b \times h$

$= \frac{1}{2} \times 4 \times 3$

$= 6 \text{ cm}^2$


Volume = area of cross-section × height

$= 6 \times 6$

$= 36 \text{ cm}^3$


Explanation

The cross-section is a triangle.




Now you try

Find the volume of this triangular prism.




9 Find the volume of these prisms.


a



b



c



Hint for Q9: First find the area of the triangular cross-section. $A = \frac{1}{2}bh$



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

6I

Problem-solving and reasoning

10, 11 | 11–13

10 A rectangular container is 10 cm wide, 20 cm long and 8 cm high. What is the capacity of the container in litres?




Hint for Q10: There are $1000\ \text{cm}^3$ in 1 L.


11 25 L of water is poured into a rectangular fish tank which is 50 cm long, 20 cm wide and 20 cm high. Will it overflow?

12 Find the volume of these solids, converting your answer to litres.

a




b



Hint for Q12c: Area of a trapezium: $A = \frac{1}{2}(a+b)h$




c




13 This diagram is a sketch of a new 25 m swimming pool to be installed in a school sports complex.

a Find the area of one side of the pool (shaded).

b Find the volume of the pool in litres.
Use $1\ \text{m}^3 = 1000$ L.





Volume of a pyramid


— | 14

14 Someone tells you that the volume of a pyramid is half of the volume of a rectangular prism with the same base. Do you think this is true?

a Make an educated guess as to what fraction of the prism's volume is the pyramid's volume.

b Use the internet to find the actual answer to part **a**.

c Draw some pyramids and find their volume using the results from part **b**.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

6J Volume of a cylinder

Learning intentions

- To know the formula for the volume of a cylinder
- To be able to find the volume of a cylinder
- To be able to find the capacity of a cylinder in litres

Key vocabulary: volume, cylinder, capacity, radius, diameter

A cylinder with its sides at right angles to its base is called a right cylinder. It has a uniform cross-section (a circle), so its volume can be calculated in a similar way to that of a prism. Cylindrical objects are often used to store gases and liquids, so working out the volume of a cylinder is an important measurement calculation.




Cylinders are essential for safely storing and transporting gases and liquids. Their robust design withstands high pressures, ensures controlled release, and offers portability, making them indispensable in various industries.

Lesson starter: How is a cylinder like a prism?


Here is a cylinder and two prisms.

- What do they all have in common?
- Do you think you can work out the volume of the cylinder in the same way as for a prism? Why?
- What do you think the formula would be for the volume of a cylinder?



Key ideas

- The volume of a cylinder is given by
  $V = \pi r^2 \times h$ or $V = \pi r^2 h$
  - r is the radius of the circular ends
  - h is the length or distance between the circular ends




Exercise 6J

Understanding


1–3(½) | 2–3(½)

1 Find the area of these circles correct to two decimal places.


a




b



c



d



2 Convert the following into the units given in the brackets. Remember: $1\text{ L} = 1000\text{ cm}^3$ and $1\text{ m}^3 = 1000\text{ L}$.


a 2000 cm^3 (L)
b 4.3 cm^3 (mL)
c 3.7 L (cm^3)
d 1 m^3 (L)
e $38\,000\text{ L}$ (m^3)
f 0.0002 m^3 (mL)

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


6J

3 State the radius (r) and the height (h) of these cylinders.


a




b




c




d



e



f





Hint for Q3: The height is perpendicular (at 90°) to the circular ends.


Fluency

4 | 4(½)


Example 23 Finding the volume of a cylinder

Find the volume of these cylinders correct to two decimal places.

a



b



Solution

a $V = \pi r^2 h$

$= \pi \times (3)^2 \times 10$

$= 282.74 \text{ cm}^3$ (to 2 d.p.)

b $V = \pi r^2 h$

$= \pi \times (0.9)^2 \times 0.3$

$= 0.76 \text{ m}^3$ (to 2 d.p.)

Explanation

πr^2 gives the area of the circular cross-section and h is the distance between the ends.

Substitute $r = 3$ and $h = 10$ into the rule.


The diameter is 1.8 m so $r = 0.9$ m.

The distance between the ends is 0.3 m, so $h = 0.3$.


Now you try

Find the volume of these cylinders correct to two decimal places.

a




b




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

4 Find the volume of these cylinders correct to two decimal places.


a




b




c




d



e



f





Hint for Q4: Substitute the values of r and h into $V = \pi r^2 h$. Note that $r = d \div 2$ for parts **d**, **e** and **f**.

Problem-solving and reasoning

5(½), 6, 7 | 5(½), 7–10

Example 24 Finding the capacity of a cylinder

Find the capacity, in litres, of a cylinder with radius 30 cm and height 90 cm. Round to the nearest litre.

Solution

$V = \pi r^2 h$

$= \pi \times (30)^2 \times 90$

$= 254\,469 \text{ cm}^3$

$= 254$ L (to nearest litre)

Explanation

Substitute $r = 30$ and $h = 90$.


There are 1000 cm^3 in 1 L, so divide by 1000 to convert to litres.

Now you try


Find the capacity, in litres, of a cylinder with diameter 20 cm and height 45 cm. Round to the nearest litre.

5 Find the capacity, in litres, of these cylinders. Round to the nearest litre.


a




b



c



d





Hint for Q5: First work out the volume in cm^3 and then use $1 \text{ L} = 1000 \text{ cm}^3$ to find the capacity in litres.


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

6J

6 A cylindrical storage drum has a radius of 0.5 m and a height of 2 m.
- **a** Find its volume, in m^3, correct to three decimal places.
- **b** Find its volume, in L, correct to the nearest litre ($1\ m^3 = 1000$ L).

7 Which has a greater capacity: a 10 cm by 10 cm by 10 cm cube or a cylinder with radius 6 cm and height 10 cm?

8 A cylindrical water tank has a radius of 2 m and a height of 2 m.
- **a** Find its capacity, in m^3, rounded to three decimal places.
- **b** Find its capacity, in L, rounded to the nearest litre.




9 How many litres of gas can a tanker carry if its tank is cylindrical with a 2 m diameter and is 12 m in length? Round to the nearest litre.





10 Draw a cylinder with its circumference equal to its height. Try to draw it to scale.


Cylinder portions — 11


11 Find the volume of these cylindrical portions correct to two decimal places.


a


b


c


d


e


f


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Vegetable and fruit growers

The importance of farming in Australia can never be underestimated. From small scale gardens to commercial agriculture, farmers and their crops are essential for our food supply. Mathematics plays a major role in the profitable management of farms, including analysis of the optimum planting times, weather cycles, irrigation rates, pricing and production costs.

The issue of water usage and storage is very important in Australia. Farmers regularly measure soil moisture content to find out what is happening around the roots of crops. Irrigation is managed according to plant watering needs, rainfall and evaporation rates.



Maths@Work

1 Farmers need to be able to convert the volume of their dams from cubic metres (m^3) to megalitres (ML) by dividing by 1000.

a Convert each of the following into ML.

i 890 m^3 **ii** 60 000 m^3 **iii** 800 000 m^3 **iv** 380 000 m^3 **v** 190 000 000 m^3

b Convert the following into cubic metres.

i 9 ML **ii** 1.2 ML **iii** 120 ML **iv** 0.98 ML **v** 12 000 ML

2 The volume of a property dam is estimated with the formula, volume = $SA \times d \times 0.4$ where SA = the surface area of the top of the dam; d = maximum depth of the dam; and the factor of 0.4 allows for the dam's sloping sides.

Calculate the volume of the following farm dams.

a Surface area = 600 m^2, maximum depth 3 m

b Surface area = 400 m^2, maximum depth 2.5 m

c Surface area = 800 m^2, maximum depth 4 m

d Surface area = 5000 m^2, maximum depth 5 m

e Surface area = 16 000 m^2, maximum depth 6 m

3 Concrete water storage tanks can be in the shape of cylinders. Calculate the capacity in litres and kilolitres for each water tank illustrated below. Recall that 1 m^3 = 1000 L = 1 kL.

a Height 1.8 m, diameter 8.2 m **b** Height 1.5 m, diameter 5 m







Hint for Q3: Recall that the volume of a cylinder is $V = \pi r^2 h$.

Essential Mathematics Core for the Victorian Curriculum Year 9 ISBN 978-1-009-59315-1 © Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Using technology

4 A soil moisture probe is a device that measures the percentage of water in soil.
- Maximum % of soil moisture is when soil is drenched.
- Minimum % of soil moisture is when plants wilt.

The difference between these percentages is the percentage of soil water that a plant uses. The depth of watering needed for plants is measured in mm, like rainfall.

Plant watering depth = percentage moisture used by plant × root depth in mm

Set up the Excel spreadsheet below to calculate watering depth for the given plants.



Hint for Q4: Format columns B, C and D as 'percentage' and enter the decimal values in each cell; e.g. enter 0.43 for 43%.

	A	B	C	D	E	F
1	Calculation of watering depths for various plants					
2	Plant	Maximum moisture percentage	Minimum moisture percentage	Percentage moisture used by plant	Root depth in mm	Plant watering depth in mm
3	Onions	41%	37%		400	
4	Carrots	43%	38%		600	
5	Tomatoes	47%	43%		1200	
6	Apple trees	47%	41%		1500	
7	Banana trees	49%	38%		600	

5 Fruit trees can be micro-irrigated by dripping or spraying water in a circular area near the trunk. The volume of water required for each tree is calculated using the formula for volume of a cylinder:

$V = \pi r^2 h = \pi r^2 \times$ plant watering depth

Note that an area of 1 m^2 covered by 1 mm of water = 1 L of water; hence area in m^2 multiplied by depth in mm gives volume in litres.
Set up the Excel spreadsheet shown below to calculate the litres of irrigation water required per 100 trees.

In the area formula, enter pi() for π.



Hint for Q5: Use the plant watering depth values calculated in Question 4.


	A	B	C	D	E	F
1	Volume of irrigation water required for 100 trees					
2	Plant	Radius in m of micro-irrigation	Area of irrigation in m^2	Plant watering depth in mm	Litres of water per plant per irrigation	Litres of water per 100 trees
3	Apple trees	0.21				
4	Banana trees	0.32				

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Pressing a new road

The construction of a new asphalt road involves pressing the surface with a roller. Two rollers are available, each with main roller wheels of varying diameter and width. The options are:

- Roller 1: Wheel diameter 1 metre and width 2 metres
- Roller 2: Wheel diameter 1.2 metres and width 2.5 metres.



The cost of hiring each roller depends on the number of revolutions in the following way:

- Roller 1: $2.20 per revolution
- Roller 2: $3 per revolution.

The rectangular section of road being built is 1 km long and 10 metres wide.

Present a report for the following tasks and ensure that you show clear mathematical workings and explanations where appropriate.

1 Preliminary task

a Write the formula for finding the curved surface area of a cylinder using the pronumerals r and h.

b What is the value of r and h for;

- **i** Roller 1
- **ii** Roller 2

c For Roller 1 find,

- **i** the surface area of the roller wheel (the curved surface area only)
- **ii** the area of ground pressed after 10 revolutions.

d Find the total surface area, in square metres, of a section of road which is 1 km long and 10 metres wide.

e How many rotations of the wheel would need to occur if roller 1 was to travel a single length of the 1 km length of road? Round to one decimal place.

2 Modelling task

Formulate

The problem is to determine the cost of hiring each roller to press the entire 1 km stretch of road.

a Write down all the relevant information that will help solve this problem with the aid of diagrams.

Solve

b Using Roller 1 determine the total number of revolutions that the wheel needs to rotate if the entire 10 metre-wide road is to be pressed. Use the following assumptions.

- It is not necessary to overlap one pressed strip with the next.
- No revolutions are counted while the roller is manoeuvring into position at the start of each strip.

c Calculate the cost of hiring Roller 1 to press the 1 km stretch of road.

d Repeat parts **b** and **c** for Roller 2.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Evaluate and verify

e Assume that each strip rolled must now overlap the previous pressed strip by 10 cm. Determine the total number of revolutions required for each of the two rollers under these conditions.

f Compare and comment on the cost of hiring the rollers if:

- **i** no overlap is required between strips
- **ii** a 10 cm overlap is required between strips.

Communicate

g Summarise your results in a table like the one below, and decide which roller should be hired and why.

Roller	Surface area of 1 revolution	Number of revolutions needed for road with no overlap	Cost of hiring with no overlap	Number of revolutions needed for road with 10 cm overlap	Cost of hiring with 10 cm overlap
1					
2					

3 Extension question

If the overlap is changed to 0.5 m instead, which roller is the cheapest to hire? Give reasons and show calculations.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


Equal volumes and different surface areas

Key technology: Spreadsheets

While some solids have the same volume, they will not necessarily have the same surface area. This will depend on the base area and the height as well as the shape and the number of faces.


1 Getting started

We will start by considering three solids including a cube of side length 1 unit, a triangular prism with equilateral base side lengths 1 unit and cylinder with diameter 1 unit.



a State the volume and surface area of the cube.

b For the base of the triangular prism (shown to the right), find the perpendicular height a.



c Find the volume and surface area of the triangular prism if the height h is 1.5 units.

d Find the volume and surface area of the cylinder if its height h is 1.5 units.

e Compare the volumes and surface areas of the solids. Which solids have similar volumes and which solids have similar surface areas?

2 Using technology

The following spreadsheet varies the height by increments of 0.2 in column A and shows the formula for the volume of the triangular prism and cylinder. The formulas for the surface areas are not given.

	A	B	C	D	E
1	Equal volumes and	different surface areas			
2					
3	Cube volume = 1	Cube surface area = 6			
4					
5		Triangular prism		Cylinder	
6	Height	Volume	Surface area	Volume	Surface area
7	0.5	=SQRT(3)/4*A7		=PI()*0.25*A7	
8	=A7+0.2				
9					

a Enter the information into a spreadsheet and fill down at cells A8, B7 and D7.

b To the nearest 0.2, find the height required so that the volume of the following solids is as close as possible to the volume of the cube of side length 1.

i Triangular prism

ii Cylinder

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

c Calculate the correct formula for the surface areas of the triangular prism and cylinder in terms of h then enter these into the spreadsheet in cells C7 and E7. Then fill down.

d For the heights found in part **b** compare the surface areas of the following solids with the surface area of the cube.

i Triangular prism **ii** Cylinder

3 Applying an algorithm

a We will repeat the questions in part **2 Using technology** but use the following steps to find a more accurate solution.

- Step 1: Change the incremental increase in cell A8 to a smaller amount. Suggest start with 0.1.
- Step 2: Fill down the columns so that the volumes for both the triangular prism and cylinder are greater than 6.
- Step 3: Find the height required so that the volume of the following solids is as close as possible to the volume of the cube of side length 1.
  i Triangular prism **ii** Cylinder
- Step 4: For the heights found in Step 3, compare the surface areas of the following solids with the surface area of the cube.
  i Triangular prism **ii** Cylinder
- Step 5: Repeat from Step 1 using a smaller incremental change and fill down.


b Comment on the shape of the triangular prism and cylinder that have the same volume as the cube. Discuss the differences in height and the surface area.




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

1 These towers are made by stacking cubes. How many cubes are needed in each case?


a



b




2 A large cube of side length 4 cm is painted then cut into 64 single 1 cm cubes. How many 1 cm cubes are not painted on any face?



3 To the nearest metre, how far will a wheel of diameter 1 m travel after 100 revolutions?

4 A goat is tethered to the centre of one side of a shed with a 10 m length of rope. In what area of grass can the goat graze?






5 A circle of radius 10 cm has a hole cut out of its centre to form a ring. Find the radius of the hole if the remaining area is 50% of the original area. Round to one decimal place.



6 Here is one net for a cube. How many different nets are possible? Do not count nets that can be rotated or reflected (flipped) to give the same net.



7 A 100 m^2 factory flat roof feeds all the water collected into a rainwater tank. If there is 1 mm of rainfall, how many litres of water go into the tank?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter summary

Measurement

Errors and accuracy

$$\%\ \text{error} = \frac{\text{Error}}{\text{Actual value}} \times 100$$

Error = recorded value – actual value

The positive (absolute) value of the error is used.

Limits of accuracy are the smallest and largest possible values of the measurement: i.e.

recorded value $\pm$ 0.5 $\times$ smallest unit

Length and perimeter

Units

$\times 1000$ $\times 100$ $\times 10$


km m cm mm

$\div 1000$ $\div 100$ $\div 10$

Circle circumference

$C = 2\pi r$ or πd

Perimeter



$P = 2 \times 10 + 2 \times 7$

$= 34$ m

Volume

Units

$\times 1000^3$ $\times 100^3$ $\times 10^3$

km^3 m^3 cm^3 mm^3

$\div 1000^3$ $\div 100^3$ $\div 10^3$

1 L = 1000 mL = 1000 cm^3

Prisms and cylinders

$V = Ah$


A = area of cross-section

h = height

$V = lwh$

$= 11 \times 6 \times 5$


$= 330$ m^3



$V = \pi r^2 h$

$= \pi(2)^2 \times 7$


$= 87.96$ cm^3



Area

Triangle

$A = \frac{1}{2}bh$



Quadrilaterals

Square $A = l^2$


Rectangle $A = lw$

Parallelogram $A = bh$

Trapezium $A = \frac{1}{2}(a + b)h$

Rhombus or kite $A = \frac{1}{2}xy$

Composite shapes



$A = lw - \frac{1}{2}\pi r^2$

$= 10 \times 8 - \frac{1}{2}\pi \times 4^2$

$= 54.87$ m^2 (to 2 d.p.)

$P = 2 \times 10 + 8 + \frac{1}{2}2\pi(4)$

$= 40.57$ m (to 2 d.p.)

Units


$\times 1000^2$ $\times 100^2$ $\times 10^2$

km^2 m^2 cm^2 mm^2

$\div 1000^2$ $\div 100^2$ $\div 10^2$

Surface area

Prisms




$A = 2 \times$ triangles $+ 3 \times$ rectangles

$= 2 \times \frac{1}{2} \times 4 \times 3 + 8 \times 5$

$+ 8 \times 4 + 8 \times 3$

$= 108$ cm^2

Cylinder



$A = 2\pi r^2 + 2\pi rh$

$= 2\pi(3)^2 + 2\pi(3)(4)$

$= 131.95$ cm^2 (to 2 d.p.)

Circle



$A = \pi r^2$






$A = \pi \times 6^2$

$= 113.10$ m^2

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.







Chapter checklist

A version of this checklist that you can print out and complete can be downloaded from your Interactive Textbook.

		✔
6A	**1 I can calculate the absolute percentage error.** e.g. Calculate the absolute percentage error if the recorded time is 20 minutes and the actual time is 26 minutes.	
6A	**2 I can find the limits of accuracy in a measurement.** e.g. A backpack is weighed using a digital scale which displays measurements in kilograms correct to one decimal place. If the measurement given is 6.7 kg, state the lower and upper limits of accuracy of this measurement.	
6B	**3 I can convert between metric units of length.** e.g. Convert these measurements to the units shown in the brackets. **a** 9.2 cm (mm) **b** 61 000 cm (m)	
6B	**4 I can find the perimeter of basic shapes including composite shapes.** e.g. Find the perimeter of these shapes. **a**  **b** 	
6C	**5 I can find the circumference of a circle.** e.g. Find the circumference of a circle with a diameter of 5 m, correct to two decimal places.	
6C	**6 I can find the perimeter of a semicircle or a quadrant.** e.g. Find the perimeter of this semicircle correct to two decimal places. 	
6D	**7 I can convert between metric units of area.** e.g. Convert these measurements to the units shown in the brackets. **a** 5.32 cm^2 (mm^2) **b** 728 000 cm^2 (m^2)	
6D	**8 I can find the area of rectangles, triangles and parallelograms.** e.g. Find the area of this triangle. 	
6D	**9 I can find the area of rhombuses and trapeziums.** e.g. Find the area of this trapezium. 	






Essential Mathematics Core for the Victorian Curriculum Year 9 ISBN 978-1-009-59315-1 © Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter checklist

		✔
6E	**10 I can find the area of a circle.** e.g. Find the area of this circle correct to two decimal places. 	
6E	**11 I can find the area of a quadrant or semicircle.** e.g. Find the area of this quadrant correct to two decimal places. 	
6F	**12 I can find the perimeter and area of simple composite shapes.** e.g. Find the perimeter and area of this composite shape correct to two decimal places. 	
6G	**13 I can find the surface area of a rectangular prism.** e.g. Find the surface area of this rectangular prism. 	
6G	**14 I can find the surface area of a triangular prism.** e.g. Find the surface area of this triangular prism. 	
6H	**15 I can find the surface area of a cylinder.** e.g. Find the surface area of this cylinder correct to two decimal places. 	

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter checklist

		✔
6H	**16** **I can find the surface area of a cylindrical portion.** e.g. Find the surface area of this half cylinder correct to two decimal places. 	
6I	**17** **I can convert between metric units of volume.** e.g. Convert these measurements to the units shown in the brackets. **a** $0.003\ m^3$ (cm^3) **b** $640\,000\ mm^3$ (cm^3)	
6I	**18** **I can find the volume of a rectangular prism.** e.g. Find the volume of this rectangular prism. 	
6I	**19** **I can find the volume of a triangular prism.** e.g. Find the volume of this triangular prism. 	
6J	**20** **I can find the volume of a cylinder.** e.g. Find the volume of this cylinder correct to two decimal places. 	
6J	**21** **I can find the capacity of a cylinder giving an answer in L or mL.** e.g. Find the volume of this cylinder in litres correct to the nearest litre. 	

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter review

Short-answer questions


6B/D/I **1** Convert the following measurements into the units given in brackets.


a 3.8 m (cm) **b** 1.27 km (m) **c** 48 mm (cm)
d 273 mm^2 (cm^2) **e** 5.2 m^2 (cm^2) **f** 0.01 m^3 (cm^3)
g 53 100 mm^3 (cm^3) **h** 3100 mL (L) **i** 0.043 L (mL)
j 2.83 kL (L) **k** 4000 cm^3 (L) **l** 1 m^3 (L)


6A **2** Calculate the absolute percentage error for the following recorded and actual measurements.

a recorded: 2.4 L actual: 2.5 L
b recorded: 3 seconds actual: 5 seconds


6B **3** Find the perimeter of each of the following shapes.


a



b



c



6D **4** Find the area of each of the following plane figures.


a


b


c



d


e


f


6F **5** Inside a rectangular lawn area of length 10.5 m and width 3.8 m, a new garden bed is to be constructed. The garden bed is to be the shape of a triangle with base 2 m and height 2.5 m. Find:

a the area of the garden bed
b the area of the lawn remaining around the garden bed.







ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

6C/E **6** Find the area and circumference/perimeter of each of the following shapes correct to two decimal places.


a



b




c




6F **7** Find the perimeter and area of each of the following composite shapes correct to two decimal places.

a




b




6G **8** Find the surface area of each of the following solid objects.

a




b




6H **9** Find the surface area of each of these cylinders, correct to two decimal places.

a




b




6I/J **10** Find the volume of each of these solid objects, rounding to two decimal places where necessary.


a




b



c



d



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter review

Multiple-choice questions

6A **1** A digital scale measures the weight of a bag of potato chips to the nearest gram to be 35 g. If the chips are sold in boxes of 10 bags, the lower limit of accuracy for the weight of a box is:

A 355 g **B** 350 g **C** 340 g **D** 360 g **E** 345 g

6B/D **2** If the area of a square field is 25 km^2, the length of one of its sides is:


A 10 km **B** 5 km **C** 4 km **D** 50 km **E** 25 km

6D **3** If 1 $m^2 = 100^2$ cm^2, 2.7 m^2 is the same as:

A 270 cm^2 **B** 0.0027 km^2 **C** 27 000 cm^2 **D** 2700 mm^2 **E** 27 cm^2

6B **4** The perimeter of this shape is:

A 21 cm **B** 14 cm **C** 24 cm
D 20 cm **E** 22 cm




6D **5** A parallelogram has area 10 m^2 and base 5 m. Using $A = bh$, its perpendicular height is:

A 50 m **B** 2 m **C** 50 m^2
D 2 m^2 **E** 0.5 m


6F **6** This composite shape could be considered as a rectangle with an area in the shape of a square removed. The shape's area is:

A 16 km^2 **B** 12 km^2 **C** 20 km^2
D 24 km^2 **E** 6 km^2




6C **7** A semicircular goal area has diameter 20 m. Its perimeter, correct to the nearest metre, is:

A 41 m **B** 36 m **C** 83 m
D 51 m **E** 52 m




6G **8** The surface area of the rectangular prism shown is:


A 30 m^2 **B** 62 m^2 **C** 31 m^2
D 60 m^2 **E** 100 m^2



6H **9** The area of the *curved surface only* of a cylinder with radius 5 mm and height 12 mm is closest to:

A 534 mm^2 **B** 94 mm^2 **C** 188 mm^2
D 754 mm^2 **E** 377 mm^2





ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

6I **10** A prism's cross-sectional area is 100 m^2. If its volume is 6500 m^3, the prism's total height would be:


A 0.65 m **B** 650 000 m **C** 65 m **D** 6.5 m **E** 650 m

6J **11** The volume of a cylinder with radius 3 cm and height 10 cm is closest to:

A 188 cm^2 **B** 251 cm **C** 141 cm^3 **D** 94 cm^3 **E** 283 cm^3

Extended-response questions

1 A kindergarten teacher collects some blocks of wood for a painting activity. Each block is a rectangular prism, as shown.



a Find the volume of each block.

b Find the total outside area to be painted for each block.

c If the paint costs \$2.50 per 100 cm^2, find the cost of painting 10 blocks.


d Another wood block is a cylinder with radius 3 cm and height 9 cm. Compared to the rectangular block:

i does it have a greater volume?

ii does it have a greater surface area?



2 An office receives five new desks with a bench shape made up of a rectangle and quarter circle as shown.



The edge of the bench is lined with a rubber strip at a cost of \$2.50 per metre.

a Find the length of the rubber edging strip for one desk, correct to two decimal places.

b Find the total cost of the rubber strip for the five desks. Round to the nearest dollar.

The manufacturer claims that the desk top area space is more than 1.5 m^2.

c Is the manufacturer's claim correct?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

7 Indices




Essential mathematics: why skills with index laws and scientific notation are important

Indices, also known as exponents or powers, and scientific notation are essential tools in many fields, simplifying complex calculations and representing very large or very small numbers efficiently.

Video game developers use indices and exponential functions to create realistic graphics and physics. They model character progression with exponential growth functions, ensuring balanced gameplay. Exponential decay represents the decrease in an object's speed due to friction, enhancing realism. Exponents in lighting calculations simulate light intensity over distance, creating immersive environments.

Astronomers and chemists use scientific notation to handle large or small numbers efficiently. An astronomer might express the distance to Proxima Centauri as 4.24×10^{13} kilometres, simplifying vast distances. A chemist uses scientific notation for precise measurements, such as creating a 0.1 M solution of hydrochloric acid by measuring 1.2×10^{-3} grams of the acid for 100 mL of water. This precision ensures accurate and consistent chemical reactions, showcasing the practical application of scientific notation in daily tasks.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.



In this chapter

7A Index notation (Consolidating)
7B Index laws for multiplication and division
7C Index laws for power of a power and the zero index
7D Index laws for brackets and fractions
7E Negative indices ★
7F Scientific notation
7G Scientific notation using significant figures

Victorian Curriculum 2.0

This chapter covers the following content descriptors in the Victorian Curriculum 2.0:

NUMBER

VC2M9N01

ALGEBRA

VC2M9A01

MEASUREMENT

VC2M9M02

Please refer to the curriculum support documentation in the teacher resources for a full and comprehensive mapping of this chapter to the related curriculum content descriptors.

© VCAA


Online resources

A host of additional online resources are included as part of your Interactive Textbook, including HOTmaths content, video demonstrations of all worked examples, auto-marked quizzes and much more.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025
Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Warm-up quiz

1 Draw a factor tree like below right for 36 and then write 36 as a product of prime factors.



2
- **a** List the factors of 24.
- **b** List the factors of 45.
- **c** List the prime factors of 24.
- **d** List the prime factors of 45.

3 Write each of the following in index form (as a power; e.g. 6^4).
- **a** $7 \times 7 \times 7$
- **b** $5 \times 5 \times 2 \times 2 \times 2$
- **c** $(3 \times 3 \times 3 \times 3) \div (3 \times 3)$
- **d** $(4 \times 4 \times 4) \div (4 \times 4)$

4 Write each of the following as 2 raised to a single power.

a $2^2 \times 2$ **b** $2^4 \times 2^2$ **c** $2^3 \times 2^2$

5 Evaluate:

a 5^2 **b** 10^2 **c** 1^4 **d** 5^1 **e** 4^0

6 Write each of the following in expanded form and evaluate.

a 3×2^2 **b** 4×3^2 **c** 5×2^3 **d** 3×10^4

7 Simplify the following by removing brackets.

a $(3^2)^2$ **b** $(2^4)^3$ **c** $(-3)^2$

8 Write each of these fractions as decimals.

a $\frac{3}{1000}$ **b** $\frac{4}{100}$ **c** $\frac{2}{1\,000\,000}$ **d** $\frac{8}{10\,000}$

9 Round the following to two decimal places.

a 3.732 **b** 24.6174 **c** 18.3654 **d** 4.3971

10 State the place value of the 4 in each of these numbers.

a 246 **b** 0.0043 **c** 4 320 000 **d** 32.48

11 State the number of significant figures in each of the following.

a 23.102 **b** 30.05 **c** 0.0012 **d** 6001

12 Complete the following.
- **a** $3.8 \times 10 =$ _____
- **b** $2.31 \times 1000 =$ _____
- **c** $17.2 \div 100 =$ _____
- **d** $0.18 \div 100 =$ _____
- **e** $3827 \div$ _____ $= 3.827$
- **f** $6.49 \times$ _____ $= 64\,900$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

7A Index notation


CONSOLIDATING

Learning intentions

- To understand index notation and know that it is a shorthand way of writing multiplication of the same base
- To be able to convert between expanded form and index form
- To be able to evaluate expressions given in index form
- To be able to express a number as a product of prime factors

Key vocabulary: index form, expanded form, power, exponent, base, index, prime, factor, substitution

When a product has the same number multiplied by itself over and over, index notation can be used to write a simpler expression.
For example, $5 \times 5 \times 5$ can be written as 5^3 and $x \times x \times x \times x \times x$ can be written as x^5. The expression 5^3 is a power and we can say '5 to the power of 3'. The 5 is called the base and the 3 is the index, power or exponent. Numbers written with indices are common in mathematics and can be applied to many types of problems.



Observe the rapid expansion of 3D squares, illustrating the concept of exponential growth.

Lesson starter: Who has the most?

A person offers you one of two prizes.

- Which offer would you take?
- Try to calculate the final amount for prize B.
- How might you use index notation to help calculate the value of prize B?
- How can a calculator help to find the amount for prize B using the power button $\boxed{\wedge}$?





Key ideas

- When a number is multiplied by itself many times, that product can be written using **index form**. For example,

Expanded form: $2 \times 2 \times 2 \times 2 \times 2$ — Index form: $2^5 = 32$

base: 2, index: 5, basic numeral: 32

$x \times x \times x \times x = x^4$

base: x, index: 4

- The **base** is the factor in the product.
- The **index** is the number of times the factor (base number) appears.
  - 2^5 reads '2 to the **power** of 5'.


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

7A

- Prime factorisation involves writing a number as a product of its prime factors.
  - A prime number has only two factors: 1 and itself.

  Prime factorisation can be completed using a factor tree or repeated division.
  For example,

  Factor tree



  Repeated division

2	24
2	12
2	6
3	3
	1

  - To write 24 in prime factor form use indices: $24 = 2 \times 2 \times 2 \times 3$
    $= 2^3 \times 3$

- Note that $a^1 = a$. For example: $5^1 = 5$
  - 3^2 does *not* mean $3 \times 2 = 6$.

Exercise 7A

Understanding

1, 2–5(½) | 1, 5

1 **a** The product $3 \times 3 \times 3 \times 3$ is called the _____ form of 81.
b 3^4 is called the _____ form of 81.
c 3^4 reads '3 to the _____ of 4'.
d In 3^4 the special name for the 3 is the _____ number.
e In 3^4 the special name for the 4 is the _____ number or _____.
f A _____ number has only two factors, itself and 1.
g Prime factorisation involves writing a number as a product of its _____ _____.

2 Evaluate:
a 5^2 **b** 2^3 **c** 3^3 **d** $(-4)^2$

3 Write the number or pronumeral that is the base in these expressions.
a 3^7 **b** $(-7)^3$ **c** $\left(\frac{2}{3}\right)^4$ **d** w^6



Hint for Q3 and Q4: In a^x, a is the base and x is the index.

4 Write the number that is the index in these expressions.
a 4^3 **b** $(-3)^7$ **c** $\left(\frac{1}{2}\right)^4$ **d** $(xy)^{13}$

5 Write the prime factors of these numbers.
a 6 **b** 15 **c** 30 **d** 77

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Fluency

6–11(½) 6–11(½)

Example 1 Writing in expanded form

Write in expanded form:

a 5^4 b a^3 c $(xy)^4$ d $2a^3b^2$

Solution

a $5^4 = 5 \times 5 \times 5 \times 5$

b $a^3 = a \times a \times a$

c $(xy)^4 = xy \times xy \times xy \times xy$

d $2a^3b^2 = 2 \times a \times a \times a \times b \times b$

Explanation

Factor 5 appears four times.

Factor a appears three times.

Factor xy appears four times.

Factor a appears three times and factor b appears twice. Factor 2 only appears once.

Now you try

Write in expanded form:

a 6^3 b b^4 c $(mn)^2$ d $3x^2y$

6 Write each of the following in expanded form.

a 4^3 b 7^4 c 3^5 d 5^3
e a^4 f b^3 g x^3 h $(xp)^6$
i $(5a)^4$ j $(3y)^3$ k $4x^2y^5$ l $(pq)^2$
m $-3s^3t^2$ n $6x^3y^5$ o $5(yz)^6$ p $4(ab)^3$



Hint for Q6:
factor → a^5 ← number of repeats
$a^5 = a \times a \times a \times a \times a$

Example 2 Expanding and evaluating

Write each of the following in expanded form and then evaluate.

a 5^3 b $(-2)^5$ c $\left(\frac{2}{5}\right)^3$

Solution

a $5^3 = 5 \times 5 \times 5$
$= 125$

b $(-2)^5 = (-2) \times (-2) \times (-2) \times (-2) \times (-2)$
$= -32$

c $\left(\frac{2}{5}\right)^3 = \frac{2}{5} \times \frac{2}{5} \times \frac{2}{5}$
$= \frac{8}{125}$

Explanation

Write in expanded form with 5 appearing three times and evaluate.

Write in expanded form with -2 appearing five times and evaluate.

Write in expanded form.

Evaluate by multiplying numerators and denominators.

Now you try

Write each of the following in expanded form and then evaluate.

a 2^3 b $(-3)^3$ c $\left(\frac{3}{5}\right)^2$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

7A

7 Write each of the following in expanded form and then evaluate.

a 6^2 **b** 2^4 **c** 3^5 **d** 12^1

e $(-2)^3$ **f** $(-1)^7$ **g** $(-3)^4$ **h** $(-5)^2$

i $\left(\frac{2}{3}\right)^3$ **j** $\left(\frac{3}{4}\right)^2$ **k** $\left(\frac{1}{6}\right)^3$ **l** $\left(\frac{5}{2}\right)^2$

m $\left(\frac{2}{-3}\right)^3$ **n** $\left(\frac{-3}{4}\right)^4$ **o** $\left(\frac{-1}{4}\right)^2$ **p** $\left(\frac{5}{-2}\right)^5$



Hint for Q7: $-2 \times (-2) \times (-2)$
$= +4 \times (-2)$
$= -8$

Example 3 Writing in index form

Write each of the following in index form.

a $5 \times 5 \times 5 \times 5$ **b** $6 \times x \times x \times x \times x$ **c** $4 \times a \times 4 \times a \times 4 \times a$

Solution | **Explanation**

a $5 \times 5 \times 5 \times 5 = 5^4$ — Factor 5 is repeated 4 times.

b $6 \times x \times x \times x \times x = 6x^4$ — Factor x is repeated 4 times; 6 appears only once.

c $4 \times a \times 4 \times a \times 4 \times a$
$= 4 \times 4 \times 4 \times a \times a \times a$
$= 4^3a^3$

Group the factors of 4 together and the factors of a together.

Write in index form.

Now you try

Write each of the following in index form.

a $9 \times 9 \times 9$ **b** $2 \times a \times a$ **c** $11 \times b \times 11 \times b \times b$

8 Write each of the following in index form.

a $3 \times 3 \times 3$ **b** $8 \times 8 \times 8 \times 8 \times 8 \times 8$

c $y \times y$ **d** $3 \times x \times x \times x$

e $4 \times c \times c \times c \times c \times c$ **f** $5 \times 5 \times 5 \times d \times d$

g $x \times x \times y \times y \times y$ **h** $7 \times b \times 7 \times b \times 7$



Hint for Q8: The index is the number of times the factor appears.

Example 4 Writing in index form with fractions

Write each of the following in index form.

a $\frac{3}{4} \times \frac{3}{4} \times \frac{3}{4}$ **b** $\frac{3}{7} \times \frac{3}{7} \times \frac{4}{5} \times \frac{4}{5} \times \frac{4}{5}$

Solution | **Explanation**

a $\frac{3}{4} \times \frac{3}{4} \times \frac{3}{4} = \left(\frac{3}{4}\right)^3$ — The fraction $\frac{3}{4}$ appears 3 times.

b $\frac{3}{7} \times \frac{3}{7} \times \frac{4}{5} \times \frac{4}{5} \times \frac{4}{5} = \left(\frac{3}{7}\right)^2 \times \left(\frac{4}{5}\right)^3$ — $\frac{3}{7}$ appears twice and $\frac{4}{5}$ three times.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Now you try

Write each of the following in index form.

a $\frac{5}{7}\times\frac{5}{7}\times\frac{5}{7}\times\frac{5}{7}$

b $\frac{2}{5}\times\frac{4}{7}\times\frac{2}{5}\times\frac{4}{7}\times\frac{4}{7}$

9 Write each of the following in index form.

a $\frac{2}{3}\times\frac{2}{3}\times\frac{2}{3}\times\frac{2}{3}$

b $\frac{3}{5}\times\frac{3}{5}\times\frac{3}{5}\times\frac{3}{5}\times\frac{3}{5}$

c $\frac{4}{7}\times\frac{4}{7}\times\frac{1}{5}\times\frac{1}{5}\times\frac{1}{5}\times\frac{1}{5}$

d $\frac{7x}{9}\times\frac{7x}{9}\times\frac{y}{4}\times\frac{y}{4}\times\frac{y}{4}$

Example 5 Writing in index form with a combination of pronumerals

Write each of the following in index form.

a $8\times a\times a\times 8\times b\times b\times a\times b$

b $3a\times 2m\times 3a\times 2m$

c $4am(4am)(4am)$

Solution

a $8\times a\times a\times 8\times b\times b\times a\times b$
$= 8\times 8\times a\times a\times a\times b\times b\times b$
$= 8^2a^3b^3$

Explanation

Group the numerals and like pronumerals and write in index form.

$64a^3b^3$ or $64(ab)^3$ are alternative answers.

b $3a\times 2m\times 3a\times 2m$
$= 2\times 2\times 3\times 3\times a\times a\times m\times m$
$= 2^2 3^2a^2m^2$

Rearrange so like factors are grouped together and write in index form.

$36a^2m^2$ or $36(am)^2$ are alternative answers.

c $4am(4am)(4am)$
$= 4\times 4\times 4\times a\times a\times a\times m\times m\times m$
$= 4^3a^3m^3$

Rearrange and write in index form.

$64a^3m^3$ or $64(am)^3$ are alternative answers.

Now you try

Write each of the following in index form.

a $7\times x\times x\times 7\times y\times 7\times x\times y$

b $2m\times 7s\times 2m\times 7s\times 2m$

c $3ab(3ab)(3ab)(3ab)$

10 Write each of the following in index form.

a $3\times x\times y\times x\times 3\times x\times 3\times y$

b $3x\times 2y\times 3x\times 2y$

c $4d\times 2e\times 4d\times 2e$

d $6by(6by)(6y)$

e $3pq(3pq)(3pq)(3pq)$

f $7mn\times 7mn\times mn\times 7$



Hint for Q10: First rearrange the factors with numbers first, then form groups of like bases.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


7A

Example 6 Finding the prime factor form

Express 48 as a product of prime factors in index form. Prime numbers are divisible only by 1 and themselves.

Solution

Tree diagram method



$48 = 2 \times 2 \times 2 \times 2 \times 3$

$= 2^4 \times 3$

Explanation

Choose a pair of factors of 48; for example, 8 and 6.

Choose a pair of factors of 8; i.e. 2 and 4.

Choose a pair of factors of 6; i.e. 2 and 3.

Continue this process until the factors are all prime numbers.

Write the prime factors of 48. Express in index notation.

Repeated division method

2	48
2	24
2	12
2	6
3	3
	1

$48 = 2 \times 2 \times 2 \times 2 \times 3$

$= 2^4 \times 3$

Start by dividing by a prime number.

$48 \div 2 = 24$

$24 \div 2 = 12$

$12 \div 2 = 6$

$6 \div 2 = 3$

$3 \div 3 = 1$

Write prime factors in ascending order.

Express in index notation.

Now you try

Express 60 as a product of prime factors.

11 Express each of the following as a product of prime factors in index form.

a 10 **b** 8 **c** 144

d 75 **e** 147 **f** 500

Problem-solving and reasoning 12, 13 12–14(½)

12 Copy and fill in the missing numbers or symbols.

a $3 \times 3 \times a \times a \times a = 3^{\square}a^{\square}$

b $\square \times \square \times k \times k \times k = 5^2k^{\square}$

c $\frac{2}{7} \times \frac{2}{7} \times \frac{2}{7} = \left(\frac{2}{7}\right)^{\square}$

d $3p^2q = 3 \times \square \times \square \times q$

e $(abc)^1 = a \square b \square c$

f $4ab^2 = \square \times (-2) \times \square \times \square \times b$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Example 7 Evaluating expressions in index form by using substitution

If $a = 5$, $b = 2$ and $c = -5$, evaluate these expressions.

a $(ab)^2$ **b** $\left(\frac{a}{c}\right)^3$ **c** $\left(\frac{b}{c}\right)^2$

Solution

a $(ab)^2 = (5 \times 2)^2$
$= 10^2$
$= 10 \times 10$
$= 100$

Explanation

Replace a with 5 and b with 2. Include the $\times$ sign. $5 \times 2 = 10$
Base of 10 appears twice.

b $\left(\frac{a}{c}\right)^3 = \left(\frac{5}{-5}\right)^3$
$= (-1)^3$
$= -1 \times (-1) \times (-1)$
$= -1$

Replace a with 5 and c with -5.
$5 \div (-5) = -1$
$-1 \times (-1) = +1, \quad +1 \times (-1) = -1$

c $\left(\frac{b}{c}\right)^2 = \left(\frac{2}{-5}\right)^2$
$= \frac{2}{-5} \times \frac{2}{-5}$
$= \frac{4}{25}$

Replace b with 2 and c with -5.

Write in expanded form.

$2 \times 2 = 4, \quad -5 \times (-5) = 25$

Now you try

If $a = 3$, $b = -2$ and $c = 7$, evaluate these expressions.

a $(ab)^2$ **b** $\left(\frac{b}{a}\right)^3$ **c** $ab^2 + ac$

13 If $a = 3$, $b = 2$ and $c = -3$, evaluate these expressions.

a $(ab)^2$ **b** $(bc)^3$ **c** $\left(\frac{a}{c}\right)^4$ **d** $\left(\frac{b}{c}\right)^3$

e $(abc)^1$ **f** $c^2 + ab$ **g** ab^2c **h** c^2ab^3



Hint for Q13: $abc = a \times b \times c$

14 Find the missing number.

a $3^{\square} = 81$ **b** $2^{\square} = 256$ **c** $\square^3 = 125$

d $\square^5 = 32$ **e** $\square^3 = -64$ **f** $\square^7 = -128$

g $\square^3 = \frac{1}{8}$ **h** $\left(\frac{2}{3}\right)^{\square} = \frac{16}{81}$ **i** $\left(\frac{\square}{4}\right)^3 = \frac{1}{64}$



Hint for Q14: For part a, ask 'How many times should 3 appear so that the product is 81?'

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

7A

Splitting cells — 15

15 Certain bacteria cells split in 2 every 1 minute. New cells also continue splitting in the same way. So, after each minute, the number of bacteria cells has doubled.

a Copy and complete this table showing the number of bacteria after each minute for 10 minutes.

Time in minutes	Number of bacteria	Number in index form
0	1	2^0
1	$1 \times 2 = 2$	2^1
2	$2 \times 2 = 4$	2^2
3	$2 \times 2 \times 2 = 8$	2^3

b How long will it take for 1 cell to divide into:

i 4 cells? **ii** 16 cells? **iii** 64 cells?

c A single cell is set aside to divide for 24 minutes. Use index form to quickly find how many cells there will be after this time.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

7B Index laws for multiplication and division

Learning intentions

- To know the index laws for multiplication and division involving a common base
- To be able to apply the index laws for multiplication and division to simplify expressions

Key vocabulary: index law, base

When multiplying or dividing numbers with the same base, index laws can be used to simplify the expression.

Consider $5^{18} \times 5^{10}$:

Using expanded form: $5^{18} \times 5^{10} = \overbrace{5 \times 5 \times 5 \times \ldots \times 5}^{18 \text{ factors of } 5} \times \overbrace{5 \times 5 \times \ldots \times 5}^{10 \text{ factors of } 5}$

$= \overbrace{5 \times 5 \times 5 \times \ldots \times 5}^{18 + 10 \text{ factors of } 5}$

$= 5^{18+10}$

$= 5^{28}$

So the total number of factors of 5 is $18 + 10 = 28$.

Also $5^{18} \div 5^{10} = \dfrac{\overbrace{5 \times 5 \times \ldots \times \cancel{5} \times \cancel{5} \times \cancel{5} \times \cancel{5}}^{18 \text{ factors of } 5}}{\underbrace{\cancel{5} \times \ldots \times \cancel{5} \times \cancel{5}}_{10 \text{ factors of } 5}}$

$= 5^{18-10}$

$= 5^{8}$

So the total number of factors of 5 is $18 - 10 = 8$.

Lesson starter: Discovering laws for multiplication and division

Consider the two expressions $2^3 \times 2^5$ and $6^8 \div 6^6$.

Complete this working. $2^3 \times 2^5 = 2 \times \square \times \square \times 2 \times \square \times \square \times \square \times \square$

$= 2^{\square}$

$6^8 \div 6^6 = \dfrac{6 \times \square \times \square \times \square \times \square \times \square \times \square \times \square}{6 \times \square \times \square \times \square \times \square \times \square}$

$= \dfrac{6 \times 6}{1}$

$= 6^{\square}$

1 What do you notice about the given expression and the answer in each case? Can you express this as a rule or law in words?

2 Repeat the type of working given above and test your laws on these expressions.

a $3^2 \times 3^7$ **b** $4^{11} \div 4^8$

Essential Mathematics Core for the Victorian Curriculum Year 9 ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

7B

Key ideas

- **Index law for multiplication**: $a^m \times a^n = a^{m+n}$
  - When multiplying terms with the same base, add the powers.
    For example, $7^3 \times 7^2 = 7^{3+2} = 7^5$
- **Index law for division**: $a^m \div a^n = \frac{a^m}{a^n} = a^{m-n}$
  - When dividing terms with the same base, subtract the powers.
    For example, $8^5 \div 8^3 = \frac{8^5}{8^3} = 8^{5-3} = 8^2$

Exercise 7B

Understanding

1–4 | 4

1 Write the missing words.

a The index law for multiplication states that if you _____ two terms with the same _____ you _____ the powers.

b The index law for division states that if you _____ two terms with the same _____ you _____ the powers.

2 Copy and complete to give an answer in index form.

a $3^2 \times 3^4 = 3 \times \square \times 3 \times \square \times \square \times \square$
$= 3^{\square}$

b $k^3 \times k^2 = k \times \square \times \square \times \square \times \square$
$= k^{\square}$



Hint for Q2: The index shows how many times the factor should appear.

3 Copy and complete to give an answer in index form.

a $5^5 \div 5^3 = \frac{5 \times \square \times \square \times \square \times \square}{5 \times \square \times \square}$
$= 5^{\square}$

b $a^6 \div a^2 = \frac{a \times \square \times \square \times \square \times \square \times \square}{\square \times \square}$
$= a^{\square}$



Hint for Q3: Show cancelling, e.g. $\frac{5 \times 5^{1} \times 5^{1}}{{}^{1}\cancel{5} \times 5^{1}}$

4 Copy and complete.

a $6^5 \times 6^7 = 6^{\square+\square} = 6^{\square}$

b $a^{13} \times a^2 = a^{\square+\square} = a^{\square}$

c $5^{12} \div 5^4 = 5^{\square-\square} = 5^{\square}$

d $\frac{m^{16}}{m^2} = m^{\square-\square} = m^{\square}$



Hint for Q4: When dividing, subtract the indices.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Fluency

5–8(½) 5–8(½)

Example 8 Using laws for multiplication and division with numerical bases

Simplify, giving your answer in index form.

a $3^6 \times 3^4$ b $4^5 \times 4$ c $7^9 \div 7^5$ d $6^8 \div 6$

Solution

a $3^6 \times 3^4 = 3^{6+4}$
$= 3^{10}$

b $4^5 \times 4 = 4^{5+1}$
$= 4^6$

c $7^9 \div 7^5 = 7^{9-5}$
$= 7^4$

d $6^8 \div 6 = 6^{8-1}$
$= 6^7$

Explanation

$a^m \times a^n = a^{m+n}$
Add powers: $6 + 4 = 10$. The base 3 is unchanged.

$4 = 4^1$
Add powers: $5 + 1 = 6$. The base 4 is unchanged.

$a^m \div a^n = a^{m-n}$
Subtract powers: $9 - 5 = 4$. The base 7 is unchanged.

$6 = 6^1$
Subtract powers: $8 - 1 = 7$. The base 6 is unchanged.

Now you try

Simplify, giving your answer in index form.

a $2^4 \times 2^7$ b $7^5 \times 7$ c $11^{12} \div 11^4$ d $5^{10} \div 5$

5 Simplify, giving your answers in index form.

a $2^4 \times 2^3$ b $5^6 \times 5^3$ c $7^2 \times 7^4$ d $8^9 \times 8$
e $3^4 \times 3^4$ f $6^5 \times 6^9$ g $3^7 \div 3^4$ h $6^8 \div 6^3$
i $5^4 \div 5$ j $10^6 \div 10^5$ k $9^9 \div 9^6$ l $(-2)^5 \div (-2)^3$



Hint for Q5: Remember: $8 = 8^1$

Example 9 Using the index law for multiplication

Simplify each of the following using the index law for multiplication.

a $x^4 \times x^5 \times x^2$ b $x^3y^4 \times x^2y$

Solution

a $x^4 \times x^5 \times x^2 = x^{4+5+2}$
$= x^{11}$

b $x^3y^4 \times x^2y = x^3x^2y^4y$
$= x^{3+2}y^{4+1}$
$= x^5y^5$

Explanation

Use the index law to add the indices, since all terms have base x.

Regroup so like bases are together. Use the index law to add the indices corresponding to each different base.
Recall that $y = y^1$.

Now you try

Simplify each of the following using the index law for multiplication.

a $a^3 \times a \times a^4$ b $m^2n^3 \times mn^4$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

7B

6 Simplify each of the following using the index law for multiplication.

a $x^4 \times x^3$ **b** $a^6 \times a^3$ **c** $t^5 \times t^3$

d $y \times y^4$ **e** $d^2 \times d$ **f** $y^2 \times y \times y^4$

g $b \times b^5 \times b^2$ **h** $q^6 \times q^3 \times q^2$ **i** $a^2m^2 \times a^3m^2$

j $k^3p^2 \times k^2p$ **k** $x^2y^3 \times x^4y^5$ **l** $m^5e^3 \times m^2e$



Hint for Q6: Check that the bases are the same before adding the indices.

Example 10 Using the index law for division

Simplify $x^{10} \div x^2$ using the index law for division

Solution

$$x^{10} \div x^2 = x^{10-2}$$
$$= x^8$$

Explanation

Use the index law to subtract the indices: $10 - 2 = 8$.
The base x is unchanged.

Now you try

Simplify $y^{11} \div y^5$ using the index law for division.

7 Simplify each of the following using the index law for division

a $a^6 \div a^4$ **b** $x^5 \div x^2$ **c** $\frac{q^{12}}{q^2}$ **d** $\frac{d^7}{d^6}$

e $\frac{b^{10}}{b^5}$ **f** $\frac{d^9}{d^4}$ **g** $\frac{a^{14}}{a^7}$ **h** $\frac{y^{15}}{y^7}$



Hint for Q7: Recall: $\frac{q^{12}}{q^2} = q^{12} \div q^2$

Example 11 Simplifying using index laws for multiplication and division

Simplify each of the following using index laws for multiplication and division.

a $3m^4 \times 2m^5$ **b** $12y^7 \div (4y^3)$ **c** $\frac{8a^6b^3}{12a^2b^2}$

Solution

a $3m^4 \times 2m^5 = 3 \times 2 \times m^4 \times m^5$
$= 6 \times m^{4+5}$
$= 6m^9$

Explanation

Regroup with numbers first then like bases together.
Multiply the numbers then use the index law for multiplication to add the indices of the base m.

b $12y^7 \div (4y^3) = \frac{12y^7}{4y^3}$
$= 3y^{7-3}$
$= 3y^4$

$12 \div 4 = 3$
Use the index law for division to subtract indices.

c $\frac{8a^6b^3}{12a^2b^2} = \frac{8}{12} \times \frac{a^6}{a^2} \times \frac{b^3}{b^2}$
$= \frac{2}{3}a^4b$ or $\frac{2a^4b}{3}$

$\frac{8}{12} = \frac{2}{3}$ in simplest form.

Use the index law for division to subtract the indices for each different base: $6 - 2 = 4$, $3 - 2 = 1$.

Recall: $b = b^1$.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Now you try

Simplify each of the following using index laws for multiplication and division.

a $4a^2 \times 5a^3$ b $20b^8 \div (5b)^3$ c $\dfrac{6x^4y^2}{8x^2y}$

8 Simplify, using index laws for multiplication and division.

a $7x^3y^3 \times x^4y^2$ b $3x^7y^3 \times x^2y$ c $5x^3y^5 \times xy^4$

d $xy^4z \times 4xy$ e $3m^3 \times 5m^2$ f $4e^4f^2 \times 2e^2f^2$

g $5c^4d \times 4c^3d$ h $9yz^2 \times 2yz^5$ i $9m^3 \div (3m^2)$

j $14x^4 \div (2x)$ k $5y^4 \div y^2$ l $6a^6 \div (2a^5)$

m $\dfrac{36m^7}{12m^2}$ n $\dfrac{5w^2}{25w}$ o $\dfrac{4a^4}{20a^3}$ p $\dfrac{7x^5}{63x}$

q $\dfrac{16x^8y^6}{12x^2y^3}$ r $\dfrac{6s^6t^3}{14s^5t}$ s $\dfrac{8m^5n^4}{6m^4n^3}$ t $\dfrac{5x^2y}{xy}$



Hint for Q8: Rearrange first, and group numbers and like bases together.

Problem-solving and reasoning 9–10(½) 9–12(½)

Example 12 Combining the index laws for multiplication and division

Simplify each of the following using index laws for multiplication and division.

a $x^2 \times x^3 \div x^4$ b $\dfrac{2a^3b \times 8a^2b^3}{4a^4b^2}$

Solution

a $x^2 \times x^3 \div x^4 = x^5 \div x^4$
$= x$

b $\dfrac{2a^3b \times 8a^2b^3}{4a^4b^2} = \dfrac{2 \times 8 \times a^3a^2 \times bb^3}{4 \times a^4 \times b^2}$
$= \dfrac{16}{4} \times \dfrac{a^5}{a^4} \times \dfrac{b^4}{b^2}$
$= 4 \times a \times b^2$
$= 4ab^2$

Explanation

Use the law for multiplication to add the indices for $x^2 \times x^3$.

Use the law for division to subtract the indices for $x^5 \div x^4$.

Rearrange the question with numbers first, then like bases grouped together.

$2 \times 8 = 16, a^3a^2 = a^{3+2} = a^5, bb^3 = b^{1+3} = b^4$

$\dfrac{16}{4} = 4, \dfrac{a^5}{a^4} = a^{5-4} = a^1 = a, \dfrac{b^4}{b^2} = b^{4-2} = b^2$

Now you try

Simplify each of the following using index laws for multiplication and division.

a $a^3 \times a^6 \div a^8$ b $\dfrac{3m^3n^2 \times 4mn^3}{2m^2n^3}$

9 Simplify each of the following using index laws for multiplication and division.

a $b^5 \times b^2 \div b$ b $y^5 \times y^4 \div y^3$ c $c^4 \div c \times c^4$

d $x^4 \times x^2 \div x^5$ e $\dfrac{t^4 \times t^3}{t^6}$ f $\dfrac{p^2 \times p^7}{p^3}$

g $\dfrac{d^5 \times d^3}{d^2}$ h $\dfrac{x^9 \times x^2}{x}$ i $\dfrac{3x^3y^4 \times 8xy}{6x^2y^2}$ j $\dfrac{9b^4}{2g^3} \times \dfrac{4g^4}{3b^2}$



Hint for Q9: Write pronumerals in alphabetical order.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

7B

10 Simplify each of the following.

a $\frac{m^4}{n^2} \times \frac{m}{n^3}$ **b** $\frac{x}{y} \times \frac{x^3}{y}$ **c** $\frac{a^4}{b^3} \times \frac{b^6}{a}$

d $\frac{12a}{3c^3} \times \frac{6a^4}{4c^4}$ **e** $\frac{3f^2 \times 8f^7}{4f^3}$ **f** $\frac{4x^2b \times 9x^3b^2}{3xb}$



Hint for Q10: Multiply numerators and denominators first.

11 Write the missing number.

a $2^7 \times 2^{\square} = 2^{19}$ **b** $6^{\square} \times 6^3 = 6^{11}$ **c** $11^6 \div 11^{\square} = 11^3$

d $19^{\square} \div 19^2 = 19$ **e** $x^6 \times x^{\square} = x^7$ **f** $a^{\square} \times a^2 = a^{20}$

g $b^{13} \div b^{\square} = b$ **h** $y^{\square} \div y^9 = y^2$ **i** $\square \times x^2 \times 3x^4 = 12x^6$

j $15y^4 \div (\square y^3) = y$ **k** $\square a^9 \div (4a) = \frac{a^8}{2}$ **l** $13b^6 \div (\square b^5) = \frac{b}{3}$

12 Evaluate without using a calculator.

a $7^7 \div 7^5$ **b** $10^6 \div 10^5$ **c** $13^{11} \div 13^9$

d $2^{20} \div 2^{17}$ **e** $101^5 \div 101^4$ **f** $200^{30} \div 200^{28}$

g $7 \times 31^{16} \div 31^{15}$ **h** $3 \times 50^{200} \div 50^{198}$ **i** $-4 \times 10^{20} \div 10^{17}$




Hint for Q12: Simplify using index laws first.


Areas and index notation — 13

13 a Write the area of each of these shapes using index notation.


i




ii



iii




iv




b Find the width of each of these shapes using index notation.

i Area = $24a^3m^2$



ii Area = $36x^2y^4$



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

7C Index laws for power of a power and the zero index

Learning intentions

- To know the index law for raising a power to another power
- To know the rule for the zero power
- To be able to simplify expressions involving a power of a power and the zero index

Key vocabulary: index law, zero index

Sometimes we find that expressions already written in index form are raised to another power, such as $(2^3)^4$ or $(a^2)^5$.

Consider $(a^2)^5$.

$$\begin{aligned}(a^2)^5 &= \overbrace{a^2} \times \overbrace{a^2} \times \overbrace{a^2} \times \overbrace{a^2} \times \overbrace{a^2} \\ &= \overbrace{a \times a} \times \overbrace{a \times a} \times \overbrace{a \times a} \times \overbrace{a \times a} \times \overbrace{a \times a} \\ &= a^{10}\end{aligned}$$

When a^2 appears 5 times there are a total of $5 \times 2 = 10$ factors of a. So when a^m appears n times, there will be a total of $m \times n = mn$ factors of a.

The power of 0 has a special property.

Consider $\dfrac{a^3}{a^3}$.

Simplify using expanded form:

$$\frac{a^3}{a^3} = \frac{\cancel{a}^1 \times \cancel{a}^1 \times \cancel{a}^1}{\cancel{a}_1 \times \cancel{a}_1 \times \cancel{a}_1} = 1$$

Simplify using the index law for division:

$$\frac{a^3}{a^3} = a^{3-3} = a^0$$

So $a^0 = 1$

Lesson starter: Discovering laws for power of a power and the zero index

Use the expanded form of 5^3 to simplify $(5^3)^2$ as shown. $(5^3)^2 = 5 \times \square \times \square \times 5 \times \square \times \square$

$= 5^{\square}$

- Repeat these steps to also simplify $(3^2)^4$ and $(x^4)^2$.
- What do you notice about the given expression and answer in each case? Can you express this as a law or rule in words?
  Now copy and complete this table.

Index form	3^5	3^4	3^3	3^2	3^1	3^0
Basic numeral	243	81				

- What pattern do you notice in the basic numerals?
- What conclusion do you come to regarding 3^0?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

7C

Key ideas

- Index law for power of a power $(a^m)^n = a^{m \times n} = a^{mn}$
  - When raising a term in index form to another power, retain the base and multiply the indices. For example: $(x^2)^3 = x^{2 \times 3} = x^6$.
  - A power outside brackets only applies to the expression inside those brackets. For example: $5(a^3)^2 = 5a^{3 \times 2} = 5a^6$.
- The **zero index**: $a^0 = 1$, where $a \neq 0$
  - Any term except 0 raised to the power of zero is 1. For example: $5^0 = 1$, $m^0 = 1$ and $(2a)^0 = 1$.

Exercise 7C

Understanding

1–4 | 4



Hint for Q1: Choose from 0, 1, 2, *divide*, *add*, *subtract* or *multiply*.

1 Write the missing word or number in these sentences.

a When raising a term or numbers in index form to another power, _____ the indices.

b Any number (except 0) raised to the power 0 is equal to _____.

2 Write the missing numbers in these tables.

a

Index form	2^6	2^5	2^4	2^3	2^2	2^1	2^0
Basic numeral	64	32					

b

Index form	4^5	4^4	4^3	4^2	4^1	4^0
Basic numeral	1024	256				

3 Copy and complete this working.

a $(4^2)^3 = 4^2 \times 4^2 \times 4^2$
$= (4 \times \square) \times (4 \times \square) \times (4 \times \square)$
$= 4^{\square}$

b $(12^3)^2 = 12^{\square} \times 12^{\square}$
$= (12 \times \square \times \square) \times (12 \times \square \times \square)$
$= 12^{\square}$

c $(x^4)^2 = x^{\square} \times x^{\square}$
$= (x \times \square \times \square \times \square) \times (x \times \square \times \square \times \square)$
$= x^{\square}$

d $(a^2)^5 = a^{\square} \times a^{\square} \times a^{\square} \times a^{\square} \times a^{\square}$
$= (a \times \square) \times (a \times \square) \times (a \times \square) \times (a \times \square) \times (a \times \square)$
$= a^{\square}$

4 Find the value of each of the following.

a 6^0 **b** 21^0 **c** 2^0 **d** 1^0

e $(3.7)^0$ **f** 582^0 **g** $\left(\frac{3}{4}\right)^0$ **h** 2760^0



Hint for Q4: $a^0 = 1$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Fluency

5–7(½) | 5–9(½)

Example 13 Using the index law for power of a power

Apply the index law for power of a power to simplify each of the following.

a $(x^5)^4$ b $3(y^5)^2$

Solution

a $(x^5)^4 = x^{5\times4}$
$= x^{20}$

Explanation

Keep x as the base and multiply the indices.

b $3(y^5)^2 = 3y^{5\times2}$
$= 3y^{10}$

Keep y and multiply the indices. The power of 2 is only applied to what is inside the brackets.

The 3 is unchanged.

Now you try

Apply the index law for power of a power to simplify each of the following.

a $(a^3)^7$ b $2(b^2)^4$

5 Apply the index law for power of a power to simplify each of the following. Leave your answers in index form.

a $(y^6)^2$ b $(m^3)^6$ c $(x^2)^5$ d $(b^3)^4$
e $(3^2)^3$ f $(4^3)^5$ g $(3^5)^6$ h $(7^5)^2$
i $5(m^8)^2$ j $4(q^7)^4$ k $-3(c^2)^5$ l $2(f^4)^6$



Hint for Q5: Keep the base. Multiply the indices.

Example 14 Using the zero index

Apply the zero index rule to evaluate each of the following.

a $(-3)^0$ b $3(5x)^0$ c $2y^0 - (3y)^0$

Solution

a $(-3)^0 = 1$

Explanation

Any number raised to the power of 0 is 1.

b $3(5x)^0 = 3 \times 1$
$= 3$

Everything in the brackets is to the power of 0 so $(5x)^0$ is 1. The 3 is not to the power of 0.

c $2y^0 - (3y)^0 = 2 \times 1 - 1$
$= 2 - 1$
$= 1$

$2y^0$ has no brackets so the power applies to the y only, so $2y^0 = 2 \times y^0 = 2 \times 1$ while $(3y)^0 = 1$.

Now you try

Apply the zero index rule to evaluate each of the following.

a $\left(\frac{2}{3}\right)^0$ b $2(7y)^0$ c $(6y)^0 + 2y^0$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

7C

6 Evaluate each of the following.

a 5^0 **b** 9^0 **c** $(-6)^0$ **d** $(-3)^0$

e $(4)^0$ **f** $\left(\frac{3}{4}\right)^0$ **g** $\left(-\frac{1}{7}\right)^0$ **h** $(4y)^0$

i $5m^0$ **j** $-3p^0$ **k** $6x^0 - 2x^0$ **l** $-5n^0 - (8n)^0$

m $(3x^4)^0$ **n** $1^0 + 2^0 + 3^0$ **o** $3a^0 + (3a)^0$ **p** $100^0 - a^0$



Hint for Q6: Any number (except zero) raised to the power 0 equals 1.

Example 15 Combining index laws for multiplication and power of a power

Simplify $(x^2)^3 \times (x^3)^5$ by applying the various index laws.

Solution

$$(x^2)^3 \times (x^3)^5 = x^{2\times 3} \times x^{3\times 5}$$
$$= x^6 \times x^{15}$$
$$= x^{21}$$

Explanation

Use the index law for power of a power to remove brackets first by multiplying indices. Then use the index law for multiplication to add indices.

Now you try

Simplify $(a^3)^3 \times (a^2)^4$ by applying the varying index laws.

7 Simplify each of the following by combining various index laws.

a $4 \times (4^3)^2$ **b** $(3^4)^2 \times 3$ **c** $x \times (x^0)^5$

d $y^5 \times (y^2)^4$ **e** $b^5 \times (b^3)^3$ **f** $(a^2)^3 \times a^4$

g $(d^3)^4 \times (d^2)^6$ **h** $(y^2)^6 \times (y)^4$ **i** $z^4 \times (z^3)^2 \times (z^5)^3$



Hint for Q7: First remove brackets by multiplying indices. Remember that $4 = 4^1$.

Example 16 Combining index laws for division and power of a power

Simplify $\frac{(m^3)^4}{m^7}$ by applying index laws.

Solution

$$\frac{(m^3)^4}{m^7} = \frac{m^{3\times 4}}{m^7}$$
$$= \frac{m^{12}}{m^7}$$
$$= m^5$$

Explanation

Remove brackets by multiplying indices then simplify using the index law for division.

$12 - 7 = 5$

Now you try

Simplify $\frac{a^8}{(a^2)^3}$ by applying index laws.

8 Simplify each of the following.

a $\frac{(b^2)^5}{b^4}$ **b** $\frac{(x^4)^3}{x^7}$ **c** $\frac{(y^3)^3}{y^3}$

d $7^8 \div (7^3)^2$ **e** $(4^2)^3 \div 4^5$ **f** $(3^6)^3 \div (3^5)^2$

g $(m^3)^6 \div (m^2)^9$ **h** $(y^5)^3 \div (y^6)^2$ **i** $(h^{11})^2 \div (h^5)^4$



Hint for Q8: First remove brackets by multiplying the powers.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Example 17 Combining index laws for multiplication, division and the zero index

Simplify $\frac{4x^2 \times 3x^3}{6x^5}$ by applying index laws.

Solution

$$\frac{4x^2 \times 3x^3}{6x^5} = \frac{4 \times 3 \times x^2 \times x^3}{6x^5}$$
$$= \frac{12x^5}{6x^5}$$
$$= 2x^0$$
$$= 2 \times 1$$
$$= 2$$

Explanation

Regroup, then simplify the numerator first by multiplying numbers and adding indices of base x. $12 \div 6 = 2$, and subtract indices: $5 - 5 = 0$

The zero index rule says $x^0 = 1$.

Now you try

Simplify $\frac{5y^4 \times 3y^2}{10y^6}$ by applying index laws.

9 Simplify each of the following using various index laws.

a $\frac{3x^4 \times 6x^8}{9x^{12}}$ **b** $\frac{5x^5 \times 4x^6}{2x^{10}}$ **c** $\frac{24(x^4)^4}{8(x^4)^2}$

d $\frac{4(d^4)^3 \times (e^4)^2}{8(d^2)^5 \times e^7}$ **e** $\frac{6(m^3)^2(n^5)^3}{15(m^5)^0(n^2)^7}$ **f** $\frac{2(a^3)^4(b^2)^6}{16(a)^0(b^6)^2}$



Hint for Q9: Remove brackets, then rearrange with numbers first and then like bases grouped together.

Problem-solving and reasoning

10 | 10, 11

10 If m and n are positive integers, in how many ways can $(a^m)^n = a^{16}$? Show each possibility.

11 Explain the error made in the following problems, then give the correct answer.

a $(a^4)^5 = a^9$ **b** $3(x^4)^2 = 9x^6$ **c** $(2x)^0 = 2$

Rabbits!

— | 12

12 There are 100 rabbits on Mt Burrow at the start of the year 2015. The rule for the number of rabbits, N, after t years (from the start of the year 2015) is $N = 100 \times 2^t$.

a Find the number of rabbits at:

i $t = 2$ **ii** $t = 6$ **iii** $t = 0$

b Find the number of rabbits at the beginning of:

i 2018 **ii** 2022 **iii** 2025

c How many years will it take for the population to first rise to more than 500 000? Give a whole number of years.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

7D Index laws for brackets and fractions

Learning intentions

- To know the index laws for removing brackets over multiplication and division
- To be able to expand and simplify expressions involving brackets
- To be able to combine a range of index laws to simplify more complex expressions

Key vocabulary: index law

It is common to find expressions such as $(2x)^3$ and $\left(\frac{x}{3}\right)^4$ in mathematical problems. These are different to most of the expressions in previous sections. They contain more than one single number or pronumeral, connected by multiplication or division, raised to a power. These expressions can also be simplified using two index laws that remove the brackets.

Consider $(a \times b)^6$.

Using expanded form: $(a \times b)^6 = \overbrace{ab \times ab \times ab \times ab \times ab \times ab}^{6 \text{ factors of } ab}$

$$= \overbrace{a \times a \times a \times a \times a \times a}^{6 \text{ factors of } a} \times \overbrace{b \times b \times b \times b \times b \times b}^{6 \text{ factors of } b}$$

$$= a^6 \times b^6$$

So this becomes a product of 6 factors of a and 6 factors of b.

Also, $\left(\frac{a}{b}\right)^6 = \overbrace{\frac{a}{b} \times \frac{a}{b} \times \frac{a}{b} \times \frac{a}{b} \times \frac{a}{b} \times \frac{a}{b}}^{6 \text{ factors of } \frac{a}{b}}$

$$= \frac{\overbrace{a \times a \times a \times a \times a \times a}^{6 \text{ factors of } a}}{\underbrace{b \times b \times b \times b \times b \times b}_{6 \text{ factors of } b}}$$

$$= \frac{a^6}{b^6}$$

So to remove the brackets we can raise each of a and b to the power 6.

Lesson starter: Discovering laws for brackets and fractions

Use the expanded form of $(2x)^3$ and $\left(\frac{x}{3}\right)^4$ to help simplify the expressions.

$(2x)^3 = 2x \times \square \times \square$

$= 2 \times 2 \times 2 \times \square \times \square \times \square$

$= 2^{\square} \times \square^{\square}$

$\left(\frac{x}{3}\right)^4 = \frac{x}{3} \times \square \times \square \times \square$

$= \frac{x \times \square \times \square \times \square}{3 \times \square \times \square \times \square}$

$= \frac{\square^{\square}}{\square^{\square}}$

- Repeat these steps to also simplify the expressions $(3y)^4$ and $\left(\frac{x}{2}\right)^5$.
- What do you notice about the given expressions and the answer in each case? Can you express this as a rule or law in words?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Key ideas

- Index law for brackets: $(a \times b)^m = (ab)^m = a^m b^m$
  - If the product of two or more numbers is raised to the power of m, raise each number in the brackets to the power of m. For example: $(2x)^2 = 2^2 x^2 = 4x^2$.
- Index law for fractions: $\left(\frac{a}{b}\right)^m = \frac{a^m}{b^m}$ and $b \neq 0$
  - If the quotient of two numbers is raised to the power of m, raise each number in the brackets to the power of m. For example: $\left(\frac{y}{3}\right)^3 = \frac{y^3}{3^3} = \frac{y^3}{27}$.

Exercise 7D

Understanding

1–4 | 4

1 Copy and complete index laws for brackets and fractions.

a $(a \times b)^m = a^m \times \square$

b $\left(\frac{a}{b}\right)^m = \frac{a^m}{\square}$.

2 Copy and complete this working.

a $(ab)^4 = ab \times \square \times \square \times \square$
$= a \times \square \times \square \times \square \times b \times \square \times \square \times \square$
$= a^4 \times \square$

b $\left(\frac{x}{6}\right)^3 = \frac{x}{6} \times \square \times \square$
$= \frac{x \times \square \times \square}{6 \times \square \times \square}$
$= \frac{x^3}{\square}$

3 Copy and complete these bracket expansions.

a $(xy)^5 = x^{\square} y^5$

b $(2m)^3 = 2^{\square} m^{\square}$

c $(abc)^4 = a^4 b^{\square} c^{\square}$

d $\left(\frac{5}{k}\right)^3 = \frac{5^3}{k^{\square}}$

e $\left(\frac{a}{m}\right)^5 = \frac{a^{\square}}{m^{\square}}$

f $\left(\frac{2}{3}\right)^4 = \frac{2^{\square}}{3^{\square}} = \frac{16}{\square}$



Hint for Q3: Raise each number or pronumeral in the brackets to the power.

4 Copy and complete these bracket expansions.

a $6(ab)^2 = 6a^{\square} b^2$

b $4(xy)^3 = \square x^{\square} y^{\square}$

c $7(2m)^2 = 7 \times 2^{\square} m^{\square}$
$= 7 \times \square \times m^2$
$= \square m^2$

d $4(3ab)^2 = \square \times 3^{\square} a^{\square} b^{\square}$
$= 4 \times \square a^{\square} b^{\square}$
$= \square a^{\square} b^{\square}$



Hint for Q4: The index outside the brackets only applies to the values inside the brackets.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

7D

Fluency

5–7(½) 5–7(½)

Example 18 Using the index law for brackets

Expand each of the following using the index law for brackets.

a $(3a)^2$ b $(9a^3)^2$

Solution

a $(3a)^2 = 3^2 \times a^2$
$= 9a^2$

b $(9a^3)^2 = 9^2(a^3)^2$
$= 81a^{3\times 2}$
$= 81a^6$

Explanation

Apply the power of 2 to each factor in the brackets.

Raise 9 and (a^3) to the power of 2.
$9^2 = 9 \times 9 = 81$, $(a^3)^2 = a^{3\times 2} = a^6$

Now you try

Expand each of the following using index law for brackets.

a $(4x)^3$ b $(2y^4)^4$

5 Expand each of the following using the index law for brackets. Evaluate numbers raised to a power.

a $(2x)^3$ b $(5y)^2$ c $(4a)^3$ d $(3r)^2$
e $(3b)^4$ f $(7r)^3$ g $(2h^2)^4$ h $(5c^2)^4$
i $(3a^2)^3$ j $(7p^4)^2$ k $(5m^3)^2$ l $(3y^{10})^2$



Hint for Q5: The index outside the brackets must be applied to each number or pronumeral inside the brackets.

Example 19 Using the index law for fractions

Apply the index law for fractions to the following.

a $\left(\frac{x}{2}\right)^3$ b $\left(\frac{m^3}{5}\right)^2$

Solution

a $\left(\frac{x}{2}\right)^3 = \frac{x^3}{2^3}$
$= \frac{x^3}{8}$

b $\left(\frac{m^3}{5}\right)^2 = \frac{(m^3)^2}{5^2}$
$= \frac{m^{3\times 2}}{5^2}$
$= \frac{m^6}{25}$

Explanation

Raise each factor in the brackets to the power of 3.

Raise m^3 and 5 to the power of 2.
$(m^3)^2 = m^{3\times 2} = m^6$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Now you try

Apply the index law for fractions to the following.

a $\left(\frac{a}{b}\right)^2$ b $\left(\frac{3a}{b^2}\right)^3$

6 Apply the index law for fractions to expand the following. Evaluate numbers raised to a power.

a $\left(\frac{p}{q}\right)^3$ b $\left(\frac{x}{y}\right)^4$ c $\left(\frac{4}{y}\right)^3$

d $\left(\frac{5}{p^2}\right)^4$ e $\left(\frac{2}{r^3}\right)^2$ f $\left(\frac{s^3}{7}\right)^2$

g $\left(\frac{7}{x^3}\right)^2$ h $\left(\frac{w^4}{10}\right)^2$ i $\left(\frac{a^2}{5}\right)^3$

j $\left(\frac{m^2}{2}\right)^4$ k $\left(\frac{2m}{n}\right)^5$ l $\left(\frac{2a^2}{3}\right)^3$



Hint for Q6: Recall: $4^3 = 4 \times 4 \times 4 = 64$

Example 20 Using the index law for brackets for more complex expressions

Expand each of the following using the index law for brackets.

a $(-2x^3y)^4$ b $4(c^2d^3)^5$

Solution	Explanation
a $(-2x^3y)^4 = (-2)^4(x^3)^4y^4$ $= 16x^{12}y^4$	Raise each value in the brackets to the power of 4. Evaluate $(-2)^4 = -2 \times (-2) \times (-2) \times (-2)$ and simplify $(x^3)^4$ using the law for power of a power.
b $4(c^2d^3)^5 = 4(c^2)^5(d^3)^5$ $= 4c^{10}d^{15}$	Raise each value in the brackets to the power of 5. Note that the coefficient (4) is not raised to the power of 5 since it is not in the brackets. Simplify using index laws.

Now you try

Expand each of the following using the index law for brackets.

a $(-4a^2b)^2$ b $6(a^3b^2)^4$

7 Expand each of the following using the index law for brackets.

a $(2x^3y^2)^5$ b $9(p^2q^4)^3$ c $2(x^3y)^2$

d $(8t^2u^9v^4)^0$ e $(-3w^3y)^3$ f $-4(p^4qr)^2$

g $(-5s^7t)^2$ h $-(-2x^4yz^3)^3$ i $-3(-2ab^2)^3$



Hint for Q7: Powers only apply to numbers or pronumerals inside the brackets.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

7D

Problem-solving and reasoning

8(½) 8–9(½)

Example 21 Using the index law for fractions for more complex expressions

Apply the index law for fractions to $\left(\frac{-2a^2}{3bc^3}\right)^4$.

Solution

$$\left(\frac{-2a^2}{3bc^3}\right)^4 = \frac{(-2)^4a^8}{3^4b^4c^{12}}$$
$$= \frac{16a^8}{81b^4c^{12}}$$

Explanation

Raise each value in the brackets to the power of 4.
Evaluate $(-2)^4$ and 3^4.

Now you try

Apply the index law for fractions to $\left(\frac{-3x^2}{2yz^3}\right)^3$.

8 Apply the index law for fractions to expand the following. Evaluate numbers that are raised to a power.

a $\left(\frac{3n^3}{2m^4}\right)^3$ **b** $\left(\frac{-2r}{n}\right)^4$ **c** $\left(\frac{-3f}{2^3g^5}\right)^2$ **d** $\left(\frac{5w^4y}{2x^3}\right)^2$

e $\left(\frac{-3x}{2y^3g^5}\right)^2$ **f** $\left(\frac{3km^3}{4n^7}\right)^3$ **g** $-\left(\frac{-5w^4y}{2zx^3}\right)^2$ **h** $\left(-\frac{3x^2y^3}{2a^5b^3}\right)^2$

Example 22 Using a variety of index laws

Simplify each of the following by applying various index laws.

a $a(-2a^2b)^3$ **b** $\left(\frac{x^2y^3}{c}\right)^3 \times \left(\frac{xc}{y}\right)^4$

Solution

a $a(-2a^2b)^3 = a(-2)^3(a^2)^3b^3$
$= a \times (-8a^6b^3)$
$= -8a^7b^3$

Explanation

Apply the power 3 to the values inside the brackets.
$(-2)^3 = -2 \times (-2) \times (-2) = -8$
$(a^2)^3 = a^{2\times3} = a^6$
Combine powers with a common base:
$a \times a^6 = a^{1+6} = a^7$

b $\left(\frac{x^2y^3}{c}\right)^3 \times \left(\frac{xc}{y}\right)^4 = \frac{x^6y^9}{c^3} \times \frac{x^4c^4}{y^4}$
$= \frac{x^{10}y^9c^4}{c^3y^4}$
$= cx^{10}y^5$

Raise each value in the brackets to the power. Multiply the numerators using the law for multiplication then divide using the law for division.

Write the answer in alphabetical order.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Now you try

Simplify each of the following by applying various index laws.

a $2(-3ab^2)^2$

b $\left(\frac{xy^2}{b}\right)^2 \times \left(\frac{b^2}{xy^4}\right)$

9 Simplify each of the following by applying the various index laws.

a $a(3b)^2$ **b** $a(3b^2)^3$ **c** $-3(2a^3b^4)^2a^2$

d $2(3x^2y^3)^3$ **e** $(-4b^2c^5d)^3$ **f** $a(2a)^3$

g $a(3a^2)^2$ **h** $5a^3(-2a^4b)^3$ **i** $-5(-2m^3pt^2)^5$

j $\left(\frac{-3x^2y^0}{5a^5b^3}\right)^3$ **k** $\left(\frac{a^3b}{c}\right)^3 \times \left(\frac{ac^4}{b}\right)^2$ **l** $\left(\frac{x^2z}{y}\right)^4 \times \left(\frac{xy^2}{z}\right)^3$



Hint for Q9: Remove brackets. Combine numerators and denominators then divide for parts j to l.

Germinating seeds — 10

10 The rule for the number of seeds germinating in a glass house over a two-week period is given by $N = \left(\frac{t}{2}\right)^3$, where N is the number of germinating seeds and t is the number of days.

a Find the number of germinating seeds after:

i 4 days **ii** 10 days

b Use the index law for fractions to rewrite the rule without brackets.

c Use your rule in part **b** to find the number of seeds germinating after:

i 6 days **ii** 4 days

d Find the number of days required to germinate:

i 64 seeds **ii** 1 seed



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

7E Negative indices

Learning intentions

- To know how negative indices can be equivalently expressed using positive indices
- To be able to express negative indices in terms of positive indices
- To be able to use the index laws with negative indices

Key vocabulary: negative, index/indices, positive

We know that $2^3 = 8$ and $2^0 = 1$ but what about 2^{-1} or 2^{-6}? Numbers written in index form using negative indices also have meaning in mathematics.

Consider the expression $= \frac{a^2}{a^5}$

Method 1: Using the law for division

$$\frac{a^2}{a^5} = a^{2-5}$$
$$= a^{-3}$$

Method 2: By cancelling

$$\frac{a^2}{a^5} = \frac{a^{\cancel{1}} \times a^{\cancel{1}}}{a \times a \times a \times \cancel{a_1} \times \cancel{a_1}}$$
$$= \frac{1}{a^3}$$

So we can see that $a^{-3} = \frac{1}{a^3}$

Lesson starter: Continuing the pattern

Explore the use of negative indices by completing this table.

Index form	2^4	2^3	2^2	2^1	2^0	2^{-1}	2^{-2}	2^{-3}
Whole number or fraction	16	8					$\frac{1}{4} = \frac{1}{2^2}$	

÷2 ÷2 ÷2 ÷2 ÷2 ÷2 ÷2

- What do you notice about the numbers with negative indices in the top row and the fractions in the second row?
- Can you describe this connection in words?
- What might be another way of writing 2^{-7} or 5^{-4}?

Key ideas

- To express a negative index as a positive index use the following rules:
  $a^{-m} = \frac{1}{a^m}$ and $a^m = \frac{1}{a^{-m}}$
  a raised to the power $-m$ is equal to the reciprocal of a raised to the power m. $(a \neq 0)$
  For example: $5^{-2} = \frac{1}{5^2}$, $\frac{1}{3^{-4}} = 3^4$.
  - Recall: the reciprocal of $\frac{a}{b}$ is $\frac{b}{a}$.
- All the index laws also apply to expressions with negative indices.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Exercise 7E

Understanding

1–4 | 4

1 Write the following using positive indices. For example: $\frac{1}{8} = \frac{1}{2^3}$.

a $\frac{1}{4}$ **b** $\frac{1}{9}$ **c** $\frac{1}{125}$ **d** $\frac{1}{27}$



Hint for Q1: Write the denominator as a power of a prime number.

2 Copy and complete these tables.

a

Index form	3^4	3^3	3^2	3^1	3^0	3^{-1}	3^{-2}	3^{-3}
Whole number or fraction	81	27					$\frac{1}{9} = \frac{1}{3^2}$	

÷3 ÷3 ÷3 ÷3 ÷3 ÷3 ÷3

b

Index form	10^4	10^3	10^2	10^1	10^0	10^{-1}	10^{-2}	10^{-3}
Whole number or fraction	10 000							$\frac{1}{1000} = \frac{1}{10^3}$

÷10 ÷10 ÷10 ÷10 ÷10 ÷10 ÷10



3 Copy and complete each of the following.

a $10^{-4} = \frac{1}{10^{\square}}$ **b** $3^{-2} = \frac{1}{3^{\square}}$ **c** $7^{-3} = \frac{1}{7^{\square}}$

Hint for Q3: $a^{-m} = \frac{1}{a^m}$

So $10^{-4} = \frac{1}{10^4}$



4 Copy and complete each of the following.

a $\frac{1}{3^{-4}} = 3^{\square}$ **b** $\frac{1}{7^{-6}} = 7^{\square}$ **c** $\frac{1}{8^{-3}} = 8^{\square}$

Hint for Q4: $\frac{1}{a^{-m}} = a^m$

So $\frac{1}{3^{-4}} = 3^4$

Fluency

5–7(½) | 5–8(½)

Example 23 Using $a^{-m} = \frac{1}{a^m}$

Rewrite the following using positive indices.

a 3^{-2} **b** 5×4^{-3}

Solution

a $3^{-2} = \frac{1}{3^2}$

b $5 \times 4^{-3} = \frac{5}{1} \times \frac{1}{4^3}$

$= \frac{5}{4^3}$

Explanation

Use $a^{-m} = \frac{1}{a^m}$.

The 5 does not have a negative power so it remains unchanged.

$4^{-3} = \frac{1}{4^3}$, $5 \times 1 = 5$, $1 \times 4^3 = 4^3$

Now you try

Rewrite the following using positive indices.

a 6^{-4} **b** 7×2^{-4}

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

7E

5 Write each of the following with positive indices.

a 5^{-2} **b** 7^{-4}
c 8^{-3} **d** 3^{-5}
e 9^{-2} **f** 10^{-3}



Hint for Q5: Remember: $a^{-m} = \frac{1}{a^m}$

So $5^{-2} = \frac{1}{5^2}$

6 Rewrite the following with positive indices only.

a 3×2^{-4} **b** 5×4^{-3} **c** 7×5^{-6}
d 2×3^{-4} **e** 4×3^{-5} **f** 9×5^{-2}
g 8×7^{-3} **h** 6×5^{-6} **i** 1×4^{-2}



Hint for Q6: $5 \times 2^{-3} = \frac{5}{2^3}$

Don't change

Example 24 Using $\frac{1}{a^{-m}} = a^m$

Rewrite the following with positive indices.

a $\frac{1}{2^{-3}}$ **b** $\frac{7}{3^{-2}}$

Solution

a $\frac{1}{2^{-3}} = 2^3$

b $\frac{7}{3^{-2}} = 7 \times \frac{1}{3^{-2}}$

$= 7 \times 3^2$

Explanation

Use $\frac{1}{a^{-m}} = a^m$.

The 7 remains unchanged.

$\frac{1}{3^{-2}} = 3^2$

Now you try

Rewrite the following with positive indices.

a $\frac{1}{10^{-3}}$ **b** $\frac{5}{9^{-4}}$

7 Write each of the following with positive powers.

a $\frac{1}{2^{-4}}$ **b** $\frac{1}{3^{-2}}$ **c** $\frac{1}{4^{-3}}$ **d** $\frac{1}{6^{-5}}$
e $\frac{1}{5^{-3}}$ **f** $\frac{1}{8^{-5}}$ **g** $\frac{1}{7^{-3}}$ **h** $\frac{1}{9^{-4}}$



Hint for Q7: Remember: $\frac{1}{a^{-m}} = a^m$

So $\frac{1}{2^{-4}} = 2^4$

8 Rewrite the following with positive indices only.

a $\frac{6}{4^{-3}}$ **b** $\frac{5}{8^{-2}}$ **c** $\frac{4}{7^{-5}}$ **d** $\frac{3}{2^{-5}}$
e $\frac{12}{5^{-4}}$ **f** $\frac{6}{10^{-3}}$ **g** $\frac{7}{9^{-4}}$ **h** $\frac{7}{10^{-6}}$



Hint for Q8:

Don't change → $\frac{7}{3^{-4}} = 7 \times 3^4$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Problem-solving and reasoning 9–10(½) 9–11(½), 12

Example 25 Changing to fractions

Rewrite 3×2^{-4} with a positive index and then as a fraction.

Solution

$$3 \times 2^{-4} = 3 \times \frac{1}{2^4}$$
$$= \frac{3}{2^4}$$
$$= \frac{3}{16}$$

Explanation

$$2^{-4} = \frac{1}{2^4}$$
$$2^4 = 2 \times 2 \times 2 \times 2 = 16$$

Now you try

Rewrite 7×5^{-2} with a positive index and then as a fraction.

9 Rewrite each of these with positive indices and then as a fraction.

a 6×5^{-2} **b** 2×3^{-2} **c** 4×5^{-3}
d 6×7^{-1} **e** 4×10^{-3} **f** 2×10^{-1}
g 5×2^{-4} **h** 4×5^{-1} **i** 1×3^{-2}



Hint for Q9:
Remember: $5^1 = 5$
So $3 \times 5^{-1} = 3 \times \frac{1}{5^1} = \frac{3}{5}$

Example 26 Changing to fractions and decimals

Rewrite 5×10^{-3} with a positive index and then as a fraction and a decimal.

Solution

$$5 \times 10^{-3} = 5 \times \frac{1}{10^3}$$
$$= \frac{5}{10^3}$$
$$= \frac{5}{1000}$$
$$= 0.005$$

Explanation

$$10^{-3} = \frac{1}{10^3}$$
$$10^3 = 10 \times 10 \times 10 = 1000$$
5 thousandths = 0.005

Now you try

Rewrite 3×10^{-2} with a positive index and then as a fraction and a decimal.

10 Rewrite each of the following with positive indices and then as a fraction and a decimal.

a 2×10^{-3} **b** 5×10^{-2}
c 7×10^{-1} **d** 3×10^{-4}
e 5×10^{-4} **f** 8×10^{-5}
g 2×10^{-6} **h** 4×10^{-8}



Hint for Q10:
$\frac{1}{10} = 0.1$, $\frac{1}{100} = 0.01$
$\frac{1}{1000} = 0.001$
$\frac{1}{10\,000} = 0.0001$
$\frac{1}{100\,000} = 0.00001$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

7E

Example 27 Evaluating without a calculator

Express using positive indices only, then evaluate without using a calculator.

a 3^{-4} b $\frac{5}{3^{-2}}$ c $\left(\frac{2}{3}\right)^{-4}$

Solution

a $3^{-4} = \frac{1}{3^4}$

$= \frac{1}{81}$

b $\frac{5}{3^{-2}} = 5 \times \frac{1}{3^{-2}}$

$= 5 \times 3^2$

$= 5 \times 9$

$= 45$

c $\left(\frac{2}{3}\right)^{-4} = \frac{2^{-4}}{3^{-4}}$

$= 2^{-4} \times \frac{1}{3^{-4}}$

$= \frac{1}{2^4} \times 3^4$

$= \frac{3^4}{2^4}$

$= \frac{81}{16}$

Explanation

Express 3^{-4} as a positive power.
$3^4 = 3 \times 3 \times 3 \times 3 = 81$

$\frac{1}{3^{-2}} = 3^2$
$3^2 = 3 \times 3 = 9$

Apply the power to each numeral in the brackets using the index law for fractions.
$\frac{a}{b} = a \times \frac{1}{b}$

$2^{-4} = \frac{1}{2^4}$, $\frac{1}{3^{-4}} = 3^4$

$3^4 = 3 \times 3 \times 3 \times 3 = 81$
$2^4 = 2 \times 2 \times 2 \times 2 = 16$

Give answer as an improper fraction.

Now you try

Express using positive indices only, then evaluate without using a calculator.

a 2^{-5} b $\frac{2}{3^{-3}}$ c $\left(\frac{4}{5}\right)^{-2}$

11 Evaluate without the use of a calculator. Express in simplified form.

a 5^{-1} b 3^{-2} c 4×10^{-2}
d -5×10^{-3} e $8 \times (2^2)^{-2}$ f $6^4 \times 6^{-6}$
g $8^{-7} \times (8^2)^3$ h $\frac{1}{8^{-1}}$ i $\frac{2}{2^{-3}}$
j $\frac{2^3}{2^{-3}}$ k $\left(\frac{3}{8}\right)^{-2}$ l $\left(\frac{4}{3}\right)^{-3}$



Hint for Q11: First write expressions using positive indices.

12 Describe the error made in these problems, then give the correct answer.

a $2x^{-2} = \frac{1}{2x^2}$ b $\frac{5}{a^4} = \frac{a^{-4}}{5}$ c $\frac{2}{(3b)^{-2}} = \frac{2b^2}{9}$

Kilograms and grams — 13

13 a The mass of a small insect is 2^{-9} kg. How many grams is this? Round to two decimal places.
b The mass of an asteroid is 3^{20} kg. How many tonnes is this?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Progress quiz

7A **1** Write the following in expanded form and evaluate parts **d–f**.

a b^4 **b** $2x^2y^3$ **c** $(3x)^3$ **d** 4^3 **e** $(-5)^4$ **f** $\left(\frac{3}{5}\right)^3$

7A **2** Write each of the following in index form.

a $5 \times 5 \times y \times y \times y \times y$

b $\frac{2}{3} \times \frac{2}{3} \times \frac{2}{3} \times \frac{5}{7} \times \frac{5}{7}$

c $4 \times a \times b \times 4 \times b \times b \times a \times b \times 4$

7A **3** Express the following as a product of their prime factors in index form.

a 45 **b** 120

7A **4** If $x = 3$, $y = 4$ and $z = -1$, evaluate these expressions.

a $(xz)^3$ **b** $\left(\frac{z}{y}\right)^2$ **c** $x(yz)^2$

7B **5** Simplify using the index laws for multiplication and division; give your answer in index form.

a $4^3 \times 4^4$ **b** $9^6 \div 9^2$ **c** $x^5 \times x^3$

d $a^2b \times a^3b^4$ **e** $y^9 \div y^3$ **f** $18a^5 \div (6a^2)$

g $3mn^4 \times 4m^2n^3$ **h** $\frac{2x^5y^3}{4x^2y}$ **i** $\frac{-6a^3b^2}{3a^2b}$

7C **6** Apply the index law for power of a power or the zero index rule to the following.

a $(x^4)^2$ **b** $5(y^3)^3$ **c** 4^0

d $(5m)^0$ **e** $(3x)^0 - 2x^0$ **f** $5(x^0)^3$

7B/C **7** Combine index laws to simplify the following.

a $\frac{3m^2 \times 4m^3n^5}{m^4n^2}$ **b** $\frac{2(x^2)^3}{12x^4}$ **c** $\frac{(3a)^0 \times 4a^3b^2}{6ab^2}$

7D **8** Expand each of the following using the index laws for brackets and fractions.

a $(2x)^2$ **b** $(2m^3)^4$ **c** $\left(\frac{x^2}{3}\right)^4$

d $4(-2ab^2)^3$ **e** $\left(\frac{3a^2}{5}\right)^3$ **f** $\left(\frac{-2a^2c}{7b^3}\right)^2$

7B/C/D **9** Combine index laws to simplify the following.

a $\frac{(4y^3)^2}{xy^2 \times 8(xy)^2}$ **b** $\left(\frac{3x^2}{4}\right)^3 \times \frac{2y^0}{x^4}$

7E **10** Rewrite the following with positive powers.

a 5^{-2} **b** 3×4^{-4} **c** $\frac{1}{3^{-2}}$ **d** $\frac{5}{2^{-3}}$

7E **11** Evaluate the following without a calculator. Express part **d** as a decimal.

a 9^{-2} **b** 5×2^{-2} **c** $\frac{6}{4^{-2}}$ **d** 6×10^{-4} **e** $\left(\frac{4}{5}\right)^{-3}$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

7F Scientific notation

Learning intentions

- To know that scientific notation is a way of representing very large and very small numbers
- To know the form of numbers written in scientific notation
- To be able to express numbers using scientific notation and as a basic numeral

Key vocabulary: scientific notation, integer

It is common in practical situations to work with very large or very small numbers. For example, the number of cubic metres of concrete used to build the Hoover Dam in the United States was $3\,400\,000$ m^3. The mass of a molecule of water is $0.000000000000000000000000299$ grams. Numbers like this can be written more efficiently using powers of 10 with positive or negative indices. This is called scientific notation. The number is written using a number between 1 and 10 that is multiplied by a power of 10. This notation is also used to show very large and very small time intervals.



The Hoover Dam, constructed with approximately 3.4×10^6 cubic metres of concrete, can hold up to 3.73×10^{10} cubic metres of water at maximum capacity. Scientific notation helps express these large quantities efficiently.

Lesson starter: Equal times

Work in pairs and help each other to complete this 'matching puzzle'.

For each time in the first column, find the equal time in the second column and in the third column.

Column 1	Column 2	Column 3
a 6 thousand seconds	6 000 000 s	6×10^2 s
b 6 millionths of a second	0.006 s	6×10^3 s
c 6 hundred seconds	0.06 s	6×10^6 s
d 6 million seconds	6000 s	6×10^{-2} s
e 6 thousandths of a second	0.000006 s	6×10^{-3} s
f 6 hundredths of a second	600 s	6×10^{-6} s

Key ideas

- Numbers written in **scientific notation** are expressed in the form $a \times 10^m$ where a is a number from 1 to less than 10 (or from greater than -10 to -1) and m is an integer.
  - To write numbers using scientific notation, place the decimal point after the first non-zero digit then multiply by a power of 10.
- Large numbers written in scientific notation will use positive powers of 10.

For example: 4 million years $= 4\,000\,000$ years

$= 4 \times 1\,000\,000$ years

$= 4 \times 10^6$ years

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

- Small numbers written in scientific notation will use negative powers of 10.
  For example: 5 thousandths of a metre = 0.005 metres

$$= \frac{5}{1000} \text{ metres}$$

$$= \frac{5}{10^3} \text{ metres}$$

$$= 5 \times 10^{-3} \text{ metres}$$

Exercise 7F

Understanding

1–4 | 3, 4

1 Copy and complete this table. The first one has been done for you.

Scientific notation	Power of 10 expanded	Basic numeral
5×10^3	5×1000	5 000
3×10^4		
2×10^5		
7×10^2		
		70 000
		400 000



Hint for Q1: The power of 10 is equal to the number of zeros in the expanded form.

2 Copy and complete this table. The first one has been done for you.

Scientific notation	Positive power	Fraction	Basic numeral
2×10^{-4}	$\frac{2}{10^4}$	$\frac{2}{10\,000}$	0.0002
3×10^{-2}			
5×10^{-3}			
7×10^{-6}			
			0.009
			0.08

3 Which of the numbers 1000, 10 000 or 100 000 completes each equation?

a $6.2 \times$ _____ $= 62\,000$
b $9.41 \times$ _____ $= 9410$
c $1.03 \times$ _____ $= 103\,000$
d $3.2 \div$ _____ $= 0.0032$
e $5.16 \div$ _____ $= 0.0000516$
f $1.09 \div$ _____ $= 0.000109$



Hint for Q3: The number of zeros tells you how many places to move the decimal point.

4 If these numbers were written using scientific notation, would positive or negative indices be used?

a 2000
b 0.0004
c 19 300
d 0.00101431

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

7F

Fluency

5–9(½) | 5–9(½)

Example 28 Writing large numbers using scientific notation

Write 4 500 000 using scientific notation.

Solution

$4\,500\,000 = 4.5 \times 10^6$

Explanation

Place the decimal point after the first non-zero digit (4).
Multiply by 10^6 since the decimal point has been moved six places to the left.

Now you try

Write 71 000 using scientific notation.

5 Write the following using scientific notation.

a 40 000	**b** 2 300 000 000 000	**c** 16 000 000 000
d 7 200 000	**e** 3500	**f** 8 800 000
g 52 hundreds	**h** 3 million	**i** 21 thousands



Hint for Q5: Large numbers: use 10 to a positive power.

Example 29 Writing small numbers using scientific notation

Write 0.0000004 using scientific notation.

Solution

$0.0000004 = 4 \times 10^{-7}$

Explanation

The first non-zero digit is 4. Multiply by 10^{-7} since the decimal point has been moved seven places to the right.

Now you try

Write 0.0000275 using scientific notation.

6 Write the following using scientific notation.

a 0.000003	**b** 0.0004	**c** 0.00876
d 0.00000000073	**e** 0.00003	**f** 0.000000000125
g 0.00000000809	**h** 0.000000024	**i** 0.0000345



Hint for Q6: Small numbers: use 10 to a negative power.

7 Write each of the following numbers using scientific notation.

a 6000	**b** 720 000	**c** 324.5	**d** 7869.03
e 8459.12	**f** 0.2	**g** 0.000328	**h** 0.00987
i 0.00001	**j** 460 100 000	**k** 17 467	**l** 128



Hint for Q7: Place the decimal point after the first non-zero digit.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Example 30 Writing basic numerals using positive powers

Express 9.34×10^6 as a basic numeral.

Solution

$9.34 \times 10^6 = 9\,340\,000$

Explanation

Move the decimal point six places to the right, so that the 9 increases in place value by six places from the units position to the millions position.

Now you try

Express 2.4×10^4 as a basic numeral.

8 Express each of the following as a basic numeral.

a 5.7×10^4 **b** 3.6×10^6 **c** 4.3×10^8
d 3.21×10^7 **e** 4.23×10^5 **f** 9.04×10^{10}
g 1.97×10^8 **h** 7.09×10^2 **i** 6.357×10^5



Hint for Q8: Move the decimal point right to increase the place value of each digit.

Example 31 Writing basic numerals using negative powers

Express 4.71×10^{-6} as a basic numeral.

Solution

$4.71 \times 10^{-6} = 0.00000471$

Explanation

Move the decimal point six places to the left and insert zeros where necessary. The 4 decreases in place value by six places from the units position to the millionths position.

Now you try

Express 9.4×10^{-3} as a basic numeral.

9 Express each of the following as a basic numeral.

a 1.2×10^{-4} **b** 4.6×10^{-6} **c** 8×10^{-10}
d 3.52×10^{-5} **e** 3.678×10^{-1} **f** 1.23×10^{-7}
g 9×10^{-5} **h** 5×10^{-2} **i** 4×10^{-1}



Hint for Q9: Move the decimal point left to decrease the place value of each digit.

Problem-solving and reasoning

10–11(½) | 10–11(½), 12

10 Express each of the following approximate numbers using scientific notation.

a The mass of Earth is 6 000 000 000 000 000 000 000 000 kg.
b The diameter of Earth is 40 000 000 m.
c The diameter of a gold atom is 0.0000000001 m.
d The radius of Earth's orbit around the Sun is 150 000 000 km.
e The universal constant of gravitation is $0.0000000000667\ \text{Nm}^2/\text{kg}^2$.
f The half-life of polonium-214 is 0.00015 seconds.
g Uranium-238 has a half-life of 4 500 000 000 years.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

7F

11 Express each of the following numbers as a basic numeral.

a Neptune is approximately 4.6×10^9 km from Earth.


b A population of bacteria contained 8×10^{12} organisms.

c Earth is approximately 3.84×10^5 km from the Moon.

d A 50c coin is approximately 3.8×10^{-3} m thick.

e The diameter of the nucleus of an atom is approximately 1×10^{-14} m.

f The population of a city is 7.2×10^5.



12 Write the answers to each of these problems using scientific notation.

a Two planets are 2.8×10^8 km and 1.9×10^9 km from their closest sun. What is the difference between these two distances?

b Two particles weigh 2.43×10^{-2} g and 3.04×10^{-3} g. Find the difference in their weights.

Scientific notation with numbers larger than 10 — 13


13 The number 47×10^4 is not written using scientific notation, since 47 is not a number between 1 and 10. The following shows how to convert to scientific notation.

$$47 \times 10^4 = 4.7 \times 10 \times 10^4$$
$$= 4.7 \times 10^5$$

Write these numbers using scientific notation.

a 32×10^3 **b** 41×10^5 **c** 0.13×10^5 **d** 0.092×10^3

e 61×10^{-3} **f** 424×10^{-2} **g** 0.02×10^{-3} **h** 0.0004×10^{-2}



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

7G Scientific notation using significant figures

Learning intentions

- To know how significant figures are counted
- To be able to round to a given number of significant figures
- To be able to write a number using scientific notation correct to a given number of significant figures

Key vocabulary: scientific notation, significant figures

The number of digits used to record measurements depends on how accurately the measurements can be recorded. The volume of Earth, for example, has been calculated as $1\,083\,210\,000\,000$ km^3 using six significant figures and could be written using scientific notation as 1.08321×10^{12}. A more accurate calculation may include more non-zero digits in the last seven places.

The mass of a single oxygen molecule is known to be $0.000000000000000000000000053$ g. This shows two significant figures and is written using scientific notation as 5.3×10^{-26}.

On many calculators you will notice that very large or very small numbers are automatically converted to scientific notation using a certain number of significant figures. Numbers can also be entered into a calculator using scientific notation.



The Earth's volume is calculated using its radius, which varies slightly because the Earth is not a perfect sphere. Using six significant figures gives us a precise measurement without overestimating our accuracy.

Lesson starter: Significant discussions

Work in pairs and sort these numbers into groups according to how many significant figures are in each number.

23	502	0.0018
7208	46	0.0001073
0.000907	1609	91
6182	6	0.0000741

Key ideas

- **Significant figures** are the important digits of a number which indicate how accurate it is.
- Significant figures are counted from left to right starting at the first non-zero digit. For example:
  - 38 041 shows five significant figures
  - 6.034 shows four significant figures
  - 0.0016 shows two significant figures
  - 0.00160 shows three significant figures.
- Zeros at the end of a number are counted for decimals (e.g. 0.00160) but not necessarily for whole numbers (e.g. 38 041 000).
- When using scientific notation, the first significant figure sits to the left of the decimal point. For example:
  3.217×10^4 shows four significant figures.
- Calculators can be used to work with scientific notation.
  - E or EE or EXP or $\times 10^x$ are common key names on calculators.
  - Pressing 2.37 EE 5 gives 2.37×10^5.
  - 2.37E5 means 2.37×10^5.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Exercise 7G

Understanding

1–4 | 3, 4

1 **a** Round each of these numbers to the nearest hundred.
i 267 **ii** 32 740 **iii** 18 350

b Round each of these numbers to the nearest tenth.
i 0.063 **ii** 0.1902 **iii** 21.04

c Round each of these numbers to the nearest thousand.
i 267 540 **ii** 38 290 **iii** 4 060 990



Hint for Q1:
Rounding rules:
Locate the first digit to the right of the required digit.
- Round down (leave it as it is) for a 4 or less
- Round up (increase by 1) for a 5 or more

2 Which of these numbers have two significant figures?
62 100, 30 500, 42, 0.0071, 0.0805, 201 000



Hint for Q2: Count significant figures from the first non-zero digit.

3 Complete the tables, rounding each number to the given number of significant figures.



Hint for Q3: In 57 263, 6 is the fourth significant figure so round to the nearest 10 for four significant figures.

a 57 263

Significant figures	Rounded number
4	
3	57 300
2	
1	

b 0.0036612

Significant figures	Rounded number
4	
3	
2	
1	0.004



Hint for Q4: The number to the left of the decimal point must be from 1 to less than 10.

4 Are the following numbers written using scientific notation with three significant figures? (Yes or no)
a 4.21×10^4 **b** 32×10^{-3} **c** 1800×10^6
d 0.04×10^2 **e** 1.89×10^{-10} **f** 9.04×10^{-6}

Fluency

5–8(½) | 5–9(½)

Example 32 Stating the number of significant figures

State the number of significant figures given in these numbers.
a 401 **b** 0.005012 **c** 3.2×10^7

Solution	Explanation
a Three significant figures	All three digits are significant.
b Four significant figures	Start counting at the first non-zero digit (5).
c Two significant figures	With scientific notation the first significant figure is to the left of the decimal point.

Now you try

State the number of significant figures given in these numbers.
a 7105 **b** 0.00016 **c** 4.21×10^{-2}

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5 State the number of significant figures given in these numbers.

a 272	**b** 1007	**c** 30 101
d 19	**e** 0.0183	**f** 0.20
g 0.706	**h** 0.00109	**i** 4.21×10^3
j 2.905×10^{-2}	**k** 1.07×10^{-6}	**l** 5.90×10^5



Hint for Q5: Start counting from the first non-zero digit.

Example 33 Writing numbers using scientific notation and significant figures

Write these numbers using scientific notation and three significant figures.

a 2 183 000 **b** 0.0019482

Solution	Explanation
a $2\,183\,000 = 2.18 \times 10^6$	Put the decimal point after the first non-zero digit. The decimal point has moved six places so multiply by 10^6. Round the third significant figure down (i.e. leave it as is) since the following digit (3) is less than 5.
b $0.0019482 = 1.95 \times 10^{-3}$	Move the decimal point three places to the right and multiply by 10^{-3}. Round the third significant figure up to 5 since the following digit (8) is greater than 4.

Now you try

Write these numbers using scientific notation and three significant figures.

a 471900 **b** 0.00032475

6 Write these numbers using scientific notation and three significant figures.

a 242 300	**b** 171 325	**c** 2829	**d** 3 247 000
e 0.00034276	**f** 0.006859	**g** 0.01463	**h** 0.001031
i 23.41	**j** 326.042	**k** 19.618	**l** 0.172046

7 Write each number using scientific notation, rounding to the number of significant figures given in the brackets.

a 47 760 (3)	**b** 21 610 (2)	**c** 4 833 160 (4)
d 37.16 (2)	**e** 99.502 (3)	**f** 0.014427 (4)
g 0.00201 (1)	**h** 0.08516 (1)	**i** 0.0001010 (1)



Hint for Q7: First round the number to the required number of significant figures.

Example 34 Using a calculator with scientific notation

Use a calculator to evaluate $3.67 \times 10^5 \times 23.6 \times 10^4$. Answer using scientific notation correct to four significant figures.

Solution	Explanation
$3.67 \times 10^5 \times 23.6 \times 10^4$	Use a calculator with an appropriate key sequence.
$= 8.661 \times 10^{10}$	Write using scientific notation with four significant figures.
	Look for a $\boxed{\times 10^x}$ or $\boxed{\text{EXP}}$ button or similar.

Now you try

Use a calculator to evaluate $62.14 \times 10^2 \times 1.6 \times 10^4$. Answer using scientific notation correct to four significant figures.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

7G

8 Use a calculator to evaluate each of the following. Write your answers using scientific notation correct to four significant figures.

a 4^{-6}

b 78^{-3}

c $(-7.3 \times 10^{-4})^{-5}$

d $\dfrac{3.185}{7 \times 10^4}$

e $2.13 \times 10^4 \times 9 \times 10^7$

f $5.671 \times 10^2 \times 3.518 \times 10^5$

g $9.419 \times 10^5 \times 4.08 \times 10^{-4}$

h $2.85 \times 10^{-9} \times 6.33 \times 10^{-3}$

i $12\,345^2$

j 87.14^8

k $\dfrac{1.83 \times 10^{26}}{4.5 \times 10^{22}}$

l $\dfrac{-4.7 \times 10^{-2} \times 6.18 \times 10^7}{3.2 \times 10^6}$

Example 35 Using a calculator with scientific notation and negative indices

Use a calculator to evaluate $7.6 \times 10^{-3} + \sqrt{2.4 \times 10^{-2}}$. Write your answer using scientific notation correct to four significant figures.

Solution

$7.6 \times 10^{-3} + \sqrt{2.4 \times 10^{-2}}$
$= 0.1625$ (to 4 sig. figs)
$= 1.625 \times 10^{-1}$

Explanation

Use a calculator with an appropriate key sequence.

Write using scientific notation with a number between 1 and 10.

Now you try

Use a calculator to evaluate $\dfrac{\sqrt{5.1 \times 10^{-7}}}{4.32 \times 10^{-2}}$. Write your answer using scientific notation correct to four significant figures.

9 Use a calculator to evaluate each of the following. Write answers using scientific notation correct to five significant figures.

a $\sqrt{8756}$

b $\sqrt{634 \times 7.56 \times 10^7}$

c $8.6 \times 10^5 + \sqrt{2.8 \times 10^{-2}}$

d $-8.9 \times 10^{-4} + \sqrt{7.6 \times 10^{-3}}$

e $\dfrac{5.12 \times 10^{21} - 5.23 \times 10^{20}}{2 \times 10^6}$

f $\dfrac{8.942 \times 10^{47} - 6.713 \times 10^{44}}{2.5 \times 10^{19}}$

g $\dfrac{2 \times 10^7 + 3 \times 10^8}{5}$


h $\dfrac{4 \times 10^8 + 7 \times 10^9}{6}$

i $\dfrac{6.8 \times 10^{-8} + 7.5 \times 10^{27}}{4.1 \times 10^{27}}$

j $\dfrac{2.84 \times 10^{-6} - 2.71 \times 10^{-9}}{5.14 \times 10^{-6} + 7 \times 10^{-8}}$



Hint for Q9: For fraction calculations, insert brackets so that the numerator and denominator are calculated before division.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Problem-solving and reasoning

10, 11 | 10, 12, 13

10 The mass of Earth is approximately $6\,000\,000\,000\,000\,000\,000\,000$ kg. The mass of the Sun is $330\,000$ times the mass of Earth. Find the mass of the Sun. Express your answer using scientific notation correct to three significant figures.

11 The diameter of Earth is approximately $12\,756\,000$ m. If the Sun's diameter is 109 times that of Earth, compute its diameter in kilometres. Express your answer using scientific notation correct to three significant figures.



12 Write these numbers from largest to smallest.
2.41×10^6, 24.2×10^5, 0.239×10^7, 2421×10^3, 0.02×10^8

Hint for Q12: First write each number using scientific notation.

13 The following output is common on a number of different calculators and computers. Write down the number that you think they represent.

a 4.26E6 **b** 9.1E−3 **c** 5.04EE11
d 1.931EE−1 **e** 2.1 ∧ 06 **f** 6.14 ∧ −11

Combining bacteria

— | 14

14 A flask of type A bacteria contains 5.4×10^{12} cells and a flask of type B bacteria contains 4.6×10^8 cells. The two types of bacteria are combined in the same flask.




Hint for Q14: Set up a table to show the number of each type of bacteria after every 8 hours.

a How many bacterial cells are there now in the flask?

b If type A bacterial cells double every 8 hours, and type B bacterial cells triple every 8 hours, how many cells are in the flask after:

i 8 hours? **ii** one day?

Express your answers to part **b** using scientific notation correct to three significant figures.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Maths@Work

Lab technician

Lab technicians work in laboratories supporting scientists by helping to carry out tests for research and to report their findings. Research can be done in medical, clinical and forensic labs as well as pharmaceutical and brand development labs.

Lab technicians require competent practical and technical skills as well as good communication skills, both verbal and written. An enquiring and analytical mind also helps. All these qualities are important in modern labs.

Mathematical skills are vital in this role. Determining growth rates of cultures can form part of a lab technician's work. Working in and with scientific notation is also required.



1 Various types of cells each have different volumes and are measured in cubic micrometres (μm^3). Write the following volumes in μm^3 using scientific notation.

- **a** Red blood cell $100\ \mu m^3$
- **b** Ear hair cell $4000\ \mu m^3$
- **c** Lymphocyte $130\ \mu m^3$
- **d** Beta cell $1000\ \mu m^3$
- **e** Fat cell $600\,000\ \mu m^3$

2 The diameter of a human egg is $120\ \mu m$. Recall that $1\ \mu m = 1 \times 10^{-6}$ m.

- **a** Write this diameter in metres as a basic numeral.
- **b** Write this diameter in metres using scientific notation.

3 The mass of one red blood cell is 1 picogram or 1×10^{-12} grams. Calculate the mass of these blood cell counts using grams and write the result using scientific notation.

- **a** 2000 red blood cells
- **b** $1\,000\,000$ red blood cells
- **c** 3×10^{12} red blood cells

4 A human being is said to be made up of $(3.7 \pm 0.8) \times 10^{13}$ cells.

- **a** What is the upper limit given here as a basic numeral?
- **b** What is the lower limit as a basic numeral?

5 A half-life is the time taken for a quantity to halve in size. Write each of the following half-lives using scientific notation.

- **a** Carbon-14 has a half-life of 5700 years. (It is used by archaeologists to date historic objects.)
- **b** Uranium-238 has a half-life of $4\,500\,000\,000$ years.
- **c** Iodine-129 has a half-life of 15.7 million years.
- **d** Polonium-214 has a half-life of 150 microseconds (state the answer in seconds).

Hint for Q5:
1 microsecond = 1 millionth of a second.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

6 Convert the half-lives in Question **5a**, **b** and **c** into days using 365.25 days per year. State answers using scientific notation to three significant figures.

7 Yeast doubles every 90–120 minutes. Taking the median value, how long would it take for a food technician to grow 1 g of yeast to $\frac{1}{2}$ kg of yeast?

8 *E. coli* bacteria can double every 20–30 minutes. Using the median time, how many grams of *E. coli* has grown from 1 mg using the following times? Write each answer in scientific notation to three significant figures.

a 50 minutes **b** $8\frac{1}{3}$ hours

9 After each 'half-life' period, 1 g of radioactive isotope halves in quantity. List the 10 fractions of radioactive isotope remaining (in grams) after each of 10 half-lives. First list these fractions with positive powers of 2 and then with negative powers of 2.

Using technology

10 Radioactive isotopes have various medical applications.

a Set up the following spreadsheet to calculate quantities remaining after multiples of half-lives for the radioactive isotopes listed. Enter formulas in the shaded cells and format cells as 'number' to four decimal places.

	A	B	C	D	E	F	G
1	**Calculation of time for multiple half-lives**						
2				**Quantity remaining in mg**			
3	Application	Radioactive isotope	**Initial quantity in mg**	After 1 half-life	After 2 half-lives	After 4 half-lives	After 10 half-lives
4	Biochemical tracer	hydrogen-3	8				
5	Biomedical imaging	carbon-11	64.128				
6	Heart system tracer	sodium-24	24.73				
7	Red blood cell lifetime tracer	iron-59	7.86				
8	Radiation therapy for cancer	radium-226	4.25				

b To help determine the correct dose, medical scientists need to calculate the length of time that a radioactive isotope lasts. Set up the following spreadsheet to calculate the total times for multiple half-lives. Enter formulas in the shaded cells and format cells as number to four decimal places.

	A	B	C	D	E	F	G	H
1	**Calculation of time for multiple half-lives**							
2			**Half-life**		**Total time**			
3	Application	Radioactive isotope	Value	Time unit	For 2 half-lives	For 4 half-lives	For 6 half-lives	For 10 half-lives
4	Biochemical tracer	hydrogen-3	12.32	hours				
5	Biomedical imaging	carbon-11	20.3	minutes				
6	Heart system tracer	sodium-24	14.951	hours				
7	Red blood cell lifetime tracer	iron-59	44.495	days				
8	Radiation therapy for cancer	radium-226	1600	years				

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Modelling

Rabbits and hares

The population of rabbits and hares in a particular part of the countryside are booming because fox numbers are low.

Present a report for the following tasks and ensure that you show clear mathematical workings, explanations and diagrams where appropriate.

1 Preliminary task

a A population of a certain colony of hares is modelled by the rule $P = 3^t$ where P is the population and t is in months. Find:
- **i** the population after 6 months
- **ii** the number of whole months for the population to reach at least 1000.

b A population of a certain colony of rabbits is modelled by the rule $P = 200 \times 2^t$ where P is the population and t is in months. Find:
- **i** the population after 6 months
- **ii** the number of whole months for the population to reach at least 20 000.

2 Modelling task

Formulate

The problem is to determine the number of months required for the hare population to overtake the population of rabbits in a particular colony with the given data.

Time in months	0	1	2	3			
Month	**Dec**	**Jan**	**Feb**	**Mar**	**Apr**	**May**	**June**
Rabbit population	500	1000	2000	4000			
Hare population	1	4	16	64			

a Write down all the relevant information that will help solve this problem.

b Copy and extend the given table showing the population of rabbits and hares in the first 6 months.

c By what factor does the rabbit population increase each month?

d The rule for the population of rabbits is of the form $P = 500 \times ?^t$. State the missing number and re-write the rule for rabbits.

e By what factor does the hare population increase each month?

f The rule for the population of hares is of the form $P = ?^t$. State the missing number and rewrite the rule for hares.

Solve

g Use the rules outlined in parts **d** and **f** to confirm the population of:
- **i** rabbits initially (at $t = 0$)
- **ii** rabbits after 6 months.
- **iii** hares after 6 months.

h Use the rules outlined in parts **d** and **f** to estimate the population of:
- **i** rabbits after 12 months
- **ii** hares after 12 months

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

i Extend your table and use your rules to calculate the population numbers for the months from August to December.

Time in months	7					
Month	July	Aug	Sep	Oct	Nov	Dec
Rabbit population						
Hare population						

Evaluate and verify

j State the first month in which the hare population overtakes the population of rabbits.

k In which month is the hare population more than twice the rabbit population?

l In which month does the rabbit population exceed 1 million?

Communicate

m Summarise your results and describe any key findings.

3 Extension question

A new population of 100 rabbits breed in a different part of the country. The population is modelled by the rule $P = 100 \times 4^t$ where t is in years from the beginning of 2021.

a What would you expect the population of rabbits be at the beginning of 2025?

b During what year will this rabbit population exceed:

i 1 million

ii 25 million?



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Technology and computational thinking

Constructing surds


Key technology: Dynamic geometry

We know that a surd is an irrational number meaning that it cannot be expressed exactly as a fraction. So to place a surd on a number line for example may be more challenging than placing a number like $\frac{3}{2}$, which is exactly half way between 1 and 2. However, the precise positions of surds on a number line can be located using geometric construction.



1 Getting started

We will start by considering the length of the diagonal of a square of side length one unit.



a Use Pythagoras' theorem to find the length of the diagonal of the given unit square. Give your answer as a surd and also as a decimal rounded to four decimal places using a calculator.

Now consider a rectangle with side lengths 1 and 2.

b Draw a diagram of the rectangle showing one of the diagonals.

c Use Pythagoras' theorem to find the surd which corresponds to the length of the diagonal of your rectangle.

2 Using technology

Open a dynamic geometry program like Desmos and complete the following to place the surd $\sqrt{2}$ on a number line.


a Show a grid using the tools menu, then place a line horizontally through two points O and A.

b Add the line segment OB which is the diagonal of a unit square.

c Construct a circle with centre O and passing through B.

d Construct the line segment OC.


e Label the segment OC with its length. You have just placed $\sqrt{2}$ on a number line at point C.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3 Applying an algorithm

In part **1b** you drew a diagonal of a rectangle with length $\sqrt{5}$.



a Use dynamic geometry to place $\sqrt{5}$ on a number line. Your construction will be similar to that of part **2 Using technology** but instead use two unit squares forming a rectangle of side lengths 1 and 2.

b Write down an algorithm (series of instructions) similar to those in part **2** which places the surd $\sqrt{5}$ on a number line.

c Note that when using dynamic geometry you can simply drag point B to a new position to construct a different surd. Try doing this as shown in this diagram.

d What size rectangle would you use to construct the surd $\sqrt{10}$?

e Use dynamic geometry to construct the following surds.

i $\sqrt{10}$

ii $\sqrt{8}$

iii $\sqrt{29}$

iv $\sqrt{34}$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Puzzles and games

1 A population of bacteria doubles every 5 minutes. What is this type of growth called? Solve this puzzle to find the answer.
Write the basic numeral for each of the following. Write the letter corresponding with each answer in the boxes below to form a word.

T $\frac{1}{10^3}$	**L** 2.15×10^3	**I** $\frac{9^5}{9^3}$
A 5×10^{-2}	**O** 2^{-2}	**N** 2^4
P $3^2 \times 2^2 - 3^0$	**X** $5^2 + 2^4 - 3^2 \times 4$	**E** $\frac{4^3}{8}$

☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐
8	5	35	0.25	16	8	16	0.001	81	0.05	2150

2 Find the answer to these calculations without using a calculator:

a $2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 5 \times 5 \times 5 \times 5 \times 5 \times 5 \times 5 \times 5$
b $5 \times 5 \times 5 \times 5 \times 5 \times 4 \times 4 \times 4 \times 4 \times 4 \times 5 \times 5 \times 5 \times 5 \times 5$

3 Show that this expression is equal to 1.
$\frac{1}{2} \times \frac{a^2b}{3am} \times \frac{6a^3}{b^2} \times \frac{10(ab)^2}{2bm^3} \times \frac{m^4}{5a^6}$

4 Insert brackets in this expression to make the given answer correct.
$4a \times 3m^3 \times \left(\frac{a}{6m}\right)^2 = 3a^3m$

5 Nick and Maddy have a big bag of 10-cent coins. They decide to make 30 piles of coins, starting with one coin in the first pile, then 2 coins in the second pile, 4 coins in the third pile, 8 in the fourth pile and so on, continuing to double the number of coins for each new pile. Given that each 10-cent coin is 2 mm thick, find the heights of the 1st, 5th, 10th, 15th, 20th, 25th and 30th piles and the value in dollars of each of these piles.

6 Each day that a bushfire burns in a National Park, the total area burnt is twice as big an area as the day before. If a National Park was totally burnt out in 15 days, on which day was the park only half burnt out?


7 It is thought that the game of chess was invented by an ancient Indian mathematician. The King was so pleased with the game that he offered the inventor any reward of his choice: rare jewels, bags of gold or even a large property.

To the King's surprise the Indian mathematician asked for some wheat! He asked for 1 grain for the first square of the chess board, 2 grains for the second square, 4 grains for the third square, 8 grains for the fourth square etc. continuing this way right up to the 64th square.

a If one grain of wheat weighs 2×10^{-8} tonnes, what weight of wheat would the inventor have received for the 64th square? Answer using scientific notation with three significant figures.
b How much money would this wheat be worth at the Australian price of $275/tonne?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter summary



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter checklist

Chapter checklist

A version of this checklist that you can print out and complete can be downloaded from your Interactive Textbook.

		✔
7A	**1 I can write expressions in expanded form and simplify.** e.g. Write the following in expanded form and simplify. **a** $(2ab)^3$ **b** $\left(\frac{2}{7}\right)^2$	
7A	**2 I can write numbers and expressions using index form.** e.g. Write the following in index form: **a** $\frac{7}{9}\times\frac{7}{9}\times\frac{7}{9}\times\frac{7}{9}$ **b** $2\times a\times a\times b\times a\times b$	
7A	**3 I can express a number as a product of its prime factors.** e.g. Express 92 as a product of its prime factors.	
7A	**4 I can evaluate expressions involving indices using substitution.** e.g. If $a=2$, $b=-3$ and $c=11$, evaluate the following. **a** $(ab)^3$ **b** $\left(\frac{b}{c}\right)^2$	
7B	**5 I can simplify expressions with numerical bases using index laws for multiplication and division.** e.g. Simplify, giving your answer in index form. **a** $4^3\times4^4$ **b** $7^5\div7$	
7B	**6 I can use the index law for multiplication.** e.g. Simplify the following using the index law for multiplication: **a** $a^3\times a^7$ **b** $9x^2\times3x^3$	
7B	**7 I can use the index law for division.** e.g. Simplify $y^6\div y^2$ using the index law for division.	
7B	**8 I can combine index laws for multiplication and division.** e.g. Simplify: $\frac{3a^2b\times4ab^3}{8a^2b^3}$.	
7C	**9 I can simplify expressions containing the zero index.** e.g. Evaluate using the zero index rule: 3^0+3a^0	
7C	**10 I can use the index law for power of a power.** e.g. Simplify using the index law for power of a power: $5(x^3)^6$	
7C	**11 I can combine index laws with the zero index rule.** e.g. Simplify the following **a** $(x^2)^3\times(x^4)^2\div x^{14}$ **b** $\frac{7a^3\times2a^4}{4a^5}$	
7D	**12 I can use index laws for brackets and fractions to simplify expressions.** e.g. Simplify, using index laws. **a** $(2a^4)^3$ **b** $\left(\frac{-2b^2}{a^3}\right)^4$	

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter checklist

		✔
7D	**13 I can combine index laws to simplify expressions.** e.g. Simplify the following. **a** $x(-xy^2)^3$ **b** $\left(\frac{a^2b^3}{2}\right)^2 \times \frac{4}{(ab)^2}$	
7E	**14 I can express negative indices in positive index form.** e.g. Rewrite the following with positive indices only: **a** $4x^{-2}$ **b** $\frac{5}{2^{-3}}$	
7E	**15 I can evaluate expressions involving negative indices.** e.g. Write the following with a positive index and then as a fraction. **a** 5×2^{-3} **b** 4×10^{-2}	
7F	**16 I can convert from scientific notation to a basic numeral.** e.g. Write these numbers as a basic numeral: **a** 4.9×10^3 **b** 3.01×10^{-6}	
7F	**17 I can write numbers using scientific notation.** e.g. Write these numbers using scientific notation: **a** 27 000 **b** 0.0000375	
7G	**18 I can write numbers using scientific notation and rounding to a given number of significant figures.** e.g. Write these numbers in scientific notation using three significant figures: **a** 9 143 000 **b** 0.00032	
7G	**19 I can use a calculator to evaluate expressions involving numbers expressed with scientific notation.** e.g. Use a calculator to evaluate $\frac{\sqrt{5.3 \times 10^{-3}}}{8.32 \times 10^{-2}}$ and express your answer using four significant figures.	

Essential Mathematics Core for the Victorian Curriculum Year 9 ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter review

Short-answer questions

7A **1** Express each of the following in index form.

a $3 \times 3 \times 3 \times 3$ **b** $2 \times x \times x \times x \times y \times y$

c $3 \times a \times a \times a \times \frac{b}{a} \times b$ **d** $\frac{3}{5} \times \frac{3}{5} \times \frac{3}{5} \times \frac{1}{7} \times \frac{1}{7}$

7A **2** Write the following as a product of prime factors in index form.

a 45 **b** 300

7B **3** Simplify using the index laws for multiplication and division.

a $x^3 \times x^7$ **b** $2a^3b \times 6a^2b^5c$ **c** $3m^2n \times 8m^5n^3$

d $a^{12} \div a^3$ **e** $x^5y^3 \div (x^2y)$ **f** $\frac{5a^8b^3}{10a^6b}$

7C/D **4** Simplify:

a $(m^2)^3$ **b** $(3a^4)^2$ **c** $(-2a^2b)^5$ **d** $3a^0b$ **e** $2(3m)^0$ **f** $\left(\frac{a^2}{3}\right)^3$

7E **5** Express each of the following with positive indices.

a 2^{-3} **b** 3×4^{-5} **c** 6×10^{-3} **d** $\frac{1}{4^{-2}}$ **e** $\frac{6}{5^{-3}}$ **f** $\frac{2}{10^{-5}}$

7B/C/D **6** Use index laws to simplify each of the following.

a $\frac{k^5 \times k^3}{k^2}$ **b** $\frac{9a^4}{2m^3} \times \frac{4m^4}{3a^2}$ **c** $(a^2)^3 \times 4a^4$

d $(h^{12})^3 \div (h^4)^5$ **e** $\frac{4(a^3)^2(b^4)^3}{20a^0(b^2)^5}$ **f** $5(3p^2q)^2$

7F **7** Write each of the following numbers as a basic numeral to arrange in ascending order.
$2.35, 0.007 \times 10^2, 0.0012, 3.22 \times 10^{-1}, 0.4, 35.4 \times 10^{-3}$.

7F **8** Write each of the following numbers as a basic numeral.

a 3.24×10^2 **b** 1.725×10^5 **c** 2.753×10^{-1} **d** 1.49×10^{-3}

7G **9** Write each of the following values using scientific notation correct to three significant figures.

a The population of Australia in 2010 was approximately 22 475 056.

b The area of the USA is 9 629 091 km^2.

c The time taken for light to travel 1 metre (in a vacuum) is 0.00000000333564 seconds.

d The wavelength of ultraviolet light is 0.000000294 m.

7F **10** Write each of the following values using scientific notation in the units given in brackets. Note: 1 tonne = 1000 kg.

a 25 years (hours) **b** 3 million years (months)

c 430 tonnes (kg) **d** 5 tonnes (grams)

Multiple-choice questions

7B **1** $3x^7 \times 4x^4$ is equivalent to:

A $12x^7$ **B** $12x^{28}$ **C** $7x^{11}$ **D** $12x^{11}$ **E** $7x^3$

7C **2** $3(2y^2)^0$ simplifies to:

A 6 **B** 3 **C** $6y^2$ **D** $3y$ **E** 12

7D **3** $(2x^2)^3$ expands to:

A $2x^5$ **B** $2x^6$ **C** $6x^6$ **D** $8x^5$ **E** $8x^6$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

7B **4** $\frac{x^6y^2}{4x^2y}$ simplifies to:

A $-4x^4y$ **B** $\frac{x^3y^2}{4}$ **C** $\frac{x^4y}{4}$ **D** $\frac{x^4y^2}{4}$ **E** $\frac{x^3y}{4}$

7D **5** $\left(\frac{2a}{k}\right)^3$ is equal to:

A $\frac{2a^3}{k}$ **B** $\frac{2a^3}{k^3}$ **C** $\frac{6a^3}{k^3}$ **D** $\frac{8a^3}{k^3}$ **E** $\frac{6a}{3k}$

7D **6** $\left(\frac{5^2}{7^3}\right)^4$ is equal to:

A $\frac{4 \times 5^2}{4 \times 7^3}$ **B** $\frac{5^8}{7^{12}}$ **C** $\frac{5^6}{7^3}$ **D** $\frac{5^8}{7^3}$ **E** $\frac{5^6}{7^7}$

7D **7** $5(2am^3)^3$ is equal to:

A $30a^3m^9$ **B** $10a^3m^9$ **C** $40a^3m^6$ **D** $40a^3m^9$ **E** $10am^6$

7E **8** 4×7^{-3} expressed with positive indices is:

A 28^3 **B** 4^37^3 **C** $\frac{4}{7^3}$ **D** $\frac{-4}{7^3}$ **E** $\frac{1}{4 \times 7^3}$

7F **9** The weight of a cargo crate is 2.32×10^4 kg. In expanded form this weight in kilograms is:

A 2 320 000 **B** 232 **C** 23 200 **D** 0.000232 **E** 2320

7G **10** 0.00032761 using scientific notation rounded to three significant figures is:

A 328×10^{-5} **B** 3.27×10^{-4} **C** 3.28×10^4 **D** 3.30×10^4 **E** 3.28×10^{-4}

Extended-response questions

1 Use a calculator to evaluate the following, giving your answer using scientific notation correct to two significant figures.

a $m_s \times m_e$ where m_s (mass of Sun) $= 1.989 \times 10^{30}$ kg and m_e (mass of Earth) $= 5.98 \times 10^{24}$ kg.

b The speed, v, in m/s of an object of mass $m = 2 \times 10^{-3}$kg and kinetic energy $E = 1.88 \times 10^{-12}$ joules where $v = \sqrt{\frac{2E}{m}}$.

2 Use the table of Solar System data below to answer these questions. Express each answer using scientific notation rounded to three significant figures.

Approximate data for some planets in the Solar System			
Planet	**Distance from the Sun in millions of km**	**Length (in Earth days) of one revolution around the Sun**	**Mass compared to Earth**
Mercury	57.9	88.0	0.553
Venus	108.2	224.7	0.815
Earth	149.6	365.25	1.00
Mars	227.9	687.0	0.1074
Jupiter	778.3	4331	317 896
Saturn	1427	10 760	95 185

a Find the distance from the Sun, in km, for:

i Mercury **ii** Earth **iii** Jupiter **iv** Saturn

b How many Earth years will have passed for these planets to complete one full revolution around the Sun?


i Mercury **ii** Mars **iii** Saturn

c The mass of Earth is approximately 5.98×10^{24} kg. Determine the mass of:

i Venus **ii** Mars **iii** Jupiter

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

8 Geometry



Essential mathematics: why geometry skills are important

Geometry is a vital skill in many fields needed to solve problems involving shapes, sizes, and spatial properties. Understanding geometric concepts helps in numerous real-life situations.

Carpenters use parallel lines and angles to ensure that furniture and structures are level and properly aligned. They also rely on geometric shapes like rectangles and triangles to create strong and stable designs, ensuring that every piece fits together perfectly.

Artists and graphic designers use congruent triangles and similar figures to create depth and perspective in their work. They apply geometric concepts to construct realistic scenes and maintain visual harmony.

Robotics engineers apply geometric principles to calculate the precise angles and distances required for a robotic arm to perform complex tasks in industries such as manufacturing, assembly, and medical surgeries with high accuracy and efficiency. They use concepts like coordinate systems and transformations to program the arm's movements

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.



In this chapter

8A Angles and triangles (Consolidating)
8B Parallel lines (Consolidating)
8C Quadrilaterals (Consolidating)
8D Polygons ★
8E Congruent triangles
8F Enlargement and similar figures
8G Similar triangles
8H Applying similar triangles

Victorian Curriculum 2.0

This chapter covers the following content descriptors in the Victorian Curriculum 2.0:

MEASUREMENT

VC2M9M03

SPACE

VC2M9SP02, VC2M9SP03

Please refer to the curriculum support documentation in the teacher resources for a full and comprehensive mapping of this chapter to the related curriculum content descriptors.

© VCAA

Online resources


A host of additional online resources are included as part of your Interactive Textbook, including HOTmaths content, video demonstrations of all worked examples, auto-marked quizzes and much more.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


Warm-up quiz

1 Name the following types of angles. Choose from *acute*, *right*, *obtuse*, *straight*, *reflex* or *revolution*.


a



b




c




d



e




f




2 Find the value of a.


a



b




c




3 Identify the following triangles as *equilateral*, *isosceles* or *scalene*.


a



b




c




4 Identify the following as *right*, *acute* or *obtuse* triangles.


a



b



c




5 Find any missing angles in these triangles.


a



b



c




6 Of the six special quadrilaterals (square, rectangle, parallelogram, rhombus, kite and trapezium), which ones match the following descriptions?

a Always two pairs of equal length sides **b** Always two pairs of parallel sides


c Always equal length diagonals

7 Find the value of the pronumerals in each of the following.


a



b




c




8 Name the following shapes.


a



b



c



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

8A Angles and triangles CONSOLIDATING

Learning intentions

- To review the various types of angles and triangles
- To know the special relationships between angles formed at a point
- To be able to find missing angles in simple geometric diagrams including angles at a point and triangles
- To be able to use the exterior angle theorem for a triangle

Key vocabulary: line, line segment, angle, complementary, supplementary, revolution, vertically opposite, acute, right, obtuse, reflex, straight, angle sum, scalene, isosceles, equilateral, exterior angle, triangle


When looking at a building or structure, we see angles formed where two lines meet. We also see triangles being used for their strength and rigidity. Angles associated with triangles and with lines that meet at a point will be revised in this section.



Steel towers use triangles for their strength and rigidity.

Lesson starter: Exterior angle discovery


Here is a triangle with one side extended to form the exterior angle $\angle BCD$ with size $c°$.



1. If $a = 50$ and $b = 85$, find $\angle ACB$. Then find $\angle BCD$. What do you notice?
2. Repeat for $a = 60$ and $b = 95$. What do you notice?
3. What is the relationship between a, b and c? This is called the exterior angle theorem.


Key ideas

- When two lines or line segments meet at a point, an **angle** is formed.
  - This angle (shown here) is named $\angle A$ or $\angle BAC$ or $\angle CAB$
  - The size of the angle is $a°$.
  - A common tool used for measuring an angle is a protractor.




- Angle types
  - **Acute**: between 0° and 90°
  - **Obtuse**: between 90° and 180°
  - **Reflex**: between 180° and 360°
  - **Right**: 90°
  - **Straight**: 180°
  - **Revolution**: 360°

- Angles at a point
  - **Complementary** (add up to 90°)




$a + 60 = 90$

  - **Supplementary** (add up to 180°)




$a + 41 = 180$

  - **Revolution** (add up to 360°)



$a + 90 + 115 = 360$






  - **Vertically opposite** (are equal)




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

8A

- Types of **triangles**


Acute-angled (all angles < 90°)	Right-angled (includes a 90° angle)	Obtuse-angled (one angle > 90°)
 **Scalene** (all sides and angles are different sizes)	 Scalene	 Scalene
 **Equilateral** (all angles 60° and all sides equal)  **Isosceles** (two angles equal and two sides equal)	 Isosceles	 Isosceles

  - The sum of the angles in a triangle (**angle sum**) is 180°.



$a + b + c = 180$

- An **exterior angle** is formed by extending one side of a shape.
  - Exterior angle theorem of a triangle: The exterior angle of a triangle is equal to the sum of the two opposite interior angles.



$c = a + b$

Exercise 8A

Understanding

1–3 | 2, 3

1 Choose a word or number to complete each sentence.

a A 90° angle is called a _____ angle.
b A _____ angle is called a straight angle.
c A 360° angle is called a _____.
d _____ angles are between 90° and 180°.
e _____ angles are between 0° and 90°.
f Reflex angles are between _____ and 360°.
g Complementary angles add up to _____.
h _____ angles add up to 180°.
i The three angles in a triangle add up to _____.
j Vertically opposite angles are _____ .


2 What type of triangle has:

a a pair of equal length sides?
b one obtuse angle?
c all angles 60°?
d one pair of equal angles?
e all angles acute?
f all sides of different length?
g one right angle?
h two 45° angles?


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3 Estimate the size of each of the following angles and use your protractor to determine an accurate measurement.


a




b




c




d



e



f



Fluency

4(½), 5, 6(½) | 4–6(½)

Example 1 Finding supplementary and complementary angles

For the angle 47°, determine the:

a supplementary angle **b** complementary angle

Solution

a $180° - 47° = 133°$

b $90° - 47° = 43°$

Explanation

Supplementary angles add up to 180°.

Complementary angles add up to 90°.

Now you try

For the angle 74°, determine the:

a supplementary angle **b** complementary angle

4 For each of the following angles determine:

i the supplementary angle

ii the complementary angle

a 55° **b** 31° **c** 74° **d** 10° **e** 89°


f 22° **g** 38° **h** 65° **i** 47° **j** 77°




Hint for Q4:
- Supplementary angles add to 180°.
- Complementary angles add to 90°.

5 Find the value of a in these diagrams.


a



b




c






Hint for Q5:
- A straight angle is 180°.
- A revolution is 360°.


d



e



f





ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


8A

Example 2 Finding unknown angles in triangles

Name the types of triangles shown here. Determine the values of the pronumerals.

a 

b 

c 

Solution

a Equilateral triangle
$x = 60$

b Obtuse isosceles triangle
$2r + 120 = 180$
$2r = 60$
$r = 30$

c Acute scalene triangle
$s + 50 + 60 = 180$
$s + 110 = 180$
$s = 70$

Explanation


All sides are equal, therefore all angles are equal.


One angle is more than 90° and two sides are equal.
Angles in a triangle add to 180° and since it is isosceles the two base angles are equal.
Subtract 120 from both sides and then divide both sides by 2.


All angles are less than 90° and all sides are of different length.
Angles in a triangle add to 180°. Simplify and solve for s.

Now you try


Name the types of triangles shown here. Determine the values of the pronumerals.


a 

b 

c 


6 Name the types of triangles shown here. Determine the values of the pronumerals.


a 


b 




Hint for Q6: The angle sum of a triangle is 180°.

c 


d 

e 


f 

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


g




h




i




j



k



l



Problem-solving and reasoning

7, 8 | 7–8(½), 9

7 For each diagram:


- **i** name the angle shown (e.g. $\angle ABC$)
- **ii** state the type of angle given
- **iii** estimate the size of the angle
- **iv** measure the angle using a protractor.




Hint for Q7: When naming an angle, put the letter at the vertex in the middle.




a




b




c



d



e



f





ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

8A

Example 3 Finding angles outside a triangle

Find the value of each pronumeral. Give reasons for your answers.


a 

b 

Solution


a $x + 47 + 47 = 180$ (angle sum)
$x + 94 = 180$
$x = 86$

Explanation



Note the isosceles triangle and vertically opposite angles and mark in those angles. Angles in a triangle add to 180° and vertically opposite angles are equal. Simplify and solve for x.

b Let a be the unknown angle.
$a + 90 + 55 = 180$ (angle sum)
$a = 35$
$y + 35 = 180$
$y = 145$




Angles in a straight line are supplementary (add up to 180°).


Alternative method: $y = 90 + 55$
$= 145$

Alternatively, use the exterior angle theorem which says that the exterior angle is equal to the sum of the two opposite interior angles.

Now you try


Find the value of each pronumeral. Give reasons for your answers.


a 

b 



8 Find the value of each pronumeral.


a 

b 


Hint for Q8: If you can't find the value of the pronumeral straight away, try finding one of the other unknown angles.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


c




d




e




f




g



h



9 Explain why $\angle OAB$ is 32° in this circle if O marks the centre of the circle.



Analogue geometry — 10, 11

10 Calculate how many degrees the minute hand of a clock rotates in:

a 1 hour **b** $\frac{1}{4}$ of an hour **c** 10 minutes **d** 15 minutes

e 72 minutes **f** 1 minute **g** 2 hours **h** 1 day

11 Find the non-reflex angle between the hour and minute hands at these times. Remember to consider how the hour hand moves between each whole number. For example, at 9.30, the hour hand is halfway between the 9 and 10.

a 3 p.m. **b** 5 a.m.
c 6:30 p.m. **d** 11:30 p.m.
e 3:45 a.m. **f** 1:20 a.m.
g 4:55 a.m. **h** 2:42 a.m.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

8B Parallel lines CONSOLIDATING

Learning intentions

- To know the special pairs of angles formed when a transversal cuts two other lines
- To know the relationship between pairs of angles formed when a pair of parallel lines is cut by a transversal
- To be able to calculate unknown angles associated with parallel lines


Key vocabulary: parallel, transversal, corresponding, alternate, cointerior, vertically opposite, supplementary

When two lines are crossed by a third line, called a transversal, many special pairs of angles are formed. If the two original lines are parallel, then the pairs of angles are either equal or they add to 180°.


Lesson starter: Are they parallel?

Here are three diagrams that show a transversal crossing two other lines.


1



2









3



Decide whether each diagram contains a pair of parallel lines. Give reasons for your answer.

Key ideas

- Parallel lines point in the same direction.
  - Arrows indicate that lines are parallel.
- A **transversal** is a line crossing two or more other lines.

Special pairs of angles	Non-parallel lines	Parallel lines
Corresponding angles • If lines are parallel, corresponding angles are equal.		
Alternate angles • If lines are parallel, alternate angles are equal.		
Cointerior angles • If lines are parallel, cointerior angles are supplementary (add up to 180°).		 $a + b = 180$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


Exercise 8B

Understanding


1, 2 | 2

1 Decide whether the pair of marked angles are corresponding, alternate or cointerior.


a




b




c




d



e



f



2 Use the word *equal* or *supplementary* to complete these sentences.

a If two lines are parallel, corresponding angles are _____.

b If two lines are parallel, alternate angles are _____.

c If two lines are parallel, cointerior angles are _____.



Hint for Q2: Supplementary angles add to 180°.


Fluency

3–4(½), 5 | 3–5(½)


Example 4 Deciding whether lines are parallel

Decide whether each diagram contains a pair of parallel lines. Give a reason.


a



b



c



Solution

a No. The two cointerior angles do not add to 180°.

b No. The two corresponding angles are not equal.

c Yes. The two alternate angles are equal.

Explanation

$60° + 114° = 174° \neq 180°$


$79° \neq 82°$

If alternate angles are equal then the lines are parallel.


Now you try

Decide whether each diagram contains a pair of parallel lines. Give a reason.


a



b



c




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


8B

3 Decide if each diagram contains a pair of parallel lines. Give a reason.


a




b




c




d




e




f




g



h



i






Hint for Q3: The reasons why lines could be parallel include:
- Corresponding angles are equal.
- Alternate angles are equal.
- Cointerior angles add to 180° (are supplementary).


Example 5 Finding angles in parallel lines

Find the value of each of the pronumerals. Give reasons for your answers.

a



b



Solution

a $a = 110$ (vertically opposite angles)

$b + 110 = 180$ (cointerior angles in parallel lines)


$b = 70$

b $b = 69$ (corresponding angles in parallel lines)


$a + 69 = 180$ (supplementary angles)

$a = 111$


Explanation



Vertically opposite angles are equal



Cointerior angles add to 180° in parallel lines



Corresponding angles are equal in parallel lines


Supplementary angles add to 180° so $a + b = 180$.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


Now you try

Find the value of each of the pronumerals. Give reasons for your answers.

a




b




4 Find the values of the pronumerals. Give a reason for each answer.

a




b






Hint for Q4: For part **a**, $x = 125$.
Reason: Alternate angles in parallel lines.
Other reasons include corresponding angles in parallel lines or cointerior angles in parallel lines.


c




d



e




f




5 Find the value of each pronumeral. Give reasons for your answers.


a




b




c




d



e



f




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


8B

Problem-solving and reasoning

6, 7 | 7, 8


6 Two pairs of train tracks meet at 40°, as shown. Find the value of a, b, c, d and e.






7 Find the value of each pronumeral.

a




b






Hint for Q7: Look for pairs of angles inside parallel lines. Mark in all known missing angles to help find the value of the pronumeral.


c




d



e




f




8 Find the value of x.


a



b



c






Hint for Q8: Use all the given information to mark in any missing angles.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


Add a new line to help — 9

9 Sometimes you can add a third parallel line to a diagram to help you find an angle. For example, to find $\angle ABC$ in this diagram you can draw a parallel line through B, then find the two alternate angles (30° and 70°). So $\angle ABC = 30° + 70° = 100°$.




Add a third parallel line to help find $\angle ABC$ in these diagrams.


a




b




c




d



e



f





ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

8C Quadrilaterals CONSOLIDATING

Learning intentions

- To know the properties of special quadrilaterals
- To know the angle sum of a quadrilateral
- To be able to calculate unknown angles inside a quadrilateral

Key vocabulary: quadrilateral, parallelogram, square, rectangle, rhombus, trapezium, kite, diagonal, parallel


Quadrilaterals are polygons with four straight sides. They include six special shapes, four of which are types of parallelograms. Like triangles, they have a special angle sum. This sum is the same for all types of quadrilaterals.



Buildings, such as the modern Library of Alexandria, has a variety of quadrilaterals in its design.

Lesson starter: What is a parallelogram?


A parallelogram is a four-sided shape made from two pairs of parallel lines.



- What are some properties of a parallelogram?
- Do you think the following shapes are parallelograms?
  - Rectangle
  - Kite
  - Rhombus
  - Square
  - Trapezium
- Discuss the properties of each of the special quadrilaterals. Consider side lengths, the lengths of diagonals and the interior angles.


Key ideas

- **Quadrilaterals** are four-sided plane figures with straight sides.
- **Parallelograms** are quadrilaterals with two pairs of parallel sides.




They include:


- **Rectangle**



- **Rhombus**




- **Square**




The red dashed lines are called the **diagonals**.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


- The kite and trapezium are also special quadrilaterals.
  - **Kite**



  - **Trapezium**




- The angle sum of a quadrilateral is $360°$.



$a + b + c + d = 360$

- Supplementary cointerior angles add to $180°$.





Exercise 8C

Understanding

1–3 | 3

1 Name the six special quadrilaterals.

2 A parallelogram has two pairs of parallel sides. Decide whether the following are parallelograms.

a Rhombus **b** Kite **c** Rectangle
d Square **e** Trapezium

3 Write the word to complete these sentences.

a A parallelogram has two pairs of _____ sides.
b The diagonals in a rhombus intersect at _____ angles.
c A quadrilateral with one pair of parallel sides is called a _____.
d The diagonals in a rectangle are _____ in length.

Fluency

4, 5–6(½) | 4, 5–6(½)

Example 6 Working with properties of quadrilaterals


Name all the types of quadrilaterals that have these properties.

a Equal length diagonals **b** Diagonals intersecting at right angles


Solution

a Rectangle and square

Explanation



b Rhombus, square and kite



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

8C

Now you try

Name all the types of quadrilaterals that have these properties.

a Equal opposite side lengths

b Exactly one pair of equal opposite angles.

4 List all the quadrilaterals that have these properties.


a 2 pairs of equal length sides
b All interior angles 90°
c Diagonals of equal length
d Diagonals intersecting at right angles
e 1 pair of parallel sides
f Diagonals of different length



Hint for Q4: Refer to the diagrams in the Key Ideas for help.

Example 7 Finding angles in quadrilaterals

Find the value of the pronumeral in this quadrilateral.



Solution

$x + 80 + 100 + 110 = 360$

$x + 290 = 360$


$x = 70$

Explanation


The angles in a quadrilateral add to 360°.
Simplify and solve for x.


Now you try


Find the value of the pronumeral in this quadrilateral.





5 Find the value of the pronumeral x in these quadrilaterals.


a 

b 

c 

d 

e 

f 




Hint for Q5: Use the angle sum of a quadrilateral, which is 360°.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


Example 8 Finding angles in special quadrilaterals

Find the value of the pronumeral in this kite and trapezium.

a



b



Solution


a $2s + 60 + 40 = 360$

$2s + 100 = 360$

$2s = 260$

$s = 130$

Explanation



The angles in a quadrilateral add to 360° and the opposite angles ($s°$) are equal in a kite.

b $y + 125 = 180$


$y = 55$

Cointerior angles inside parallel lines are supplementary (add to 180°).


Now you try

Find the value of the pronumeral in this kite and trapezium.

a




b




6 Find the value of the pronumerals.

a




b






Hint for Q6: Remember: A kite has one pair of equal opposite angles. Cointerior angles inside parallel lines are supplementary (add to 180°).


c




d



e



f



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

8C

Problem-solving and reasoning 7, 8 7, 9, 10

7 Decide whether it is possible to draw a quadrilateral with:
- **a** 1 interior reflex angle
- **b** 2 interior reflex angles
- **c** 3 acute angles
- **d** 4 acute angles




Hint for Q7: A reflex angle is between 180° and 360°.

8 A shopping centre's floor is a quadrilateral with one 120° angle and two 90° angles. What is the size of the fourth angle?


9 Explain why a rectangle, square and rhombus are all parallelograms.

10 These quadrilaterals include exterior angles. Find the value of x in each.


a



b




c




Quad *x* challenge — 11

11 Find the value of x in these diagrams.


a




b




c




d



e



f





Essential Mathematics Core for the Victorian Curriculum Year 9 ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

8D Polygons ★

Learning intentions

- To know the rule for the angle sum of a polygon
- To be able to calculate unknown angles inside a polygon
- To be able to calculate the interior angle of regular polygons

Key vocabulary: polygon, regular polygon, convex, non-convex

Closed two-dimensional shapes with straight sides are called polygons. They are classified by their number of sides. We have already studied triangles (3 sides) and quadrilaterals (4 sides), which have been classified further by their special properties. The properties of other polygons will be studied here.



Buildings aren't limited to triangles and quadrilaterals in their design. This structure features hexagons.


Lesson starter: Can you draw a polygon?

Use your knowledge of polygons to draw each of the following shapes.


- a convex quadrilateral (all angles less than $180°$)
- a non-convex pentagon (at least one angle greater than $180°$)
- a regular hexagon (all sides and all angles equal)

Key ideas

- A **polygon** is a closed two-dimensional shape with straight sides.
- **Convex** polygons have all interior angles less than $180°$. A **non-convex** polygon has at least one interior angle greater than $180°$.



Convex quadrilateral



Non-convex hexagon

- The sum of the interior angles, S, in a polygon with n sides is given by $S = (n - 2) \times 180°$.


Polygon	Number of sides	Angle sum
Triangle	3	$180°$
Quadrilateral	4	$360°$
Pentagon	5	$540°$
Hexagon	6	$720°$
Heptagon	7	$900°$
Octagon	8	$1080°$
Nonagon	9	$1260°$
Decagon	10	$1440°$
Undecagon	11	$1620°$
Dodecagon	12	$1800°$
n-gon	n	$(n - 2) \times 180°$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

8D

- **Regular polygons** have equal length sides and equal interior angles.

A regular pentagon



Exercise 8D

Understanding

1–4 | 4

1 Write the missing word/rule.

a A _____ has 6 sides.

b A _____ polygon has equal sides and equal angles.

c A _____ polygon has at least one reflex angle.

d The rule for the angle sum of a polygon is $S =$ _____.

2 How many sides do these polygons have?

a Pentagon

b Heptagon

c Quadrilateral

d Undecagon

e Nonagon

f Dodecagon



Hint for Q2: Refer to the Key Ideas for help.

3 Use $S = (n - 2) \times 180°$ to find the angle sum, S, of these polygons. n is the number of sides.

a Hexagon


b Octagon

c Undecagon

d Heptagon

4 Name each of these shapes as convex or non-convex, and write their polygon name.


a




b



c



d





Hint for Q4: A convex polygon has all internal angles less than $180°$.


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Fluency

5–7(½) 5–7(½)


5 Find the value of x using the given angle sum.

a



Angle sum = 540°

b




Angle sum = 720°




Hint for Q5: All the angles must sum to the given angle sum.

c



Angle sum = 900°


d



Angle sum = 720°

Example 9 Finding angles in polygons

Find the angle sum using $S = (n - 2) \times 180°$, then find the value of a.



Solution

$n = 6$ and $S = (n - 2) \times 180°$
$= (6 - 2) \times 180°$
$= 4 \times 180°$
$= 720°$

Explanation


The shape is a hexagon with 6 sides so $n = 6$.

$a + 90 + 115 + 100 + 135 + 40 = 720$
$a + 480 = 720$
$a = 240$

The sum of all angles is 720°. Simplify and solve for a.

Now you try

Find the angle sum using $S = (n - 2) \times 180°$, then find the value of a.




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


8D

6 For each polygon find the angle sum using $S = (n - 2) \times 180°$. Then find the value of a.


a



b




c






Hint for Q6: First find the angle sum using $S = (n - 2) \times 180°$, where n is the number of sides.


d



e




f



Example 10 Working with regular polygons

Find the size of an interior angle of a regular octagon.



Solution

$n = 8$ and $S = (n - 2) \times 180°$

$= (8 - 2) \times 180°$

$= 1080°$

$a = 1080 \div 8$

$= 135$


Explanation

The regular octagon has 8 sides so use $n = 8$.

Each interior angle in a regular polygon is equal so divide by 8 to get a single angle.


Now you try

Find the size of an interior angle of a regular decagon.




7 Find the size of an interior angle of these regular polygons. Round to two decimal places in part **e**.


a




b




c




d



e



f



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Problem-solving and reasoning 8, 9 8–10

8 Calculate the number of sides if a polygon has the given angle sum.
Suggestion: Use the rule $S = (n - 2) \times 180°$.

a $2520°$ **b** $4140°$ **c** $18\,000°$

9 Decide whether the following are possible. If so, make a drawing.


a A hexagon with all angles equal but not all sides equal.

b A hexagon with all sides equal but not all angles equal.




10 These polygons include exterior angles. Find the value of x in each.


a




b



c



d





Hint for Q10: The interior angle sum will first need to be calculated.

Maximum reflex — 11

11 Recall that a non-convex polygon has at least one reflex interior angle.

a Use drawings to decide whether the following are possible.

i A quadrilateral with 2 reflex angles.

ii A pentagon with 2 reflex angles

iii A hexagon with 4 reflex angles

b What is the maximum number of interior reflex angles possible for these polygons?

i Quadrilateral **ii** Pentagon **iii** Octagon

c Write a rule for the maximum number of interior reflex angles for a polygon with n sides.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

8E Congruent triangles

Learning intentions

- To understand the four tests for congruence of triangles
- To be able to recognise a pair of congruent triangles using one of the four tests
- To be able to find unknowns in a pair of congruent triangles

Key vocabulary: congruent/congruent figures, corresponding sides, corresponding angles, congruence statement

Congruent shapes and objects have the same shape and size. Their matching angles are equal and their matching sides have the same length.

It is possible to determine whether or not two triangles are congruent by using four tests, which we call SSS, SAS, AAS and RHS.



The Petronas Twin Towers in Kuala Lumpur look congruent.

Lesson starter: Constructing congruent triangles

For this task you will need a ruler, pencil and protractor. (You might also use compasses.) Divide these constructions up equally among the members of the class. Each group is to draw one of the following triangles, with the given properties.

1. Triangle ABC with $AB = 8$ cm, $AC = 5$ cm and $BC = 4$ cm
2. Triangle DEF with $DE = 7$ cm, $DF = 6$ cm and $\angle EDF = 40°$
3. Triangle GHI with $GH = 6$ cm, $\angle IGH = 50°$ and $\angle IHG = 50°$
4. Triangle JKL with $\angle JKL = 90°$, $JL = 5$ cm and $KL = 4$ cm


- Compare all triangles with the vertices ABC. What do you notice? What does this say about two triangles that have three pairs of equal side lengths?
- Compare all triangles with the vertices DEF. What do you notice? What does this say about two triangles that have two pairs of equal side lengths and the included angles equal?
- Compare all triangles with the vertices GHI. What do you notice? What does this say about two triangles that have two equal corresponding angles and one corresponding equal length side?
- Compare all triangles with the vertices JKL. What do you notice? What does this say about two triangles that have one right angle, the hypotenuse and one other corresponding equal length side?


Key ideas

- **Congruent figures** have the same shape and size.



- If triangle ABC (ΔABC) is congruent to triangle DEF (ΔDEF), we write $\Delta ABC \equiv \Delta DEF$. This is called a **congruence statement**.
  - Letters are usually written in matching order.






Corresponding sides	Corresponding angles
$AB = DE$	$\angle A = \angle D$
$BC = EF$	$\angle B = \angle E$
$AC = DF$	$\angle C = \angle F$


- **Corresponding sides** are opposite equal corresponding angles.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


- Tests for triangle congruence:
  - Side, Side, Side (SSS)
    Three pairs of corresponding sides are equal.




  - Angle, Angle, Side (AAS)
    Two angles and any pair of corresponding sides are equal.



  - Side, Angle, Side (SAS)
    Two pairs of corresponding sides and the included angle are equal.



  - Right angle, Hypotenuse, Side (RHS)
    A right angle, the hypotenuse and one other pair of corresponding sides are equal.




Exercise 8E

Understanding

1–3 | 2, 3(½)

1 These two triangles are congruent.



a Name the point on ΔXYZ that matches:
- **i** point B
- **ii** point A
- **iii** point C

b Name the side on ΔXYZ that corresponds to (matches):
- **i** AB
- **ii** AC
- **iii** BC

c Name the angle in ΔABC that corresponds to (matches):
- **i** $\angle X$
- **ii** $\angle Y$
- **iii** $\angle Z$

2 Copy and complete the sentences below.

a Congruent figures are exactly the same shape and _____.

b If triangle ABC is congruent to triangle STU then we write $\Delta ABC \equiv$ _____.

c The short names of the four congruence tests for triangles are SSS, ____, ____ and ____.

3 Write a congruence statement if:

a triangle ABC is congruent to triangle FGH

b triangle DEF is congruent to triangle STU



Hint for Q3: Example:
$\Delta DEF \equiv \Delta GHI$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


8E


Fluency


4, 5(½) | 4–5(½)


Example 11 Choosing a congruence test

Which congruence test (SSS, SAS, AAS or RHS) would be used to show that these pairs of triangles are congruent?

a 

b 


c 


d 


Solution	Explanation
a SAS	Two pairs of corresponding sides and the included angle are equal.
b RHS	A right angle, hypotenuse and one pair of corresponding sides are equal.
c AAS	Two angles and a pair of corresponding sides are equal.
d SSS	Three pairs of corresponding sides are equal.


Now you try

Which congruence test (SSS, SAS, AAS or RHS) would be used to show that these pairs of triangles are congruent?

a 

b 


c 

d 


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

4 Which congruence test (SSS, SAS, AAS or RHS) would be used to show that these pairs of triangles are congruent?


a




b




c




d




e




f



g



h






Hint for Q4: Choose SSS for three matching sides, SAS for two matching sides and the included angle, AAS for two equal angles and one matching side or RHS for a right angle, hypotenuse and one other matching side.


Example 12 Finding missing side lengths and angles using congruence

Find the values of the pronumerals in these pairs of congruent triangles.

a



b



Solution

a $x = 5$

$y = 7$

b $a = 25$

$b = 180 - 100 - 25$

$= 55$

Explanation

The side of length x and the side of length 5 are in corresponding positions (opposite the □).

The longest side on both triangles must be equal.


The angle marked $a°$ corresponds to the 25° angle in the other triangle.

The sum of three angles in a triangle is 180°.


Now you try

Find the values of the pronumerals in these pairs of congruent triangles.

a



b




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


8E

5 Find the values of the pronumerals in these pairs of congruent triangles.

a




b






Hint for Q5: You already know they are congruent!


c




d




e




f



g




h



Problem-solving and reasoning 6–8 7–10

6 Measurements were taken for three matching triangular tiles and are shown here.




a Which two triangles are congruent?


b What reason (SSS, SAS, AAS, RHS) is used to explain their congruence?

7 For each set of three triangles, choose the two that are congruent. Give a reason (SSS, SAS, AAS or RHS) and write a congruence statement (e.g. $\Delta ABC \equiv \Delta FGH$).

a



b








Hint for Q7: Write matching vertices in the same order.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


c









d







8 Explain why RHS is not the first reason you would use to explain why these two triangles are congruent.






Hint for Q9: Draw examples to justify your decision.

9 Are all triangles with three pairs of equal corresponding angles congruent? Explain why or why not.

10 $ABCD$ is a parallelogram.


a Give the reason why $\Delta ABC \equiv \Delta CDA$.


b What does this say about $\angle B$ and $\angle D$?





Fishing for congruence — 11


11 Identify all pairs of congruent triangles from those below by writing a congruence statement. Angles with the same mark are equal.



































ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


Progress quiz

8A **1** For the angle 63°, determine the:


a complementary angle **b** supplementary angle

8A **2** Name the types of triangles shown and find the value of each pronumeral. Part **c** includes an exterior angle.


a



b




c




8B **3** Explain why the pairs of lines shown below are *not* parallel.

a




b




8B **4** Find the values of the pronumerals in the parallel lines below. Give reasons for your answers.


a



b




c




8C **5** Find the values of the pronumerals in these quadrilaterals and name the type of quadrilateral in parts **b** and **c**.


a



b




c




8D **6** Find the angle sum of each of the following polygons using $S = (n - 2) \times 180°$ and then find the value of a. The polygon in part **b** is a regular polygon.

a




b




8E **7** Which congruence test (SSS, SAS, AAS or RHS) would be used to show that these pairs of triangles are congruent?


a



b



8E **8** Find the value of the pronumerals in this pair of congruent triangles.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

8F Enlargement and similar figures

Learning intentions

- To know that enlargement of a figure produces a similar figure
- To be able to recognise that two shapes are similar
- To be able to construct a similar figure using the enlargement transformation
- To be able to calculate a scale factor for similar figures
- To be able to find missing angles and side lengths in similar figures

Key vocabulary: similar figures, enlargement, scale factor, ratio, corresponding sides, corresponding angles, image

Whenever we look at a map, TV screen, microscope, model, movie screen or computer document we are looking at similar figures.


Similar figures have the same shape but not necessarily the same size. If two figures are similar then one of them can be enlarged or reduced so that it is identical (congruent) to the other. If a figure is enlarged by a scale factor greater than 1, the image will be larger than the original. If the scale factor is between 0 and 1, the image will be smaller.





The image on the screen is similar to the original; it has the same shape but a different size.

Lesson starter: Are they similar?

Here are three quadrilaterals, not drawn to scale.








- Which pair of quadrilaterals do you think are similar?
- Why do you think they are similar? Discuss your reasons.
- Give reasons why the other quadrilateral is not similar.

Key ideas

- **Enlargement** is a transformation that involves the increase or decrease in size of an object.
  - The 'shape' of the object is unchanged.
  - Enlargement uses a centre of enlargement and an enlargement factor or **scale factor**.
  - The scale factor is the number by which you multiply each side length to enlarge or reduce the size of a shape.
  - The **image** of a point A can be labelled as A'.




- Two figures are **similar** if one can be enlarged to be congruent to the other. They are of the same shape but not the same size.
  - Corresponding angles are equal.
  - Pairs of corresponding sides are in the same proportion or ratio.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

8F

- The **scale factor** = $\frac{\text{image length}}{\text{original length}}$
- The symbols ||| and ~ are used to describe similarity.
  - For example, $ABCD \mid\mid\mid EFGH$ or $ABCD \sim EFGH$
  - The letters are usually written in matching order.
  - Scale factor = $\frac{EF}{AB} = \frac{10}{5} = 2$
  - The scale factor will be the same for all matching pairs of sides.



Exercise 8F

Understanding

1–4 | 1, 2, 4

1 Write the missing word or symbol.

a Scale factor = $\frac{\text{_____ length}}{\text{original length}}$

b Similar shapes are the same _____ but different size.

c If $ABCD$ is similar to $EFGH$, then we write $ABCD$ _____ $EFGH$.

2 The two figures below are similar.

a Name the angle in the larger figure that corresponds to (matches) $\angle A$.

b Name the angle in the smaller figure that corresponds to $\angle I$.


c Name the side in the larger figure that corresponds to BC.

d Name the side in the smaller figure that corresponds to FJ.

e Use FG and AB to find the scale factor.



Hint for Q2: For part e, use scale factor = $\frac{\text{image length}}{\text{original length}}$



3 This diagram shows ΔABC enlarged to give the image $\Delta A'B'C'$.


a Measure the lengths OA and OA'. What do you notice?

b Measure the lengths OB and OB'. What do you notice?

c Measure the lengths OC and OC'. What do you notice?


d What is the scale factor?

e Is $A'B'$ twice the length of AB? Measure to check.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

4 This diagram shows rectangle $ABCD$ enlarged (in this case reduced) to rectangle $A'B'C'D'$.




a Measure the lengths OA and OA'. What do you notice?
b Measure the lengths OD and OD'. What do you notice?
c What is the scale factor?
d Compare the lengths AD and $A'D'$. Is $A'D'$ one quarter of the length of AD?

Fluency

5, 6 | 5(½), 6, 7

Example 13 Using the scale factor

These figures are similar.
a Find the scale factor.
b Find the value of x.
c Find the value of y.



Solution

a Scale factor $= \frac{6}{4} = 1.5$

b $x = 3 \times 1.5$
$= 4.5$

c $y \times 1.5 = 7.5$
$y = 7.5 \div 1.5$
$= 5$

Explanation


Choose two corresponding sides and use
$\text{scale factor} = \frac{\text{image length}}{\text{original length}}$

Multiply by the scale factor to get the length of the side on the larger image.

When y is multiplied by the scale factor it gives a length of 7.5 on the image. Divide by 1.5 to solve for y.

Now you try

These figures are similar.
a Find the scale factor.
b Find the value of x.
c Find the value of y.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

8F

5 Each of the pairs of figures shown here are similar. For each pair:

i Find the scale factor.


ii Find the value of x.

iii Find the value of y.




Hint for Q5: Scale factor = $\frac{\text{larger}}{\text{smaller}}$ Make sure you choose a matching pair of sides.


a




b




c




d



e



f




Example 14 Enlarging figures

Copy the given diagram using plenty of space, and use the given centre of enlargement (O) and these scale factors to enlarge ΔABC.


a Scale factor $\frac{1}{2}$

b Scale factor 3



Solution

a




Explanation

Connect dashed lines between O and the vertices A, B and C.

Since the scale factor is $\frac{1}{2}$, place A' so that OA' is half of OA.

Repeat for B' and C'. Join vertices A', B' and C'.

b



Draw dashed lines from O through A, B and C. Place A' so that OA' is 3 times OA. Repeat for B' and C' and form $\Delta A'B'C'$.


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Now you try

Copy the given diagram using plenty of space, and use the given centre of enlargement (O) and these scale factors to enlarge ΔABC.

a Scale factor 2


b Scale factor $\frac{1}{2}$



6 Copy the given diagram, leaving plenty of space around it. Use the given centre of enlargement (O) and given scale factors to enlarge ΔABC.

a Scale factor $\frac{1}{3}$

b Scale factor 2




7 This diagram includes a square with centre O and vertices $ABCD$.

a Copy the diagram, leaving plenty of space around it.

b Enlarge square $ABCD$ by these scale factors and draw the image. Use O as the centre of enlargement.

i $\frac{1}{2}$ **ii** 1.5



Problem-solving and reasoning


8, 9 | 10, 11

8 A square is enlarged by a scale factor of 4.

a Are the internal angles the same for both the original and the image?

b If the side length of the original square was 2 cm, what would be the side length of the image square?

c If the side length of the image square was 100 cm, what would be the side length of the original square?



9 A photo of a sunflower is enlarged by a scale factor of 2. If the larger flower has a diameter of 7.4 cm, what is the diameter of the smaller flower?



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


8F

10 These diagrams show a shape and its image after enlargement. For each part, find:


i the scale factor

ii the coordinates (x, y) of the centre of enlargement


a




b



c



d





Hint for Q10: For parts **c** and **d**, first draw dashed red lines connecting A with A' etc.

11 Explain why:

a any two squares are similar

b any two equilateral triangles are similar

c any two rectangles are not necessarily similar

d any two isosceles triangles are not necessarily similar


Actual lengths from drawings and maps — 12, 13

12 The container of this truck is 12.7 m long.

a Measure the length of the container on the truck in the drawing.

b Measure the height of the container on the truck in the drawing.

c Estimate the actual height of the container on the truck.





13 A map has a scale ratio of $1 : 50\,000$.

a What length on the ground is represented by 2 cm on the map?

b What length on the map is represented by 12 km on the ground?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

8G Similar triangles

Learning intentions

- To understand the four tests for similarity of triangles
- To be able to recognise a pair of similar triangles using one of the four tests
- To be able to calculate and use the scale factor to find an unknown length

Key vocabulary: similar, scale factor, ratio, corresponding sides, corresponding angles, hypotenuse


As with congruent triangles, there are mathematical tests that can be used to see if two triangles are similar. Once we know that two triangles are similar, we can work out any missing angles and lengths.


Lesson starter: Why are AA and AAA the same test for triangles?

If all three angles are the same inside two triangles then we know that the triangles are similar.

Here are a pair of triangles.


- Do you think they are similar?
- What is the missing angle in each triangle?
- Can you explain why the AA test is the same as AAA?






Key ideas


- Two triangles are similar if:
  - corresponding angles are equal
  - corresponding sides are in proportion (the same ratio).
- The similarity statement for two similar triangles ΔABC and ΔDEF is:
  - $\Delta ABC \text{ ||| } \Delta DEF$ or
  - $\Delta ABC \sim \Delta DEF$
- Tests for similar triangles (not to be confused with the congruence tests for triangles):
  - Side, Side, Side (SSS):
    All three pairs of corresponding sides are in the same ratio.
    $$\frac{12}{6} = \frac{8}{4} = \frac{14}{7}$$




  - Side, Angle, Side (SAS):
    Two pairs of corresponding sides are in the same ratio and the included angles are equal.
    $$\frac{22}{11} = \frac{10}{5}$$



  - Angle, Angle, Angle (AAA or AA):
    All three corresponding angles are equal
    (If there are two equal pairs then the third pair must be equal.)



  - Right angle, Hypotenuse, Side (RHS):
    The hypotenuses of right-angled triangles and another corresponding pair of sides are in the same ratio.
    $$\frac{15}{5} = \frac{6}{2}$$



Essential Mathematics Core for the Victorian Curriculum Year 9
ISBN 978-1-009-59315-1
© Greenwood et al. 2025
Photocopying is restricted under law and this material must not be transferred to another party.
Cambridge University Press

Exercise 8G

Understanding

1, 2 | 2

1 Copy and complete the following sentences.


a The abbreviated tests for similar triangles are SSS, ___, ___ and ___.

b Similar figures have the same _____ but are not necessarily the same _____.

c ΔABC is a _____ (*line*, *triangle* or *quadrilateral*).

d If $\Delta ABC \mid\mid\mid \Delta DEF$, then ΔABC is _____ to ΔDEF.

2 These two triangles are similar.



a Which vertex on ΔDEF corresponds to (matches) vertex B on ΔABC?

b Which vertex on ΔABC corresponds to (matches) vertex F on ΔDEF?

c Which side on ΔDEF corresponds to (matches) side AC on ΔABC?

d Which side on ΔABC corresponds to (matches) side EF on ΔDEF?

e Which angle on ΔABC corresponds to (matches) $\angle D$ on ΔDEF?

f Which angle on ΔDEF corresponds to (matches) $\angle B$ on ΔABC?


Fluency

3, 4(½), 5, 6 | 3–4(½), 6, 7


Example 15 Choosing a similarity test for triangles

Choose the similarity test that proves that these pairs of triangles are similar.


a




b



c



d



Solution	Explanation
a AAA	Three pairs of angles are equal.
b RHS	Both are right-angled triangles and the hypotenuses and another pair of sides are in the same ratio $\left(\frac{9}{3} = \frac{6}{2}\right)$.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


c SAS — Two pairs of corresponding sides are in the same ratio $\left(\frac{21}{7} = \frac{18}{6}\right)$ and the included angles are equal.

d SSS — Three pairs of corresponding sides are in the same ratio $\left(\frac{15}{5} = \frac{12}{4} = \frac{6}{2}\right)$.


Now you try

Choose the similarity test that proves that these pairs of triangles are similar.


a




b



c




d




3 Choose the similarity test which proves that these pairs of triangles are similar.

a




b






Hint for Q3: Choose from SSS, SAS, AAA or RHS.


c




d




e




f



g



h




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


8G

4 Write similarity statements for these pairs of similar triangles. Write letters in matching order.


a




b




c




d



e



f






Hint for Q4: Here is an example of a similarity statement: $\Delta ABC \text{ ||| } \Delta DEF$



Hint for Q4: For parts **e** and **f**, use angles in parallel lines.


5 What is the scale factor of this pair of similar triangles that enlarges ΔABC to ΔDEF?



Example 16 Finding a missing length using similarity

For this pair of triangles:

a give a reason (SSS, SAS, AAA or RHS) why the two triangles are similar
b find the scale factor
c find the value of x



Solution

a AAA or just AA.

b Scale factor $= \frac{3}{2} = 1.5$.

c $x = 5 \times 1.5$
$= 7.5$

Explanation

$\angle EAC = \angle DBC$ since AE is parallel to BD and $\angle C$ is common to both triangles. (Also $\angle AEC = \angle BDC$ since AE is parallel to BD).

$\frac{AE}{BD} = \frac{3}{2}$.


Multiply CD by the scale factor to find the length of the corresponding length CE.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Now you try


For this pair of triangles:

- **a** give a reason (SSS, SAS, AAA or RHS) why the two triangles are similar
- **b** find the scale factor
- **c** find the value of x



6 For this pair of triangles:

- **a** give a reason (SSS, SAS, AAA or RHS) why the two triangles are similar
- **b** find the scale factor
- **c** find the value of x






Hint for Q6: To use SSS, SAS or RHS, you need more than one pair of matching sides.

7 For this pair of triangles:


- **a** give a reason (SSS, SAS, AAA or RHS) why the two triangles are similar
- **b** find the scale factor
- **c** find the value of x



Problem-solving and reasoning


8, 9 | 8–11

8 Two people create a shadow in the Sun, as shown. What is the height of the taller person?



9 Name the triangle that is not similar to the other two in each group of three triangles.


a





Hint for Q9: Name a triangle like this: ΔABC or ΔSTU.


b





ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

8G


c









d







10 For each pair of similar triangles:


i give a reason (SSS, SAS, AAA or RHS) why the two triangles are similar

ii find the value of x




Hint for Q10: Consider angle properties in parallel lines.


a




b




c



11 Give reasons why these two triangles are similar.





Area ratio — 12

12 Consider these three similar triangles (not drawn to scale).

Original




Image 1




Image 2





Image 3



a Complete this table, comparing each image to the original.

Triangle	Original	Image 1	Image 2	Image 3
Length scale factor	1	2		
Area	4	16		
Area scale factor	1			

b What do you notice about the area scale factor compared to the length scale factor?

c What would be the area scale factor if the length scale factor is:

i 10? **ii** 20? **iii** 100?

Essential Mathematics Core for the Victorian Curriculum Year 9 ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

8H Applying similar triangles

Learning intentions

- To be able to identify a pair of similar triangles in a given context
- To be able to give a reason why two triangles are similar
- To be able to calculate and use the scale factor to find an unknown length in a real situation

Key vocabulary: similar figures, scale factor

Similar triangles can be used in many mathematical and practical problems. If two triangles are proved to be similar then the properties of similar triangles can be used to find unknown lengths or angles. The approximate height of a tall object, or the width of a projected image, can be found using similar triangles.




By knowing the girl's height and the length of the kite string to the girl and the ground, you can calculate the kite's height.

Lesson starter: How high is it?

A 1 m vertical stick gives a shadow 80 cm long. At the same time, a tall tree has a shadow 24 m long.

- How can AAA be used to explain the two similar triangles?
- What is the scale factor?
- What is the height of the tree?



Key ideas

- To apply similarity in practical problems involving triangles:
  - Prove that two triangles are similar by stating one of the tests: SSS, SAS, AAA or RHS.
  - Find a scale factor.
  - Find the value of any unknowns.

Exercise 8H

Understanding

1–3 | 3


1 Here are two triangles.


a How many pairs of equal matching angles are given?

b Which test (SSS, SAS, AAA or RHS) explains why they are similar?

c What is the scale factor if the smaller triangle is enlarged to the size of the larger triangle?

d Find the value of x.






ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

8H


2 In this diagram, name the angle that is common to both ΔACE and ΔBCD.





Hint for Q2: You can name the angle like this: $\angle S$.


3 In this diagram:
- **a** name the pair of vertically opposite angles
- **b** name the two pairs of equal alternate angles



Fluency


4–7 | 5–8

4 Two similar triangles are created by cables supporting a yacht's mast.
- **a** Find a scale factor for the two triangles.
- **b** Find the height of the mast.



Example 17 Applying similarity

Chris' shadow is 1.2 m long while a 1 m vertical stick has a shadow 0.8 m long.
- **a** Give a reason why the two triangles are similar.
- **b** Determine Chris' height.



Solution

a All angles are the same (AAA).

b Scale factor $= \frac{1.2}{0.8} = 1.5$

$\therefore$ Chris' height $= 1 \times 1.5$

$= 1.5$ m

Explanation


The Sun's rays will pass over Chris and the stick and hit the ground at approximately the same angle.

First find the scale factor.

Multiply the height of the stick by the scale factor to find Chris' height.

Now you try

Two ladders lean against a vertical wall at the same angle as shown. The distances that the base of the ladders are from the wall are 3 m and 6 m.
- **a** Give a reason why the two triangles are similar.
- **b** Find how high the longer ladder reaches up the wall if the shorter ladder reaches up 5 m.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5 A tree's shadow is 20 m long while a 2 m vertical stick has a shadow 1 m long.

a Give a reason why the two triangles contained within the objects and their shadows are similar.

b Find the height of the tree.




Hint for Q5: Construct a diagram as in Example 17.

6 John stands 6 m from a lamp post and casts a 2 m shadow. The shadow from the pole and from John end at the same place. If John is 1.5 m tall, what is the height of the lamp post?

7 Two cables support a steel pole at the same angle as shown. The two cables are 4 m and 10 m in length and the shorter cable reaches 3 m up the pole.

a Give a reason why the two triangles are similar.

b Find the height of the pole.






8 Ali is at the beach and decides to estimate how far an exposed rock is from the seashore. He places four pegs in the sand as shown, and measures the distance between them.

a Why do you think Ali has placed the four pegs in the way that is shown in the diagram?


b Are the two triangles similar? Which test (SSS, SAS, AAA or RHS) could be used?

c How far is the rock from the beach?



Problem-solving and reasoning 9, 10 9, 11


9 A deep chasm has a large rock (R) sitting on its side, as shown. Find the width of the chasm.





Hint for Q9: Don't forget to find the scale factor.


10 Joanne wishes to determine the width of the river shown without crossing it. She places four pegs as shown. Calculate the river's width.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

8H

11 A person 1.8 m tall stands in front of a light that sits on the floor.
The person casts a shadow on the wall behind them.




a Find how tall the shadow will be if the distance between the wall and the light is:
i 4 m
ii 10 m
iii 3 m

b How tall will the shadow be if the distance between the wall and the person is:
i 4 m? **ii** 5 m?

c Find the distance from the wall to the person if the shadow is:
i 5.4 m **ii** 7.2 m

Visual challenge — 12, 13


12 Find the length AB in this diagram if the two triangles are similar.





Hint for Q12: Remember to line up corresponding sides for the scale factor.

13 In this diagram, AC is perpendicular (at right angles) to BD and ΔABD is right-angled.



a How many similar triangles are there?

b Find the lengths:
i BD **ii** AC **iii** AB

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Animator

Animators are creative people who use their skills to hand-draw, model or computer-generate images to create animated effects.

While animators are artistic, many other skills are necessary to excel in this field. Animators must be patient, have an eye for detail, and be able to work in groups as well as independently. They need excellent communication skills and IT skills.

Geometry and linear programming are important in this field as they allow the animator to illustrate how an image can be rotated and shifted. Similarity and congruency of objects are used to ensure consistency between scenes.

Animating is not just restricted to the movies. Computer and console computer games, television series, advertising and even corporate training organisations often directly employ or out-source animators.



1 Being able to accurately copy drawings by hand is a skill used in some forms of animation. Use your knowledge of angles and proportion to carefully copy these logos into your workbook.

a



b




c




2 Using graph paper, enlarge each of the following logos by a factor of two.

a




b




Essential Mathematics Core for the Victorian Curriculum Year 9 ISBN 978-1-009-59315-1 © Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3 How many pairs of congruent triangles are in the symmetrical logo below? Can you see any pairs of similar triangles?



4 Compare each of the following figures and state how each has been rotated relative to the preceding one. You will need to use a protractor.



Using technology

5 Animators enjoy creating original designs. Use computer drawing software, such as Geometer's Sketchpad, to design a geometrical logo from the initials of your name. Some ideas for 'D and W' logos are shown below, but your design can be original if you wish.




Essential Mathematics Core for the Victorian Curriculum Year 9 ISBN 978-1-009-59315-1 © Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Ocean Ironman

The organisers of an ocean ironman event want to estimate the distance from the beach to a floating buoy in the sea. Flags are placed on the beach by the organisers in an effort to create similar triangles as shown.





The flag positions are at A, C, D and E and the line segments AB and DE are assumed to be at right angles with AD at the beach.

Present a report for the following tasks and ensure that you show clear mathematical workings and explanations where appropriate.

1 Preliminary task

Once the flags are in position one of the organisers measures the following;

- $AC = 200$ metres
- $CD = 5$ metres
- $DE = 4$ metres

a Find the scale factor between the two similar triangles.

b Hence, estimate the distance from the beach to the buoy (distance AB).

2 Modelling task

The problem is to determine the distance between the beach and the buoy using similar triangles.

a Write down all the relevant information that will help solve this problem, and illustrate using diagrams.

b Explain why the positions of the flags placed by the organisers produce a pair of similar triangles. (That is, triangle DCE is similar to triangle ACB).

c Two more attempts at positioning flags and measuring distances are undertaken by the organisers.
Jimmy's placement:

- $AC = 360$ metres
- $CD = 8$ metres
- $DE = 3$ metres.

Olivia's placement:

- $AC = 90$ metres
- $CD = 5$ metres
- $DE = 8$ metres.

i Use Jimmy's model to estimate the distance between the beach and the buoy.
ii Use Olivia's model to estimate the distance between the beach and the buoy.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Evaluate and verify

d Another organiser attempts to place the flags such that:

- $AC = 100$ metres
- $CD = 50$ centimetres
- $DE = 1.2$ metres.

Do you think that arranging the flags in this way is a good idea? Explain why.

e If the actual distance from the beach to the buoy is 200 metres, determine an arrangement for the flags that would deliver the correct distance. Illustrate with a diagram including the measurements.

Communicate


f Summarise your results in a table.

	AC	CD	DE	AB
PERSON 1	200	5	4	
JIMMY				
OLIVIA				
PERSON 4				

Comment on any findings and recommendations.

3 Extension questions

a Another way to arrange the flags is to use the model shown here. Using this model, identify the pair of similar triangles and explain why they are similar.



b Using the following measurements, estimate the distance from the beach to the buoy.

- $AC = 100$ metres
- $AE = 60$ metres
- $DE = 140$ metres

c Compare the two models studied in this task using the different diagrams. Do you have a preferred model? Explain why.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Geometric bisection

Key technology: Dynamic geometry

To generate drawings with a high degree of accuracy it is important not just to rely on guess work or trial and error. Instead, we use construction techniques using the properties of lines and shapes to pinpoint positions accurately. These construction techniques could be achieved using a pair of compasses, pencil and ruler; however, dynamic geometry software could also be used with the added benefit that points in the construction can be dragged to test multiple cases.



1 Getting started

We will start by considering constructing the perpendicular bisector of a line segment. Use dynamic geometry like Desmos to complete the following.


a Construct a line segment AB.

b Construct a circle with centre A and radius AB and a circle with centre B and radius AB.

c Construct points C and D at the intersection of the circles.

d Construct CD, which is the perpendicular bisector of AB, and the point M, which is the midpoint of AB.


e Check that your construction is correct by dragging either of points A or B. The segment CD should move with the construction as the perpendicular bisector of AB.




2 Using technology

a Use dynamic geometry to construct the following. The given diagrams will provide clues as to how to complete the construction.

i Equilateral triangle




ii Isosceles triangle




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

iii Angle bisector



iv Perpendicular line




b For each of the above constructions, drag one of the initial points to see if the properties of the construction are retained. For the angle bisector, for example, the ray OE should always bisect the angle AOB when point A is dragged.

3 Applying an algorithm

a Choose one of the constructions above and write a simple algorithm (sequence of steps) which tells the user how to create the construction in the correct order.

b Follow this algorithm to construct a square. The special tool 'Perpendicular line' inside the dynamic geometry software is used.

- Step 1: Construct segment AB.
- Step 2: Construct lines perpendicular to AB passing through A and perpendicular to AB passing through B.
- Step 3: Construct the circle with centre A and radius AB.
- Step 4: Construct point C.
- Step 5: Construct the line perpendicular to AC passing through C.
- Step 6: Construct point D.




c In your square construction, drag one of points A or B. ABCD should always be a square regardless of the position of A or B.


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

1 Use 12 matchsticks to make 6 equilateral triangles.


2 Find the value of a in the diagram.




3 How many acute angles are there in this diagram?




4 A circle is divided using chords (one chord is shown here, giving two regions).
What is the maximum number of regions that can be formed if the circle is divided with 4 chords?



5 Explore (using dynamic geometry) where the points A,B and C should be on the sides of Δ DEF so that the perimeter of Δ ABC is a minimum.




6 The Sierpinski triangle shown is a mathematically generated pattern.





It is created by repeatedly enlarging triangles by a factor of $\frac{1}{2}$.
The steps are:

- Start with an equilateral triangle, as in figure 1.
- Enlarge the triangle by a factor of $\frac{1}{2}$.
- Arrange three copies of the image, as in figure 2.
- Continue repeating steps 2 and 3 with each triangle.

a Make a large copy of figures 1 to 3 and then draw the next two figures in the pattern.


Figure 1


Figure 2


Figure 3

b If the original triangle (figure 1) had side length l, what is the side length of the smallest triangle in:
i figure 2? **ii** figure 3?
iii figure 8 (assuming figure 8 is the 8th diagram in the pattern)?

c What fraction of the area is shaded in:
i figure 2? **ii** figure 3?
iii figure 6 (assuming figure 6 is the 6th diagram in the pattern)?

d The Sierpinski triangle is one where the process of enlargement and copying is continued forever. What is the shaded area of a Sierpinski triangle?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter summary

Geometry

Angles

Acute $0° < \theta < 90°$
Right $90°$
Obtuse $90° < \theta < 180°$
Straight $180°$
Reflex $180° < \theta < 360°$
Revolution $360°$
Complementary angles add up to $90°$
Supplementary angles add up to $180°$
Vertically opposite angles are equal

Parallel lines

If the lines are parallel:

- corresponding angles are equal
- alternate angles are equal
- cointerior angles are supplementary $a + b = 180$



Quadrilaterals

Angle sum = $360°$


Parallelogram
Rhombus
Rectangle
Square
Kite
Trapezium

Polygons

The sum of the interior angles in a polygon with n sides is $S = (n - 2) \times 180°$. Regular polygons have all sides and angles equal.

Triangles

Sum of angles is $180°$.
Exterior angle equals sum of two opposite interior angles $a = b + c$



Types:
Acute-angled – all angles $< 90°$
Obtuse-angled – 1 angle $> 90°$
Right-angled – 1 angle $90°$
Equilateral – all angles $60°$ – all sides are equal
Isosceles – 2 angles and 2 sides are equal
Scalene – all sides and angles are different sizes

Congruent triangles

These are identical in shape and size. They may need to be rotated or reflected. We write $\Delta ABC \equiv \Delta DEF$ (congruent to)

Tests for congruence:
SSS – three pairs of corresponding sides equal
SAS – two pairs of corresponding sides and included angle are equal
AAS – two angles and any pair of corresponding sides are equal
RHS – a right angle, hypotenuse and one other pair of corresponding sides are equal

Similar figures

These are the same shape but different size.
Scale factor $= \frac{\text{image length}}{\text{original length}}$

Similar triangles

Similar triangles have corresponding angles equal and corresponding sides are in the same ratio.
We say $\Delta ABC \,|||\, \Delta DEF$ or $\Delta ABC \sim \Delta DEF$ (similar to)

Tests for similar triangles:
SSS – three pairs of corresponding sides in the same ratio
SAS – two pairs of corresponding sides in the same ratio and included angle equal
AAA or AA – all three pairs of corresponding angles are equal; two pairs is enough to prove this
RHS – the hypotenuses of right-angled triangles and another pair of corresponding sides are in the same ratio


Applying similar triangles

In practical problems:

- look to identify and prove pairs of similar triangles
- find a scale factor
- use this to find the value of any unknowns.

e.g. The shadow cast by a tree is 10.5 m while a person 1.7 m tall has a 1.5 m shadow. How tall is the tree?

Similar AAA
Scale factor $= \frac{10.5}{1.5} = 7$
$\therefore\ h = 1.7 \times 7$
$= 11.9$ m
Tree is 11.9 m tall.








(Not to scale)

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.






Chapter checklist

A version of this checklist that you can print out and complete can be downloaded from your Interactive Textbook.



		✔
8A	**1 I can find supplementary and complementary angles.** e.g. State the supplementary and complementary angle to 25°.	
8A	**2 I can name triangles and find the size of unknown angles.** e.g. Name the type of triangle in this diagram and find the values of the pronumerals. 	
8B	**3 I can find unknown angles in parallel lines.** e.g. Find the values of the pronumerals in this diagram and give a reason. 	
8B	**4 I can explain why two lines are parallel.** e.g. Decide, with reasons, whether the given pair of lines are parallel. 	
8C	**5 I can identify properties of various quadrilaterals.** e.g. Name the special quadrilaterals which have diagonals which intersect at 90°.	
8C	**6 I can find an unknown angle in a quadrilateral.** e.g. Find the value of a in this quadrilateral. 	
8C	**7 I can find an unknown angle in a special quadrilateral.** e.g. Find the value of a in this kite. 	

Essential Mathematics Core for the Victorian Curriculum Year 9
ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter checklist

		✔
8D	**8 I can find an angle sum of a polygon and an unknown angle in a polygon.** e.g. Find the value of a in this heptagon after finding the angle sum. 	
8D	**9 I can find the internal angle in a regular polygon.** e.g. Find the size of an internal angle inside a regular octagon.	
8E	**10 I can choose a test and write a congruence statement for a pair of congruent triangles.** e.g. Write a congruence statement and choose a test to prove congruence for this pair of triangles. 	
8E	**11 I can find missing side lengths and angles in congruent triangles.** e.g. Find the value of the pronumerals given the following pair of congruent triangles. 	
8F	**12 I can find the scale factor for a pair of similar figures and find the length of unknown sides.** e.g. Find the scale factor for these similar figures then find the value of the pronumerals. 	
8F	**13 I can draw a similar figure using the enlargement transformation and a given scale factor.** e.g. Copy the given diagram and enlarge the given shape using the centre of enlargement (O) and the scale factors of 3 and also $\frac{1}{2}$. 	

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

		✔
8G	**14 I can recognise when two triangles are similar using one of the four tests.** e.g. Give the reason why these triangles are similar. 	
8G	**15 I can find a scale factor and use this to find an unknown length.** e.g. If the given pair of triangles is known to be similar, find the value of x. 	
8H	**16 I can identify similar triangles in a real context and use a scale factor to find an unknown length.** e.g. A tree's shadow is 20 m long and a 4 m high vertical stick has a 5 m shadow at the same time of day. **a** Use a diagram to form a pair of similar triangles and give a reason why they are similar. **b** Use the scale factor to find the height of the tree.	


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter review


Short-answer questions

8A **1** Find the value of the pronumerals.


a



b




c




8A **2** Name the following triangles and find the value of the pronumerals.


a



b




c




8B **3** Decide whether the following contain a pair of parallel lines. Give a reason.


a



b




c




8B **4** State the values of the pronumerals.


a



b



c



8C **5** Name the quadrilaterals that have:


a equal length diagonals

b diagonals intersecting at right angles


c two pairs of equal length sides

8C **6** Find the value of each pronumeral in the following quadrilaterals.


a



b




c




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

8D **7** Find the value of the pronumeral in these polygons. Use $S = (n - 2) \times 180°$ for the angle sum.


a



b




c




8E **8** Determine whether each pair of triangles is congruent. If congruent, give the abbreviated reason (SSS, SAS, AAS or RHS) and state the value of any pronumerals.


a




b




c



d




8F **9** Copy the given diagram using plenty of space. Using the centre of enlargement (O) and a scale factor of 3, enlarge ΔABC.




8G **10** Determine whether the following pairs of triangles are similar, and state the similarity test (SSS, SAS, AAS or RHS) that proves this.


a




b





c





d






ISBN 978-1-009-59315-1
© Greenwood et al. 2025
Photocopying is restricted under law and this material must not be transferred to another party.


Chapter review

8G **11** For the following pairs of similar triangles, find the value of x.


a



b




c



8H **12** A conveyor belt loading luggage onto a plane is 12 m long. A vertical support 1.6 m high is placed under the belt. It is 4 m along the conveyor belt, as shown in the diagram.

a Find the scale factor for the two similar triangles.

b Find the height (AE) of the luggage door above the ground.




Multiple-choice questions

8A **1** The angle that is supplementary to an angle of 55° is:

A 35° **B** 55° **C** 95° **D** 125° **E** 305°

8B **2** What is the value of x if AB is parallel to CD?

A 110 **B** 70 **C** 20 **D** 130 **E** 120



8A **3** If two angles in a triangle are 30° and 80°, then the third angle is:

A 110° **B** 70° **C** 90° **D** 80° **E** 30°

8B **4** The values of x and y in the diagram are:


A $x = 35, y = 85$

B $x = 45, y = 45$

C $x = 50, y = 60$

D $x = 55, y = 65$

E $x = 65, y = 55$



8D **5** Using $S = (n - 2) \times 180°$, the sum of the interior angles in a hexagon is:

A 1078° **B** 360° **C** 720° **D** 900° **E** 540°

8C **6** The quadrilateral with all sides equal, two pairs of opposite parallel sides and no right angles is:

A a kite **B** a trapezium **C** a parallelogram

D a rhombus **E** a square

8C **7** The value of a in the diagram is:


A 45

B 310

C 75


D 60

E 50




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

8E **8** The abbreviated reason for congruence in the two triangles shown is:




A AA **B** SAS **C** SSS
D AAS **E** RHS

8F **9** The scale factor in the two similar figures shown that enlarges the original figure to its image is:



A $\frac{2}{3}$ **B** 2 **C** 1.2
D 1.5 **E** 0.5


8G **10** The value of x in the diagram is:



A 6 **B** 9 **C** 10
D 8 **E** 7.5


Extended-response questions

1 In this complicated diagram there are two triangles and one quadrilateral.



a Name the types of triangles.
b Find the values of the pronumerals in alphabetical order. Give a reason at each step.

2 A buoy (E) is floating in the sea at some unknown distance from the beach, as shown. The points A, B, C and D are measured and marked out on the beach, as shown.



a Name the angle that is vertically opposite to $\angle ACB$.
b Explain, with reasons, why $\Delta ABC \,|||\, \Delta EDC$.
c Find the distance from the buoy to the beach (ED) to one decimal place.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

9

Algebraic techniques



Essential mathematics: why skills with algebra are important

Algebraic techniques simplify and solve complex problems in various fields.

Retail managers use algebra to manage inventory and decide on discounts. By analysing past sales data, they can predict periods when items will be popular, such as summer for ice creams. To prepare, they use a seasonal index for each month and adjust their orders based on expected demand. They also use algebra to determine the best time to put items on sale by analysing sales trends and calculating the break-even point where extra sales from a discount offset the lower price.

Music producers use algebra to design synthesizers and mix audio. Synthesizers generate sound waves using algebraic expressions like $A \times \sin(2\pi ft + \varphi)$, where A is amplitude, f is frequency, t is time, and φ is phase. These variables allow producers to shape sound precisely, creating different tones and effects. Algebraic techniques also help adjust volume levels, balance frequencies, and apply effects, crafting high-quality tracks.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025
Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.



In this chapter

9A Reviewing algebra (Consolidating)
9B Expanding binomial products
9C Expanding perfect squares
9D Forming a difference of two squares
9E Factorising algebraic expressions
9F Factorising the difference of two squares
9G Factorising by grouping in pairs
9H Factorising monic quadratic trinomials
9I Simplifying algebraic fractions: multiplication and division ★
9J Simplifying algebraic fractions: addition and subtraction ★

Victorian Curriculum 2.0

This chapter covers the following content descriptors in the Victorian Curriculum 2.0:

ALGEBRA

VC2M9A01, VC2M9A02, VC2M9A06

Please refer to the curriculum support documentation in the teacher resources for a full and comprehensive mapping of this chapter to the related curriculum content descriptors.

© VCAA

Online resources

A host of additional online resources are included as part of your Interactive Textbook, including HOTmaths content, video demonstrations of all worked examples, auto-marked quizzes and much more.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025
Photocopying is restricted under law and this material must not be transferred to another party.
Cambridge University Press

Warm-up quiz

1 Write down the coefficient of x in these expressions.

a $4x - 1$ **b** $3y + 6x$ **c** $3xy - 2x$ **d** $-4x^2 - 7x$

2 What is the constant term in these expressions?

a $5 - 2x$ **b** $-16a + 2$

c $5xy - 2x + 1$ **d** $101x - 6$

3 Evaluate the following if $a = 2$ and $b = -5$.

a ab **b** $-3b + 1$ **c** $4 - b + a$ **d** $5a^2b$

e $\sqrt{a^2}$ **f** $2b^2 - a$ **g** $5 - b^2$ **h** $a^2b - 2b^2$

4 Expand the following using $a(b + c) = ab + ac$.

a $2(x + 3)$ **b** $3(a - 5)$ **c** $4x(3 - 2y)$ **d** $-3(2b - 1)$

5 Write down the highest common factor of:

a 4 and 6 **b** 12 and 18 **c** $2x$ and $4x$

d $3xy$ and $9y$ **e** $10x$ and $15x^2$ **f** $3a^2b$ and $4ab^2$

6 Factorise by taking out the highest common factor.

a $2a + 6$ **b** $3x + 12y$ **c** $5x^2 - 15x$ **d** $4m - 6mn$

7 Add or subtract these fractions. You will need a common denominator for parts **c** and **d**.

a $\frac{3}{7} + \frac{2}{7}$ **b** $\frac{4}{9} - \frac{5}{9}$ **c** $\frac{1}{2} + \frac{2}{3}$ **d** $\frac{3}{8} - \frac{1}{2}$

8 Multiply or divide these fractions.


a $\frac{2}{3} \times \frac{4}{5}$ **b** $\frac{3}{4} \times \frac{2}{3}$ **c** $\frac{7}{14} \times \frac{2}{3}$ **d** $\frac{6}{11} \times \frac{22}{12}$

e $\frac{2}{3} \div \frac{1}{3}$ **f** $\frac{7}{8} \div \frac{14}{24}$ **g** $\frac{3}{2} \div \frac{4}{3}$ **h** $\frac{7}{9} \div \frac{4}{3}$

9 Expand and simplify.

a $3(x - 1) + 5$ **b** $4(1 - x) + 5x$ **c** $-2(5 + x) - x$

10 Write two expressions for the area of this rectangle, one with brackets and one without.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

9A Reviewing algebra

CONSOLIDATING

Learning intentions

- To know the names of the parts of an algebraic expression
- To be able to form algebraic expressions from simple word phrases
- To know that only like terms can be combined under addition and subtraction
- To be able to simplify algebraic expressions using the four operations: +, −, × and ÷
- To be able to expand expressions involving brackets
- To be able to evaluate expressions by substituting given values

Key vocabulary: expression, pronumeral, variable, term, like terms, constant term, coefficient, distributive law, evaluate

A high level of skill in algebra is required to solve more complex mathematical problems. Skills include adding, subtracting, multiplying and dividing algebraic expressions, as well as expanding and factorising expressions.

In this section we will review some basic concepts in algebra.

Lesson starter: Vocab review

Consider the expression $3x^2 + 4xy - 2 - xy$.

- How many terms are given in the expression?
- What letters are used as pronumerals?
- What is the coefficient of x^2?
- What is the constant term?
- Are there any like terms?
- Can the expression be simplified? If so, how?
- What would be the value of the expression if $x = 2$ and $y = -1$?

Key ideas

- An **expression** is a combination of numbers and pronumerals connected by mathematical operations.
  - A **term** is part of an expression with numbers and pronumerals connected only by multiplication and division
  - A **coefficient** is the number part in front of a term
  - A **constant term** is a term that does not contain any pronumerals
  - This is an example of a 3-term expression

$$7y^2 + 2xy - 4$$

coefficient of y^2 (7) constant term (−4)

- **Like terms** have the same pronumeral part.
  - They can be collected using addition and subtraction.
    For example: $5a - 7a = -2a$ and $3xy^2 + 2y^2x = 5xy^2$
- For multiplication and division, the symbols × and ÷ are not usually shown.
  For example: $(-3 \times a \times b) \times (2 \times b) = -3ab \times 2b = -6ab^2$

$$14a^2b \div (7ab) = \frac{14a^2b}{7ab} = 2a$$

- The **distributive law** is used to expand brackets.

  For example: $-3(x - 2) = -3x + 6$

$$\begin{aligned} 4x + 2(1 - x) &= 4x + 2 - 2x \\ &= 2x + 2 \end{aligned}$$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

9A

- To evaluate an expression, substitute a value for each pronumeral and simplify.
  For example: if $a = 2$ and $b = -3$ then

$$\begin{aligned} b^2 - a &= (-3)^2 - 2 \\ &= 9 - 2 \\ &= 7 \end{aligned}$$

Exercise 9A

Understanding

1–3 | 3

1 Write expressions for each of the following.

a The cost of:

i x movie passes at \$11 each

ii n apples at 50 cents each

b The number of tickets purchased for:

i x adults and y children

ii b boys and g girls

c The cost of hiring an electrician for n hours if the callout fee is \$50 and the cost of labour is \$60 per hour

2 Consider the expression $5x - 2xy - y^2 - 7$.

a What is the coefficient of x?

b What is the coefficient of xy?

c What is the constant term?

d Are there any like terms?

e Evaluate the expression if $x = 1$ and $y = 2$.

f Evaluate the expression if $x = 2$ and $y = -1$.



Hint for Q2: Remember to include the negative sign in the coefficient of xy.

3 Write expressions for the following.

a The sum of 5 and $2x$.

b 7 less than $4a$.

c 1 less than the square of y.

d The product of x and z.

e The sum of the squares of a and b.

f The square root of 8 more than x.



Hint for Q3: 'Product' means multiply.

Fluency

4–6(½) | 4–6(½)

Example 1 Collecting like terms

Collect like terms to simplify the following.

a $5y - x + 2y$

b $4a^2b + ab - ba^2$

Solution

a $5y - x + 2y = 7y - x$

b $4a^2b + ab - ba^2 = 3a^2b + ab$

Explanation

$5y$ and $2y$ are like terms.

$4a^2b$ and ba^2 (or a^2b) are like terms and $4 - 1 = 3$.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Now you try

Collect like terms to simplify the following.

a $7m + 2n - 5m - 3n$ **b** $2x^2y - xy - 4yx^2$

4 Simplify by collecting like terms.

a $3a - 4a$	**b** $4x + 7x$	**c** $6ab + 2ab$
d $5xy - 9xy$	**e** $3a - 1 - 2a$	**f** $7y + 2x - 11y$
g $5x - y - 3y + x$	**h** $ab + ba$	**i** $a^2b - 5a^2b - 3$
j $11xy - 14yx$	**k** $10r^2a + 2ar^2$	**l** $9st^2 - st - 8t^2s$



Hint for Q4: Only collect 'like' terms. Note that $ab = ba$.

Example 2 Multiplying and dividing terms

Simplify these expressions.

a $-4a \times 2ab$ **b** $21a^2b \div (3abc)$

Solution

a $-4a \times 2ab = -8a^2b$

b $21a^2b \div (3abc) = \frac{21a^2b}{3abc}$

$= \frac{7a}{c}$

Explanation

$-4 \times 2 = -8$ and $a \times a = a^2$

Write as a fraction then cancel where possible.

Note: $\frac{a^2}{a} = \frac{a \times \cancel{a}^1}{\cancel{a}^1} = a$

Now you try

Simplify these expressions.

a $5xy \times (-3y)$ **b** $6a^2b \div (12ab)$

5 Simplify the following.

a $5 \times 3x$	**b** $7a \times 2$	**c** $-4x \times 2y$
d $6a \times (-7a)$	**e** $-3ab \times b$	**f** $-6a^2b \times 2b$
g $4x \div 2$	**h** $7a \div a$	**i** $12a \div (4a)$
j $22a^2 \div (11a)$	**k** $40a \div (4a^2)$	**l** $100x^2y \div (25xy)$
m $\frac{7ab^2}{7b}$	**n** $\frac{48a^2bc}{16abc^2}$	**o** $\frac{12xy}{36xy^2}$

Example 3 Expanding brackets

Use the distributive law to expand and simplify:

a $-x(2 + x)$ **b** $4x - 2(x - 1)$

Solution

a $-x(2 + x) = -2x - x^2$

b $4x - 2(x - 1) = 4x - 2x + 2$

$= 2x + 2$

Explanation

$-x \times 2 = -2x$ and $-x \times x = -x^2$

First expand the brackets then collect like terms.

Note: $-2 \times (-1) = 2$, not -2.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

9A

Now you try

Use the distributive law to expand and simplify:

a $-3(x+2)$ **b** $-4x(1-x)+2x^2$

6 Use the distributive law to expand and simplify:

a $3(x+2)$ **b** $2(4+x)$
c $-3(x+4)$ **d** $-6(x+1)$
e $-2(x-3)$ **f** $-x(x+1)$
g $-3y(2-3y)$ **h** $-7a(a-b)$
i $3+2(x-1)$ **j** $3x+4(1-x)$
k $3(x+1)-7x$ **l** $4(x+2)-2(x+1)$
m $7(x-3)-3(x-4)$ **n** $-4(1-x)-21x+1$
o $-6(2-x)-7(4-x)$ **p** $-2(3-x)-(5-x)$




Hint for Q6: Remember: a negative times a negative is a positive.

Problem-solving and reasoning


7, 8(½), 9 | 8(½), 9–11

7 For these shapes, write expressions for **i** perimeter and **ii** area.


a



b



c



8 Evaluate the following if $a=3$, $b=-2$ and $c=-4$.

a $2a+b$ **b** abc **c** $2a \div (3b)$
d $-3c+b$ **e** a^2+b^2 **f** $c^2 \div b$
g $\dfrac{a^2+c^2}{2}$ **h** $\dfrac{a-c}{7}$ **i** $\dfrac{1}{a}(b+c)$
j a^3-b^3 **k** $2b^3-c$ **l** $\sqrt{a^2+b^2}$



Hint for Q8: Substitute pronumeral values and work out the answer.

9 You can hire a sports car for an upfront fee of $100 plus $80 per hour after that.

a What is the cost of hiring a sports car for:
i 2 hours? **ii** 9 hours?

b Write an expression for the cost of hiring a sports car for n hours.




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10 Find the area of these shapes in expanded form. All angles are $90°$.


a



b



c



11 Identify the errors, then correct to find the answer.

a $-3(x-1)$
$= -3x - 3$

b $\frac{3ab}{6b} = 2a$

c $\frac{4a}{2a^2} = 2a$

d $3(x+2) - 3(x-1)$
$= 3x + 6 - 3x - 3$
$= 3$

Jake vs Lucas — 12

12 Jake and Lucas operate separate computer consulting services.

- Jake charges $\$60$ per hour.
- Lucas charges a $\$40$ callout fee plus $\$50$ per hour after that.

a What is the cost of hiring Jake for:

i 2 hours? **ii** 10 hours?

b What is the cost of hiring Lucas for:

i 2 hours? **ii** 10 hours?

c Write an expression for the cost of hiring Jake for n hours.

d Write an expression for the cost of hiring Lucas for n hours.

e After what time is the cost of hiring Jake or Lucas the same?



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

9B Expanding binomial products

Learning intentions

- To understand the distributive law for expanding binomial products
- To be able to expand and simplify binomial products

Key vocabulary: binomial product, distributive law, expand

A binomial is an expression with two terms such as $x + 5$ or $x^2 + 3$. You will recall from the previous section that we looked at the product of a single term with a binomial expression; e.g. $2(x - 3)$ or $x(3x - 1)$.


The product of two binomial expressions can also be expanded using the distributive law. This involves multiplying every term in one expression by every term in the other expression.



Expanding the product of two expressions can be applied to problems involving the expansion of rectangular areas, such as a farmer's paddocks.

Lesson starter: Rectangular expansions

If $(x + 1)$ and $(x + 2)$ are the side lengths of a rectangle as shown, the total area can be found as an expression in two different ways.



- Write an expression for the total area of the rectangle using length = $(x + 2)$ and width = $(x + 1)$.
- Now find the area of each of the four parts of the rectangle and combine to give an expression for the total area.
- Compare your two expressions above and complete this equation:

$$(x + 2)(___) = x^2 + ___ + ___.$$

- Can you explain a method for expanding the left-hand side to give the right-hand side?

Key ideas

- A binomial is an expression with two terms.
- Expanding **binomial products** uses the distributive law.

$$\begin{aligned}(a + b)(c + d) &= a(c + d) + b(c + d) \\ &= ac + ad + bc + bd\end{aligned}$$

- Diagrammatically

$$(a + b)(c + d) = ac + ad + bc + bd$$

For example: $(x + 1)(x + 5) = x^2 + 5x + x + 5$

$= x^2 + 6x + 5$

	a	b
c	ac	bc
d	ad	bd

	x	5
x	x^2	$5x$
1	$1x$	5


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Exercise 9B

Understanding

1–3 | 2, 3

1 The given diagram shows the area $(x+2)(x+3)$.




a Write down an expression for the area of each of the four regions inside the rectangle.

b Copy and complete: $(x+2)(x+3) = _____ + 3x + _____ + 6$

$= _____ + 5x + _____$

2 The given diagram shows the area $(2x+3)(x+1)$.



a Write down an expression for the area of each of the four regions inside the rectangle.

b Copy and complete: $(2x+3)(_____) = 2x^2 + _____ + 3x + _____$

$= _____ + _____ + _____$

3 Copy and complete these expansions

a $(x+1)(x+5) = _____ + 5x + _____ + 5$

$= _____ + 6x + _____$

b $(x-3)(x+2) = _____ + _____ - 3x - _____$

$= _____ - x - _____$

c $(3x-2)(7x+2) = _____ + 6x - _____ - _____$

$= _____ - _____ - _____$

d $(4x-1)(3x-4) = _____ - _____ - 3x + _____$

$= _____ - 19x + _____$



Hint for Q3: The product is negative if there are opposite signs $(+, -)$ or $(-, +)$ and positive if they are the same sign $(+, +)$ or $(-, -)$.

Fluency

4–5(½) | 4–5(½)

Example 4 Expanding binomial products

Expand $(x+3)(x+5)$.

Solution

$(x+3)(x+5) = x^2 + 5x + 3x + 15$

$= x^2 + 8x + 15$

Explanation

Use the distributive law to expand the brackets and then collect the like terms $5x$ and $3x$.

Now you try

Expand $(x+2)(x+9)$.

4 Expand the following.

a $(x+2)(x+5)$ **b** $(b+3)(b+4)$

c $(t+8)(t+7)$ **d** $(p+6)(p+6)$

e $(x+9)(x+6)$ **f** $(d+15)(d+4)$

g $(a+1)(a+7)$ **h** $(y+10)(y+2)$

i $(m+4)(m+12)$



Hint for Q4: First expand with the distributive law then collect the two like terms.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

9B

Example 5 Expanding binomial products involving subtraction signs

Expand the following.

a $(x-4)(x+7)$ **b** $(2x-1)(x-6)$ **c** $(5x-2)(3x+7)$

Solution

a $(x-4)(x+7) = x^2+7x-4x-28$
$= x^2+3x-28$

b $(2x-1)(x-6) = 2x^2-12x-x+6$
$= 2x^2-13x+6$

c $(5x-2)(3x+7) = 15x^2+35x-6x-14$
$= 15x^2+29x-14$

Explanation

After expanding to get the four terms, collect the like terms $7x$ and $-4x$.
Note: $x \times 7 = 7x$ and $-4 \times x = -4x$.

Remember: $2x \times x = 2x^2$ and $-1 \times (-6) = 6$.

Recall: $5x \times 3x = 5 \times 3 \times x \times x = 15x^2$.

Now you try

Expand the following.

a $(x+3)(x-6)$ **b** $(3x-2)(x-4)$ **c** $(6x+5)(3x-4)$

5 Expand the following.


a $(x+3)(x-4)$ **b** $(x+5)(x-2)$ **c** $(x+4)(x-8)$
d $(x-6)(x+2)$ **e** $(x-1)(x+10)$ **f** $(x-7)(x+9)$
g $(x-2)(x+7)$ **h** $(x-1)(x-2)$ **i** $(x-4)(x-5)$
j $(4x+3)(2x+5)$ **k** $(3x+2)(2x+1)$ **l** $(3x+1)(5x+4)$
m $(2x-3)(3x+5)$ **n** $(8x-3)(3x+4)$ **o** $(3x-2)(2x+1)$
p $(5x+2)(2x-7)$ **q** $(2x+3)(3x-2)$ **r** $(4x+1)(4x-5)$
s $(3x-2)(6x-5)$ **t** $(5x-2)(3x-1)$ **u** $(7x-3)(3x-4)$



Hint for Q5: First expand with the distributive law then collect the two like terms.

Problem-solving and reasoning 6–8, 9(½) 7–11

6 A 10 m by 7 m rectangular factory shed is expanded by x metres on two sides.




a Write expressions for:
i the new length of the shed (horizontal length)
ii the new width of the shed (vertical length)

b Using your results from part **a**, expand brackets to find an expression for the new total area.

c What would be the new area if $x = 2$?

7 A rectangular room in a house with dimensions 4 m by 5 m is to be extended. Both the length and width are to be increased by x m.



a Find an expanded expression for the area of the new room.


b **i** If $x = 3$, find the area of the new room.
ii By how much has the area increased?



Hint for Q7: First label the length and width of the new room.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

8 A rectangular trampoline of length 3 m and width 2 m is to be surrounded by padding of width x metres.





a Write expressions for:
- **i** the total length of the trampoline and padding
- **ii** the total width of the trampoline and padding

b Find an expression for the total area by expanding brackets.

c What would be the total area if $x = 1$?



Hint for Q8: Don't forget to count the x on both sides.

9 Write the missing terms in these expansions.

a $(x+2)(x+___) = x^2+5x+6$
b $(x+___)(x+5) = x^2+7x+10$
c $(x+1)(x+___) = x^2+7x+___$
d $(x+___)(x+9) = x^2+11x+___$
e $(x+3)(x-___) = x^2+x-___$
f $(x-5)(x+___) = x^2-2x-___$

10 Expand these binomial products.

a $(a+b)(a+c)$
b $(a-b)(a+c)$
c $(b-a)(a+c)$
d $(2x+y)(x-2y)$
e $(2a+b)(a-b)$
f $(3x-y)(2x+y)$


11 A picture frame 5 cm wide has a length that is twice the width, x cm.

a Find an expression for the total area of the frame and picture.

b Find an expression in expanded form for the area of the picture only.




Hint for Q11: Length of picture = $2x - 5 - 5 = 2x - 10$



Paving the pool — 12

12 The outside edge of a path around a rectangular swimming pool is 15 m long and 10 m wide. The path is x metres wide.



a Write expressions for:
- **i** the length of the pool
- **ii** the width of the pool

b Find an expression for the area of the pool in expanded form.

c Find the area of the pool if $x = 2$.

d What value of x makes the pool area 50 m^2? Use trial and error.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

9C Expanding perfect squares

Learning intentions

- To be able to recognise a perfect square
- To understand that expressions that are perfect squares can be expanded
- To be able to expand and simplify perfect squares

Key vocabulary: perfect square, distributive law, expand

A special type of binomial product involves perfect squares. Examples of perfect squares are $2^2 = 4$, $15^2 = 225$, x^2 and $(a+b)^2$. To expand $(a+b)^2$ we multiply $(a+b)$ by $(a+b)$ and use the distributive law:

$$\begin{aligned}(a+b)^2 &= (a+b)(a+b)\\ &= a(a+b) + b(a+b)\\ &= a^2 + ab + ba + b^2\\ &= a^2 + 2ab + b^2\end{aligned}$$

A similar result is obtained for the square of $(a-b)$:

$$\begin{aligned}(a-b)^2 &= (a-b)(a-b)\\ &= a(a-b) - b(a-b)\\ &= a^2 - ab - ba + b^2\\ &= a^2 - 2ab + b^2\end{aligned}$$

Lesson starter: Seeing the pattern

Using $(a+b)(c+d) = ac + ad + bc + bd$, expand and simplify the binomial products below.

$(x+1)(x+1) = x^2 + x + x + 1$
$=$

$(x+3)(x+3) =$
$=$

$(x-5)(x-5) =$
$=$

$(x-7)(x-7) =$
$=$

- Describe the patterns you see in the expansions above.
- Generalise your observations by completing the following expansions.

$(a+b)(a+b) = a^2 + ___ + ___ + ___$
$= a^2 + ___ + ___$

$(a-b)(a-b) =$
$=$

Key ideas

- $3^2 = 9$, a^2, $(2y)^2$, $(x-1)^2$ and $(3-2y)^2$ are all examples of **perfect squares**. They are expressions that can be written as a single square.
- Expanding perfect squares:

$$\begin{aligned}(a+b)^2 &= (a+b)(a+b)\\ &= a(a+b) + b(a+b)\\ &= a^2 + ab + ba + b^2\\ &= a^2 + 2ab + b^2\end{aligned}$$

$$\begin{aligned}(a-b)^2 &= (a-b)(a-b)\\ &= a(a-b) - b(a-b)\\ &= a^2 - ab - ba + b^2\\ &= a^2 - 2ab + b^2\end{aligned}$$


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Exercise 9C

Understanding

1–3 | 3

1 The side lengths of this square are $(x + 3)$ units.



a What are the areas of each of the four regions? Write expressions each time.

b Add up all the area expressions to find an expression for the total area.

c Complete the following: $(x + 3)(x + 3) = x^2 + 3x + __ + __$

$= x^2 + 6x + __$

2 Complete these expansions.

a $(x + 4)(x + 4) = x^2 + 4x + __ + __$

$= _____$

b $(x + 5)(x + 5) = x^2 + 5x + __ + __$

$= _____$

c $(x - 2)(x - 2) = x^2 - 2x - __ + __$

$= _____$

d $(x - 7)(x - 7) = x^2 - 7x - __ + __$

$= _____$



Hint for Q2: Expand using the distributive law: $(a + b)(c + d) = ac + ad + bc + bd$

3 **a** Substitute the given value of b into $x^2 + 2bx + b^2$ and simplify.

i $b = 3$ **ii** $b = 11$

b Substitute the given value of b into $x^2 - 2bx + b^2$ and simplify.

i $b = 2$ **ii** $b = 9$

Fluency

4–5(½) | 4–6(½)

Example 6 Expanding perfect squares

Expand each of the following.

a $(x + 3)^2$ **b** $(x - 2)^2$

Solution

a $(x + 3)^2 = (x + 3)(x + 3)$

$= x^2 + 3x + 3x + 9$

$= x^2 + 6x + 9$

Alternative solution:

$(x + 3)^2 = x^2 + 2 \times x \times 3 + 3^2$

$= x^2 + 6x + 9$

Explanation

Write in expanded form.

Use the distributive law.

Collect like terms.

Expand using $(a + b)^2 = a^2 + 2ab + b^2$, where $a = x$ and $b = 3$.

b $(x - 2)^2 = (x - 2)(x - 2)$

$= x^2 - 2x - 2x + 4$

$= x^2 - 4x + 4$

Alternative solution:

$(x - 2)^2 = x^2 - 2 \times x \times 2 + 2^2$

$= x^2 - 4x + 4$

Write in expanded form.

Use the distributive law.

Collect like terms.

Expand using $(a - b)^2 = a^2 - 2ab + b^2$ where $a = x$ and $b = 2$.

Now you try

Expand each of the following.

a $(x + 8)^2$ **b** $(x - 5)^2$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

9C

4 Expand each of the following perfect squares.

a $(x+1)^2$	**b** $(x+3)^2$	**c** $(x+2)^2$	**d** $(x+5)^2$
e $(x+4)^2$	**f** $(x+9)^2$	**g** $(x+7)^2$	**h** $(x+10)^2$
i $(x-2)^2$	**j** $(x-6)^2$	**k** $(x-1)^2$	**l** $(x-3)^2$
m $(x-9)^2$	**n** $(x-7)^2$	**o** $(x-4)^2$	**p** $(x-12)^2$



Hint for Q4: Recall
$(a+b)^2 = (a+b)(a+b)$
$(a-b)^2 = (a-b)(a-b)$

Example 7 Expanding more perfect squares

Expand $(2x+3)^2$.

Solution

$(2x+3)^2 = (2x+3)(2x+3)$

$= 4x^2 + 6x + 6x + 9$

$= 4x^2 + 12x + 9$

Alternative solution:

$(2x+3)^2 = (2x)^2 + 2 \times 2x \times 3 + 3^2$

$= 4x^2 + 12x + 9$

Explanation

Write in expanded form.

Use the distributive law.

Collect like terms.

Expand using $(a+b)^2 = a^2 + 2ab + b^2$

where $a = 2x$ and $b = 3$.

Recall $(2x)^2 = 2x \times 2x = 4x^2$.

Now you try

Expand $(5x-3)^2$.

5 Expand each of the following perfect squares.

a $(2x+1)^2$	**b** $(2x+5)^2$	**c** $(3x+2)^2$	**d** $(3x+1)^2$
e $(5x+2)^2$	**f** $(4x+3)^2$	**g** $(7+2x)^2$	**h** $(5+3x)^2$
i $(2x-3)^2$	**j** $(3x-1)^2$	**k** $(4x-5)^2$	**l** $(2x-9)^2$



Hint for Q5:
$(2x)^2 = 2x \times 2x$
$= 4x^2$

6 Expand each of the following perfect squares.


a $(3-x)^2$	**b** $(5-x)^2$	**c** $(1-x)^2$	**d** $(6-x)^2$
e $(11-x)^2$	**f** $(4-x)^2$	**g** $(7-x)^2$	**h** $(12-x)^2$
i $(8-2x)^2$	**j** $(2-3x)^2$	**k** $(9-2x)^2$	**l** $(10-4x)^2$



Hint for Q6: $-x \times (-x) = x^2$

Problem-solving and reasoning 7, 8 8–10

7 A farmer extends a 12 m square sheep pen on two sides by x metres.





a Write the expression for the side length of the new pen.

b Write an expression for the area of the new pen and expand this expression.

c Use your result to find the new area if:


i $x = 2$ **ii** $x = 5$



Hint for Q7: An example of an expression is $10 - x$ or $x^2 - 20x + 100$.


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

8 A child at pre-school cuts off a strip of width x cm from two sides of a square piece of paper of side length 10 cm.



- **a** Write an expression for the new side length of the remaining paper.
- **b** Find the area of the new piece of paper in expanded form.
- **c** Use your result to find the new area if:
  - **i** $x = 2$ **ii** $x = 6$
- **d** What value of x makes the new area one quarter of the original?

9 A square piece of tin of side length 20 cm has four squares of side length x cm removed from each corner. The sides are folded up to form a tray. The centre square forms the tray base.




- **a** Write an expression for the side length of the base of the tray.
- **b** Write an expression for the area of the base of the tray. Expand your answer.
- **c** Find the area of the tray base if $x = 3$.
- **d** Find the volume of the tray if $x = 3$.



Hint for Q9: For part **d**, what is the height of the tray?


10 A square of side length b is removed from a square of side length a.



- **a** Using subtraction, write down an expression for the remaining area.
- **b** Write expressions for the area of the regions in expanded form:
  - **i** A **ii** B **iii** C
- **c** Add all the expressions from part **b** to see if you get your answer from part **a**.

Arranging tennis courts — 11

11 Four tennis courts are arranged as shown, with a square storage space in the centre. Each court area has the same dimensions, $a \times b$.



- **a** Write an expression for the side length of the total area.
- **b** Write an expression for the area of the total area.
- **c** Write an expression for the side length of the inside storage space.
- **d** Write an expression for the area of the storage space in expanded form.
- **e** Subtract your answer to part **d** from your answer to part **b** to find the area of the four courts.
- **f** Find the area of one court. Does your answer confirm that your answer to part **e** is correct?



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

9D Forming a difference of two squares

Learning intentions

- To understand how a difference of two squares is formed
- To be able to expand and simplify to form a difference of two squares

Key vocabulary: difference of two squares, distributive law, expand

Another type of expansion deals with the product of the sum and difference of the same two terms. The result is the difference of two squares:

$$\begin{aligned}(a+b)(a-b) &= a(a-b)+b(a-b) \\ &= a^2-ab+ba-b^2 \\ &= a^2-b^2\end{aligned} \quad \text{or} \quad \begin{aligned}(a+b)(a-b) &= a^2-ab+ba-b^2 \\ &= a^2-b^2\end{aligned}$$

Lesson starter: How is 16×14 a difference of two squares?

Using the fact that $15^2 = 225$, follow these steps to show how 16×14 can be calculated mentally using a difference of two squares.

- $16 \times 14 = (15 + ___) \times (15 - ___)$ Rewrite.
  $= 15^2 - ___ + ___ - ___$ Expand.
  $= ___ - 1$ Simplify.
  $= ___$ Evaluate.
- Now try this technique on 17×13 and 19×21.

Key ideas

- A **difference of two squares** is formed when one square is subtracted from another.
- This is formed when $(a+b)(a-b)$ is expanded and simplified.

$$\begin{aligned}(a+b)(a-b) &= a^2-ab+ba-b^2 \\ &= a^2-b^2\end{aligned}$$

$(a-b)(a+b)$ also expands to a^2-b^2
The result is a difference of two squares.

Exercise 9D

Understanding

1–3 | 2, 3

1 Why do the two middle terms in an expansion of $(x+a)(x-a)$ (i.e. $x^2+ax-xa-a^2$) cancel out?

2 Decide whether each of the following shows a single perfect square or a difference of two squares.

a 4^2 **b** 7^2-3^2 **c** a^2-b^2 **d** x^2

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3 Complete these expansions.

a $(x+4)(x-4) = x^2 - 4x + ___ - ___$
$= _____$

b $(2x-1)(2x+1) = 4x^2 + ___ - ___ - ___$
$= _____$



Hint for Q3:
$(a+b)(a-b) = a^2 - ab + ab - b^2$
$= a^2 - b^2$

Fluency

4-5(½) 4-5(½)

Example 8 Forming a difference of two squares

Expand and simplify the following.

a $(x+2)(x-2)$ **b** $(x-7)(x+7)$

Solution

a $(x+2)(x-2) = x^2 - 2x + 2x - 4$
$= x^2 - 4$

Alternate solution:
$(x+2)(x-2) = (x)^2 - (2)^2$
$= x^2 - 4$

Explanation

Expand using the distributive law.
$-2x + 2x = 0$.

$(a+b)(a-b) = a^2 - b^2$. Here $a = x$ and $b = 2$.

b $(x-7)(x+7) = x^2 + 7x - 7x - 49$
$= x^2 - 49$

Alternate solution:
$(x-7)(x+7) = (x)^2 - (7)^2$
$= x^2 - 49$

Expand, then note that $7x - 7x = 0$.

$(a-b)(a+b) = a^2 - b^2$, with $a = x$ and $b = 7$.

Now you try

Expand and simplify the following.

a $(x+5)(x-5)$ **b** $(x-15)(x+15)$

4 Expand and simplify the following to form a difference of two squares.

a $(x+1)(x-1)$ **b** $(x+3)(x-3)$
c $(x+8)(x-8)$ **d** $(x+4)(x-4)$
e $(x+12)(x-12)$ **f** $(x+11)(x-11)$
g $(x-9)(x+9)$ **h** $(x-5)(x+5)$
i $(x-6)(x+6)$ **j** $(5-x)(5+x)$
k $(2-x)(2+x)$ **l** $(7-x)(7+x)$



Hint for Q4: Recall
$(a+b)(a-b) = a^2 - b^2$

Example 9 Forming more differences of two squares

Expand and simplify $(3x-2y)(3x+2y)$.

Solution

$(3x-2y)(3x+2y) = 9x^2 + 6xy - 6xy - 4y^2$
$= 9x^2 - 4y^2$

Explanation

Expand using the distributive law.
$6xy - 6xy = 0$.

Continued on next page

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

9D

Alternate solution:
$(3x - 2y)(3x + 2y) = (3x)^2 - (2y)^2$
$= 9x^2 - 4y^2$

$(a + b)(a - b) = a^2 - b^2$, with $a = 3x$ and $b = 2y$ here. Recall that $(3x)^2 = 3x \times 3x$.

Now you try

Expand and simplify $(12x - 5y)(12x + 5y)$.

5 Expand and simplify the following.

a $(3x - 2)(3x + 2)$ **b** $(5x - 4)(5x + 4)$ **c** $(4x - 3)(4x + 3)$
d $(7x - 3y)(7x + 3y)$ **e** $(9x - 5y)(9x + 5y)$ **f** $(11x - y)(11x + y)$
g $(8x + 2y)(8x - 2y)$ **h** $(10x - 9y)(10x + 9y)$ **i** $(7x - 5y)(7x + 5y)$
j $(6x - 11y)(6x + 11y)$ **k** $(8x - 3y)(8x + 3y)$ **l** $(9x - 4y)(9x + 4y)$



Hint for Q5:
$(3x)^2 = 3x \times 3x$
$= 9x^2$

Problem-solving and reasoning

6(½), 7 | 6(½), 8

6 To calculate 21×19, here is a method using the difference of two squares.
$21 \times 19 = (20 + 1) \times (20 - 1)$
$= 20^2 - 1^2$
$= 400 - 1$
$= 399$

Use this technique to evaluate the following (mentally if you can).

a 31×29 (Note: $30^2 = 900$) **b** 41×39 (Note: $40^2 = 1600$) **c** 26×24 (Note: $25^2 = 625$)
d 51×49 (Note: $50^2 = 2500$) **e** 22×18 (Use $20^2 = 400$) **f** 23×17 (Use $20^2 = 400$)
g 35×25 (Use $30^2 = 900$) **h** 54×46 (Use $50^2 = 2500$)



7 Lara is x years old and her two best friends are $(x - 2)$ and $(x + 2)$ years old.

a Write an expression for:
i the square of Lara's age
ii the product of the ages of Lara's best friends

Hint for Q7: Expand in part **a ii**.

b Are the answers from parts **a i** and **ii** equal? If not, by how much do they differ?

8 A square of side length x has one side reduced by 1 unit and the other increased by 1 unit.

a Find an expanded expression for the area of the resulting rectangle.
b Is the area of the original square the same as the area of the resulting rectangle? Explain why/why not.




Hint for Q8: Determine expressions for the length and width of the rectangle first.

Classroom renovation

— | 9

9 A square classroom is to be shortened on the south side by 3 m and extended on the east side by 3 m. The original side length of the classroom was x m.

a Write expressions for:
i the original area of the classroom
ii the new length of the classroom
iii the new width of the classroom



b Expand to find an expression for the new area of the classroom.

c Is the new area the same as the original area? If not, by how much do they differ?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


9E Factorising algebraic expressions

Learning intentions

- To understand that factorising and expanding are reverse processes
- To be able to identify the highest common factor
- To know the form of a factorised expression
- To be able to factorise algebraic expressions involving a common factor

Key vocabulary: highest common factor, factorise


The process of factorisation is a key step in the simplification of many algebraic expressions and in the solution of equations. It is the reverse process of expansion and involves writing an expression as a product of its factors.



Lesson starter: Factorised areas


Here is a rectangle of length $(x + 3)$ and width x.

- Write an expression for the total area using the given length and width.
- Write an expression for the total area by adding up the area of the two smaller regions.
- Are your two expressions equivalent? How could you work from your second expression (expanded) to the first expression (factorised)?



Key ideas

- To **factorise** means to write an expression as a product of its factors.
- When factorising expressions with common factors, take out the highest common factor (HCF). The HCF could be:
  - a number
    For example: $2x + 10 = 2(x + 5)$
  - a pronumeral
    For example: $x^2 + 5x = x(x + 5)$
  - the product of numbers and pronumerals
    For example: $2x^2 + 10x = 2x(x + 5)$



- A factorised expression can be checked by using expansion.
  - For example: $2x(x + 5) = 2x^2 + 10x$.

Essential Mathematics Core for the Victorian Curriculum Year 9 ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Exercise 9E

Understanding

1–3 | 3

1 Write down the highest common factor (HCF) of these pairs of numbers.

a 8, 12 **b** 10, 20 **c** 5, 60 **d** 24, 30

2 Write down the missing factor.

a $5 \times ___ = 5x$
b $7 \times ___ = 7x$
c $2a \times ___ = 2a^2$
d $5a \times ___ = 10a^2$
e $___ \times 3y = -6y^2$
f $___ \times 12x = -36x^2$

3 **a** Write down the missing factor in each part.

i $___(x^2 + 2x) = 6x^2 + 12x$
ii $___(2x + 4) = 6x^2 + 12x$
iii $___(x + 2) = 6x^2 + 12x$

b Which equation above uses the HCF of $6x^2$ and $12x$?

Fluency

4–6(½) | 4–6(½)

Example 10 Finding the highest common factor (HCF)

Determine the HCF of the following.

a $6a$ and $8ab$ **b** $3x^2$ and $6xy$

Solution

a HCF of $6a$ and $8ab$ is $2a$

b HCF of $3x^2$ and $6xy$ is $3x$

Explanation

HCF of 6 and 8 is 2.
HCF of a and ab is a.

HCF of 3 and 6 is 3.
HCF of x^2 and xy is x.

Now you try

Determine the HCF of the following.

a $10ab$ and $5b$ **b** $7xy^2$ and $21xy$

4 Determine the HCF of the following.

a $6x$ and $14xy$ **b** $12a$ and $18a$
c $10m$ and 4 **d** $12y$ and 8
e $15t$ and $6s$ **f** 15 and p
g $9x$ and $24xy$ **h** $6n$ and $21mn$
i $10y$ and $2y$ **j** $8x^2$ and $14x$
k $4x^2y$ and $18xy$ **l** $5ab^2$ and $15a^2b$



Hint for Q4: HCF stands for highest common factor.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Example 11 Factorising expressions

Factorise the following.

a $4x + 12$ **b** $10y - 25y^2$

Solution

a $4x + 12 = 4(x + 3)$

Explanation

4 is the HCF of $4x$ and 12.
Place 4 in front of the brackets and divide each term by 4:
$4x \div 4 = x$ and $12 \div 4 = 3$.
Check your answer using expansion.

b $10y - 25y^2 = 5y(2 - 5y)$

The HCF of $10y$ and $25y^2$ is $5y$. Place $5y$ in front of the brackets and divide each term by $5y$.

Now you try

Factorise the following.

a $15a + 20$ **b** $2x^2 - 6x$

5 Factorise the following.

a $7x + 7$	**b** $3x + 3$	**c** $4x - 4$	**d** $5x - 5$
e $4 + 8y$	**f** $10 + 5a$	**g** $3 - 9b$	**h** $6 - 2x$
i $12a + 3b$	**j** $6m + 6n$	**k** $10x - 8y$	**l** $4a - 20b$
m $x^2 + 2x$	**n** $a^2 - 4a$	**o** $y^2 - 7y$	**p** $x - x^2$
q $3p^2 + 3p$	**r** $8x - 8x^2$	**s** $4b^2 + 12b$	**t** $6y - 10y^2$



Hint for Q5: Always take out the highest common factor (HCF).

Example 12 Taking out the common negative sign

Factorise $-8x^2 - 12x$.

Solution

$-8x^2 - 12x = -4x(2x + 3)$

Explanation

The HCF of the terms is $-4x$, including the common negative. Place the factor in front of the brackets and divide each term by $-4x$.
Note: $-8x^2 \div (-4x) = 2x$ and $-12x \div (-4x) = 3$.

Now you try

Factorise $-11a - 22a^2$.

6 Factorise the following by including the negative sign in the common factor.

a $-8x - 4$	**b** $-4x - 2$		
c $-10x - 5y$	**d** $-7a - 14b$		
e $-9x - 12$	**f** $-6y - 8$		
g $-10x - 15y$	**h** $-4m - 20n$		
i $-3x^2 - 18x$	**j** $-8x^2 - 12x$		
k $-16y^2 - 6y$	**l** $-5a^2 - 10a$		



Hint for Q6: Take a negative factor out of both terms.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

9E

Problem-solving and reasoning

7, 8(½) | 8(½), 9–11


7 Write the missing number or expression.

a $3x + 9 = ___(x + 3)$ **b** $xy + x = x(___ + 1)$ **c** $a^2 - a = ___(a - 1)$


d $5xy + 10x = ___(y + 2)$ **e** $-7a - 14 = ___(a + 2)$ **f** $-24a^2 - 36a = ___(2a + 3)$

8 Write down the perimeter of these shapes in factorised form.


a



b




c






Hint for Q8: First find an expression for the perimeter, then factorise it.


d




e



f



9 The expression for the area of a rectangle is $(4x^2 + 8x)$. Find an expression for its width if the length is $(x + 2)$.



10 The height, in metres, of a ball thrown in the air is given by $5t - t^2$, where t is the time in seconds.

a Write an expression for the ball's height in factorised form.

b Find the ball's height at these times:

i $t = 0$ **ii** $t = 2$ **iii** $t = 4$

c How long does it take for the ball's height to return to 0 metres? Use trial and error if required.

11 $7 \times 9 + 7 \times 3$ can be evaluated by first factorising to $7(9 + 3)$. This gives $7 \times 12 = 84$. Use a similar technique to evaluate the following.

a $9 \times 2 + 9 \times 5$ **b** $6 \times 3 + 6 \times 9$ **c** $-2 \times 4 - 2 \times 6$

d $-5 \times 8 - 5 \times 6$ **e** $23 \times 5 - 23 \times 2$ **f** $63 \times 11 - 63 \times 8$

Further factorisation

— | 12, 13

12 Common factors can also be removed from expressions with more than two terms.
For example: $2x^2 + 6x + 10xy = 2x(x + 3 + 5y)$

Factorise these expressions by taking out the HCF.

a $3a^2 + 9a + 12$ **b** $5z^2 - 10z + zy$ **c** $x^2 - 2xy + x^2y$

d $4by - 2b + 6b^2$ **e** $-12xy - 8yz - 20xyz$ **f** $3ab + 4ab^2 + 6a^2b$



13 You can factorise some expressions by taking out a binomial factor.
For example: $3(x - 2) + x(x - 2) = (x - 2)(3 + x)$

Factorise the following by taking out a binomial common factor.

Hint for Q13: $x - 2$ is the common factor.

a $4(x + 3) + x(x + 3)$ **b** $3(x + 1) + x(x + 1)$ **c** $7(m - 3) + m(m - 3)$

d $x(x - 7) + 2(x - 7)$ **e** $8(a + 4) - a(a + 4)$ **f** $5(x + 1) - x(x + 1)$

g $y(y + 3) - 2(y + 3)$ **h** $a(x + 2) - x(x + 2)$ **i** $t(2t + 5) + 3(2t + 5)$

j $m(5m - 2) + 4(5m - 2)$ **k** $y(4y - 1) - (4y - 1)$ **l** $(7 - 3x) + x(7 - 3x)$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Progress quiz

9A **1** Simplify the following.

a $3x - 2y + 5x - 4y$ **b** $3ab^2 - ab + 4ab^2$

c $4a \times 5ab$ **d** $\frac{3xy^2}{9xy}$

9A **2** Expand and simplify the following.

a $3(x - 4)$ **b** $2y(3y + 5)$ **c** $5 - 2(3x + 1)$

9B **3** Expand the following binomial products.

a $(x + 4)(x + 6)$ **b** $(x + 3)(x - 2)$


c $(2x - 3)(x + 4)$ **d** $(3x - 2)(2x - 5)$

9C **4** Expand the following perfect squares.


a $(x + 6)^2$ **b** $(x - 8)^2$ **c** $(3x + 7)^2$ **d** $(5 - x)^2$

9B/C **5** Find the area of these figures in expanded form.

a



b



9D **6** Expand and simplify the following.

a $(x + 10)(x - 10)$ **b** $(x - 1)(x + 1)$

c $(3x + 2)(3x - 2)$ **d** $(4x - 5y)(4x + 5y)$

9C/D **7** Expand and simplify $(x + 1)^2 - (x + 1)(x - 1)$.

9E **8** Determine the HCF of the following.

a $8x$ and 12 **b** $6a$ and $15a$

c $5ab$ and $10ab^2$ **d** $12x^2$ and $9x$


9E **9** Factorise the following. Take a negative factor out in parts **e** and **f**.

a $6x + 12$ **b** $15y - 20$ **c** $8ab + 20a$


d $4x^2 - 2x$ **e** $-10x - 5$ **f** $-4x^2 - 10x$

9E **10** Give the perimeter of these shapes in factorised form

a



b



Essential Mathematics Core for the Victorian Curriculum Year 9 ISBN 978-1-009-59315-1 © Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

9F Factorising the difference of two squares

Learning intentions

- To be able to recognise a difference of two squares
- To be able to express a difference of two squares in factorised form
- To know to check for a common factor first before further factorisation

Key vocabulary: difference of two squares

Recall that a difference of two squares is formed when expanding the product of the sum and difference of two terms. For example, $(x+2)(x-2) = x^2 - 4$. Reversing this process means that a difference of two squares can be factorised into two binomial expressions of the form $(a+b)$ and $(a-b)$.



Engineers use algebra when designing products used for bridges, buildings and machinery, such as this steel turbine for a power station. For example, a circle area difference:

$$\begin{aligned} A &= \pi R^2 - \pi r^2 \\ &= \pi(R^2 - r^2) \\ &= \pi(R-r)(R+r). \end{aligned}$$

Lesson starter: Expanding to understand factorising

Complete the steps in these expansions then write the conclusion.

- $(x+3)(x-3) = x^2 - 3x + ___ - ___$
  $= x^2 - ___$
  $\therefore x^2 - 9 = (___ + ___)(___ - ___)$
- $(a+b)(a-b) = a^2 - ab + ___ - ___$
  $= ___ - ___$
  $\therefore a^2 - ___ = (___ + ___)(___ - ___)$
- $(2x-5)(2x+5) = 4x^2 + 10x - ___ - ___$
  $= ___ - ___$
  $\therefore 4x^2 - ___ = (___ + ___)(___ - ___)$

Key ideas

- Factorising the difference of two squares uses the rule $a^2 - b^2 = (a+b)(a-b)$.
  - $x^2 - 16 = x^2 - 4^2$
    $= (x+4)(x-4)$
  - $9x^2 - 100 = (3x)^2 - 10^2$
    $= (3x+10)(3x-10)$
  - $25 - 4y^2 = 5^2 - (2y)^2$
    $= (5+2y)(5-2y)$
- First take out common factors where possible.
  - $2x^2 - 18 = 2(x^2 - 9)$
    $= 2(x+3)(x-3)$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Exercise 9F

Understanding

1, 2, 3(½) | 1, 3

1 Expand these binomial products to form a difference of two squares.

a $(x+2)(x-2)$

b $(x-7)(x+7)$

c $(3x-y)(3x+y)$



Hint for Q1: $a^2 - b^2$ is a difference of two squares.

2 State the missing term. Assume it is a positive number.

a $(___)^2 = 9$

b $(___)^2 = 121$

c $(___)^2 = 25b^2$

d $(___)^2 = 49y^2$

3 State the missing terms to complete these factorisations.

a $x^2 - 16 = x^2 - 4^2$
$= (x+4)(___ - ___)$

b $x^2 - 144 = x^2 - (___)^2$
$= (___ + 12)(x - ___)$

c $16x^2 - 1 = (___)^2 - (___)^2$
$= (4x + ___)(___ - 1)$

d $9a^2 - 4b^2 = (___)^2 - (___)^2$
$= (3a + ___)(___ - 2b)$

Fluency

4–6(½) | 4–6(½)

Example 13 Factorising a simple difference of two squares

Factorise each of the following.

a $x^2 - 4$

b $144 - y^2$

Solution

a $x^2 - 4 = x^2 - 2^2$
$= (x+2)(x-2)$

Explanation

Write as a difference of two squares (4 is the same as 2^2).

Write in factorised form:
$a^2 - b^2 = (a+b)(a-b)$

Here $a = x$ and $b = 2$.

b $144 - y^2 = 12^2 - y^2$
$= (12+y)(12-y)$

$12^2 = 144$
Use $a^2 - b^2 = (a+b)(a-b)$

Now you try

Factorise each of the following.

a $x^2 - 16$

b $49 - y^2$

4 Factorise each of the following.

a $x^2 - 9$ **b** $y^2 - 25$ **c** $y^2 - 1$ **d** $x^2 - 64$

e $x^2 - 16$ **f** $b^2 - 49$ **g** $a^2 - 81$ **h** $x^2 - y^2$

i $a^2 - b^2$ **j** $16 - a^2$ **k** $25 - x^2$ **l** $1 - b^2$

m $36 - y^2$ **n** $121 - b^2$ **o** $x^2 - 400$ **p** $900 - y^2$



Hint for Q4: $a^2 - b^2 = (a+b)(a-b)$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

9F

Example 14 Factorising move complex differences of two squares

Factorise each of the following.

a $9a^2 - 25$ b $81x^2 - y^2$

Solution

a $9a^2 - 25 = (3a)^2 - 5^2$
$= (3a + 5)(3a - 5)$

Explanation

Write as a difference of two squares. $9a^2$ is the same as $(3a)^2$.

Write in factorised form.

b $81x^2 - y^2 = (9x)^2 - y^2$
$= (9x + y)(9x - y)$

$81x^2 = (9x)^2$

Use $a^2 - b^2 = (a + b)(a - b)$.


Now you try

Factorise each of the following.

a $4a^2 - 9$ b $25a^2 - b^2$

5 Factorise each of the following.

a $4x^2 - 25$ b $9x^2 - 49$
c $25b^2 - 4$ d $4m^2 - 121$
e $100y^2 - 9$ f $81a^2 - 4$
g $1 - 4x^2$ h $25 - 64b^2$
i $16 - 9y^2$ j $36x^2 - y^2$
k $4x^2 - 25y^2$ l $64a^2 - 49b^2$
m $4p^2 - 25q^2$ n $81m^2 - 4n^2$
o $25a^2 - 49b^2$ p $100a^2 - 9b^2$



Example 15 Factorising a difference of two squares with a common factor

Factorise $2b^2 - 32$ by first taking out a common factor.

Solution

$2b^2 - 32 = 2(b^2 - 16)$
$= 2(b^2 - 4^2)$
$= 2(b + 4)(b - 4)$

Explanation

First, factor out the common factor of 2.
Write as a difference of two squares and then factorise.

Now you try

Factorise $3a^2 - 27$ by first taking out a common factor.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

6 Factorise each of the following by first taking out the common factor.

a $3x^2 - 108$ **b** $10a^2 - 10$ **c** $6x^2 - 24$

d $4y^2 - 64$ **e** $98 - 2x^2$ **f** $32 - 8m^2$

g $5x^2y^2 - 5$ **h** $3 - 3x^2y^2$ **i** $63 - 7a^2b^2$

Problem-solving and reasoning 7–9 7–10

7 The height above ground (in metres) of an object thrown off the top of a building is given by $36 - 4t^2$ where t is in seconds.

a Factorise the expression for the height of the object by first taking out the common factor.

b Find the height of the object:

i initially ($t = 0$) **ii** at 2 seconds ($t = 2$).

c How long does it take for the object to hit the ground? Use trial and error if you wish.

8 Initially it may not appear that an expression such as $-4 + 9x^2$ is a difference of two squares. However, swapping the position of the two terms makes $-4 + 9x^2 = 9x^2 - 4$, which can be factorised to $(3x + 2)(3x - 2)$. Use this idea to factorise these difference of two squares.

a $-9 + x^2$ **b** $-121 + 16x^2$ **c** $-25a^2 + 4$ **d** $-y^2 + x^2$

9 Olivia factorises $16x^2 - 4$ to get $(4x + 2)(4x - 2)$, but the answer says $4(2x + 1)(2x - 1)$.

a What should Olivia do to get from her answer to the actual answer?

b What should Olivia have done initially to avoid this issue?

10 By using $a^2 - b^2 = (a + b)(a - b)$ we can factorise $(x + 1)^2 - 4$ in the following way:

$$\begin{aligned}(x + 1)^2 - 4 &= (x + 1)^2 - 2^2 \\ &= ((x + 1) + 2)((x + 1) - 2) \\ &= (x + 3)(x - 1)\end{aligned}$$


Use this technique to factorise the following.

a $(x + 5)^2 - 9$ **b** $(x + 3)^2 - 4$ **c** $(x + 10)^2 - 16$

d $(x - 3)^2 - 25$ **e** $(x - 7)^2 - 1$ **f** $(x - 3)^2 - 36$

The multisize frame — 11

11 This 'multisize' square picture frame has side length 30 cm and can hold a square picture with any side length less than 26 cm.



a If the side length of the picture is x cm, write an expression for:

i the area of the picture

ii the area of the frame (in factorised form).

b Use your result from part **a ii** to find the area of the frame if:

i $x = 20$

ii the area of the picture is 225 cm^2.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

9G Factorising by grouping in pairs

Learning intentions

- To know that a common factor can include a binomial term
- To be able to use the grouping method to factorise some four-term expressions
- To know that expressions can be rearranged to find common factors to use in the grouping process

Key vocabulary: binomial, grouping in pairs

When an expression can be written using four terms, such as $x^2 + 2x - x - 2$, it may be possible to factorise it into a product of two binomial terms such as $(x - 1)(x + 2)$. In such situations the method of grouping in pairs is often used.



Algebra is an essential tool for the engineers who work together making renewable energy possible, including: aerospace, civil, electrical, electronics, environmental, industrial, materials, mechanical and solar engineers.

Lesson starter: Two methods - Same result

The four-term expression $x^2 - 3x - 3 + x$ is written on the board.

Tommy chooses to rearrange the terms to give $x^2 - 3x + x - 3$ then factorises by grouping in pairs.

Sharon chooses to rearrange the terms to give $x^2 + x - 3x - 3$ then also factorises by grouping in pairs.

- Complete Tommy and sharon's factorisation working.

Tommy	Sharon
$x^2 - 3x + x - 3 = x(x - 3) + 1(_____)$	$x^2 + x - 3x - 3 = x(_____) - 3(_____)$
$= (x - 3)(_____)$	$= (x + 1)(_____)$

- Discuss the differences in the methods. Is there any difference in their answers?
- Whose method do you prefer?

Key ideas

- **Factorisation by grouping in pairs** is a method that is often used to factorise a four-term expression.
  - Terms are grouped into pairs and factorised separately.
  - The common binomial factor is then taken out to complete the factorisation.
  - Terms can be rearranged to assist in the search of a common factor.

$$x^2 + 3x - 2x - 6$$
$$= x(x + 3) - 2(x + 3)$$
$$= (x + 3)(x - 2)$$

which could also be written as $(x - 2)(x + 3)$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Exercise 9G

Understanding

1, 2 | 1(½), 2, 3

1 Expand each expression.

a $2(x-1)$
b $-5(1-a)$
c $a(a+5)$
d $y(4-y)$
e $x(a+1)+2(a+1)$
f $b(x-2)-3(x-2)$

2 State the missing information.

a $2(x+1)+x(x+1)=(x+1)$(_____)
b $5(x+5)-x(x+5)=(x+5)$(_____)
c $a(x-3)+(x-3)=(x-3)$(_____)
d $(x-3)-a(x-3)=(x-3)$(_____)



Hint for Q2: What is left after dividing by the common factor?

3 Identify the common binomial term then factorise each expression.

a $x(x-3)-2(x-3)$
b $x(x+4)+3(x+4)$
c $3(2x+1)-x(2x+1)$
d $4(3x-2)-x(3x-2)$

Fluency

4–6(½) | 4–6(½)

Example 16 Factorising by grouping in pairs

Use the method of grouping in pairs to factorise $x^2+2x+3x+6$.

Solution

$$\begin{aligned} x^2+2x+3x+6 &= (x^2+2x)+(3x+6) \\ &= x(x+2)+3(x+2) \\ &= (x+2)(x+3) \end{aligned}$$

Explanation

Group the first and second pair of terms.

Factorise each group.

Take the common factor $(x+2)$ out of both groups.

Now you try

Use the method of grouping in pairs to factorise $x^2+4x+2x+8$.

4 Use the method of grouping in pairs to factorise these expressions.

a $x^2+3x+2x+6$
b $x^2+4x+3x+12$
c $x^2+7x+2x+14$
d $x^2-6x+4x-24$
e $x^2-4x+6x-24$
f $x^2-3x+10x-30$



Hint for Q4: Factorise the first pair then the second pair.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

9G

Example 17 Factorising by grouping in pairs using a negative common factor

Use the method of grouping in pairs to factorise $x^2 + 3x - 5x - 15$.

Solution

$$\begin{aligned} x^2 + 3x - 5x - 15 &= (x^2 + 3x) + (-5x - 15) \\ &= x(x + 3) - 5(x + 3) \\ &= (x + 3)(x - 5) \end{aligned}$$

Explanation

Group the first and second pair of terms.

Factorise each group, including the common negative in the second part.

Take out the common factor $(x + 3)$.

Now you try

Use the method of grouping in pairs to factorise $x^2 + 2x - 3x - 6$.

5 Use the method of grouping in pairs to factorise these expressions.

a $x^2 + 4x - 3x - 12$ **b** $x^2 + 2x - 6x - 12$

c $x^2 + 2x - 18x - 36$ **d** $x^2 + 3x - 14x - 42$

e $x^2 + 4x - 18x - 72$ **f** $x^2 - 2x - xa + 2a$



Hint for Q5: You can take out a negative common factor if necessary.

6 Factorise these expressions. Use a factor of 1 where necessary, for example, $x^2 - ax + x - a = x(x - a) + 1(x - a)$.

a $x^2 - bx + x - b$ **b** $x^2 - cx + x - c$

c $x^2 + bx + x + b$ **d** $x^2 + cx - x - c$

e $x^2 + ax - x - a$ **f** $x^2 - bx - x + b$

Problem-solving and reasoning 7–8(½) 7–9(½), 10

Example 18 Rearranging an expression to factorise by grouping in pairs

Factorise $2x^2 - 9 - 18x + x$ using grouping in pairs.

Solution

$$\begin{aligned} 2x^2 - 9 - 18x + x &= 2x^2 + x - 18x - 9 \\ &= x(2x + 1) - 9(2x + 1) \\ &= (2x + 1)(x - 9) \end{aligned}$$

Explanation

Rearrange so that each group has a common factor.

Factorise each group then take out $(2x + 1)$.

Alternative method:

$$\begin{aligned} 2x^2 - 9 - 18x + x &= 2x^2 - 18x + x - 9 \\ &= 2x(x - 9) + 1(x - 9) \\ &= (x - 9)(2x + 1) \end{aligned}$$

Alternatively, you can group in another order where each group has a common factor. Then factorise.

The answer will be the same.

Now you try

Factorise $2x^2 - 15 - 10x + 3x$ using grouping in pairs.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

7 Factorise these expressions by first rearranging the terms.

a $2x^2 - 7 - 14x + x$
b $5x + 2x + x^2 + 10$
c $2x^2 - 3 - x + 6x$
d $3x - 8x - 6x^2 + 4$
e $11x - 5a - 55 + ax$
f $12y + 2x - 8xy - 3$
g $6m - n + 3mn - 2$
h $15p - 8r - 5pr + 24$

8 Note that $x^2+5x+6 = x^2+2x+3x+6$, which can be factorised by grouping in pairs. Use a similar method to factorise the following.

a $x^2 + 7x + 10$
b $x^2 + 8x + 15$
c $x^2 + 10x + 24$
d $x^2 - x - 6$
e $x^2 + 4x - 12$
f $x^2 - 11x + 18$

9 Use the method of grouping in pairs to factorise these expressions. The HCF for each pair includes a pronumeral. For example, $2ab - 2ad + bc - cd = 2a(b - d) + c(b - d) = (b - d)(2a + c)$

a $3ab + 5bc + 3ad + 5cd$
b $4ab - 7ac + 4bd - 7cd$
c $2xy - 8xz + 3wy - 12wz$
d $5rs - 10r + st - 2t$
e $4x^2 + 12xy - 3x - 9y$
f $2ab - a^2 - 2bc + ac$

10 Make up at least three of your own four-term expressions that factorise to a binomial product. Describe the method that you used to make up each four-term expression.

Comparing two methods — 11

11 $xa - 21 + 7a - 3x$ could be rearranged in two different ways before factorising.

Method 1

$xa + 7a - 3x - 21 = a(x + 7) - 3(_____)$

$= _____$

Method 2

$xa - 3x + 7a - 21 = x(a - 3) + 7(_____)$

$= _____$

a Copy and complete both methods for the above expression.

b Use different arrangements of the four terms to complete the factorisation of the following in two ways. Show working using both methods.

i $xb - 6 - 3b + 2x$
ii $xy - 8 + 2y - 4x$
iii $4m^2 - 15n + 6m - 10mn$
iv $2m + 3n - mn - 6$
v $4a - 6b^2 + 3b - 8ab$
vi $3ab - 4c - b + 12ac$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


9H Factorising monic quadratic trinomials

Learning intentions

- To know the form of a monic quadratic trinomial
- To understand how a quadratic trinomial is formed from a binomial expansion
- To know how to find the numbers required to express a quadratic trinomial in factorised form

Key vocabulary: monic quadratic trinomial

An expression that takes the form $x^2 + bx + c$, where b and c are constants, is an example of a monic quadratic trinomial that has the coefficient of x^2 equal to 1. To factorise a quadratic expression, we need to use the distributive law in reverse. Consider the expansion shown below:





Quadratic equations are widely used in engineering, business and science, and factorisation is a key step when solving such equations. Self-driving cars are programmed to solve equations, including quadratic equations of motion.

If we examine the expansion above we can see how each term of the product is formed.

Product of x and x is x^2

$(x + 2)(x - 4) = x^2 - 2x - 8$

Product of 2 and -4 is -8
$(2 \times (-4) = -8$, the constant term)

$x \times (-4) = -4x$

$(x + 2)(x - 4) = x^2 - 2x - 8$

$2 \times x = 2x$

Add $-4x$ and $2x$ to give the middle term, $-2x$
$(-4 + 2 = -2$, the coefficient of $x)$

Lesson starter: So many choices

Mia says that as $-2 \times 3 = -6$ then $x^2 + 5x - 6$ must equal $(x - 2)(x + 3)$.

- Expand $(x - 2)(x + 3)$ to see if Mia is correct.
- What other pairs of numbers multiply to give -6?
- Which pair of numbers should Mia choose to correctly factorise $x^2 + 5x - 6$?
- What advice can you give Mia when trying to factorise these types of trinomials?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Key ideas

- To factorise a **monic quadratic trinomial** of the form $x^2 + bx + c$, find two numbers that:
  - multiply to give c and
  - add to give b.

  For example: $x^2 - 3x - 10 = (x - 5)(x + 2)$ choose -5 and $+2$ since $-5 \times 2 = -10$ and $-5 + 2 = -3$
- Check factorisation steps by expanding.

  check: $(x - 5)(x + 2) = x^2 + 2x - 5x - 10$
  $= x^2 - 3x - 10$
- Write the factors in any order.

  write $(x - 5)(x + 2)$ or $(x + 2)(x - 5)$
- If c is a negative number, the signs in the factorised expression will be different from each other.
- If c is a positive number, the signs in the factorised expression will be the same as each other.

Exercise 9H

Understanding

1, 2 | 1, 2

1 Give the result of expanding these binomial products.

a $(x + 2)(x + 7)$ **b** $(x + 3)(x + 11)$
c $(x + 12)(x - 5)$ **d** $(x - 2)(x - 6)$

2 Decide which two numbers multiply to give the first number and add to give the second number.

a $6, 5$ **b** $12, 13$ **c** $20, 9$
d $-15, 2$ **e** $-30, -1$ **f** $18, -11$

Fluency

3–5(½) | 3–6(½)

Example 19 Factorising monic quadratic trinomials

Factorise $x^2 + 7x + 10$.

Solution

$x^2 + 7x + 10 = (x + 5)(x + 2)$

Explanation

Need two numbers that × to 10 and + to 7.
Factors of 10 include $(10, 1)$ and $(5, 2)$. The pair that adds to 7 is $(5, 2)$.

Now you try

Factorise $x^2 + 9x + 14$

3 Factorise each of the following quadratic expressions.

a $x^2 + 8x + 12$ **b** $x^2 + 10x + 9$ **c** $x^2 + 8x + 7$
d $x^2 + 15x + 14$ **e** $x^2 + 7x + 12$ **f** $x^2 + 10x + 16$
g $x^2 + 8x + 15$ **h** $x^2 + 9x + 20$ **i** $x^2 + 11x + 24$



Hint for Q3: Look for factors of c which add to b.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

9H

Example 20 Factorising monic quadratic trinomials with a negative c value

Factorise $x^2 + 2x - 8$.

Solution

$x^2 + 2x - 8 = (x + 4)(x - 2)$

Explanation

Need two numbers that $\times$ to -8 and $+$ to 2.
Factors of -8 are $(-8, 1)$ or $(8, -1)$ or $(4, -2)$ or $(-4, 2)$ and $4 + (-2) = 2$, so choose $(4, -2)$.

Now you try

Factorise $x^2 + 4x - 12$

4 Factorise each of the following quadratic expressions.

a $x^2 + 3x - 4$	**b** $x^2 + x - 2$	**c** $x^2 + 4x - 5$
d $x^2 + 5x - 14$	**e** $x^2 + 2x - 15$	**f** $x^2 + 8x - 20$
g $x^2 - 3x - 18$	**h** $x^2 - 7x - 18$	**i** $x^2 - x - 12$



Hint for Q4: Remember that sometimes b and c are negative.

Example 21 Factorising monic quadratic trinomials with a negative b value

Factorise $x^2 - 7x + 10$

Solution

$x^2 - 7x + 10 = (x - 2)(x - 5)$

Explanation

Need two numbers that $\times$ to 10 and $+$ to -7.
Factors of 10 are $(10, 1)$ or $(-10, -1)$ or $(5, 2)$ or $(-5, -2)$.

To add to a negative (-7), both factors must then be negative: $-5 + (-2) = -7$, so choose $(-5, -2)$.

Now you try

Factorise $x^2 - 7x + 12$

5 Factorise each of the following quadratic expressions.

a $x^2 - 6x + 5$	**b** $x^2 - 2x + 1$	**c** $x^2 - 5x + 4$
d $x^2 - 9x + 8$	**e** $x^2 - 4x + 4$	**f** $x^2 - 8x + 12$
g $x^2 - 11x + 18$	**h** $x^2 - 10x + 21$	**i** $x^2 - 5x + 6$

6 Factorise each of the following quadratic expressions.

a $x^2 - 7x - 8$	**b** $x^2 - 3x - 4$	**c** $x^2 - 5x - 6$
d $x^2 - 6x - 16$	**e** $x^2 - 2x - 24$	**f** $x^2 - 2x - 15$
g $x^2 - x - 12$	**h** $x^2 - 11x - 12$	**i** $x^2 - 4x - 12$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Problem-solving and reasoning 7–8(½) 7–9(½)

Example 22 Factorising with a common factor

Factorise the quadratic expression $2x^2 - 2x - 12$.

Solution

$2x^2 - 2x - 12 = 2(x^2 - x - 6)$
$= 2(x - 3)(x + 2)$

Explanation

First take out common factor of 2.

Factors of -6 are $(-6, 1)$ or $(6, -1)$ or $(-3, 2)$ or $(3, -2)$.

$-3 + 2 = -1$ so choose $(-3, 2)$.

Now you try

Factorise the quadratic expression $3x^2 - 6x - 45$.

7 Factorise each of the following quadratic expressions by first taking out a common factor.

a $2x^2 + 10x + 8$ **b** $2x^2 + 22x + 20$ **c** $3x^2 + 18x + 24$
d $2x^2 + 14x - 60$ **e** $2x^2 - 14x - 36$ **f** $4x^2 - 8x + 4$
g $2x^2 + 2x - 12$ **h** $6x^2 - 30x - 36$ **i** $5x^2 - 30x + 40$
j $3x^2 - 33x + 90$ **k** $2x^2 - 6x - 20$ **l** $3x^2 - 3x - 36$

8 Find the missing term in these trinomials if they are to factorise using positive integers. For example: the missing term in $x^2 + \square + 10$ could be $7x$ because $x^2 + 7x + 10$ factorises to $(x + 5)(x + 2)$ and 5 and 2 are integers. There may be more than one answer in each case.


a $x^2 + \square + 5$ **b** $x^2 - \square + 9$ **c** $x^2 - \square - 12$ **d** $x^2 + \square - 12$
e $x^2 + \square + 18$ **f** $x^2 - \square + 18$ **g** $x^2 - \square - 16$ **h** $x^2 + \square - 25$

9 The expression $x^2 - 6x + 9$ factorises to $(x - 3)(x - 3) = (x - 3)^2$, which is a perfect square. Factorise these perfect squares. A common factor may also be involved.

a $x^2 + 8x + 16$ **b** $x^2 + 10x + 25$ **c** $x^2 + 30x + 225$
d $x^2 - 2x + 1$ **e** $x^2 - 14x + 49$ **f** $x^2 - 26x + 169$
g $2x^2 + 4x + 2$ **h** $5x^2 - 30x + 45$ **i** $-3x^2 + 36x - 108$

The paved backyard — 10

10 A rectangular backyard has a length 2 metres more than its width (x metres). Inside the rectangle are three paved squares each of area 5 m^2 as shown. The remaining area is lawn.



a Find an expression for:
- **i** the total backyard area
- **ii** the area of lawn in expanded form
- **iii** the area of lawn in factorised form.

b Find the area of lawn if:
- **i** $x = 10$ **ii** $x = 7$.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

9I Simplifying algebraic fractions: multiplication and division ★

Learning intentions

- To know that expressions must be factorised before common factors can be cancelled
- To be able to simplify algebraic fractions by cancelling common factors
- To be able to multiply and divide algebraic fractions

Key vocabulary: algebraic fraction, common factor, factorise, numerator, denominator, reciprocal

Fractions such as $\frac{4}{6}$ and $\frac{24}{16}$ can be simplified by cancelling common factors. This is also true of algebraic fractions such as $\frac{3x}{6}$, $\frac{5a^2}{10a}$ and $\frac{2x+4}{2}$.

Lesson starter: Does $\frac{2x+6^3}{2^1} = 2x+3$?

For the expression $\frac{2x+6}{2}$, a student attempts to cancel the 6 with the 2 in the denominator.

They write $\frac{2x+6^3}{2^1} = 2x+3$.

- Is this a correct method?
- Substitute $x = 1$ into the left hand side and then the right hand side. Are they equal?
- Can you show a correct method? Does it involve factorisation?

Key ideas

- Simplify **algebraic fractions** by cancelling **common factors**

$$\frac{3^1x}{6^2} = \frac{x}{2} \qquad \frac{5a^2}{3a} = \frac{5a}{3} \qquad \frac{2x+4}{2} = \frac{2^1(x+2)}{2^1} = x+2$$

  - This is an incorrect cancellation: $\frac{2x+4^2}{2^1} = 2x+2$

- To multiply algebraic fractions, first cancel any numerator with any denominator.

$$\frac{2^1x^1}{5_1} \times \frac{15^3}{4_2x^1y} = \frac{1 \times 3}{1 \times 2y} = \frac{3}{2y}$$

- To divide algebraic fractions, multiply by the reciprocal of the fraction following the division sign.
  - The **reciprocal** of $\frac{a}{b} = \frac{b}{a}$.

$$\frac{3x}{4} \div \frac{9}{8y} = \frac{3^1x}{4_1} \times \frac{8^2y}{9^3} = \frac{2xy}{3}$$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Exercise 9I

Understanding

1–3 | 3

1 Cancel to simplify these fractions.

a $\frac{4}{2}$ **b** $\frac{4x}{2}$ **c** $\frac{4(x+2)}{2}$ **d** $\frac{4(x-1)}{2}$



2 Write down the reciprocal of these fractions.

a $\frac{2}{3}$ **b** $\frac{4}{3}$ **c** $\frac{7x}{2}$

d $\frac{5}{4a^2}$ **e** 7 **f** $6x$

Hint for Q2:
Recall: $7 = \frac{7}{1}$
for part e.



3 Divide these simple fractions.

a $\frac{2}{3} \div \frac{3}{4}$ **b** $\frac{7}{8} \div \frac{3}{4}$ **c** $\frac{6}{7} \div \frac{4}{7}$

d $\frac{5}{14} \div \frac{10}{7}$ **e** $\frac{21}{4} \div \frac{7}{8}$ **f** $\frac{4}{9} \div \frac{4}{9}$

Hint for Q3:
- Copy the first fraction.
- Change the ÷ to ×.
- Flip over the second fraction.

Fluency

4–7(½) | 4–7(½)

Example 23 Cancelling common factors

Simplify by cancelling common factors.

a $\frac{7x}{21}$ **b** $\frac{5(x+2)(x-1)}{x+2}$

Solution

a $\frac{\cancel{7}^{1}x}{\cancel{21}_{3}} = \frac{x}{3}$

b $\frac{5\cancel{(x+2)}^{1}(x-1)}{\cancel{x+2}^{1}} = 5(x-1)$

Explanation

The HCF of 7 and 21 is 7.

Treat $(x+2)$ as a common factor to both the numerator and denominator.

Now you try

Simplify by cancelling common factors.

a $\frac{-2ab^2}{4b}$ **b** $\frac{-3(x-2)(x+7)}{3(x-2)}$

4 Simplify by cancelling common factors.

a $\frac{4x}{8}$ **b** $\frac{8x}{16}$ **c** $\frac{-4ab}{12}$ **d** $\frac{7a}{14b}$

e $\frac{4(x-1)}{4}$ **f** $\frac{-6(2x+1)}{3}$ **g** $\frac{-7x}{21x}$ **h** $\frac{4a^2}{40a}$

i $\frac{3(x+2)}{x+2}$ **j** $\frac{-2(x-7)}{x-7}$ **k** $\frac{5(x+3)}{10(x+3)}$ **l** $\frac{-16(x-14)}{24(x-14)}$

m $\frac{(x+2)(x-4)}{x+2}$ **n** $\frac{2(x-1)(x+3)}{x+3}$ **o** $\frac{-4(x+2)(x-5)}{2(x-5)}$ **p** $\frac{-7(x+1)(x-9)}{7(x+1)(x-9)}$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

91

Example 24 Simplifying by factorising

Simplify these fractions by factorising first.

a $\frac{2x+6}{2}$ b $\frac{-3x-9}{x+3}$

Solution

a $\frac{2x+6}{2} = \frac{2^{1}(x+3)}{2^{1}}$

$= x+3$

Explanation

First factorise the numerator, then cancel the 2.

b $\frac{-3x-9}{x+3} = \frac{-3(x+3)^{1}}{x+3^{1}}$

$= -3$

-3 is common to both $-3x$ and -9. $(x+3)$ can now be cancelled.

Now you try

Simplify these fractions by factorising first.

a $\frac{5a-30}{5}$ b $\frac{-7x-14}{x+2}$

5 Simplify these fractions by factorising first.

a $\frac{4x+8}{4}$ b $\frac{2x-6}{2}$ c $\frac{3x-12}{3}$

d $\frac{3}{3x+3}$ e $\frac{-7}{14x+21}$ f $\frac{5a}{15a-10}$

g $\frac{2x+8}{x+4}$ h $\frac{7x+14}{x+2}$ i $\frac{-5x-15}{x+3}$

j $\frac{-4x-10}{2x+5}$ k $\frac{18+12x}{3+2x}$ l $\frac{-25-15x}{5+3x}$



Hint for Q5: First factorise the numerator or denominator, then cancel.

Example 25 Multiplying algebraic fractions

Simplify these products.

a $\frac{3x}{4} \times \frac{8}{9x}$ b $\frac{4(x+1)}{3} \times \frac{12}{x+1}$

Solution

a $\frac{3x^{1}}{4^{1}} \times \frac{8^{2}}{9x^{3}} = \frac{1 \times 2}{1 \times 3}$

$= \frac{2}{3}$

Explanation

Cancel common factors from any numerator to any denominator, then multiply.

b $\frac{4(x+1)^{1}}{3_{1}} \times \frac{12^{4}}{x+1^{1}} = \frac{4 \times 4}{1 \times 1}$

$= 16$

Cancel all common factors before multiplying.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Now you try

Simplify these products.

a $\frac{2a^2}{3} \times \frac{9}{a}$

b $\frac{3x-1}{5} \times \frac{20}{6x-2}$

6 Simplify these products.

a $\frac{4x}{5} \times \frac{10}{8x}$

b $\frac{3x}{7} \times \frac{7}{6x}$

c $\frac{15a}{4} \times \frac{8}{25}$

d $\frac{16}{3a} \times \frac{9}{32}$

e $\frac{4y^2}{3} \times \frac{7x}{4y}$

f $\frac{8}{3x} \times \frac{5x^2}{16}$

g $\frac{2(x-1)}{3} \times \frac{6}{x-1}$

h $\frac{5(2x+1)}{4} \times \frac{4}{2x+1}$

i $\frac{3(x-7)}{7} \times \frac{21}{2(x-7)}$

j $\frac{x-3}{4} \times \frac{x+1}{x-3}$

k $\frac{3x+7}{x-2} \times \frac{x-2}{3x+7}$

l $\frac{5(x-2)}{3(x+7)} \times \frac{3(x+6)}{10(x-2)}$



Hint for Q6: Remember to cancel first before multiplying.

Example 26 Dividing algebraic fractions

Simplify the following.

a $\frac{3x}{4} \div \frac{6}{7}$

b $\frac{5(x+1)}{6} \div \frac{2(x+1)}{9}$

Solution

a $\frac{3x}{4} \div \frac{6}{7} = \frac{3^1x}{4} \times \frac{7}{6^2}$

$= \frac{7x}{8}$

Explanation

Multiply by the reciprocal of the second fraction.
Cancel before multiplying.

b $\frac{5(x+1)}{6} \div \frac{2(x+1)}{9} = \frac{5(x+1)^1}{6_2} \times \frac{9^3}{2(x+1)^1}$

$= \frac{15}{4}$

Invert the second fraction and multiply.
Cancel before multiplying.

Now you try

Simplify the following.

a $\frac{-2x}{3} \div \frac{4}{9}$

b $\frac{2}{3(4a-1)} \div \frac{5}{4a-1}$

7 Simplify the following.

a $\frac{2x}{3} \div \frac{4}{7}$

b $\frac{5x}{4} \div \frac{15}{8}$

c $\frac{-3a}{4} \div \frac{2a}{7}$

d $\frac{-6a}{7} \div \frac{12}{7a}$

e $\frac{4(x+1)}{3} \div \frac{2(x+1)}{9}$

f $\frac{2(x-1)}{5} \div \frac{3(x-1)}{10}$

g $\frac{11(x+4)}{x+1} \div \frac{x+4}{x+1}$

h $\frac{-3(2x-1)}{2x+1} \div \frac{2(2x-1)}{2x+1}$



Hint for Q7: Multiply by the reciprocal of the fraction following the division sign.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

91

Problem-solving and reasoning

8–9(½) | 8–10(½)

8 Find the error in these problems and then find the correct answer.

a $\frac{2x+2^{1}}{2^{1}} = 2x+1$

b $\frac{-3x-6}{3} = \frac{-3(x-2)}{3}$
$= -(x-2)$

c $\frac{2x}{3} \div \frac{4x}{9} = \frac{8x^2}{27}$

d $\frac{x+2^{1}}{3^{1}} \times \frac{6^{2}x}{2^{1}} = 2x^2+1$

9 Simplify the following.

a $\frac{2x+4}{3} \times \frac{6}{x+2}$

b $\frac{5x-15}{7} \times \frac{7}{x-3}$

c $\frac{-6x-12}{x} \times \frac{3x}{x+2}$

d $\frac{5x-5}{2} \div \frac{x-1}{6}$

e $\frac{-17x-34}{x+1} \div \frac{x+2}{2x+2}$

f $\frac{-7x+14}{3x+2} \div \frac{2-x}{6x+4}$



Hint for Q9: Factorise expressions before cancelling; e.g. $2x+4$ can be written as $2(x+2)$.

10 We know that $\frac{a^2}{a}$ cancels to $\frac{a}{1} = a$. Similarly, $\frac{(x+1)^2}{x+1} = \frac{(x+1)(x+1)}{x+1} = x+1$. Use this idea to cancel the following.

a $\frac{(x+3)^2}{x+3}$

b $\frac{(2x-1)^2}{2x-1}$

c $\frac{-5(x-3)^2}{x-3}$

d $\frac{-2(x-4)}{6(x-4)^2}$

The negative 1 factor

— | 11

11 Note that $3-2x = -1(2x-3)$. Use this idea to cancel the following.

a $\frac{3-2x}{2x-3}$

b $\frac{5-7x}{7x-5}$

c $\frac{4(2-x)}{3(x-2)}$

d $\frac{-2(x-5)}{5-x}$

e $\frac{2-x}{(x-2)^2}$

f $\frac{16(x-6)^2}{4(6-x)}$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

9J Simplifying algebraic fractions: addition and subtraction ★

Learning intentions

- To know that the steps for adding and subtracting algebraic fractions are the same as for numerical fractions
- To be able to find the lowest common denominator of fractions
- To be able to add and subtract algebraic fractions

Key vocabulary: lowest common denominator, equivalent fraction, algebraic fraction, numerator, denominator

The process required for adding or subtracting algebraic fractions is similar to that used for fractions without pronumerals.

To simplify $\frac{2}{3}+\frac{4}{5}$, for example, you would find the lowest common multiple of the denominators (15) then express each fraction with this denominator. Adding the numerators completes the task.

Lesson starter: Compare the working

Here is the working for the simplification of the sum of a pair of numerical fractions and the sum of a pair of algebraic fractions.

$$\frac{2}{5}+\frac{3}{4}=\frac{8}{20}+\frac{15}{20}$$
$$=\frac{8+15}{20}$$
$$=\frac{23}{20}$$

$$\frac{2x}{5}+\frac{3x}{4}=\frac{8x}{20}+\frac{15x}{20}$$
$$=\frac{8x+15x}{20}$$
$$=\frac{23x}{20}$$

- What steps taken to simplify the algebraic fractions were the same as those used for the numerical fractions?
- Write down the steps required to add (or subtract) algebraic fractions.

Key ideas

- To add or subtract algebraic fractions:
  - determine the lowest common denominator (LCD)
  - express each fraction using the LCD
  - add or subtract the numerators.

$$\frac{2x}{3}-\frac{x}{5}=\frac{10x}{15}-\frac{3x}{15}$$
$$=\frac{10x-3x}{15}$$
$$=\frac{7x}{15}$$

Exercise 9J

Understanding

1–4 | 3, 4

1 Find the lowest common multiple (LCM) of these pairs of numbers.

a (6, 8) **b** (3, 5) **c** (11, 13) **d** (12, 18)

2 Evaluate:

a $\frac{1}{2}+\frac{1}{3}$ **b** $\frac{3}{5}+\frac{1}{4}$ **c** $\frac{7}{8}+\frac{3}{4}$

d $\frac{9}{10}-\frac{1}{5}$ **e** $\frac{2}{3}-\frac{4}{5}$ **f** $\frac{11}{12}-\frac{5}{6}$



Hint for Q2: First find the LCM. Recall that the LCM of 8 and 4 is 8, not 32.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

9J

3 Write equivalent fractions by stating the missing expression.

a $\frac{2x}{5} = \frac{\square}{10}$ **b** $\frac{7x}{3} = \frac{\square}{9}$ **c** $\frac{x+1}{4} = \frac{\square(x+1)}{12}$

d $\frac{3x+5}{11} = \frac{\square(3x+5)}{22}$ **e** $\frac{4}{x} = \frac{\square}{2x}$ **f** $\frac{30}{x+1} = \frac{\square}{3(x+1)}$

4 Copy and complete these simplifications.

a $\frac{x}{4} + \frac{2x}{3} = \frac{\square}{12} + \frac{\square}{12} = \frac{\square}{12}$

b $\frac{5x}{7} - \frac{2x}{5} = \frac{\square}{35} - \frac{\square}{35} = \frac{\square}{35}$



Hint for Q4: Remember to multiply the numerator by the same number that you multiplied the denominator.

Fluency

5, 6(½) | 5, 6–7(½)

5 Write down the LCD (lowest common denominator) for these pairs of fractions.

a $\frac{x}{3}, \frac{2x}{5}$ **b** $\frac{3x}{7}, \frac{x}{2}$ **c** $\frac{-5x}{4}, \frac{x}{8}$ **d** $\frac{2x}{3}, \frac{-5x}{6}$ **e** $\frac{7x}{10}, \frac{-3x}{5}$

Example 27 Adding and subtracting with a numeral in the denominator

Simplify:

a $\frac{7x}{3} + \frac{x}{6}$ **b** $\frac{x}{4} - \frac{2x}{5}$

Solution

a $\frac{7x}{3} + \frac{x}{6} = \frac{7x \times 2}{3 \times 2} + \frac{x}{6}$

$= \frac{14x}{6} + \frac{x}{6}$

$= \frac{14x + x}{6}$

$= \frac{15x}{6}$

$= \frac{5x}{2}$

b $\frac{x}{4} - \frac{2x}{5} = \frac{5x}{20} - \frac{8x}{20}$

$= \frac{5x - 8x}{20}$

$= -\frac{3x}{20}$

Explanation

Note that the LCM of 3 and 6 is 6, not $3 \times 6 = 18$.

For $\frac{7x}{3}$, multiply the numerator and denominator by 2. $\frac{x}{6}$ already has a denominator of 6.

$\frac{15}{6} = \frac{5}{2}$ after cancelling.

Determine the LCM of 4 and 5; i.e. 20. Express each fraction as an equivalent fraction with a denominator of 20. $2x \times 4 = 8x$.
Then subtract numerators.

Now you try

Simplify:

a $\frac{2a}{5} + \frac{3a}{10}$ **b** $\frac{7x}{8} - \frac{2x}{3}$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

6 Simplify:

a $\frac{x}{7}+\frac{x}{2}$ **b** $\frac{x}{3}+\frac{x}{15}$ **c** $\frac{x}{4}-\frac{x}{8}$

d $\frac{x}{9}+\frac{x}{5}$ **e** $\frac{y}{7}-\frac{y}{8}$ **f** $\frac{a}{2}+\frac{a}{11}$

g $\frac{b}{3}-\frac{b}{9}$ **h** $\frac{m}{3}-\frac{m}{6}$ **i** $\frac{m}{6}+\frac{3m}{4}$

j $\frac{a}{4}+\frac{2a}{7}$ **k** $\frac{2x}{5}+\frac{x}{10}$ **l** $\frac{p}{9}-\frac{3p}{7}$

m $\frac{b}{2}-\frac{7b}{9}$ **n** $\frac{9y}{8}+\frac{2y}{5}$ **o** $\frac{4x}{7}-\frac{x}{5}$



Hint for Q6:
The LCM of 7 and 2 is 14, so write $\frac{x}{7}+\frac{x}{2}=\frac{2x}{14}+\frac{7x}{14}$

$=\frac{2x+7x}{14}$

$=\frac{9x}{14}$

Example 28 Adding more algebraic fractions

Simplify: $\frac{x+3}{2}+\frac{x-2}{5}$

Solution

$\frac{(x+3)}{2}+\frac{(x-2)}{5}=\frac{5(x+3)}{10}+\frac{2(x-2)}{10}$

$=\frac{5x+15+2x-4}{10}$

$=\frac{7x+11}{10}$

Explanation

The LCM of 2 and 5 is 10. Write as equivalent fractions with denominator 10.
Expand the brackets and simplify the numerator by adding and collecting like terms.

Now you try

Simplify: $\frac{x-3}{4}+\frac{x-2}{3}$

7 Simplify:

a $\frac{x+1}{2}+\frac{x+3}{5}$ **b** $\frac{x+3}{3}+\frac{x-4}{4}$

c $\frac{a-2}{7}+\frac{a-5}{8}$ **d** $\frac{y+4}{5}+\frac{y-3}{6}$

e $\frac{m-4}{8}+\frac{m+6}{5}$ **f** $\frac{x-2}{12}+\frac{x-3}{8}$

g $\frac{2b-3}{6}+\frac{b+2}{8}$ **h** $\frac{3x+8}{6}+\frac{2x-4}{3}$

i $\frac{2y-5}{7}+\frac{3y+2}{14}$ **j** $\frac{2t-1}{8}+\frac{t-2}{16}$

k $\frac{4-x}{3}+\frac{2-x}{7}$ **l** $\frac{2m-1}{4}+\frac{m-3}{6}$



Hint for Q7: Remember to include brackets when multiplying a numeral by these binomial numerators.

Problem-solving and reasoning 8, 9 9–11

8 A student thinks that the LCD to use when simplifying $\frac{x}{2}+\frac{3x}{4}$ is 8.

a Complete the simplification using a common denominator of 8.
b Now complete the simplification using the actual LCD of 4.
c How does your working for parts **a** and **b** compare? Which method is preferable and why?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

9J

9 Find and describe the error in each set of working. Then give the correct answer.

a $\frac{4x}{5} - \frac{x}{3} = \frac{3x}{2}$

b $\frac{x+1}{5} + \frac{x}{2} = \frac{2x+1}{10} + \frac{5x}{10}$

$= \frac{7x+1}{10}$

10 Simplify:

a $\frac{3}{x} + \frac{5}{2x}$

b $\frac{7}{3x} - \frac{2}{x}$

c $\frac{7}{4x} - \frac{5}{2x}$

d $\frac{4}{3x} + \frac{2}{9x}$



Hint for Q10: Your common denominator will contain x. The LCM of x and $2x$ is $2x$.

11 Simplify by first finding the LCD:

a $\frac{2x}{5} - \frac{3x}{2} - \frac{x}{3}$

b $\frac{x}{4} - \frac{2x}{3} + \frac{5x}{6}$

c $\frac{5x}{8} - \frac{5x}{6} + \frac{3x}{4}$

d $\frac{x+1}{4} + \frac{2x-1}{3} - \frac{x}{5}$

e $\frac{2x-1}{3} - \frac{2x}{7} + \frac{x-3}{6}$

f $\frac{1-2x}{5} - \frac{3x}{8} + \frac{3x+1}{2}$



Hint for Q11: Choose a common denominator that works for all three fractions.

The missing algebraic fractions — 12

12 Find the missing algebraic fraction. The fraction should be in simplest form.

a $\frac{x}{2} + \frac{\square}{\square} = \frac{5x}{6}$

b $\frac{\square}{\square} + \frac{x}{4} = \frac{3x}{8}$

c $\frac{2x}{5} + \frac{\square}{\square} = \frac{9x}{10}$

d $\frac{2x}{3} - \frac{\square}{\square} = \frac{7x}{15}$

e $\frac{\square}{\square} - \frac{x}{3} = \frac{5x}{9}$

f $\frac{2x}{3} - \frac{\square}{\square} = \frac{5x}{12}$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Automotive technology

The automotive industry is now more technically complex with the increased use of computer technology in cars. Training for automotive trades requires students to have mechanical ability as well as problem-solving and mathematical skills.

The machinists and technicians constructing and installing parts must have an eye for detail and excellent measurement skills.


Formulas are widely used in all trades to calculate quantities such as area, volume and maximum loads and friction. The universal tool for simplifying such formulas is algebra.




Maths@Work

1 Write the algebraic formula for the surface area of each solid below using the given pronumerals. Simplify each formula by taking out any common factors.


a




b



c



d



2 Match each of the following volume formulae to one of the solids shown in Question **1**.

i $V = \pi r^2 h$

ii $V = lwh$


iii $V = \frac{1}{2}abh$

iv $V = \frac{\pi d^2 h}{4}$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3 The cross-sectional area of a pipe is used when calculating the weight of a given length of pipe. To simplify the area formula, the algebraic difference of two squares rule is used.

a Copy and complete this procedure.



$$\begin{aligned}\text{Area of cross-section} &= \pi R^2 - \square \\ &= \pi(\square - r^2) \\ &= \pi\left(\frac{\square}{2^2} - \frac{d^2}{2^2}\right) \\ &= \frac{\pi}{4}(D^2 - \square) \\ &= \frac{\pi}{4}(D+d)(\square - \square)\end{aligned}$$

b Use the above formula to calculate the exact area of cross-section, as a multiple of π, given the following diameters of various pipes.

i $D = 40$ mm and $d = 36$ mm
ii $D = 32$ mm and $d = 30$ mm
iii $D = 2.7$ cm and $d = 2.3$ cm
iv $D = 6.5$ cm and $r = 5.5$ cm

Using technology



Hint for Q4: When entering the formula for cross-sectional area use pi() for π. Link all the length cells' formulae to cell F2 by using $ signs, i.e. =$F$2. Format area and weight cells to 'number' with two decimal places.

4 The approximate weight of a length of pipe can be determined by this formula:

$$\text{weight} = \text{cross-section area (cm}^2\text{)} \times \text{length (cm)} \times \text{density (g/cm}^3\text{)}$$

Develop the Excel spreadsheet shown below to calculate the weights of various lengths of pipe. Enter appropriate formulas into the shaded cells.

	A	B	C	D	E	F	G	H
1	Weight of pipe lengths							
2						100		
3	Pipe material	Outer diameter in cm	Inner diameter in cm	Area of cross-section in cm^2	Density in g/cm^3	Length in cm	Weight in g	Weight in kg
4	Poly pipe a	4.3	3.7		0.955			
5	Poly pipe b	12.5	11.9		0.955			
6	Poly pipe c	31.5	28.5		0.955			
7	Copper pipe a	1.2	1		8.94			
8	Copper pipe b	2	1.9		8.94			
9	Copper pipe c	5	4.8		8.94			
10	Steel pipe a	4.8	4.4		7.85			
11	Steel pipe b	6	5		7.85			
12	Steel pipe c	16.5	15.5		7.85			


a Find the difference in kg/m between copper pipe c and steel pipe b.
b Find the difference in weight between 3.5 m of poly pipe a and 3.5 m of steel pipe a.
c Find the difference in weight between 2.75 m of poly pipe c and 5 m of steel pipe c.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Square castle moats

Moats were constructed around castles to help protect its occupants from invaders. A square castle of side length 40 metres is to be surrounded by a square moat of a certain width as shown.

Present a report for the following tasks and ensure that you show clear mathematical workings, explanations and diagrams where appropriate.



1 Preliminary task

a Find the area of land occupied by the square castle of side 40-metres.

b If the width of the moat is 10 metres, find the surface area of the water in the moat.

c Find the outside perimeter of the moat if the width of the moat is 8.5 metres.

2 Modelling task

The problem is to determine a width for the moat so that the surface area of water is at least 10 times the area of the castle.

Formulate

a Write down all the relevant information that will help solve this problem with the aid of a diagram.

b If the width of the moat is x metres, find expressions in terms of x for the following:

i the perimeter of the outside of the moat

ii the total area occupied by the moat and the castle.

Solve

c Use your expressions from part **b** to show that the area occupied by the moat, not including the castle, can be written in the form $4x(40 + x)$.

d Using the expression from part **c**, find the area occupied by the moat for the following values of x.

i 10 **ii** 40 **iii** 50

e For each value of x in part **d**, calculate the value of $\frac{\text{Moat area}}{\text{Castle area}}$.

f Using trial and error, determine the **smallest integer** value of x such that the area of the moat is at least 10 times the area of the castle.

Evaluate and verify

g Assuming that the lord of the castle wants the area of the moat to be as close as possible to 10 times the area of the castle, investigate if this is possible using non-integer values of x to one decimal place.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Communicate

h Summarise your results using a table like the one below and describe any key findings.

Moat width x metres					
Area of moat in m^2					
Area of castle in m^2					
$\frac{\text{Moat area}}{\text{Castle area}}$					

3 Extension questions

a Instead, the lord wants to build a square castle of side length x metres inside a square lake of side length 50 metres.

i Draw a labelled diagram showing the square castle within the square moat.

ii Show that the remaining lake area forming the moat is given by the expression $(50 + x)(50 - x)$.

iii Calculate the value of $\frac{\text{Moat area}}{\text{Castle area}}$ to one decimal place, when:

I $x = 13$

II $x = 17$

iv Determine the smallest integer width of the castle that delivers a moat area which is at least 10 times the area of the castle.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Generating framing pictures


Key technology: Spreadsheets and programming

Area problems can often be solved using quadratic equations because the multiplication of a length and a width, both of which depend on the variable x, will result in an expression including a x^2 term. For example, a painting surrounded by a picture frame of width x cm will have an area which is a quadratic expression in terms of x. The coefficients will depend on the dimensions of the overall framed painting.



1 Getting started

The given diagram shows a picture frame of equal width x cm surrounding a square painting. The total width of the framed painting including the frame is 50 cm.



a Write an expression in terms of x for the side length of the painting inside the frame.

b Give the realistic values of x and give reasons.


c Write an expression for the area of the painting. Expand and simplify your expression.

d Use your expression from part **c** to find the area of the painting for the following values of x.
i 3 **ii** 5 **iii** 10

2 Applying an algorithm

Rather than solving for x for a given painting area algebraically, we can search for a solution numerically. Consider an algorithm which searches for a value of x which makes the area of the painting equal to 1000 cm^2.

This flowchart describes a simple algorithm which asks for a starting value of x. This should be a small integer value for which the area of the painting is greater than 1000 cm^2. $x = 5$ is chosen.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

a Starting with an x value of 5 run through the algorithm and fill in the table of values for the output of x and A.

b Use your result above to decide between which two integers lies the x value that makes the area of the painting equal to 1000 cm^2.

c Modify the algorithm and complete parts **a** and **b** above if the area of the painting is instead 2000 cm^2.

x	A
5	1600
6	

3 Using technology

a Set up a spreadsheet which calculates the area of the painting for a given value of x as shown.

b Fill down at cells A7 and B6 and locate the integer value of x which gives the area closest to 1000 cm^2.

c Now try to find a more accurate value of x for which the area is equal to 1000 cm^2 by changing the following. Fill down to locate the value of x.
- Alter the formula in cell A7 to be $= A6 + 0.1$
- Alter the starting value in cell A6 to 8

	A	B
1	**Picture frames**	
2		
3	**Painting area**	**1000**
4		
5	**x value**	**Area**
6	1	=4*A6^2-200*A6+2500
7	=A6+1	

d Repeat part **c** using a smaller increment until you are satisfied that you have found the correct value of x to one decimal place.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

1 'You use me when dividing fractions.' Simplify each expression, then match the answers to the letters to solve the riddle.

P	L	C	R
$-2x \times 3x$	$(x+1)(x-1)$	$\frac{2x-6}{2}$	$\frac{5x^2}{10x}$


O	A	E	I
$\frac{x}{2} - \frac{x}{3}$	$\frac{x+3}{2} \times \frac{6}{x+3}$	$\frac{2x}{3} \div \frac{4}{9}$	$7ab - 6ba$

☐	☐	☐	☐	☐	☐	☐	☐	☐	☐
$\frac{x}{2}$	$\frac{3x}{2}$	$x-3$	ab	$-6x^2$	$\frac{x}{2}$	$\frac{x}{6}$	$x-3$	3	x^2-1

2 A square framed photograph has side length 20 cm and frame width x cm.


a Find an expanded expression for the area of the picture.

b What value of x makes the picture $\frac{1}{4}$ of the total area?




3 Write an expression that gives the total number of dots required for the nth diagram.


a



b



c



4 Simplify these algebraic fractions. Don't be fooled by the double negative.

a $\frac{x+1}{3} - \frac{x-2}{4}$ **b** $\frac{3x-4}{5} - \frac{4x-2}{3}$

5 **a** The difference between the squares of two consecutive numbers is 97. What are the two numbers?

b The difference between the squares of two consecutive odd numbers is 136. What are the two numbers?

c The difference between the squares of two consecutive multiples of 3 is 81. What are the two numbers?

6 In a race over 4 km, Ryan ran at a constant speed. Sophie ran the first 2 km at a speed 1 km/h faster than Ryan. She ran the second 2 km at a speed 1 km/h slower than Ryan. Who won the race?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter summary

Algebraic techniques

Difference of two squares

$a^2 - b^2 = (a - b)(a + b)$

e.g. $x^2 - 16 = x^2 - 4^2$
$= (x - 4)(x + 4)$

e.g. $4x^2 - 9 = (2x)^2 - 3^2$
$= (2x - 3)(2x + 3)$

By grouping in pairs

If there are four terms, it may be possible to group into two binomial terms.

e.g. $5x + 10 - 3x^2 - 6x$
$= 5(x + 2) - 3x(x + 2)$
$= (x + 2)(5 - 3x)$

Monic quadratic trinomials

These are of the form:
$x^2 + bx + c$

Require two numbers that multiply to c and add to b

e.g. $x^2 - 7x - 18 = (x - 9)(x + 2)$
since $-9 \times 2 = -18$
and $-9 + 2 = -7$

Multiplying and dividing terms

$-3ab \times 2a = -6a^2b$

$15xy \div (3x^2y) = \frac{15xy}{3x^2y} = \frac{5}{x}$

Factorisation

The reverse process of expansion. Always remove the highest common factor first.

E.g. $2ab + 8b$ HCF $= 2b$
$= 2b(a + 4)$

E.g. $3x - 9 = 3(x - 3)$ HCF $= 3$
$-4x - 8 = -4(x + 2)$ HCF $= -4$

Expansion

The process of removing brackets

E.g. $2(x + 5) = 2x + 10$



$(x + 3)(2x - 4) = 2x^2 - 4x + 6x - 12$
$= 2x^2 + 2x - 12$

Review

3-term expression

$\overbrace{7xy - 4yx + 6}$

like terms (can collect) — constant term

The coefficient of x in $4 - 2x$ is -2.

Difference of two squares

$(a - b)(a + b) = a^2 - b^2$

E.g. $(x - 5)(x + 5) = x^2 + 5x - 5x - 25$
$= x^2 - 25$

Perfect squares

$(a + b)^2 = a^2 + 2ab + b^2$
$(a - b)^2 = a^2 - 2ab + b^2$

E.g. $(2x + 3)^2 = (2x + 3)(2x + 3)$
$= 4x^2 + 6x + 6x + 9$
$= 4x^2 + 12x + 9$

Simplifying by cancelling

E.g. $\frac{7(x+1)}{2(x+1)} = \frac{7}{2}$

E.g. $\frac{5x - 10}{5} = \frac{{}^{1}\cancel{5}(x - 2)}{\cancel{5}_{1}} = x - 2$

Algebraic fractions ⭐

These involve algebraic expressions in the numerator and/or denominator.

E.g. $\frac{2x}{x - 7}$, $\frac{3}{5x}$, $\frac{7}{(x - 1)^2}$

Add/subtract ⭐

Must find lowest common denominator (LCD) before applying operation.

E.g. $\frac{2x}{3} - \frac{x}{4} = \frac{8x}{12} - \frac{3x}{12}$
$= \frac{5x}{12}$

E.g. $\frac{x - 1}{2} + \frac{x - 3}{3} = \frac{3(x - 1)}{6} + \frac{2(x - 3)}{6}$
$= \frac{3x - 3 + 2x - 6}{6}$
$= \frac{5x - 9}{6}$

Multiplying ⭐

Cancel any numerator with any denominator.

E.g. $\frac{\cancel{3x + 2}^{1}}{\cancel{5}_{1}} \times \frac{\cancel{15}^{3}}{\cancel{2(3x + 2)}_{1}}$
$= \frac{3}{2}$

Dividing ⭐

To divide, multiply by the reciprocal. Factorise all expressions, cancel and then multiply.

E.g. $\frac{3x}{4} \div \frac{9x}{8}$
$= \frac{{}^{1}\cancel{3x}}{\cancel{4}_{1}} \times \frac{\cancel{8}^{2}}{\cancel{9x}_{3}}$
$= \frac{2}{3}$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter checklist

A version of this checklist that you can print out and complete can be downloaded from your Interactive Textbook.

		✔
9A	**1 I can identify and collect like terms.** e.g. Simplify: **a** $4a^2 - ab - 2a^2 + 1$ **b** $5x^2y - xy - 6yx^2$	
9A	**2 I can evaluate an algebraic expression using substitution.** e.g. If $x = 2$ and $y = -5$, evaluate $y^2 - xy$.	
9A	**3 I can multiply and divide algebraic terms.** e.g. Simplify: **a** $2ab \times 7a$ **b** $8xy \div (24x)$	
9A	**4 I can expand expressions with brackets.** e.g. Expand the following: **a** $-3(x - 2)$ **b** $-2a(4a - 5)$	
9B	**5 I can expand binomial products.** e.g. Expand: **a** $(x - 3)(2x + 5)$ **b** $(3x - 7)(2x - 5)$	
9C	**6 I can expand perfect squares.** e.g. Expand: **a** $(x + 7)^2$ **b** $(4x - 3)^2$	
9D	**7 I can expand to form a difference of two squares.** e.g. Expand: **a** $(x - 4)(x + 4)$ **b** $(3x + 5)(3x - 5)$	
9E	**8 I can determine the HCF.** e.g. Determine the HCF of the terms $6xy$ and $18xy^2$.	
9E	**9 I can factorise expressions with common factors.** e.g. Factorise the following: **a** $5a + 15$ **b** $11x^2 - 33xy$ **c** $-14mn - 7m$ (including common negative)	
9F	**10 I can factorise a difference of two squares.** e.g. Factorise $x^2 - 100$ and $4a^2 - 9$.	
9G	**11 I can factorise by grouping in pairs.** e.g. Use the method of grouping in pairs to factorise $x^2 + 6x - 3x - 18$.	
9G	**12 I can factorise using grouping in pairs by first rearranging.** e.g. Factorise $2x^2 - 9 - 6x + 3x$.	

Essential Mathematics Core for the Victorian Curriculum Year 9 ISBN 978-1-009-59315-1 © Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter checklist

		✔
9H	**13 I can factorise a quadratic trinomial.** e.g. Factorise $x^2 + 4x - 21$.	
9F/H	**14 I can factorise with trinomials or difference of two squares by first taking out a common factor.** e.g. Fully factorise $2x^2 - 18$ and $3x^2 - 21x + 36$.	
9I	**15 I can simplify algebraic fractions.** e.g. Simplify this fraction by factorising first: $\frac{2x-4}{x-2}$.	
9I	**16 I can multiply algebraic fractions.** e.g. Simplify $\frac{x+3}{6} \times \frac{9}{x+3}$.	
9I	**17 I can divide algebraic fractions.** e.g. Simplify $\frac{5(x+2)}{8} \div \frac{10(x+2)}{4}$.	
9J	**18 I can add and subtract simple algebraic fractions.** e.g. Simplify: **a** $\frac{2x}{5} - \frac{3x}{10}$ **b** $\frac{x-3}{4} + \frac{2x-5}{3}$	

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Short-answer questions

9A **1** Simplify these expressions.

a $3x \times 2x$ **b** $-3ab \times 6b$ **c** $7ab \div (14a)$ **d** $-40x^2y \div (10xy)$

9A **2** Expand the following.

a $3(x-4)$ **b** $-2(x+6)$ **c** $-x(x-1)$ **d** $-4x(2x-3)$

9A **3** Collect like terms to simplify.

a $5x - 1 + 2x$ **b** $7ab - 2ba$ **c** $4a^2b - 7a^2b$ **d** $5x^2 - xy + 3x^2$

9B **4** Expand the following binomial products.

a $(x-3)(x+4)$ **b** $(x-7)(x-2)$ **c** $(2x-3)(3x+2)$ **d** $(x-1)(3x+4)$

9C/D **5** Expand the following.

a $(x+3)^2$ **b** $(x-4)^2$ **c** $(3x-2)^2$

d $(x-5)(x+5)$ **e** $(7-x)(7+x)$ **f** $(11x-4)(11x+4)$

9E **6** Write the following in fully factorised form by removing the highest common factor.

a $4a + 12b$ **b** $x^2 + x$

c $6x - 9x^2$ **d** $-3 - 9x$

9F **7** Factorise the following difference of two squares.

a $x^2 - 100$ **b** $25x^2 - y^2$

c $49 - 9x^2$ **d** $3x^2 - 48$

9G **8** Factorise the following by grouping in pairs.

a $x^2 - 3x + 6x - 18$ **b** $4x^2 + 10x - 2x - 5$ **c** $3x - 8b + 2bx - 12$

9H **9** Factorise the following trinomials.

a $x^2 + 8x + 15$ **b** $x^2 - 3x - 18$

c $x^2 - 7x + 6$ **d** $3x^2 + 15x - 42$

9I **10** Simplify the following.

a $\frac{3(x-1)}{6}$ **b** $\frac{(x+1)(x-3)}{x-3}$

c $\frac{3x+12}{3}$ **d** $\frac{-5x-15}{10}$

e $\frac{2x-16}{3x-24}$ **f** $\frac{2(x+2)(x-1)}{x+2}$

9I **11** Simplify the following algebraic fractions by first factorising and cancelling where possible.

a $\frac{3}{2x} \times \frac{x}{6}$ **b** $\frac{7a^2}{x+1} \times \frac{x+1}{14a}$ **c** $\frac{x(x-4)}{8(x+1)} \times \frac{4(x+1)}{x}$

d $\frac{3a}{5} \div \frac{9}{10}$ **e** $\frac{2(x-1)}{3} \div \frac{x-1}{6}$ **f** $\frac{2x}{5x+20} \div \frac{x}{x+4}$

9J **12** Simplify the following by first finding the lowest common denominator.

a $\frac{x}{4} + \frac{2x}{3}$ **b** $\frac{5x}{6} - \frac{7x}{8}$ **c** $\frac{3a}{4} - \frac{a}{2}$

d $\frac{x}{2} + \frac{x-1}{3}$ **e** $\frac{2x-3}{4} + \frac{x+1}{3}$ **f** $\frac{x-1}{6} + \frac{x+3}{8}$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter review

Multiple-choice questions

9A **1** The constant term in the expression $2x^2 - 3$ is:

A 3 **B** 2 **C** -2 **D** $2x^2$ **E** -3

9A **2** The expanded form of $-2(x + 3)$ is:

A $2x + 6$ **B** $-2x - 3$ **C** $-2x - 6$ **D** $-2x + 6$ **E** $-2x + 3$

9A **3** $x^2 - 2xy + 2yx$ is equal to:

A xy **B** x^2 **C** $x^2 - 4xy$ **D** 0 **E** $4xy$

9A **4** $-3ab \times 4b$ is equal to:

A $-7ab^2$ **B** $-12ab^2$ **C** $-7a^2b$ **D** $-12a^2b$ **E** $12ab^2$

9B **5** $(x - 3)(x - 4)$ expands to:

A 12 **B** $x^2 + 12$ **C** $x^2 - 7x + 12$ **D** $x^2 - x - 12$ **E** $x^2 + 7x - 12$

9B **6** $(2x - 1)(x + 5)$ in expanded and simplified form is:

A $2x^2 + 9x - 5$ **B** $x^2 + 11x - 5$ **C** $4x^2 - 5$ **D** $3x^2 - 2x + 5$ **E** $2x^2 + 4x - 5$

9C **7** $(3a + 2b)^2$ is equivalent to:

A $9a^2 + 6ab + 4b^2$ **B** $9a^2 + 4b^2$ **C** $3a^2 + 6ab + 2b^2$
D $3a^2 + 12ab + 2b^2$ **E** $9a^2 + 12ab + 4b^2$

9F **8** $16x^2 - 49$ in factorised form is:

A $(4x - 7)^2$ **B** $(16x - 49)(16x + 49)$ **C** $(2x - 7)(8x + 7)$
D $(4x - 7)(4x + 7)$ **E** $4\left(4x^2 - 49\right)$

9G **9** The factorised form of $x^2 + 3x - 2x - 6$ is:

A $(x - 3)(x + 2)$ **B** $x - 2(x + 3)^2$ **C** $(x + 3)(x - 2)$
D $x - 2(x + 3)$ **E** $x(x + 3) - 2$

9H **10** If $(x - 2)$ is a factor of $x^2 + 5x - 14$ the other factor is:

A x **B** $x + 7$ **C** $x - 7$
D $x - 16$ **E** $x + 5$

9I **11** $\frac{5x}{3} \div \frac{15x}{12}$ is equal to:

A $\frac{4}{3}$ **B** $\frac{4x}{3}$ **C** $\frac{4}{3x}$ **D** $\frac{25x^2}{12}$ **E** $\frac{25x}{12}$

9J **12** $\frac{4x}{3} - \frac{2x}{9}$ is equal to:

A $\frac{10x}{27}$ **B** $\frac{30x}{9}$ **C** $\frac{2x}{9}$ **D** $\frac{2x}{3}$ **E** $\frac{10x}{9}$


9J **13** $\frac{x + 2}{5} + \frac{2x - 1}{3}$ written as a single fraction is:

A $\frac{11x + 1}{15}$ **B** $\frac{11x + 9}{8}$ **C** $\frac{3x + 1}{8}$ **D** $\frac{11x + 7}{15}$ **E** $\frac{13x + 1}{15}$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Extended-response questions

1 A pig pen for a small farm is being redesigned. It is originally a square of side length x m.



a In the planning, the length is initially kept as x m. The width is altered such that the area of the pen is $(x^2 + 3x)$ square metres. What is the new width? (Consider factorising.)

b Instead, it is decided that the original length will be increased by 1 metre and the original width will be decreased by 1 metre.

i What effect does this have on the perimeter of the pig pen compared with the original size?

ii Work out an expression for the new area of the pig pen in expanded form. How does this compare to the original area?


c For the pig pen described in part **b**:

i what would be the area if $x = 2$?

ii what value of x gives an area of 15 m^2? Use trial and error.



2 The security tower for a palace is on a small square piece of land 20 m by 20 m. It has a moat of width x metres the whole way around it, as shown.



a State the area of the piece of land.

b **i** Give an expression for the side length of the combined moat and land.

ii Find an expression, in expanded form, for the entire area occupied by the moat and the land.

c Use your answers to parts **a** and **b** to give an expression for the area occupied by the moat alone. Answer in factorised form.

d Use trial and error to find the value of x such that the area of the moat alone is 500 m^2.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.



10 Statistics and probability

Essential mathematics: why skills in statistics and probability are important

Statistics and probability are useful tools for analysing data, making predictions, and understanding the likelihood of events. Many industries apply these mathematical fields practically.

Meteorologists use probability and data from weather stations, satellites, and radars to predict weather. They analyse atmospheric pressure, humidity, temperature, and wind speed, then input this data into computer models. If models show moist air converging with a low-pressure system, meteorologists might predict a 70% chance of rain tomorrow, helping people plan activities.

Epidemiologists use probability and mathematical models to track disease spread and plan interventions. They analyse infection rate data and use the basic reproduction number R_0 to see how many others one sick person can infect. This involves calculating the probability of infection spreading. If R_0 is greater than 1, the disease can spread quickly.

Manufacturers use statistics in quality control to monitor production and maintain high-quality products. They analyse defect data and use probability to predict future issues, scheduling maintenance before problems occur, reducing waste, and increasing efficiency.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.



In this chapter

10A Review of probability (Consolidating)
10B Venn diagrams and two-way tables
10C Using arrays for two-step experiments
10D Using tree diagrams
10E Using relative frequencies to estimate probabilities
10F Data and sampling
10G Summarising data: range and measures of centre
10H Interpreting data from tables and graphs
10I Stem-and-leaf plots
10J Grouped data

Victorian Curriculum 2.0

This chapter covers the following content descriptors in the Victorian Curriculum 2.0:

STATISTICS

VC2M9ST01, VC2M9ST02, VC2M9ST03, VC2M9ST04, VC2M9ST05

PROBABILITY

VC2M9P01, VC2M9P02, VC2M9P03

Please refer to the curriculum support documentation in the teacher resources for a full and comprehensive mapping of this chapter to the related curriculum content descriptors.

© VCAA

Online resources

A host of additional online resources are included as part of your Interactive Textbook, including HOTmaths content, video demonstrations of all worked examples, auto-marked quizzes and much more.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025
Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Warm-up quiz

1 Write as decimals:

a $\frac{1}{10}$ **b** $\frac{2}{8}$ **c** 30% **d** 85% **e** 23.7%

2 Express in simplest form:

a $\frac{4}{8}$ **b** $\frac{7}{21}$ **c** $\frac{20}{30}$ **d** $\frac{100}{100}$ **e** $\frac{0}{4}$

f $\frac{12}{144}$ **g** $\frac{36}{72}$ **h** $\frac{36}{58}$ **i** $\frac{72}{108}$ **j** $\frac{2}{7}$

3 A six-sided die is tossed.

a List the possible outcomes.

b How many of the outcomes are:

i even? **ii** less than 3? **iii** less than or equal to 3?


iv at least 2? **v** not a 6? **vi** not odd?

4 From the group of the first 10 integers $\{1, 2, \ldots, 10\}$, how many of the numbers are:

a odd? **b** less than 8? **c** greater than or equal to 5?

d no more than 7? **e** prime? **f** not prime?

5 Several cards were randomly selected from a pack of playing cards. The suit of each card was noted, the card was replaced and the pack was shuffled. The frequency of each suit is shown in the column graph.



a How many times was a heart selected?

b How many times was a card selected in total?

c In what fraction of the trials was a diamond selected?

6 Consider this simple data set: 1, 2, 3, 5, 5, 7, 8, 10, 13.

a State the number of scores, n.

b Find the mean (the sum of scores divided by n).

c Find the median (the middle value).

d Find the mode (the most common value).

e Find the range (the difference between the highest and lowest value).

f Find the probability of randomly selecting:

i a 5

ii a number that is not 5

iii a number that is no more than 5

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10A Review of probability CONSOLIDATING

Learning intentions

- To understand the concept of chance and how to describe it numerically
- To be able to interpret the language of probability
- To be able to find the probability of an event for equally likely outcomes

Key vocabulary: probability, experiment, sample space, outcome, event, complement, chance

The mathematics used to calculate chance is called probability. We use it to compare the number of favourable outcomes to the total number of outcomes. This shows us how likely it is that the favourable event will occur. The probability of an event occurring is a number between 0 and 1. An impossible event has a probability of 0. An event that is certain to occur has a probability of 1.




We can use probability to find the likelihood of rolling a particular total score with two dice.

Lesson starter: Events and probabilities

As a class group, write down and discuss at least three events that have the following chance of occurring.

- impossible chance
- very low chance
- medium to low chance
- even (50 : 50) chance
- medium to high chance
- very high chance
- certain chance

For each decimal marked on the number line below, choose the probability description from the list above that best matches it.



Key ideas

- A random **experiment** has various possible **outcomes** that occur without interference.
- An outcome is a possible result from a **chance** experiment.
- The **sample space** is the list of all possible outcomes of an experiment.
- An **event** is a collection of outcomes resulting from an experiment.
  For example, rolling a die is a random experiment with six possible outcomes: 1, 2, 3, 4, 5 and 6.
  The event 'rolling a number greater than 4' includes the outcomes 5 and 6.
  The event 'rolling a number at least 4' includes the outcomes 4, 5 and 6.
- **Probability** is a measure of the likelihood that an event will occur.
- The probability of an event where all outcomes are equally likely is given by:


$$\text{Pr(Event)} = \frac{\text{Number of outcomes where event occurs}}{\text{Total number of outcomes}}$$

- Probabilities are numbers between 0 and 1 and can be written as a decimal, fraction or percentage.
  For example: 0.55 or $\frac{11}{20}$ or 55%

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10A

- For all events, $0 \leq \Pr(\text{Event}) \leq 1$.



- The **complement** of event A is written A′ (or not A). A′ is the event that A does not occur.

$\Pr(A') = 1 - \Pr(A)$ or $\Pr(\text{not A}) = 1 - \Pr(A)$

Exercise 10A

Understanding

1–4 | 3,4

1 Write the missing words in each statement.

a The _____ _____ is the list of all possible outcomes in an experiment.

b An _____ is a collection of the outcomes from an experiment.

c Pr stands for _____.

d An _____ event has a probability of 0, an event that is _____ to occur has a probability of 1.

e An event with probability of 0.5 has _____ or 50 : 50 chance of occurring.

f The _____ of an event, A, is the event where A does not occur.



2 Jim believes that there is a 1 in 4 chance that the flower on his prized rose will bloom tomorrow.

Hint for Q2: '1 in 4' means $\frac{1}{4}$.

a Write the chance '1 in 4' as:



i a fraction **ii** a decimal **iii** a percentage



b Draw a number line from 0 to 1 and mark the level of chance described by Jim.

Hint for Q3: Percentage ÷ 100 = decimal

3 Copy and complete this table.

	Percentage	Decimal	Fraction	Number line
a	50%	0.5	$\frac{1}{2}$	
b	25%			
c			$\frac{3}{4}$	
d				

4 Ten people make the following guesses of the chance that they will get a salary bonus this year.



Hint for Q4: First write each chance as a decimal.

$0.7, \frac{2}{5}, 0.9, \frac{1}{3}$, 2 in 3, $\frac{3}{7}$, 1 in 4, 0.28, $\frac{2}{9}$, 0.15

Order their chances from lowest to highest.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


Fluency

5(½), 6–9 | 5–7(½), 8, 9(½)

Example 1 Finding probabilities of events

This spinner has five equally divided sections.

a List the sample space using the different numbers.
b Find Pr(3).
c Find Pr(not a 3).
d Find Pr(a 3 or a 7).
e Find Pr(a number that is at least a 3).



Solution

a $\{1, 2, 3, 7\}$

b $\Pr(3) = \frac{2}{5}$ or 0.4

c $\Pr(\text{not a } 3) = 1 - \Pr(3)$
$= 1 - \frac{2}{5}$ or $1 - 0.4$
$= \frac{3}{5}$ or 0.6

d $\Pr(\text{a 3 or a 7}) = \frac{2}{5} + \frac{1}{5}$
$= \frac{3}{5}$

e $\Pr(\text{at least a 3}) = \frac{3}{5}$

Explanation

Use set brackets, {}, and list all the possible outcomes in any order.

$\Pr(3) = \frac{\text{number of sections labelled 3}}{\text{number of equal sections}}$

'Not a 3' is the complementary event of obtaining a 3.

Alternatively, count the number of sectors that are not 3.


There are two 3s and one 7 in the five sections.

Three of the sections have the numbers 3 or 7, which are at least 3; i.e. 3 or more.

Now you try

This spinner has six equally divided sections.

a List the sample space using the different numbers.
b Find Pr(2).
c Find Pr(not a 1).
d Find Pr(a 2 or a 3).
e Find Pr(a number which is at most 2).



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10A


5 The spinners below have equally divided sections. Complete the following for each spinner. Write all probabilities as simplified fractions.

- **i** List the sample space using the given numbers.
- **ii** Find Pr(2).
- **iii** Find Pr(not a 2).
- **iv** Find Pr(a 2 or a 3).
- **v** Find Pr(a number which is at least a 2).




Hint for Q5: Pr(event) means 'find the probability of the event'. 'At least 2' means '2 or more'.


a




b




c




d



e



f



6 Find the probability of obtaining a blue ball if a ball is selected at random from a box that contains:

- **a** 4 blue balls and 4 red balls
- **b** 3 blue balls and 5 red balls
- **c** 1 blue ball, 3 red balls and 2 white balls
- **d** 8 blue balls, 15 black balls and 9 green balls
- **e** 15 blue balls only
- **f** 5 yellow balls and 2 green balls



Hint for Q6: $\Pr(\text{blue}) = \frac{\text{number of blue balls}}{\text{total number of balls}}$

7 Find the probability of *not* selecting a blue ball if a ball is selected at random from a box containing the balls described in Question **6**, parts **a** to **f**, above.



Hint for Q7: Pr(not blue) = 1 − Pr(blue)

8 If a swimming pool has eight lanes and each of eight swimmers has an equal chance of being placed in lane 1, find the probability that a particular swimmer:

- **a** will swim in lane 1
- **b** will not swim in lane 1

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Example 2 Choosing letters from a word

A letter is randomly chosen from the word PROBABILITY. Find the following probabilities.

a Pr(L)
b Pr(not L)
c Pr(vowel)
d Pr(consonant)
e Pr(vowel or a B)
f Pr(vowel or consonant)

Solution	Explanation
a $\Pr(\text{L}) = \frac{1}{11}$	One of the 11 letters is an L.
b $\Pr(\text{not L}) = 1 - \frac{1}{11}$ $= \frac{10}{11}$	The event 'not L' is the complement of the event selecting an L. Complementary events add up to 1. Pr(not L) = 1 − Pr(L).
c $\Pr(\text{vowel}) = \frac{4}{11}$	The vowels of the alphabet are A, E, I, O and U. There are 4 vowels in PROBABILITY: O, A and two letter Is.
d $\Pr(\text{consonant}) = 1 - \frac{4}{11}$ $= \frac{7}{11}$	The events 'vowel' and 'consonant' are complementary. Alternatively, the other 7 letters are consonants; thus $\frac{7}{11}$.
e $\Pr(\text{vowel or a B}) = \frac{6}{11}$	There are 4 vowels and 2 letter Bs.
f $\Pr(\text{vowel or consonant}) = 1$	This event includes all possible outcomes, since a letter is either a vowel or a consonant.

Now you try

A letter is chosen from the word EXPERIMENT. Find the following probabilities.

a Pr(M)
b Pr(not an E)
c Pr(vowel)
d Pr(consonant)
e Pr(E or consonant)
f Pr(letter from the first half of the alphabet)

9 A letter is chosen at random from the word ALPHABET. Find the following probabilities.

a Pr(L)
b Pr(A)
c Pr(A or L)
d Pr(vowel)
e Pr(consonant)
f Pr(vowel or consonant)
g Pr(Z)
h Pr(A or Z)
i Pr(not an A)
j Pr(letter from the first half of the alphabet)



Hint for Q9: Recall that vowels are A, E, I, O and U.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10A

Problem-solving and reasoning

10, 11 | 11–13

10 The school captain is to be chosen at random from four candidates. Two are girls (Hayley and Alisa) and two are boys (Rocco and Stuart).

a List the sample space.

b Find the probability that the school captain will be:

i Hayley

ii male

iii neither Stuart nor Alisa



Hint for Q10: The sample space is a list of the possible students.

11 A card is drawn at random from a pack of 52 playing cards. Find the probability that the selected card will be:

a the queen of diamonds

b an ace

c a red king


d a red card

e a jack or a queen

f any card except a 2

g any card except a jack or a black queen

h not a black ace



12 A six-sided die is tossed and the upper-most face is observed and recorded. Find the following probabilities.

a Pr(6)

b Pr(3)

c Pr(not a 3)

d Pr(1 or 2)

e Pr(a number less than 5)

f Pr(even number or odd number)

g Pr(square number)

h Pr(not a prime number)

i Pr(a number greater than 1)



Hint for Q12: 'Less than 5' means that 5 is not included. Square numbers are $1^2 = 1$, $2^2 = 4$, $3^2 = 9$ etc.

13 A letter is chosen at random from the word PROBABILITY. Find the probability that the letter will be:

a a B

b not a B

c a vowel

d not a vowel

e a consonant

f a letter belonging to one of the first five letters in the alphabet

g a letter from the word RABBIT

h a letter that is not in the word RABBIT

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Faulty record player — 14

14 A vinyl record contains eight tracks. The time length for each track is shown in the table on the right.

A vinyl record collector places the record on a faulty record player, which begins playing randomly at an unknown place somewhere on the record, not necessarily at the beginning of a track.

Track	Time (minutes)
1	3
2	4
3	4
4	5
5	4
6	3
7	4
8	4

a Find the total number of minutes of music available on the record.

b Find the probability that the record player will begin playing somewhere on track 1.

c Find the probability that the record player will begin somewhere on:

i track 2 **ii** track 3 **iii** a track that is 4 minutes long

iv track 4 **v** track 7 or 8 **vi** a track that is not 4 minutes long



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10B Venn diagrams and two-way tables

Learning intentions

- To understand how Venn diagrams and two-way tables are used to show how the sample space is distributed among events
- To be able to use a Venn diagram to display the distribution of two sets
- To be able to fill out a two-way table either from a problem or from a Venn diagram
- To be able to use a Venn diagram or two-way table to calculate probabilities of events


Key vocabulary: Venn diagram, two-way table, probability

When the results of an experiment involve overlapping categories it can be very helpful to organise the information into a Venn diagram or two-way table. Probabilities can easily be calculated from these types of diagrams.

Lesson starter: Solving puzzles

Work with a partner to find the answer to each puzzle. For Puzzles 1 – 4, draw some Venn diagrams, like the ones shown, to help you. For Puzzle 5, use a two-way table.


Puzzle 1



12 students travel to school by bus only and 10 students don't travel by either bus or train.

- How many students don't travel by train?


Puzzle 2



14 students in total have blue eyes, and of these students 5 have both blue eyes and blonde hair.

- How many have blue eyes but not blonde hair?


Puzzle 3



11 students own a bicycle only and 4 students own both a skateboard and a bicycle.

- How many students in total own a bicycle?

Puzzle 4



12 students in total like sport, 14 students in total like reading, and 9 students like both sport and reading.

- How many students altogether like either sport or reading or both?

Puzzle 5

A survey of 40 students found that a total of 22 play basketball, 9 play both volleyball and basketball and 6 do not play either basketball or volleyball.

- How many basketball players don't play volleyball?
- How many students in total don't play volleyball?
- How many students in total do play volleyball?


	Basketball	Not Basketball	Total
Volleyball			
Not Volleyball			
Total			

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Key ideas

- A **Venn diagram** and a **two-way table** help to organise outcomes into different categories. This example shows the types of computers owned by 100 people.

Venn diagram



Two-way table

	Mac	No Mac	Total
PC	12	50	62
No PC	31	7	38
Total	43	57	100

These diagrams show, for example, that:


- 12 people own both a Mac and a PC
- 62 people own a PC
- 57 people do not own a Mac
- $\Pr(\text{Mac}) = \frac{43}{100}$
- $\Pr(\text{only Mac}) = \frac{31}{100}$
- $\Pr(\text{Mac or PC}) = \frac{93}{100}$
- $\Pr(\text{Mac and PC}) = \frac{12}{100} = \frac{3}{25}$

Exercise 10B

Understanding

1–4 | 3, 4

1 This Venn diagram shows the number of people who enjoy riding and running.



a How many people in total are represented by this Venn diagram?

b Find how many people enjoy:

- **i** riding only
- **ii** riding (in total)
- **iii** running only
- **iv** running (in total)
- **v** both riding and running
- **vi** neither riding nor running
- **vii** riding or running or both?

c How many people do not enjoy:

i riding? **ii** running?



Hint for Q1b: 'Riding only' does not include the overlapping section.




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


10B

2 Each statement is about the shaded area in the Venn diagram. State the missing number in each statement.


a _____ people in total like pineapple.




b _____ people like pineapple only.




c _____ people like both pineapple and mango.




d _____ people like pineapple or mango or both.



e _____ people don't like mango.



f _____ person likes neither mango nor pineapple.




3 Match the diagrams A, B, C or D with the given description.

a own a cat
b own a cat only
c own both a cat and a dog
d own a cat or a dog or both




Hint for Q3: 'Both a cat and a dog' is where the circles overlap.


A




B



C



D



4 Fill in the missing numbers in these two-way tables.

a

	A	Not A	Total
B	7	8	
Not B		1	
Total	10		

b

	A	Not A	Total
B	2		7
Not B		4	
Total			20



Hint for Q4: Start with the column or row that has two numbers in it.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Fluency 5–8 5–9

Example 3 Constructing a Venn diagram


A survey of 30 people found that 21 like AFL and 12 like soccer. Also, 7 people like both AFL and soccer and 4 like neither AFL nor soccer.

a Construct a Venn diagram for the survey results.
b How many people:
  i like AFL or soccer?
  ii do not like soccer?
  iii like only AFL?
c If one of the 30 people was randomly selected, find:
  i Pr(like AFL and soccer)
  ii Pr(like neither AFL nor soccer)
  iii Pr(like only soccer)

Solution

Explanation

a



Place the appropriate number in each category.
Place the 7 in the overlap first, then ensure that:
- the total that like AFL is 21 $(14 + 7 = 21)$.
- the total that like soccer is 12 $(5 + 7 = 12)$.

The 4 people that like neither are placed outside the circles.

b i 26 like AFL or soccer or both

The total number of people who like AFL, soccer or both is $14 + 7 + 5 = 26$.

ii $30 - 12 = 18$ don't like soccer

12 like soccer, so 18 do not.

iii 14 like AFL but not soccer

21 like AFL but 7 of these also like soccer.

c i $\Pr(\text{like AFL and soccer}) = \frac{7}{30}$

7 out of 30 people like AFL and soccer.

ii $\Pr(\text{like neither AFL nor soccer})$

$= \frac{4}{30}$

$= \frac{2}{15}$

The 4 people who like neither AFL nor soccer sit outside both categories.

iii $\Pr(\text{like soccer only})$

$= \frac{5}{30}$

$= \frac{1}{6}$

5 people like soccer but not AFL.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10B

Now you try

A survey of 50 people found that 25 watch live TV and 32 people watch streamed shows. Also 14 people watch both live TV and streamed shows and 7 watch neither.


a Construct a Venn diagram for the survey results.

b How many people:
- i watch live TV or streamed shows?
- ii do not watch live TV?
- iii watch only streamed shows?

c If one of the 50 people was randomly selected, find:
- i Pr(watch live TV and streamed shows)
- ii Pr(watch neither live TV nor streamed shows)
- iii Pr(watch only live TV)

5 In a class of 30 students, 22 carried a phone and 9 carried an iPad. Three carried both a phone and an iPad and 2 students carried neither.

a Copy and complete this Venn diagram.





b How many students:
- **i** carried a phone or an iPad (includes carrying both)?
- **ii** do not carry an iPad?
- **iii** carry only an iPad?

c If one of the 30 students was selected at random, find the following probabilities.
- **i** Pr(carry a phone and an iPad)
- **ii** Pr(carry neither a phone nor an iPad)
- **iii** Pr(carry only a phone)



Hint for Q5: Start by writing a number in the overlapping section. '22 with a phone' is the total for the phone circle, including the overlap.

6 Match each diagram with the correct statement.
- **i** Not A
- **ii** A or B
- **iii** A and B
- **iv** B
- **v** B only
- **vi** Neither A nor B

Diagram P




Diagram Q




Diagram R




Diagram S




Diagram T





Diagram U






Hint for Q6: 'A or B' means 'either A or B or both'. 'A and B' means 'both A and B'.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


7 For each Venn diagram, find the following probabilities.

i Pr(A) **ii** Pr(A only)
iii Pr(not B) **iv** Pr(A and B)
v Pr(A or B) **vi** Pr(neither A nor B)

a




b






Hint for Q7:

A and B



A or B





Hint for Q7: First calculate the total number in each sample.

Example 4 Constructing a two-way table

At a car yard, 24 cars are tested for fuel economy. Eighteen of the cars run on petrol, 8 cars run on gas and 3 cars can run on both petrol and gas.

a Illustrate the situation using a two-way table.
b How many of the cars:
  i do not run on gas?
  ii run on neither petrol nor gas?
c Find the probability that a randomly selected car:
  i runs on gas
  ii runs on only gas
  iii runs on gas or petrol

Solution

a

	Gas	Not gas	Total
Petrol	3	15	18
Not petrol	5	1	6
Total	8	16	24

Explanation

Set up a table as shown and enter the numbers (in black) from the given information.

Fill in the remaining numbers (in red) ensuring that each column and row adds to the correct total.

b i 16

The total at the base of the 'Not gas' column is 16.

ii 1

The number at the intersection of the 'Not gas' column and the 'Not petrol' row is 1.

c i $\Pr(\text{gas}) = \frac{8}{24}$

$= \frac{1}{3}$

8 cars in total run on gas out of the 24 cars.

ii $\Pr(\text{only gas}) = \frac{5}{24}$

Of the 8 cars that run on gas, 5 of them do not also run on petrol.

iii $\Pr(\text{gas or petrol}) = \frac{15+5+3}{24}$

$= \frac{23}{24}$

Of the 24 cars, some run on petrol only (15), some run on gas only (5) and some run on gas and petrol (3).

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10B

Now you try

Of 100 clothing items in a small shop, 65 contained natural fibres, 45 contained artificial fibres and 20 contained both types of fibres.

a Illustrate the situation using a two-way table.

b How many items:
- i do not have artificial fibres?
- ii have neither type of fibre?

c Find the probability that a randomly selected item:
- i has natural fibres
- ii has only natural fibres
- iii has artificial or natural fibres

8 Of 50 desserts served at a restaurant one evening, 25 were served with ice cream, 21 were served with cream and 5 were served with both cream and ice cream.



Hint for Q8: 25 is the total for the ice cream row.



a Copy and complete this two-way table.

	Cream	Not cream	Total
Ice cream			
Not ice cream			
Total			

b How many of the desserts:
- i did not have cream?
- ii had neither cream nor ice cream?

c Find the probability that a chosen dessert:
- i had cream
- ii had only cream
- iii had cream or ice cream

9 Find the following probabilities using each of the given tables. First copy and complete each two-way table.

i Pr(A) ii Pr(not A) iii Pr(A and B)
iv Pr(A or B) v Pr(B only) vi Pr(neither A nor B)



Hint for Q9: 'Neither A nor B' is the same as 'Not A and Not B'.

a

	A	Not A	Total
B	3	1	
Not B	2		4
Total			

b

	A	Not A	Total
B		4	15
Not B	6		
Total			26


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Problem-solving and reasoning 10–12 11–14

10 For each two-way table, fill in the missing numbers then transfer the information to a Venn diagram.


a

	A	Not A	Total
B	2		8
Not B			
Total		7	12



b

	A	Not A	Total
B		4	
Not B	9		13
Total	12		



11 In a group of 17 people, 13 rented their house, 6 rented a car and 3 did not rent either a car or their house.

a Draw a Venn diagram, showing circles for 'Rents house' and 'Rents car'.

b How many people rented both a car and their house?

c Find the probability that one of them rented only a car.



Hint for Q11: 14 people will be in the circles, but $13 + 6 = 19$. How many must be in the overlap?

12 One hundred citizens were surveyed regarding their use of water in their garden. Of these, 23 said that they used tank water, 48 said that they used tap water and 41 said that they did not use water on their garden at all.

a Copy and complete this two-way table.

	Tank water	Not tank water	Total
Tap water			
Not tap water			
Total			

b How many people used both tank and tap water?

c What is the probability that one of the people uses only tap water?


d What is the probability that one of the people uses tap water or tank water?




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10B

13 All members of a ski club enjoy either skiing and/or snowboarding.
Seven enjoy only snowboarding, 16 enjoy skiing and 4 enjoy both snowboarding and skiing.




- **a** Copy and complete the Venn diagram shown.
- **b** How many people are in the club?
- **c** If a person from the club is randomly selected, what is the probability of choosing a snowboarder?
- **d** If a person is randomly selected out of the group that likes skiing, what is the probability of choosing a snowboarder?
- **e** What is the probability of choosing a skier out of the group that likes snowboarding?




14 Of a group of 30 cats, 24 like either tinned or dry food or both, 10 like only dry food and 5 like both tinned and dry food.

- **a** Find the probability that a selected cat likes only tinned food.
- **b** Out of the group of cats that eat dry food, what is the probability of selecting a cat that also likes tinned food?



Numbers challenge — 15

15 One hundred people were surveyed and it was found that 55 were males and 30 were doctors. The number of male doctors was 17. Copy and complete this Venn diagram and then determine the number of people in each question below.



- **a** The number who are neither male nor a doctor
- **b** The number who are not males
- **c** The number who are not doctors
- **d** The number who are male but not a doctor
- **e** The number who are a doctor but not male
- **f** The number who are female and a doctor
- **g** The number who are female or a doctor



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10C Using arrays for two-step experiments

Learning intentions

- To know how to list the sample space of a two-step experiment in an array/table
- To understand the difference between experiments carried out with replacement and without replacement
- To be able to construct arrays for two-step experiments with and without replacement and find associated probabilities

Key vocabulary: array, with replacement, without replacement, two-step experiment, sample space

Sometimes an experiment consists of two steps, such as tossing a single coin twice or rolling two dice. Or perhaps a card is pulled from a hat and then a spinner is spun. We can use tables to list the sample space for such experiments.

Lesson starter: The maths cup

This activity can be run in small groups or as a class.



- **a** Draw the table below on the whiteboard. Each student should also have a copy.
- **b** Each student selects one competitor as their 'own' (choose a winner!).
- **c** Students take turns to roll 2 dice and state the sum of the uppermost faces of the dice.
- **d** The total of each roll refers to the competitor number. When its number is rolled, that competitor moves another 100 m towards the finish line. A cross is placed in the cell to show the move.
- **e** The winning competitor is the first to reach the finish at 1000 m.
- **f** Keep rolling the dice until first, second and third places are decided.

	Competitor	100 m	200 m	300 m	400 m	500 m	600 m	700 m	800 m	900 m	1000 m
1	SCRATCHED										
2	Greased Lightning										
3	Flying Eagle										
4	Quick Stix										
5	Break a Leg										
6	Slow and Steady										
7	The Donkey										
8	The Wombat										
9	Tooting Tortoise										
10	Ripper Racer										
11	Speedy Gonzo										
12	Pharaoh										

Discussion questions

- Do you think that the competitor that won your maths cup race would always win?
- In this game, do all the competitors have the same chance of winning?
- Why is Competitor 1 scratched?
- When two dice are thrown, what are all the possible outcomes for the sum of the two uppermost faces?
- What are the competitor numbers that are highly likely to win this race?
- What are the competitor numbers that are unlikely to win this race?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10C

Key ideas

- An **array** (or table) can be used to list the sample space for experiments involving two steps.
- When listing outcomes, it is important to be consistent with the order for each outcome. For example: the outcome (heads, tails) is different from the outcome (tails, heads). So when two coins are tossed, the Coin 1 outcome is written first in each cell of the table.

		Coin 2	
		H	T
Coin 1	H	HH	HT
	T	TH	TT

  The sample space is shown in the central part of the table; i.e. {HH, HT, TH, TT}.
- The probability is still given by:

$$\text{Pr(Event)} = \frac{\text{number of outcomes where the event occurs}}{\text{total number of possible outcomes}}$$

  For example, when two coins are tossed, Pr(a tail and a head) $= \frac{2}{4} = \frac{1}{2}$
- Some experiments are conducted **without replacement**, which means that each individual trial outcome cannot be achieved again.
  For example: Two letters are chosen from the word CAT. 'Without replacement' means that the first letter chosen is removed and it is not possible to choose it again.

With replacement

		2nd		
		C	A	T
1st	C	(C, C)	(C, A)	(C, T)
	A	(A, C)	(A, A)	(A, T)
	T	(T, C)	(T, A)	(T, T)

9 outcomes

Without replacement

		2nd		
		C	A	T
1st	C	×	(C, A)	(C, T)
	A	(A, C)	×	(A, T)
	T	(T, C)	(T, A)	×

6 outcomes

Exercise 10C

Understanding

1–4 | 3, 4

1 A letter is chosen from the word HI and a letter is chosen from the word BYE.
Copy and complete this table to show the sample space.

	B	Y	E
H	HB		
I			



Hint for Q1: Write 'H' in each cell of the top srow. Write 'B' in each cell of the first column.

2 A coin and a four-sided die are tossed.

a Copy and complete this table to show the sample space.

	1	2	3	4
H				
T				

Hint for Q2: The sample space is all the outcomes in the central part of the table.



b How many outcomes are in the sample space?

c What is the probability of tossing a 'H3'?

d What is the probability of tossing a 'T2'?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


3 A coin is flipped and then a spinner with numbers 1 to 5 is spun. The possible outcomes are listed in the table below.

	1	2	3	4	5
H	H1	H2	H3	H4	H5
T	T1	T2	T3	T4	T5

a How many outcomes are possible?
b List the four outcomes in which an even number is displayed on the spinner.
c Hence, state the probability that an even number is displayed.
d List the outcomes for which the coin shows tails and the spinner shows an odd number.
e What is Pr(T, odd number)?

4 Two coins are tossed and the four possible outcomes are shown below.

		20-cent coin	
		H	T
50-cent coin	H	HH	HT
	T	TH	TT



a What is the probability that the 50-cent coin will be heads and the 20-cent coin will be tails?
b For which outcomes are the two coins displaying the same face?
c What is the probability of the two coins displaying the same face?

Fluency

5, 6 | 5–7

Example 5 Using a table for two-step experiments

A spinner with the numbers 1, 2 and 3 is spun, and then a card is chosen at random from cards containing the letters of ATHS.

a Draw a table to list the sample space of this experiment.
b How many outcomes does the experiment have?
c Find the probability of the combination 2S.
d Find the probability of an odd number being spun and the letter H being chosen.

Solution

a

	A	T	H	S
1	1A	1T	1H	1S
2	2A	2T	2H	2S
3	3A	3T	3H	3S

b There are 12 outcomes.

c $\Pr(2S) = \frac{1}{12}$

d $\Pr(\text{odd, H}) = \frac{2}{12} = \frac{1}{6}$

Explanation

The sample space of the spinner (1, 2, 3) is put into the left column.

The sample space of the cards (A, T, H, S) is put into the top row.

The table has $4 \times 3 = 12$ items in it.

All 12 outcomes are equally likely. Spinning 2 and choosing an S is one of the 12 outcomes.

Possible outcomes are 1H and 3H, so probability = $2 \div 12$.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10C

Now you try

A coin is flipped and an 8-sided die is rolled.

a Draw a table to list the sample space of this experiment.
b How many outcomes does the experiment have?
c Find the probability of the outcome H7.
d Find the probability of obtaining a tail and a number less than 4.

5 A coin is flipped and then a regular die is rolled.

a Copy and complete this table to list the sample space for this experiment.

	1	2	3	4	5	6
H						
T						

b How many possible outcomes are there?
c Find the probability of the pair H3.
d Find the probability of 'heads' on the coin with an odd number on the die.

6 A letter is chosen from the word LINE and another is chosen from the word RIDE.

a Copy and complete this table to list the sample space.


	R	I	D	E
L				
I				
N				
E				

b How many possible outcomes are there?
c Find Pr(N, R); i.e. the probability that N is chosen from LINE and R is chosen from RIDE.
d Find Pr(L, D).
e Find the probability that two vowels are chosen.
f Find the probability that two consonants are chosen.
g Find the probability that the two letters chosen are the same.



Hint for Q6e: Recall that the vowels are A, E, I, O and U.

7 The spinners shown below are each spun.



Spinner 1 Spinner 2

a Copy and complete this table to list the sample space. Use R for red, P for purple and so on.

		Spinner 2		
		R	P	B
Spinner 1	R			
	P			
	G			
	B			



Hint for Q7: List the colour from spinner 1 first in each cell.

b Find the probability that spinner 1 will display red and spinner 2 will display blue.
c Find the probability that both spinners will display red.
d What is the probability that one of the spinners displays red and the other displays blue?
e What is the probability that both spinners display the same colour?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Problem-solving and reasoning 8–10 8, 10, 11

8 Two dice are rolled for a board game. The numbers showing are then added together to get a number between 2 and 12.



Hint for Q8: Write the sum in each cell. For example, for (1, 1), write 2.

		Die 1					
		1	2	3	4	5	6
Die 2	1						
	2						
	3						
	4						
	5						
	6						

a Copy and complete a table like the one above to show the sample space. State the number of outcomes.

b Find the probability that the two dice add to 5.

c Find the probability that the two dice add to an even number.

d What is the most likely sum to occur?

e What are the two least likely sums to occur between 2 and 12?



9 In Rosemary's left pocket she has two orange marbles and one white marble. In her right pocket she has a yellow marble, a white marble and 3 blue marbles. She chooses a marble at random from each pocket.



Hint for Q9: The left-pocket outcomes are W, O, O.

a Draw a table to describe the sample space.

b Find the probability that she will choose an orange marble and a yellow marble.

c What is the probability that she chooses a white marble and a yellow marble?

d What is the probability that she chooses a white marble and an orange marble?

e Find the probability that a white and a blue marble are selected.

f What is the probability that the two marbles selected are the same colour?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10C

Example 6 Finding the sample space for events without replacement

The letters T R E K are written on cards. A letter is chosen and not replaced so that this letter is now unavailable. Then a second letter is chosen.

a In a table, list the outcomes from choosing two letters without replacement.
b Find the probability that the two letters chosen are (E, R), in that order.
c Find the probability of obtaining an outcome with an E in it.

Solution

Explanation

a

		1st choice			
		T	R	E	K
2nd choice	T	×	(R, T)	(E, T)	(K, T)
	R	(T, R)	×	(E, R)	(K, R)
	E	(T, E)	(R, E)	×	(K, E)
	K	(T, K)	(R, K)	(E, K)	×

List all the outcomes, writing the '1st choice' letter first in each cell. Note that the same letter cannot be chosen twice.

b $\Pr(\text{E, R}) = \frac{1}{12}$

Without replacement there are 12 outcomes.

c $\Pr(\text{includes E}) = \frac{6}{12} = \frac{1}{2}$

6 of the 12 outcomes contain an E.

Now you try

Two digits are selected from the set {3, 5, 6} without replacement.

a Draw a table to list the sample space.
b Find the probability that the outcome (5, 6) will be the result.
c Find the probability that (5, 6) or (6, 5) will be the result.

10 Two letters are chosen from the word DOG *without replacement*.

a Complete the given table.

		1st		
		D	O	G
2nd	D	×	(O, D)	(G, D)
	O		×	
	G			×



Hint for Q10: 'Without replacement' means the first letter chosen is removed, so it is not possible to choose it again.

b Find the probability of obtaining the (G, D) outcome.
c Find the probability of obtaining an outcome with an O in it.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

11 Two digits are selected *without replacement* from the set $\{1, 2, 3, 4\}$.

a Draw a table to show the sample space.

b Find:

i Pr(1, 2)

ii Pr(4, 3)

c Find the probability that:

i both numbers will be at least 3

ii the outcome will contain a 1 or a 4

iii the outcome will contain a 1 and a 4

iv the outcome will not contain a 3



Hint for Q11a: Remember that doubles such as (1, 1), (2, 2) etc. are not allowed.



Hint for Q11c: 'At least 3' includes 3.

Game show — 12


12 A wheel is spun during a game show to determine the prize money, and then a six-sided die is rolled. The prize money shown on the wheel is multiplied by the number on the die to give the total winnings.

a What is the probability that a contestant will win $6000?

b What is the probability that they win more than $11 000?



Hint for Q12: Draw a table to show all outcomes. List the prize in brackets under each outcome.





ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10D Using tree diagrams


Learning intentions

- To know how to use a tree diagram to list the sample space from experiments with two or more steps
- To be able to construct the sample space using tree diagrams for experiments with replacement and those without replacement
- To be able to use a tree diagram to find the probability of certain events

Key vocabulary: tree diagram, with replacement, without replacement, sample space

When two coins are flipped, we can draw a table to list the sample space. Yet if three coins are flipped, then we would need a three-dimensional table to list all outcomes. Imagine trying to find probabilities when five coins are flipped!

Another tool that mathematicians use for probability is the tree diagram. This tree diagram describes the four outcomes when two coins are flipped.



It is important to be able to read a tree diagram correctly. The first row (HH) represents the outcome where the first coin flipped was heads and the second coin flipped was heads. The third row (TH) represents the outcome where the first coin was tails and the second was heads.

Lesson starter: Coin puzzle

If two coins are flipped, rank these outcomes from most likely to least likely:


- Exactly two heads are obtained.
- Exactly one head and exactly one tail are obtained.
- At least one coin shows tails.
- Three tails are shown.

How might the order change if three coins are flipped? Compare your answers with other students.


Key ideas

- **Tree diagrams** are diagrams used to list the sample space for multistage experiments with two or more steps.
  - The outcomes for each stage of the experiment are listed vertically and each stage is connected with branches.
    For example:

Tossing a coin 3 times (with replacement)



Selecting 2 letters from {A, B, C} (without replacement)



In these examples, each set of branches produces outcomes that are all equally likely. Tree diagrams can also be used where all outcomes are not equally likely.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


Exercise 10D

Understanding


1–3 | 1, 3

1 These two tree diagrams show the selection of two letters from the word CAT.

A



B



a Which tree diagram shows selection 'with replacement'?

b Which tree diagram shows selection 'without replacement'?


2 The letters S, K and Y are written on cards and placed in a hat. One letter is randomly selected and then returned to the container. A second letter is then selected.

a Write the missing word or number from each statement about this event.

i 'With _____' means the first letter is returned to the hat before the second letter is selected.

ii When the first letter chosen is replaced, there are still _____ letters to choose from for the second letter.

b Copy and complete this tree diagram for selecting two letters *with replacement*.




3 The letters T, W and O are written on cards and placed in a container. One letter is randomly selected and not returned to the container. A second letter is then selected.

a Write the missing word or number from each statement about this event.

i _____ _____ means the first letter chosen is placed aside and the second letter is selected from the remaining letters.

ii Since the first letter chosen is not replaced, there will be only _____ letters to choose from for the second letter.

b Complete the tree diagram for selecting two letters *without replacement* from the letters T, W and O.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10D

Fluency 4–6 5–7


Example 7 Constructing a tree diagram

An experiment involves tossing two coins.

a Complete a tree diagram to show all possible outcomes.
b What is the total number of outcomes?
c Find the probability of tossing:
i two tails ii one tail iii at least one head

Solution

a



b The total number of outcomes is 4.

c i $\Pr(\text{TT}) = \frac{1}{4}$

ii $\Pr(1 \text{ tail}) = \frac{2}{4} = \frac{1}{2}$

iii $\Pr(\text{at least 1 head}) = \frac{3}{4}$

Explanation

Tree diagram shows two coin tosses one after the other, resulting in $2 \times 2 = 4$ outcomes.

There are four possibilities in the outcomes column.

One out of the four outcomes is TT.

Two outcomes have one tail: {HT, TH}

Three outcomes have at least one head: {HH, HT, TH}


Now you try

An experiment involves tossing two 4-sided dice.

a Complete a tree diagram to show all possible outcomes.
b What is the total number of outcomes?
c Find the probability of tossing:
i two 3's ii one 4 iii at most one 2

4 A coin is tossed twice.

a Complete this tree diagram to show all the possible outcomes.



b What is the total number of outcomes?
c Find the probability of obtaining:
i two heads
ii exactly one head
iii at least one head
iv at least one tail




Hint for Q4: 'At least one' means one or two in this case.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5 A spinner with numbers 1, 2 and 3 is spun twice.

- **a** Draw a tree diagram and list the outcomes.
- **b** Find Pr(1 then 1).
- **c** Find Pr(1 then 2).
- **d** Find Pr(1 and 2 spun in either order).
- **e** Find Pr(both show the same number).
- **f** Find Pr(numbers add to 4).





Hint for Q5: Tree diagram headings: 'First spin', 'Second spin', 'Outcomes'.


Example 8 Constructing a tree diagram without replacement

Two people are selected without replacement from a group of three: Annabel (A), Brodie (B) and Chris (C).

- **a** List all the possible combinations for the selection using a tree diagram.
- **b** Find the probability that the selection will contain:
  - **i** Annabel and Brodie
  - **ii** Chris
  - **iii** Chris or Brodie

Solution

a



Explanation

On the first choice there are three options (A, B or C) but on the second choice there are only two remaining.

b i $\Pr(\text{Annabel and Brodie}) = \frac{2}{6}$

$= \frac{1}{3}$

2 of the 6 outcomes contain Annabel and Brodie (AB) and (BA).

ii $\Pr(\text{Chris}) = \frac{4}{6}$

$= \frac{2}{3}$

4 out of the 6 outcomes contain Chris.

iii $\Pr(\text{Chris or Brodie}) = \frac{6}{6}$

$= 1$

All of the outcomes contain at least one of Chris or Brodie.

Now you try

A drawer contains two red and one blue sock. Two socks are selected at random without replacement.

- **a** List all the possible combinations for the selection using a tree diagram.
- **b** Find the probability that the selection will contain:
  - **i** two red socks
  - **ii** two socks of different colour
  - **iii** at least one red sock.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10D

6 Two people are selected without replacement from a group of three: Donna (D), Elle (E) and Fernando (F).

a List all the possible combinations for the selection using a tree diagram.

b Find the probability that the selection will contain:

i Donna and Elle

ii Fernando

iii Fernando or Elle



Hint for Q6: The combinations are the outcomes.

7 A drawer contains 2 red socks (R), 1 blue sock (B) and 1 yellow sock (Y) and two socks are selected at random without replacement.

a Complete this tree diagram.


b Find the probability of obtaining:

i a red sock and a blue sock

ii two red socks

iii any pair of socks of the same colour

iv any pair of socks of different colour



Problem-solving and reasoning 8 8, 9

8 A coin is tossed three times.

a Draw a tree diagram and list the outcomes.

b Find Pr(3 tails).

c Find Pr(2 tails then 1 head).

d Find Pr(2 tails and 1 head, in any order).

e Which is more likely: getting exactly 3 tails or getting exactly 2 tails?



Hint for Q8: Use headings: Coin 1', 'Coin 2', 'Coin 3', 'Outcomes'.

9 There are three bottles of white wine and two bottles of red wine on a shelf in a cellar. It is too dark to read the labels so two bottles are randomly selected, one at a time. Find the probability that:

a two bottles of different colour wine are selected

b two bottles of the same colour wine are selected

c one or more bottles of red wine are selected



Hint for Q9: Draw a tree diagram using the 'without replacement' method.

Selecting matching clothes — 10

10 A man randomly selects a tie from his collection of one green and two red ties. He selects a shirt from a collection of one red and two white. He then selects either a red or black hat. Use a tree diagram to help find the probability that the man selects a tie, shirt and hat according to the following descriptions:

a a red tie, red shirt and black hat

b exactly one item red

c at least two items red

d a green tie and a black hat

e not a red item

f all three items red

g exactly two items red

h a green hat

i a green tie or a black hat

j a red tie or white shirt or black hat



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10E Using relative frequencies to estimate probabilities

Learning intentions

- To understand how experimental probability (relative frequency) is calculated
- To be able to calculate experimental probability
- To be able to calculate the expected number of occurrences given the experimental probability

Key vocabulary: expected number of occurrences, experimental probability, experiment, favourable outcome, survey, long-run proportion

The sample space is used to calculate theoretical probability. Experimental probability, however, is calculated from the results of a survey, an experiment or a simulation. A larger number of trials make experimental probability calculations more accurate. Experimental probability is used to predict an expected number of results.



Fishermen use experimental data to decide on the bait, place and time of day that gives the highest probability of catching a fish.

Lesson starter: Newspaper theories

A tabloid newspaper reports that of 10 people interviewed in the street, 5 had a dose of the flu. At a similar time a medical student tested 100 people and found that 21 had the flu.

- What is the experimental probability of having the flu, according to the newspaper's survey?
- What is the experimental probability of having the flu, according to the medical student's results?
- Which of the two sets of results would be most reliable and why? Discuss the reasons.
- Using the results from the medical student, how many people would you expect to have the flu in a group of 1000, and why?

Key ideas

- **Experimental probability** or **relative frequency** is calculated using the results of an experiment or **survey**.
  - Experimental probability is a relative frequency calculated from a number of repeated trials.

$$\text{Experimental probability} = \frac{\text{number of times the outcome occurs}}{\text{total number of trials in the experiment}}$$

- The **long-run proportion** is the experimental probability for a sufficiently large number of trials.
- The **expected number of occurrences** is the expected number of favourable outcomes from an experiment.
  - Expected number of occurrences = probability $\times$ number of trials

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Exercise 10E

Understanding

1, 2 | 2

1 Insert the word *experimental*, *theoretical* or *expected* to complete each statement.

a _____ probability is calculated from the results of an experiment or survey.

b _____ probability is calculated from the sample space of an event.

c When a coin is tossed, there are 2 possible equally likely outcomes. So the _____ probability of obtaining a head is $\frac{1}{2}$.

d Jayce tossed a coin 20 times and obtained a head 12 times. So the _____ probability of obtaining a head was $\frac{12}{20} = \frac{3}{5}$.

e A survey found that 3 out of every 5 students have a sister. Out of 1000 students, the _____ number of students with a sister would be $\frac{3}{5} \times 1000 = 600$.

2 Write the experimental probability from each set of results listed here.

a Out of 20 people surveyed, 18 preferred cereal for breakfast.

b 50 students were surveyed and it was found that 40 had pets.

c There were 25 boys and 20 girls at the school bus stop one afternoon. If one child is selected from those at the bus stop, state the experimental probability that the child is a boy.



Hint for Q2: In part **c**, first find the total number of students at the bus stop.

Fluency

3–7 | 4–8

Example 9 Using experimental probability

Kris plays rugby and has a record of kicking a goal from a penalty kick 4 times out of every 7 attempts.

a State the experimental probability for Kris to achieve a penalty goal.

b Calculate the expected number of goals from 28 penalty kicks that Kris takes.

Solution	Explanation
a $\Pr(\text{goal}) = \frac{4}{7}$	Experimentally, we know that Kris can kick 4 goals out of every 7 kicks.
b Expected number $= \frac{4}{7} \times 28 = 16$ goals	Expected number = probability × number of trials

Now you try

In the last 10 school days Mia caught the correct bus in the morning 8 times.

a State the experimental probability of Mia catching the correct bus in the morning on the next school day.

b Calculate the expected number of times Mia will catch the correct morning bus in the next 50 school days.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3 Ashleigh has found that she can shoot a basketball through the hoop 4 times out of 10 from the '3 point' area.

a State the experimental probability for Ashleigh to shoot a basketball through the hoop from the '3 point' area.

b Calculate the expected number of times the ball would go through the hoop from 100 shots Ashleigh makes from the '3 point' area.

4 The experimental probability of Jess hitting a bullseye on a dartboard is 0.05 (or $\frac{5}{100}$). How many bullseyes would you expect Jess to get if he threw the following number of darts?

a 100 darts **b** 200 darts

c 1000 darts **d** 80 darts



Hint for Q4: Expected number = probability × number of trials



5 A bus company surveyed a random selection of 90 people in one suburb. They found that 35 of these people regularly used a bus service from that suburb to the centre of the city.

a State the experimental probability for a person in that suburb to regularly use a bus service from that suburb to the city.

b If there are 2700 residents in that suburb, find the expected number who would regularly use a bus service from that suburb to the city.

Example 10 Finding the experimental probability

A box contains an unknown number of coloured balls and a ball is drawn from the box and then replaced. The procedure is repeated 100 times and the colour of the ball drawn is recorded each time. Twenty-five red balls were recorded.

a Find the experimental probability for selecting a red ball.

b Find the expected number of red balls if the box contained 500 balls in total.

Solution

a $\Pr(\text{red balls}) = \frac{25}{100}$

$= 0.25$

b Expected number of red balls in 500

$= 0.25 \times 500$

$= 125$

Explanation

$\Pr(\text{red balls}) = \frac{\text{number of red balls drawn}}{\text{total number of balls drawn}}$

There are 25 red balls and 100 balls in total.

Expected number of occurrences = probability × number of trials

Now you try

120 people were surveyed and asked if they had contracted the flu in the last 12 months. 25 responded Yes, 90 responded No and the remainder responded Not sure.

a Find the experimental probability that a person contracted the flu in the last 12 months.

b Find the expected number of people who contracted the flu in the last 12 months if 960 people are surveyed.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10E

6 A bag contains an unknown number of counters.
A counter is selected from the bag and then replaced.
The procedure is repeated 100 times and the colour of the counter is recorded each time. Sixty of the counters drawn were blue.



Hint for Q6: Experimental probability $= \frac{\text{number of blue counters}}{\text{total number of counters selected}}$

a Find the experimental probability for selecting a blue counter.

b Find the expected number of blue counters if the bag contained:
i 100 counters **ii** 200 counters **iii** 600 counters

7 In an experiment involving 200 people chosen at random, 175 people said that they owned a home computer.



Hint for Q7: Expected number = probability × number of trials

a Calculate the experimental probability of choosing a person who owns a home computer.

b Find the expected number of people who would own a home computer in the following group sizes:
i 400 people **ii** 5000 people **iii** 40 people

8 By calculating the experimental probability, estimate the chance that each of the following events will occur.

a Nat will walk to work today, given that she walked to work five times in the last 75 working days.

b Mike will win the next game of cards if, in the last 80 games, he has won 32.

c Brett will hit the bullseye on the dartboard with his next attempt if, in the last 120 attempts, he was successful 22 times.

Problem-solving and reasoning 9–11 10–13

9 This table shows the results of three different surveys of people in Perth about their use of public transport (PT).

Survey	Number who use PT	Survey size	Experimental probability
A	2	10	$\frac{2}{10} = 0.2$
B	5	20	_____
C	30	100	_____



Hint for Q9: A large survey size makes experimental probability calculations more accurate.

a What are the two missing numbers in the experimental probability list?

b Which survey should be used to estimate the probability that a person uses public transport and why?

10 The results of tossing a drawing pin and observing how many times the pin lands with the spike pointing up are shown in the table. Results are recorded at different stages of the experiment.

Number of throws	Frequency (spike up)	Experimental probability
1	1	1.00
5	2	0.40
10	5	0.50
20	9	0.45
50	18	0.36
100	41	0.41



Experimental probability shows that the chance of a drawing pin landing spike-up or spike-down is not 50 : 50.

Which experimental probability would you choose to best represent the probability that the pin will land spike up? Why?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

11 A six-sided die is rolled 120 times. State how many times you would expect the following events to occur?

a a 6 **b** a 1 or a 2

c a number less than 4 **d** a number which is at least 5



Hint for Q11: Use the theoretical probability to calculate the expected numbers.

12 The colour of cars along a highway was noted over a short period of time and summarised in this frequency table.

Colour	White	Silver	Blue	Green
Frequency	7	4	5	4



a How many cars had their colour recorded?

b Find the experimental probability that a car's colour is:

i blue **ii** white

c If the colour of 100 cars was recorded, find the expected number of:

i blue cars

ii green cars

iii blue or green cars

13 A spinner is divided into three regions not necessarily of equal size. The regions are numbered 1, 2 and 3 and the spinner is spun 50 times. The table shows the results.

Number	1	2	3
Frequency	26	11	13

a Find the experimental probability of obtaining:

i a 1 **ii** at least a 2 **iii** a 1 or a 3

b Based on these results, how many 3s would you expect if the spinner is spun 300 times?

c In fact, the spinner is divided up using simple and common fractions. Draw and label how you think the spinner regions are divided up.



Hint for Q13: The angles of the spinner regions are multiples of 90°.


Different-shaped dart boards — 14

14 One hundred darts are randomly thrown at the given dartboards. No darts miss the dartboard entirely. How many darts do you expect to hit the blue shaded region? Give reasons.




Hint for Q14: First find the proportion of the entire dartboard taken up by each blue region.


a




b



c




d



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Progress quiz

10A **1** This spinner has six equally divided sections.



a List the sample space using the given numbers.

b Find:

i Pr(4) **ii** Pr(not a 4)

iii Pr(4 or a 5) **iv** Pr(at most 5)

10A **2** A letter is randomly chosen from the word PROGRESS. Find the following probabilities.

a Pr(G) **b** Pr(S) **c** Pr(not S)

d Pr(vowel) **e** Pr(S or R)

10B **3** **a** Complete the following two-way table.

	A	Not A	Total
B	4		
Not B		7	11
Total		12	

b Use your two-way table to find the following probabilities.

i Pr(A) **ii** Pr(A and B) **iii** Pr(B only) **iv** Pr(A or B)

10B **4** A survey of 20 people found that 8 people attended fitness classes and 12 people ran for fitness. 5 of these people did both.

a Construct a Venn diagram for the survey results.

b How many people:

i do not run?

ii run or attend a fitness class?

iii only attend a fitness class?

c If one of the 20 people was randomly selected, find:

i Pr(run and attend a class)

ii Pr(neither run nor attend a class)

iii Pr(only run)

d Turn your Venn diagram into a two-way table.

10C **5** A coin is tossed and then a 4-sided die is rolled.

a Copy and complete the table to list the sample space of this experiment.

	1	2	3	4
H				
T				

b How many possible outcomes are there?

c Find the probability of:

i the combination H4

ii tails on the coin and a factor of 6 on the die

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10C **6** The letters of the word BEE are written on cards. One card (letter) is chosen and not replaced and then a second card is chosen.

a Complete the table below to list the possible outcomes.

		1st choice		
		B	E	E
2nd choice	B	×		
	E		×	
	E			×

b Find the probability that the word BE is formed from the cards chosen (in order)

c Find the probability of choosing a B.

10D **7** Two snake lollies are selected without replacement from a bag containing three snakes: 1 red (R), 1 blue (B) and 1 orange (O).

a List all the possible combinations for the selection using a tree diagram.

b Find the probability that the selection will contain:

i red and orange **ii** blue

10E **8** In a survey involving 80 people chosen at random, 48 said they catch public transport to work.

a Calculate the experimental probability of choosing a person who catches public transport to work.

b Find the expected number of people who would catch public transport to work in a group of 400 people.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10F Data and sampling

Learning intentions

- To know the key words associated with data and sampling and what they mean
- To be able to classify the different types of data
- To know the common types of sampling methods
- To be able to recognise when bias may exist when collecting data

Key vocabulary: population, sample, bias, census, categorical data, numerical data, random sampling, systematic sampling, stratified sampling

Statistics is an essential part of how science collects, analyses and interprets data. We use data to support ideas and uncover new theories. We also use data using tables and graphs to help display information in an organised and understandable way. Data can also be used to influence decision making and help sell products, which is why it is very important to think about how data is collected, how it is analysed and the conclusions that might be communicated.



We use technology to help collect and analyse large data samples.

Lesson starter: Is there bias?

A data collection company has designed a survey to collect information about people's preferences regarding how they communicate with friends and family. The company is interested in feedback from the entire population of a town. Three employees, Jemma, Karri and Moshi, are asked to survey a range of individuals and report back with their data samples.

- Jemma's method involves messaging all her friends on her contact list.
- Karri surveys people in-person in a busy street at around 3:30 p.m. once school is finished for the day.
- Moshi uses a list of mobile phone numbers which belong to people who have agreed to complete surveys for a small fee.
- Discuss each of the methods employed by Jemma, Karri and Moshi. Do you think any of the survey methods would lead to bias? Give reasons.
- Discuss alternative methods to collect the data that might be less biased.


Key ideas

- Key words associated with statistics
  - **Population** is a complete group with at least one characteristic in common. This group could be people, animals, objects or any event.
  - **Sample** is a selection or subset of the population.
  - **Survey** is an investigation or tool to collect information about a given sample or population.
    - A survey of the entire population is called a **census**.
  - **Bias** is a tendency which causes a difference in the results of a survey and the true facts.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Types of data

- **Categorical** (or qualitative) data can be sorted into groups or categories. For example: people's favourite types of animals.
- **Numerical** (or quantitative) data are expressed as numbers. For example: the heights of twenty 15-year-old students.



Types of sampling methods

- **Simple random sampling:** Each individual in the population has an equal chance of being chosen. For example, each individual is assigned a number and a random number generator selects a number.
- **Systematic sampling:** Individuals are selected at regular intervals. For example, every 20th car is stopped on a highway and the driver is surveyed regarding alcohol consumption.
- **Stratified sampling:** The population is divided into subgroups and individuals are surveyed within that group. For example, a population is surveyed regarding their heart health but are first divided into age groups.

Errors may occur in sampling. Examples include:

- Sample size is too small.
- The sample does not represent the entire population.
- Poor or inaccurate measurements are recorded.
- An inappropriate sampling method is chosen.

Exercise 10F

Understanding

1–3 | 1–3

1 Give the key statistical word which matches each definition,

a An investigation or tool to collect information about a given sample or population.

b A tendency which causes a difference in the results of a survey and the true facts.

c A complete group with at least one characteristic in common.

d A selection or subset of the population.

2 Give the name of the type of data matching the description.

a Data that can be sorted into groups or categories.

b Data which can be expressed as numbers.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10F

3 Give the name of the type of sampling method matching the description.

a The population is divided into subgroups and individuals are surveyed within that group.

b Each individual in the population has an equal chance of being chosen.

c Individuals are selected at regular intervals.



Hint for Q3: Choose from simple random, systematic or stratified.

Fluency

4, 5, 7 | 5–8

Example 11 Classifying data

Classify the data obtained from the following survey as either numerical discrete, numerical continuous, categorical ordinal or categorical nominal.

a The mass of each shell in a collection of sea shells
b The results of a survey asking people to rate a TV program as good, satisfactory or poor
c The colours of the next 10 cars that pass on a busy road
d The number of computers in a household

Solution	Explanation
a Numerical continuous	The data includes numbers which could take on any number within a given range. The recordings may depend on the accuracy of the measuring instrument.
b Categorical ordinal	The data includes words which are categories and have a clear order.
c Categorical nominal	The data includes colours which are categories with no particular order.
d Numerical discrete	The data includes numbers from a limited range of values, in this case, the natural numbers $1, 2, 3, \ldots$

Now you try

Classify the data obtained from the following survey as either numerical discrete, numerical continuous, categorical ordinal or categorical nominal.

a The number of students in all the academic classes in a school
b The type of caterpillar found in a number of vegetable garden beds
c The time taken to complete a task
d The type of safety vest used with one of three sizes available including large, medium or small

4 Classify the following as numerical or categorical data.

a The number of pets in a household
b The colour of hair on the heads of 13 models
c The type of mark on a test as either answer, method or consequential
d The number of litres of drink in 10 bottles



Hint for Q4: Categorical data can be sorted into categories and numerical data are expressed as numbers.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5 Classify the data obtained from the following survey as either numerical discrete, numerical continuous, categorical ordinal or categorical nominal.

- **a** The style of jacket on a wall in a shop
- **b** The level of firmness of a bed described as hard, medium or soft
- **c** The weight of 25 packets of chips delivered to a shop
- **d** The shirt size of a range of clothing using the integers 36, 37, 38, ...



Hint for Q5: Discrete data has a limited number of values. Continuous data can take on any value in a given range. Nominal data has no order. Ordinal data has order.

6 Classify the data obtained from the following survey as either numerical discrete, numerical continuous, categorical ordinal or categorical nominal.

- **a** The type of food purchased at a supermarket
- **b** The time taken to complete a 100 m sprint
- **c** The rating of a guide's use of the English language with the options: excellent, good, fair and poor
- **d** The scores out of 20 using whole numbers only for an assignment



Example 12 Choosing a sampling method

a Identify the sampling method used in each of these surveys.

- **i** A car factory assembly line chooses every fifth car to check the quality of paint.
- **ii** One hundred email addresses are randomly selected from a list of all email addresses in a workplace.
- **iii** Fifty dogs are first grouped into breed type then tested for fleas.

b Eighty people on the street at 7 p.m. are surveyed about how often they go out for dinner. This data is then used to make conclusions about the population of that suburb and how often they go out to dinner. Do you think the sample will contain bias? If so, explain why.

Solution	Explanation
a i Systematic	Individual cars are selected at regular intervals.
ii Simple random	Each individual in the workplace has an equal chance of being chosen.
iii Stratified	The dogs are divided into subgroups (breeds) and then dogs are tested for fleas within that breed.
b Yes, it is perhaps more likely that the people surveyed tend to go out for dinner more often.	People who are already out in the street at 7 p.m. are perhaps more likely to go out to dinner on any night of the week.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10F

Now you try

a Identify the sampling method used in each of these surveys.
- i Athletes are divided into sport type then tested for heart fitness.
- ii Every fifth competition paper ordered by surnames is selected for analysis regarding the working shown.
- iii Every family in an apartment block is assigned a number and then five random numbers are chosen so that families can be surveyed regarding their thoughts on the standard of the cleaning in the block.

b Wasram is to seek opinions about what people in the community think of the most recent Soccer World Cup. He surveys his team-mates at the soccer club using a questionnaire. Do you think the sample will contain bias? If so, explain why.

7 a Identify the sampling method used in each of these surveys.
- i A random selection of passport numbers are selected from a government list then checked for people who travelled to the United States.
- ii People at a hospital are sorted into age groups then tested for their reaction to a particular type of medicine.
- iii Every tenth person in the queue at a football match is surveyed regarding their thoughts about who will win the match.



Hint for Q7: Choose from simple random sampling, systematic sampling or stratified sampling.

b A lecturer wants to learn about the opinions of his students and asks them in a lecture to put their hand up and respond verbally to a couple of questions. Do you think the sample will contain bias? If so, explain why.

8 a Identify the sampling method used in each of these surveys.
- i Every eighth Christmas tree sold at a shop is tested for bugs.
- ii A school surveys students about the new uniform and selects a random sample of 20 students from each year level.
- iii A random selection of people chosen from the electoral role are surveyed regarding their opinion on how the election was conducted.



b Matty records information about the bird population on her farm and sits by the lake to wait for the birds to come along before recording the bird type and colour. Do you think the sample will contain bias? If so, explain why.

Problem-solving and reasoning

9–11 | 10–12

9 In a survey involving stratified sampling, 10 people are to be selected from a company including 30 people working in the sales department and 70 people working in the administration department. How many people should be selected from each department?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10 The following statistics were collected about Wally's pet cat.

- Length: 81.45 cm
- Weight: 14.72 kg
- Favourite food: Dine
- Number of toys: 8
- Type of scratching post: short, standard and tall

Describe the type of data for each of the five characteristics.



11 What advice would you give to the following people who are trying to generate unbiased data?

a Hudson wants to form a view as to the difficulty of a test and talks to his five closest friends.

b Molly, a vet, wants to review the time that it takes to serve customers and looks at the data for her most experienced vet in the practice.

c Prabhdeep is researching the cost of bread in a town and records the cost of all the types of bread in a supermarket.

d Zach wants to assess the amount of study students put into their mathematics. He surveys students exiting the Extension maths exam for the Accelerated students.

12 Explain why stratified sampling might be used to explore people's resting heart rates.

The Challenger disaster — 13

13 In 1985 the Challenger space shuttle disintegrated soon after launch and killed all seven crew members. The cause of the disaster was an O-ring which failed and allowed pressurised gasses to escape and cause a chain reaction.

Statistics had been collected and analysed before the launch which collected data regarding the variables *temperature* and *the number of times the O-ring had failed*. The conclusion was that there was no correlation between the two variables. Unfortunately, they did not look at the data collected from when the O-rings had not failed and after including this data there proved to be a more significant correlation between the variables and if discovered earlier, may have stopped the launch.

Use the internet to research a different statistical error which led to a disaster or embarrassing mistake made by a company or researcher. Comment on the following:

a The statistical error made

b The consequences of the error

c Suggestions regarding what could have been done to avoid the error.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10G Summarising data: range and measures of centre

Learning intentions

- To know that measures of centre are used to summarise a data set
- To be able to find the mean, median, mode and range of a set of data
- To know how the median is found for data sets with an odd or even number of values
- To understand that an outlier is a value much smaller or larger than the rest of the data

Key vocabulary: mode, mean, range, median, outlier, bimodal

Statistics involves collecting and summarising data. It also involves drawing conclusions and making predictions. The type and amount of product stocked on supermarket shelves, for example, is determined by the sales statistics and other measures such as average cost and price range.

Lesson starter: Game purchase

Arathi purchases 7 computer games at a sale. 3 games cost $20 each, 2 games cost $30, 1 game costs $50 and the last game cost $200.

- Recall and discuss the meaning of the words mean, median and mode.
- Can you work out the mean, median or mode for the cost of Arathi's games?
- Which of the mean, median or mode gives the best 'average' for the cost of Arathi's games?
- Why is the mean greater than the median in this case?

Key ideas

Mean $\bar{x}$

The mean, $\bar{x}$, of a set of numbers is given by

$$\bar{x} = \frac{\text{sum of all the values}}{\text{number of scores}}$$

For example, mean $= \frac{6+7+10+12+13}{5}$

$= 9.6$

Median

The median is the middle value if the data is placed in order.

- If there are two middle values, the median is calculated as the mean of these two values.

odd number of values

1 3 5 (5) 6 7 10

median

even number of values

13 17 17 20 21 27 27 28

20.5

median

- The **mode** is the most common value.
  - There can be more than one mode.
  - If there are two modes, we say that the data set is **bimodal**.
- An **outlier** is a score that is much larger or smaller than the rest of the data.
- The **range** is the difference between the highest and lowest values.

  Range = maximum value − minimum value

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Exercise 10G

Understanding

1–5 | 1–5

1 **a** To calculate the _____, you add up all the values and divide by the number of values.

b Find the mean of these data sets (round each answer to one decimal place).

i 3 6 8 10 12

ii 6 8 4 12 9 11 16 13



Hint for Q1: First find the sum of the scores, then divide by the number of scores. Mean = $\frac{39}{5}$

2 **a** The mode is the most _____ value.

b State the mode of these data sets.

i 1 2 2 2 4 5 6

ii 1 4 8 8 9 10 10 10 12

3 **a** The median is the _____ value when the scores are listed in order.

b Find the median of these data sets.

i 1 3 5 6 8

ii 4 5 7 9 10 11

4 **a** The _____ is the difference between the highest and lowest values.

b Find the range of these data sets.

i 3 6 7 12 15 24

ii 5 12 14 18 23 27



Hint for Q4: Range = highest value − lowest value

5 **a** An _____ is a score that is much larger or smaller than the rest of the data.

b For each of these data sets, if there is an outlier, state its value.

i 2 4 5 4 6 2 3 36

ii 21 25 3 27 28 24 29 30

Fluency

6(½), 7, 8 | 6(½), 7–9

Example 13 Finding measures of centre

For the given data sets, find the following:

i the mean **ii** the median **iii** the mode **iv** the range

a 5 2 4 10 6 1 2 9 6

b 17 13 26 15 9 10

Solution	Explanation
a i Mean $= \frac{45}{9}$	$5+2+4+10+6+1+2+9+6=45$
$= 5$	Find the sum of all the numbers and divide by the number of values.
ii 1 2 2 4 (5) 6 6 9 10	First, order the data.
Median = 5	The median is the middle value.
iii Mode = 2 and 6	The data set is bimodal since there are two numbers with the highest frequency.
iv Range = 10 − 1	The range is the highest score minus the lowest score.
$= 9$	

Continued on next page

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10G

b i Mean $= \frac{90}{6}$
$= 15$

$17 + 13 + 26 + 15 + 9 + 10 = 90$
There are 6 values.

ii 9 10 13 15 17 26
14

Median $= \frac{13 + 15}{2}$
$= 14$

First, order the data.

Since there are two values in the middle, find the mean of them.

iii No mode

None of the values are repeated so there is no mode.

iv Range $= 26 - 9$
$= 17$

The highest score is 26 and the lowest score is 9.

Now you try

For the given data sets, find the following:

i the mean **ii** the median
iii the mode **iv** the range

a 8 4 2 7 7 3 4 7 3

b −4 2 0 1 −1 0 3 5

6 For the given data sets, find:

i the mean **ii** the median
iii the mode **iv** the range

a 7 2 3 8 5 9 8
b 6 13 5 4 16 10 3 5 10
c 12 9 2 5 8 7 2 3
d 10 17 5 16 4 14
e 3.5 2.1 4.0 8.3 2.1
f 0.7 3 2.9 10.4 6 7.2 1.3 8.5
g 6 0 −3 8 2 −3 9 5
h 3 −7 2 3 −2 −3 4



Hint for Q6: Recall that to subtract a negative, you add a positive; e.g. $7 - (-2) = 7 + 2 = 9$.

7 These data sets include an outlier. Calculate the mean and the median and write down the outlier. Include the outlier in your calculations.

a 5 7 7 8 12 33
b 2 40 42 45 47
c 1.3 1.1 1.0 1.7 1.5 1.6 −1.1 1.5
d −58 −60 −59 −4 −64



Hint for Q7: An outlier is much larger or smaller than the other scores.

8 Decide whether the following data sets are bimodal.

a 2 7 9 5 6 2 8 7 4
b 1 6 2 3 3 1 5 4 1 9
c 10 15 12 11 18 13 9 16 17
d 23 25 26 23 19 24 28 26 27



Hint for Q8: Bimodal means that there are two modes.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

9 This ordered data set shows the number of fish Daniel caught in the 12 weekends that he went fishing during the year.

1 2 3 4 4 5 6 7 7 9 11 13

a Find the range.
b Find the median.
c Find the mean.
d Find the mode.



Problem-solving and reasoning

10–12 | 11–13

10 In three running races, Paula recorded the times 25.1 seconds, 24.8 seconds and 24.1 seconds.

a What is the mean time of the races? Round to two decimal places.
b Find the median time.

11 This is a data set of six house prices in Darwin.

$324 000 $289 000 $431 000 $295 000 $385 000 $1 700 000

a Which price would be considered the outlier?
b If the outlier was removed from the data set, by how much would the median change? (First work out the original median house price.)
c If the outlier was removed from the data set, by how much would the mean change, to the nearest dollar? (First work out the original mean house price.)



Hint for Q11: To find the median, list the prices in order, then find the middle value.

Example 14 Finding a data value for a required mean

The hours a shop assistant spends cleaning the store in eight successive weeks are:
8, 9, 12, 10, 10, 8, 5, 10

a Calculate the mean for this set of data.
b Determine the number of hours that needs to be added to this data to make the mean equal to 10.

Solution

a Mean $= \frac{72}{8}$

$= 9$

Explanation

$8 + 9 + 12 + 10 + 10 + 8 + 5 + 10 = 72$.
Sum of the 8 data values is 72.

b Let a be the new score.

Require $\frac{72 + a}{8 + 1} = 10$

$\frac{72 + a}{9} = 10$

$72 + a = 90$

$a = 18$

The new score would need to be 18.

$72 + a$ is the total of the new data and $8 + 1$ is the new total number of scores. Set this equal to the required mean of 10.

Solve for a.

9 scores have a mean of 10, so the sum of the scores $= 9 \times 10 = 90$.

Write the answer.

Now you try

Amanda received scores of 85, 91, 94 and 78 on her last four maths tests.

a Calculate the mean for this set of data.
b Determine the score that needs to be added to this data to make the mean equal to 89.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10G

12 A netball player scored the following number of goals in her 10 most recent games:
15 14 16 14 15 12 16 17 16 15

a What is her mean score?

b What number of goals does she need to score in the next game for the mean of her scores to be 16?



13 Stevie obtained the following scores on her first five Maths tests: 92 89 94 82 93

a What is her mean test score?

b If there is one more test left to complete, and she wants to achieve an average of at least 85, what is the lowest score Stevie can obtain for her final test?

Aiming for an A — 14

14 A school gives grades in Mathematics each semester according to this table. Raj has scored the following results for four topics this semester, and has one topic to go: 75 68 85 79

Average score	Grade
90–100	A+
80–	A
70–	B+
60–	B
50–	C+
0–	C

a What is Raj's mean score so far?

b What grade will Raj get for the semester if his fifth score is:
i 50? **ii** 68? **iii** 94?

c Find the maximum average score Raj can receive for the semester. Is it possible for him to get an A+?

d Find the least score that Raj needs in his fifth topic for him to receive an average of:
i B+ **ii** A



Using a calculator 10G: Finding measures of centre
The activity can be found in the More Resources section of the Interactive Textbook in the form of a printable PDF.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10H Interpreting data from tables and graphs

Learning intentions

- To know the common types of statistical graphs including those that are skewed or symmetrical
- To be able to interpret common types of statistical graphs
- To be able to recognise when a graph may be misleading or drawn to give a false impression


Key vocabulary: histogram, dot plot, pie chart, symmetrical data, skewed data, mean, median, mode, range

In our everyday lives, it is important to be able to understand many forms of information. Data that is presented in a table or a graph is much easier to interpret than a long list of data. The headings in a table add detail and graphs give a visual comparison between values or categories.


Lesson starter: An unfair comparison

Each pair of graphs below shows the same data. Identify which one of the pair is misleading and discuss the reasons why.


Graph A1




Graph A2




Graph B1




Graph B2



Graph C1



Graph C2




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10H


Key ideas

- Graphs can be presented in different forms. Some common examples are:



  - The titles, scales and column heights or sector angles tell us something about the data that is used to make the graphs.


- Misleading graphs give a false impression about data.
  - When only part of the scale is shown, the difference between results may be exaggerated.
  - If one column has a different size or colour, it may appear to have greater value than the other columns.
  - When pictograph symbols are not the same size, the larger symbol appears to have a higher value.
  - A measure of centre can be misleading if it is not a fair representation of the centre of the data values.
    For example, for the data in the graph below, the mean wage of $850 is actually larger than all other wages except the outlier of $2000. So the median (middle) wage of $620 is a better representation of the 'average' wage in this case.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

- Graphs can help to determine if data is symmetrical or skewed.

Symmetrical **Skewed**



- **Symmetrical data** will appear like the data has a mirror line at its centre.
- **Skewed data** will have values bunched to one side of the middle.

Exercise 10H


Understanding

1–3 | 2, 3


1 Which of the following are not examples of graphs?
dot plot, mean, column graph, pie chart, range and histogram.

2 Which of the following graphs is symmetrical and which is skewed?

A




B




3 Which of the two graphs might a newspaper use if it wanted to give the impression that the Panthers beat the Jaguars by a large margin? Give a reason.

A



B




Fluency

4–6, 8, 9 | 4, 6, 7, 9, 10

Example 15 Interpreting histograms or column graphs

This column graph shows the results of a spelling test out of 10.

a How many results are shown in this histogram?
b List the results in ascending order.
c Calculate the mean.
d Calculate the median.
e What is the range of results from this test?
f Is this data skewed or symmetrical?
g What proportion of results is greater than or equal to 7?



Continued on next page

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10H

Solution	Explanation
a $1+1+2+3+4+4=15$	The frequency shows how many times each score occurred. Add the frequency values to find the total number of scores.
b 4, 5, 7, 7, 8, 8, 8, 9, 9, 9, 9, 10, 10, 10, 10	1 lot of 4, 1 lot of 5, 2 lots of 7, 3 lots of 8, 4 lots of 9, 4 lots of 10.
c Mean $= \frac{123}{15}$ $= 8.2$	$4+5+7+7+8+8+8+9+9+9+9+10+10+10+10=123$
d Median $= 9$	The 8th score is the middle score.
e Range $= 10-4$ $= 6$	The range is the highest score minus the lowest score.
f Skewed	Results are bunched to the higher scores.
g $\frac{13}{15}$	13 scores are greater than or equal to 7 out of a total of 15 scores.

Now you try


This pie chart shows the proportion of households in a particular town that have a type of pet. There are 500 households in the town.

a How many of the households owned:

i dogs? **ii** cats? **iii** other?

b How many more households own cats compared to dogs?

c What is the sector angle for cats?



4 This graph shows the results of a number of maths tests out of 10.

a How many results are shown in this graph?

b List the results in ascending order.


c Calculate the mean.

d Calculate the median.

e What is the range of results from this test?


f Is this data skewed or symmetrical?

g What proportion (fraction) of results is greater than or equal to 8?



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5 This dot plot shows the heights, in cm, of the students in a Year 9 class.






Hint for Q5: Each dot represents a student.

a How many students have their heights recorded on this dot plot?
b What is the range of heights?
c What is the mode of these heights?
d What is the median height for this class?
e What is the mean height for this class?
f What value is the outlier?

6 This pie chart shows the proportions of deaths in Australia from diseases caused by smoking.

Deaths from smoking-related diseases





Hint for Q6:
Number of deaths
$= \frac{\text{percentage}}{100} \times \text{total deaths}$
Sector angle = percentage $\times$ 360°

Around 15 000 Australians in total die from smoking-related illness per year.

a What is the total percentage of smoking deaths caused by cancer?
b How many Australian smokers die of cancer in a year?
c How many Australian smokers die of cancer each day? Round to the nearest whole number.
d How many Australian smokers die from various breathing diseases each day? Round to one decimal place.
e Calculate the sector angle for lung cancer. Recall that 100% represents 360°.

7 The following table shows tide times and heights in December for Yamba, on the NSW north coast. The tide heights shown are red for low tide and blue for high tide. The times are in 24-hour time.

Saturday 17		*Sunday* 18		*Monday* 19		*Tuesday* 20		*Wednesday* 21	
Time	Height	Time	Height	Time	Height	Time	Height	Time	Height
0038	1.14	0145	1.18	0255	1.26	0401	1.37	0.502	1.49
0619	0.45	0729	0.49	0847	0.50	1008	0.46	1123	0.38
1243	1.42	1340	1.34	1445	1.26	1556	1.21	1703	1.18
1925	0.29	2018	0.29	2115	0.28	2211	0.26	2307	0.22

Write all time answers in both 24-hour time and 12-hour time.



Hint for Q7: 1520 in 24-hour time is 3.20 p.m. in 12-hour time.

a How high is the second high tide on Saturday 17 December?
b What time is the first low tide on Monday 19 December?
c How much later in the morning is the low tide on Tuesday 20 December than the low tide on Monday 19 December?
d What is the difference in height between the two high tides on Wednesday 21 December?
e How long is it between the two high tides on Sunday 18 December?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10H

8 The following is a table of life expectancy estimates for people in various countries.

Life expectancy in years			
Country	**Overall average**	**Male**	**Female**
Japan	83	79	86
Switzerland	82	80	84
Australia	81	79	84
Malaysia	74	72	77
Vietnam	74	72	77
Indonesia	71	69	73
Papua New Guinea	57	55	60
South Africa	49	49	50
Mozambique	39	38	39


a How many years more do Australian males expect to live compared to Mozambique males?

b How many years more do Australian females expect to live compared to Papua New Guinea females?

c If the overall world average of life expectancy is 67 years, how far above the average is Japan's overall average life expectancy?

d Find the increase in life expectancy between Indonesian males and Australian males. Now calculate this increase as a percentage of Indonesian male life expectancy. Round to one decimal place.

9 The line graphs below show the mean monthly minimum and maximum temperatures and also the monthly rainfall in mm for Hobart.
The temperature values are read from the scale on the left and the rainfall values are read from the scale on the right. For example, in January the mean maximum temperature is 22°C and the rainfall is 44 mm.



a What was the mean maximum temperature in April?

b What was the rainfall in April?

c What was the mean minimum temperature in July?

d What was the rainfall in July?

e During what months was the mean maximum temperature greater than 19°C?

f During what months was the rainfall less than 40 mm per month?

g Which were the wettest two months in Hobart?

h What was the temperature range in December?




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10 The following graph shows impact speeds (reaction time and distance travelled while braking) for cars when driving at various speeds on a dry road.



Hint for Q10: Reaction time is shown by the purple bar, braking time is shown by the orange bar.



a The red line on the graph represents an object or person in front of the braking car. How far in front of the car is the object or person at the start of the driver's reaction time?

b How many metres are travelled during *reaction time* when driving at 60 km/h?

c How many metres are travelled during *braking time* when driving at 60 km/h?

d How much distance is needed, overall, to stop when driving at 80 km/h?


e By how much does the braking distance increase when driving at 80 km/h compared to 50 km/h?

Problem-solving and reasoning


11, 12 | 11, 13, 14

11 Here are two column graphs, each showing the same results of a survey that asked people which TV channel they preferred.

Graph A



Graph B



a From Graph A, write down how many viewers preferred each channel.

b Which graph could be titled 'Channel 15 is clearly most popular'?

c Which graph could be titled 'All TV channels have similar popularity'?

d What is the difference between the two graphs?

e Which graph is misleading, and why?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10H


12 This column graph shows the number of sales for one month of four popular breakfast cereals. The height of each column represents the sales.

a List the breakfast cereals in order of sales.

b Did Yum-crunch have more sales than the other cereals?


c List 3 changes that have been made to the Yum-crunch column to make it look like it has better sales than any of the other cereals.

d How should the columns be drawn so the graph is not misleading?



13 This pictograph shows the number of TV viewers for the final matches of various sports.

Size of TV audience for finals matches



a Which sport *appears* to have had the greatest TV audience?

b List the sports in order, according to their *length* on the pictograph.

c Using the key, determine the audience sizes for the rugby and soccer finals.

d Which sport actually had the largest TV audience? What was the size of its audience?

e List the sports in order according to the audience size calculated using the key.


f In what way is this graph misleading?

g How should a pictograph be drawn so it is not misleading?



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

14 This column graph shows the price of some houses that were sold during one November in a Sydney suburb.





a How many house sales are shown in this column graph?
b List the price of each house in ascending order.
c Find the mean house price.
d Find the median house price.
e A newspaper headline read 'House prices now average over $900 000'. Is this 'average' referring to the mean or median price?
f What proportion of house prices were less than the mean value?
g Do you think the mean or the median would better represent the 'average' house price? Give a reason for your answer.

Viewing distance for TV — 15

15 a Draw a line graph showing the Minimum and Maximum viewing distances for each TV screen size. Include a key.

TV screen size (length of diagonal in inches)	Maximum viewing distance (cm)	Minimum viewing distance (cm)
26	200	100
30	220	120
34	260	130
42	320	160
47	360	180
50	380	200
55	420	220
60	470	230
65	490	250

b If a lounge chair is 1.5 m in front of the TV, which sizes of TV would be suitable?
c If a lounge chair is 3 m in front of the TV, which sizes of TV would be suitable?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10I Stem-and-leaf plots

Learning intentions

- To be able to construct and interpret a stem-and-leaf plot
- To be able to construct and interpret a back-to-back stem-and-leaf plot
- To be able to interpret the shape of a stem-and-leaf plot to describe the distribution of the data as symmetrical or skewed

Key vocabulary: stem-and-leaf plot, symmetrical data, skewed data, back-to-back stem-and-leaf plot

Stem-and-leaf plots (or stem plots) are commonly used to display a single data set or two related data sets. They help to show how the data is distributed. They retain all the individual data elements so no detail is lost. The median and mode can be easily read from a stem-and-leaf plot because all the data sits in order.

Lesson starter: Ships vs Chops

At a school, Ms Ships' class and Mr Chops' class sit the same exam. The scores are displayed using this back-to-back stem-and-leaf plot. Discuss the following.

- Which class had the most students?
- What were the lowest and highest scores from each class?
- What were the median scores from each class?
- Which class could be described as symmetrical and which as skewed?
- Which class had the better results?

Leaf Ms Ships' class	Stem	Leaf Mr Chops' class
3 1	5	0 1 1 3 5 7
8 8 7 5	6	2 3 5 5 7 9 9
6 4 4 2 1	7	8 9 9
7 4 3	8	0 3
6	9	1

7 | 8 means 78



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Key ideas

- A **stem-and-leaf plot** uses a stem number and leaf number to represent data.
  - The data is shown in two parts: a stem and a leaf.
  - The 'key' tells you how the plot is to be read.

Ordered stem-and-leaf plot

Stem	Leaf
1	2 6
2	2 3 4 7
3	1 2 4 7 8 9
4	2 3 4 5 8
5	7 9

2 | 4 means 24 people

The leaf numbers are written in order.

A Key is added to show the place value of the stems and leaves.

- **Back-to-back stem-and-leaf plots** can be used to compare two sets of data. The stem is drawn in the middle, with the leaves on either side.

Scores for the last 30 football games

Winning scores		Losing scores
	7	4 5 8 8 9
1	8	0 0 3 3 6 7
7 5	9	1 2 3 6
8 4 4 1	10	3 9
9 5 0	11	1
3 1	12	

10 | 9 means 109

81 lowest winning score

Symmetrical data

Skewed data

111 highest losing score

- **Symmetrical data** will produce a graph that is symmetrical about the centre.
- **Skewed data** will produce a graph that includes data bunched to one side of the centre.

Exercise 10I

Understanding

1–3 | 3

1 List the statistical data that would produce these stem-and-leaf plots. Use the key to help.



Hint for Q1: The same stem goes with each leaf along each row.

a

Stem	Leaf
3	5 7
4	1 3 8

3 | 1 means 3.1

b

Stem	Leaf
5	2
6	0 1 7
7	3 5

6 | 3 means 63

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

101

2 This stem-and-leaf plot shows the number of minutes Alexis spoke on her phone for a number of calls.



Stem	Leaf
0	8
1	5 9
2	1 1 3 7
3	4 5

2 | 1 means 21 minutes

a How many calls are represented by the stem-and-leaf plot?

b What is the length of the:
- **i** shortest phone call?
- **ii** longest phone call?

c What is the mode (the most common) call time?

d What is the median call time (middle value)?



Hint for Q2: Count the leaves to find the number of calls.

3 This back-to-back stem-and-leaf plot shows the thickness of tyre tread on a selection of cars from the city and country.

City		Country
8 7 3 1 0 0 0	0	6 8 8 9
8 6 3 1 0	1	0 4 5 5 6 9
1	2	3 4 4

1 | 3 means 13 mm



Hint for Q3: 06 is written just as 6.

a How many car tyres were tested altogether?

b What was the smallest tyre tread thickness in:
- **i** the city?
- **ii** the country?

c What was the largest tyre tread thickness in:
- **i** the city?
- **ii** the country?

d Find the median tyre tread thickness for tyres in:
- **i** the city
- **ii** the country



Hint for Q3: Skewed distributions are not symmetrical.

e Is the distribution of tread thickness for city cars more symmetrical or skewed?

f Is the distribution of tread thickness for country cars more symmetrical or skewed?

Fluency

4(½), 5, 6 | 4(½), 6, 7

Example 16 Constructing and using a stem-and-leaf plot

For this set of data:

0.3 2.5 4.1 3.7 2.0 3.3 4.8 3.3 4.6 0.1 4.1 7.5 1.4 2.4
5.7 2.3 3.4 3.0 2.3 4.1 6.3 1.0 5.8 4.4 0.1 6.8 5.2 1.0

a Organise the data into an ordered stem-and-leaf plot.

b Find the median.

c Find the mode.

d Describe the data as symmetrical or skewed.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Solution

a

Stem	Leaf
0	1 1 3
1	0 0 4
2	0 3 3 4 5
3	0 3 3 4 7
4	1 1 1 4 6 8
5	2 7 8
6	3 8
7	5

3 | 4 means 3.4

Explanation

The minimum is 0.1 and the maximum is 7.5 so stems range from 0 to 7.

Place leaves in order from smallest to largest. Some numbers appear more than once; e.g. two instances of 0.1 means that the leaf 1 appears twice: 0.1, 0.1, …

b $\text{Median} = \frac{3.3 + 3.4}{2}$

$= 3.35$

There are 28 data values. The median is the average of the two middle values (the 14th and 15th values).

c Mode is 4.1.

The most common value is 4.1.

d Data is approximately symmetrical.

The distribution of numbers is approximately symmetrical about the stem containing the median.

Now you try

For this set of data:

24 37 52 16 11 29 13 26

34 42 9 29 21 35 46 50

a Organise the data into an ordered stem-and-leaf plot.
b Find the median.
c Find the mode.
d Describe the data as symmetrical or skewed.

4 For each of the following data sets:

i organise the data into an ordered stem-and-leaf plot
ii find the median
iii find the mode
iv describe the data as symmetrical or skewed

a 41 33 28 24 19 32 54 35

19 23 32 26 28 26 28

b 31 33 23 35 15 23 48 50 35 42 45 15 21

51 31 34 23 42 50 26 30 45 37 39 45

c 34.5 34.9 33.7 34.5 35.8 33.8 34.3 35.2 37.0 34.7

35.2 34.4 35.5 36.5 36.1 33.3 35.4 32.0 36.3 34.8

d 167 159 159 193 161 164 167 157 158 175 177

185 177 202 185 187 159 189 167 159 173 198 200



Hint for Q4: Draft version: Enter the leaves in the same order as in the question. Final version: Rewrite the stem-and-leaf plot with leaves in order. Remember to include the key.



Hint for Q4: In part **d**, use 16 as the stem for 167.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

101

5 The number of vacant rooms in a motel each week over a 20-week period is shown below.

12 8 11 10 21 12 6 11 12 16
14 22 5 15 20 6 17 8 14 9

a Draw a stem-and-leaf plot of this data.
b In how many weeks were there fewer than 12 vacant rooms?
c Find the median number of vacant rooms.



Example 17 Constructing back-to-back stem-and-leaf plots

A shop owner has two jeans shops. The daily sales in each shop over a 16-day period are monitored and recorded as follows.

Shop A
3 12 12 13 14 14 15 15 21 22 24 24 24 26 27 28

Shop B
4 6 6 7 7 8 9 9 10 12 13 14 14 16 17 27

a Draw a back-to-back stem-and-leaf plot.
b Compare and comment on differences between the sales made by the two shops.

Solution

a

Shop A		Shop B
3	0	4 6 6 7 7 8 9 9
5 5 4 4 3 2 2	1	0 2 3 4 4 6 7
8 7 6 4 4 4 2 1	2	7

1 | 3 means 13

Explanation

The data for each shop is already ordered. Stems are in intervals of 10. Record leaf digits for Shop A on the left and Shop B on the right, ordered from the middle to the outside.

b Shop A sales are generally between 12 and 28, with one low value of 3. Shop B sales are generally between 4 and 17, with one high value of 27. Shop A has a lot more high values than Shop B. Shop B has more low values than Shop A.

Look at both sides of the plot for the highest and lowest values and whether there are a few or many of the small and large numbers.

Now you try

At a school, two reading classes in year 6 read books for homework one night. The number of pages read by the students are recorded as follows.

Class A 21 36 19 15 19 31 17 24 29
Class B 3 16 14 20 11 18 12 13

a Draw a back-to-back stem-and-leaf plot.
b Compare and comment on the differences between the amount of reading completed by the different class groups.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

6 For each of the following sets of data:

i Draw a back-to-back stem-and-leaf plot.

ii State the smallest and largest value in each set and compare the numbers of small and large values in each set.



Hint for Q6: Order the leaves with the smallest on the inside and largest on the outside. State whether each set has a few or many of the small numbers and large numbers.

a

Set A:	46	32	40	43	45	47	53
	33	48	39	43	54	40	54
Set B:	48	49	31	40	43	47	48
	44	46	53	44	41	49	51

b

Set A:	0.7	0.8	1.4	8.8	9.1	2.6	3.2	0.3	1.7	1.9	2.5	4.1	4.3	3.3	3.4
	3.6	3.9	3.9	4.7	1.6	0.4	5.3	5.7	2.1	2.3	1.9	5.2	6.1	6.2	8.3
Set B:	0.1	0.9	0.6	1.3	0.9	0.1	0.3	2.5	0.6	3.4	4.8	5.2	8.8	4.7	5.3
	2.6	1.5	1.8	3.9	1.9	0.1	0.2	1.2	3.3	2.1	4.3	5.7	6.1	6.2	8.3

7 **a** Draw a back-to-back stem-and-leaf plot for the final scores of St Kilda and Collingwood in the 24 AFL games given here.

St Kilda:	126	68	78	90	87	118
	88	125	111	117	82	82
	80	66	84	138	109	113
	122	80	94	83	106	68
Collingwood:	104	80	127	88	103	95
	78	118	89	82	103	115
	98	77	119	91	71	70
	63	89	103	97	72	68



Hint for Q7:
Percentage = fraction × 100

b In what percentage of their games did each team score more than 100 points?

c Comment on the symmetry of the distribution of the scores for each team.

d Which team has scores that are more consistent? Which team has more higher scores?

Problem-solving and reasoning 8, 9 8–10

8 This stem-and-leaf plot shows the time taken, in seconds, by Helena to run 100 m in her last 25 races.

Stem	Leaf
14	9
15	4 5 6 6 7 7 7 8 9
16	0 0 1 1 2 2 3 4 4 5 5 5 7 7
17	2

14 | 9 means 14.9 seconds

a Find Helena's median time.

b What is the difference between the slowest and fastest time?

c If in her 26th race her time was 14.8 seconds and this was added to the stem-and-leaf plot, would her median time change? If so, by how much?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10I

9 Two brands of batteries were tested to determine their lifetime in hours. The data below shows the lifetime of 20 batteries of each brand.

Brand A: 7.3 8.2 8.4 8.5 8.7 8.8 8.9 9.0 9.1 9.2
9.3 9.4 9.4 9.5 9.5 9.6 9.7 9.8 9.9 9.9

Brand B: 7.2 7.3 7.4 7.5 7.6 7.8 7.9 7.9 8.0 8.1
8.3 9.0 9.1 9.2 9.3 9.4 9.5 9.6 9.8 9.8



Hint for Q9: The stems will be the 'units' value and each leaf the 'tenths' value.

a Draw a back-to-back stem-and-leaf plot for this data.

b How many batteries from each brand lasted more than 9 hours?

c Which brand shows the best performance?

10 Find the median if all the data in each back-to-back stem-and-leaf plot was combined.

a

5	3	8 9
9 7 7 1	4	0 2 2 3 6 8
8 6 5 2 2	5	3 3 7 9
7 4 0	6	1 4

4 | 2 means 42

b

3	16	0 3 3 6 7 9
9 6 6 1	17	0 1 1 4 8 8
8 7 5 5 4 0	18	2 2 6 7
2	19	0 1

16 | 3 means 16.3



Hint for Q10: First combine each back-to-back plot into just one stem-and-leaf plot.

Birth weights and smoking — 11

11 The back-to-back stem-and-leaf plot below shows the birth weight in kilograms of babies born to mothers who do or don't smoke.



Birth weight of babies

Smoking mothers		Non-smoking mothers
4 3 2 2	2	4
9 9 8 7 6 6 5 5	2*	8 9
4 3 2 1 1 1 0 0 0	3	0 0 1 2 2 3
6 5 5	3*	5 5 5 6 6 7 7 8
1	4	
	4*	5 5 6

2 | 4 means 2.4 kg
2* | 5 means 2.5 kg

a What percentage of babies born to smoking mothers have a birth weight of less than 3 kg?

b What percentage of babies born to non-smoking mothers have a birth weight of less than 3 kg?

c Compare and comment on the differences between the birth weights of babies born to mothers who smoke and those born to mothers who don't smoke.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10J Grouped data

Learning intentions

- To be able to construct a frequency table from a set of data including a percentage frequency column
- To be able to construct and analyse a histogram or percentage frequency histogram

Key vocabulary: frequency table, percentage frequency histogram, histogram, class interval


For some data, especially large sets, it makes sense to group the data and then record the frequency for each group to produce a frequency table. For numerical data, a graph generated from a frequency table gives a histogram. Like a stem-and-leaf plot, a histogram shows how the data is distributed across the full range of values. A histogram, for example, is used to display the level of exposure of the pixels in an image in digital photography. It uses many narrow columns to show how the light values are distributed across the scale from black to white.



A luminance value histogram used in digital photography software.

Lesson starter: Baggage check

This histogram shows the distribution of the weight of a number of bags checked at an airport.



- How many bags had a weight in the range 10–15 kg?
- How many bags were checked in total?
- Is it possible to determine the exact mean, median or mode of the weight of the bags by looking at the histogram? Discuss.
- Describe the distribution of checked bag weights for the given graph.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10J

Key ideas


- A **frequency table** shows the number of values within a set of categories or **class intervals**.
- Grouped numerical data can be illustrated using a **histogram**.
  - The height of a column corresponds to the frequency of values in that class interval.
  - There are usually no gaps between columns.
  - The scales are evenly spread, with each bar spreading across the boundaries of the class interval.
  - A **percentage frequency histogram** shows the frequencies as percentages of the total.
- Like a stem-and-leaf plot, a histogram shows whether the data is skewed or symmetrical.

Frequency table

Class interval	Frequency	Percentage frequency
60–	3	12
65–	4	16
70–	6	24
75–	8	32
80–85	4	16
Total	25	100


70– includes numbers from 70 to less than 75.

Frequency histogram



This gap may be used when the intervals do not start at zero.

Percentage frequency histogram



Exercise 10J

Understanding

1–4 | 4

1 Write the missing word for each of these statements.

a A group of scores such as $10 - 15$ kg is called a _____ _____.

b A _____ table shows the class intervals and the number of values in each class interval.

c A column graph that shows frequencies and scores is called a _____.

d A _____ _____ histogram shows each frequency as a percentage of the total number of scores.


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

2 Write the total number of scores in each of these displays.

a

Class interval	Frequency
0–	3
5–	5
10–	8
15–	4

b






Hint for Q2: The sum of the frequencies equals the number of scores.



Hint for Q3: The frequency shows the number of years.

3 The frequency histogram below shows the number of years for which the number of international visitors to South Australia was within a given range for the decade from 2000 to 2009.






a How many years in the decade were there less than 330 000 international visitors?

b Which range of visitor numbers had the highest frequency?

4 Some Year 9 students selected a sample of textbooks from the library and recorded the number of pages in each book. The frequency histogram below shows their results.



a How many textbooks had between 100 and 200 pages?

b How many textbooks were selected from the library?

c What percentage of textbooks had between:

i 200 and 300 pages? **ii** 200 and 400 pages?



Hint for Q4: The frequency shows the number of textbooks.
Percentage $= \frac{\text{frequency}}{\text{total}} \times 100$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10J

Fluency

5, 7, 8 | 6–8

Example 18 Constructing frequency tables and histograms

The data below shows the number of hamburgers sold each hour by a 24-hour fast-food store during a 50-hour period.

1	10	18	14	20	11	19	10	17	21
5	16	7	15	21	15	10	22	11	18
12	12	3	12	8	12	6	5	14	14
14	4	9	15	17	19	6	24	16	17
14	11	17	18	19	19	19	18	18	20

a Set up and complete a grouped frequency table, using class intervals 0–, 10–, 20–, etc. Include a percentage frequency column.

b Construct a frequency histogram.

c For how many hours did the fast-food store sell:

i fewer than 10 hamburgers?

ii at least 15 hamburgers?

Solution


Explanation

a

Class interval	Frequency	Percentage frequency
0–	3	6
5–	7	14
10–	15	30
15–	19	38
20–24	6	12
Total	50	100

Create class intervals of 5 from 0 up to 25, since 24 is the maximum number. Record the number of data values in each interval in the frequency column. Convert to a percentage by dividing by the total (50) and multiplying by 100.

b



Create a frequency histogram with frequency on the vertical axis and the class intervals on the horizontal axis. The height of the column shows the frequency of that interval.

c i $3 + 7 = 10$ hours

Fewer than 10 hamburgers covers the 0–4 and 5–9 intervals.

ii $19 + 6 = 25$ hours

At least 15 hamburgers covers the 15–19 and 20–24 intervals.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Now you try

The data below shows the number of cars sold each month at a large city car yard.

18	26	31	12	52	46
38	19	24	40	57	54

a Set up and complete a grouped frequency table, using class intervals 0–, 5–, 10–, etc. Include a percentage frequency column.
b Construct a frequency histogram.
c For how many months did the car yard sell more than 50 cars?
d What percentage of months were less than 20 cars sold?

5 The data below shows the number of ice creams sold from an ice cream van over a 50-day period.

0	5	0	35	14	15	18	21	21	36	45	2	8
2	2	3	17	3	7	28	35	7	21	3	46	47
1	1	3	9	35	22	7	18	36	3	9	2	
11	37	37	45	11	12	14	17	22	1	2	2	



Hint for Q5a: The interval '0–' means scores from 0 to 9.



Hint for Q5c: 'Fewer than 20' doesn't include 20. 'At least 30' includes 30.

a Set up and complete a grouped frequency table using class intervals 0–, 10–, 20– etc. Include a percentage frequency column.
b Construct a frequency histogram.
c How many days did the ice cream van sell:
  i fewer than 20 ice creams?
  ii at least 30 ice creams?
d What percentage of days were 20 or more ice creams sold?

6 The data below shows the mark out of 100 on the Science exam for 60 Year 9 students.

50	67	68	89	82	81	50	50	89	52	60	82	52	60	87
89	71	73	75	83	86	50	52	71	80	95	87	87	87	74
60	60	61	63	63	65	82	86	96	88	50	94	87	64	64
72	71	72	88	86	89	69	71	80	89	92	89	89	60	83



Hint for Q6: The frequency for '50–' will be the number of scores from 50 to 59.

a Set up and complete a grouped frequency table, using class intervals 50–, 60–, 70– etc. Include a percentage frequency column.
b Construct a frequency histogram.
c **i** How many marks were less than 70 out of 100?
  ii What percentage of marks were at least 70 out of 100?

7 The number of goals kicked by a country footballer in each of his last 30 football matches is given below.

8	9	3	6	12	14	8	3	4	5	2	5	6	4	13
8	9	12	11	7	12	14	10	9	8	12	10	11	4	5



Hint for Q7: The frequency for '0–2' will be the number of scores that are 0, 1 or 2. Label the horizontal scale 0, 3, 6, 9 etc.


a Organise the data into a grouped frequency table using class intervals of 0–2, 3–5 etc.
b Draw a frequency histogram for the data.
c In how many games did the player kick fewer than six goals?
d In how many games did he kick more than 11 goals?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


10J

8 Which one of these histograms illustrates a symmetrical data set and which one shows a skewed data set?

a



b





Hint for Q8: Skewed data is bunched either below or above the middle.

Problem-solving and reasoning 9–11 9, 11, 12

9 Write down the missing numbers in these frequency tables; i.e. find the values of the pronumerals.

a

Class interval	Frequency	Percentage frequency
0–4	1	10
5–9	3	c
10–14	4	d
a–19	b	e
Total	10	f

b

Class interval	Frequency	Percentage frequency
40–	20	20
a–	28	b
60–	12	c
70–	d	40
Total	100	e

10 The data below shows the length of overseas phone calls (in minutes) made by a particular household over a six-week period.

1.5	1	1.5	1	4.8	4	4	10.1	9.5	1	3
8	5.9	6	6.4	7	3.5	3.1	3.6	3	4.2	4.3
4	12.5	10.2	10.3	4.5	4.5	3.4	3.5	3.5	5	3.5
3.6	4.5	4.5	12	11	12	14	14	12	13	10.8
12.1	2.4	3.8	4.2	5.6	10.8	11.2	9.3	9.2	8.7	8.5

What percentage of phone calls were more than 3 minutes in length? Answer to one decimal place.



Hint for Q10: Percentage $= \frac{\text{frequency}}{\text{total}} \times 100$

11 This percentage frequency histogram shows the heights of office towers in a city.

a What percentage of office towers have the following heights?

- **i** between 50 m and 100 m
- **ii** less than 150 m
- **iii** no more than 200 m
- **iv** at least 100 m
- **v** between 100 m and 150 m or greater than 200 m

b If the city had 100 office towers, how many would be expected to have a height of:


- **i** between 100 m and 150 m?
- **ii** at least 150 m?

c If the city had 40 office towers, how many would be expected to have a height of:

- **i** between 0 m and 50 m?
- **ii** no more than 150 m?




Hint for Q11: In parts **b** and **c**, number of office towers = percentage × total



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

12 Two students show different histograms for the same set of data.





Hint for Q12: A histogram that shows a lot of detail is more useful.

a Which histogram is more useful in helping to analyse the data?

b What would you advise student B to do differently when constructing the histogram?

The distribution of weekly wages — 13

13 The data below shows the weekly wages of 50 people in dollars.

400	500	552	455	420	424	325	204	860	894	464	379	563
940	384	370	356	345	380	720	540	654	678	628	656	670
740	750	730	766	760	700	700	768	608	576	890	920	874
450	674	725	612	605	600	548	670	230	725	860		

a What is the minimum weekly wage and the maximum weekly wage?

b **i** Organise the data into about 10 class intervals: 200–, 300– etc.

ii Draw a frequency histogram for the data.

c **i** Organise the data into about five class intervals.

ii Draw a frequency histogram for the data.

d Discuss the shapes of the two graphs. Which graph represents the data better and why?



Using a calculator 10J: Graphing grouped data

The activity can be found in the More Resources section of the Interactive Textbook in the form of a printable PDF.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Maths@Work

Personal trainer

Personal training is a growing trend in our society. Personal trainers, while being fit themselves, must have an understanding of mathematics. The mathematics involved in personal training includes calculations of body mass index (BMI) using body mass and weight, calories and kilojoules, business-related mathematics and clients' pulse rates.



1 Calculate the mean pulse rate (measured in beats per minute, bpm) for each of the following clients, over their first five training sessions. Round to a whole number of bpm.
- **a** 85, 92, 86, 90, 88
- **b** 78, 90, 79, 82, 91
- **c** 98, 100, 102, 94, 98
- **d** 120, 122, 134, 116, 104

2 The resting heart rate of a person aged from 6 to 15 is 70 bpm to 100 bpm. Decide if each of the following children aged between 6 and 15 years has a healthy resting heart rate.
- **a** Ella 88 bpm
- **b** Caleb 102 bpm
- **c** Whitney 72 bpm
- **d** Mia 120 bpm
- **e** Kevin 92 bpm

3 As a personal trainer, monitoring a client's resting pulse rate over time is important. Luke's resting pulse rates are displayed in the following stem-and-leaf plot.
Find the mean, mode, median and range to one decimal place of Luke's resting heart rates for the month of February.

Resting heart rate in bpm

Stem	Leaf
7	8 9
8	2 4 4 6 7 8 8
9	0 0 0 1 3 4 5 5 8 8 9
10	4 5 6 6 7 9
11	0 3

8|4 means 84

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

4 A personal trainer needs to be able to compare resting pulse rates to pulse rates after exercise. Look at the back-to-back stem-and-leaf plot below for Luke, and find the mean, mode and median bpm after exercise and compare to **Question 3**.

Heart rate in bpm

Leaf (after exercise)	Stem	Leaf (resting)
8 6 4	7	8 9
6 6 5	8	2 4 4 6 7 8 8
9 8 8 8 8 7 6	9	0 0 0 1 3 4 5 5 8 8 9
6 6 5 4	10	4 5 6 6 7 9
8 6 2	11	0 3
7 6 6 5 4	12	
3 2 2	13	

Using technology


5 Wal, a personal fitness trainer, keeps a monthly record of the PBs (personal best times) for his clients. One popular routine is performing the following three exercises:

- 20 step-ups
- 10 burpees
- 10 squats.

Here is a list of PBs, in seconds, for Wal's clients when doing this routine one October.

94	63	103	84	98	89	73	105	85	115
77	96	87	107	64	90	88	102	91	78
99	119	82	76	82	71	83	80	92	117

Enter these 30 times into a graphics calculator and construct a histogram using interval widths of 10, starting with 60 seconds.
Note: Refer to the PDF in the More Resources section of the Interactive Textbook, which has instructions for graphing grouped data using a graphics calculator.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Modelling

The missing data

Gregory is a marine biologist and is collecting and analysing data regarding shark and dolphin numbers in a particular area of the Great Barrier Reef.

He collects data for 14 consecutive days, but unfortunately two pieces of data ($\boldsymbol{D_1}$ and $\boldsymbol{D_2}$) are lost due to a computer glitch as shown in the table below.

Day	1	2	3	4	5	6	7	8	9	10	11	12	13	14
Sharks	15	12	4	D_1	15	11	19	5	1	10	12	6	24	11
Dolphins	4	7	2	3	2	6	4	2	D_2	4	7	2	4	6

Present a report for the following tasks and ensure that you show clear mathematical workings and explanations where appropriate.

1 Preliminary task

a If Gregory assumes that $D_1 = 10$, find the following:

- **i** The mean and median number of sharks recorded, correct to one decimal place where necessary.
- **ii** The range for the number of sharks recorded over the two weeks.

b If Gregory assumes that $D_2 = 5$, find the following:

- **i** The mean and median number of dolphins recorded, correct to one decimal place where necessary.
- **ii** The range for the number of dolphins recorded over the two weeks.

c Which set of data for the two species (sharks or dolphins) appears to be more inconsistent and spread out?

2 Modelling task

Formulate

The problem is to determine the possible valves of D_1 and D_2 so that Gregory can complete the report based on the statistics he can remember.

a Write down all the relevant information that will help solve this problem, and illustrate using diagrams.

Solve

b Gregory seems to remember that the range for the number of dolphins was 7. Determine the value of D_2 assuming Gregory remembers the range correctly.

c From his memory, Gregory thought that the mean number of sharks over the 14 day period was approximately 12. Based on this value, determine the likely value of D_1.

d From his memory, Gregory thought that the mean number of dolphins over the 14-day period was approximately 5. Based on this value, determine the likely value of D_2.

e If the median number of sharks was 11, find the possible value(s) of D_1.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Evaluate and verify

f If in fact the mean number of dolphins over the 14-day period was 4 after rounding to the nearest integer, find all possible values of D_2.

g If the mean of the first 7 days was the same as the mean of the next 7 days for the dolphin data collected, determine the value of D_2.

h Is it possible for the mean of the first 7 days to be the same as the mean of the next 7 days for the shark data collected? Give reasons.

Communicate

i Summarise your results, showing the calculations that lead to each possible value of D_1 and D_2 and decide which values you believe are mostly likely to represent the missing data.

3 Extension questions

a Draw a dot plot for each species and decide on a set of likely values for D_1 and D_2

b Decide if it is possible for the median of either set of data to be anything other than an integer. If so, determine the values of D_1 and D_2 to allow this to happen.

c Estimate the probability on any given day the shark numbers exceed 12 at the area of the reef Gregory is studying?



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Technology and computational thinking

Simulating candidates

Key technology: Spreadsheets and programming

When selecting more than one item from a group, we are working with compound or multiple events for which tables and tree diagrams could be used to list possible outcomes and their associated probabilities. For more complex problems it may become too cumbersome to use such techniques to calculate probabilities. Hence, computer simulations are used to find suitable approximations.



1 Getting started

Consider a simple selection of two people from a group of five of which two support the Red (R) political party and three support the Blue (B) political party. The two selections are made to fill two key roles on a committee. Assume at this stage that the same person can hold both positions and so the experiment is to be run with replacement.


R/B	R	R	B	B	B
R	(R, R)	(R, R)	(B, R)		
R					
B					
B					
B					

a Complete this table above to show all the outcomes for the two committee positions.

b Use your table results to find the following probability of obtaining:
- **i** 2 Reds
- **ii** 2 Blues
- **iii** 1 Red and 1 Blue in any order.

2 Applying an algorithm

The given flowchart simulates the selection of two people from a group of 5 of which two support the Red (R) political party and three support the Blue (B) political party. Two random integers from 1 to 5 are generated with the numbers 1 and 2 meaning that a Red is chosen. The number of times two reds are chosen is represented by i and the number of trials is represented by n.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

a Run through the algorithm using a random number generator to generate the values of a and b. Suggestions include Excel: = RandBetween(1, 5) or CAS: RandInt(1, 5). Complete the following table for the different n values.

n	a	b	i
1			
2			
3			
4			
5			

b Why does the algorithm stop after 5 trials?

c Which variable counts how many times there are two Reds? How many times were two Red candidates selected in your simulation using the algorithm?

3 Using technology

We will use a spreadsheet to apply the above algorithm simulating the selection of two Red or Blue candidates with replacement.

a Enter the following formulas into a spreadsheet as shown.

	A	B	C	D	E
1	Two-step simulation with replacement			Frequency	=COUNTIF(E5:E24, "YES")
2				Probability	=E1/20
3					
4	Outcome 1	Red or Blue?	Outcome 2	Red or Blue?	Two red?
5	=RANDBETWEEN(1,5)	=IF(OR(A5=1,A5=2), "Red", "Blue")	=RANDBETWEEN(1,5)	=IF(OR(C5=1,C5=2), "Red", "Blue")	=IF(AND(B5="Red", D5="Red"), "YES","NO")

b Explain what the formulas in row 5 do.

c Fill down at cells A5, B5, C5, D5 and E5 to row 24. This should generate results for 20 trials.

d Now fill in the formulas at cells E1 and E2. Explain what you think these formulas do.

e Press Shift F9 to rerun the simulation and note the results in E1 and E2. Keep a record of the probability of obtaining two Reds (cell E2) for 10 reruns of the simulation and find the average of these results. Compare this with the theoretical result obtained in part **1bi**.

The following spreadsheet is similar to the one above, but this time assumes that selections are made without replacement. This means that a person from the group of five cannot be chosen for both positions on the committee.

	A	B	C	D	E	F
1	Two-step simulation without replacement			Frequency	=COUNTIF(F5:F24, "YES")	
2				Probability	=E1/20	
3						
4	Outcome 1	Red or Blue?	Outcome 2	Red or Blue?	Allowed?	Two red?
5	=RANDBETWEEN(1,5)	=IF(OR(A5=1,A5=2), "Red", "Blue")	=RANDBETWEEN(1,5)	=IF(OR(C5=1,C5=2), "Red", "Blue")	=IF(A5=C5,"NO","YES")	=IF(AND(B5="Red", D5="Red",E5="YES"), "YES","NO")
6						

f Enter this information into a new sheet and fill down the cells A5, B5, C5, D5 and E5 to row 24.

g Compare the probability result (cell E2) in this new spreadsheet with the same result from the previous spreadsheet. What does this tell you about the probability of obtaining two Reds by making selections with replacement compared to making selections without replacement?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Puzzles and games

1 One coin says to another coin: *How does my face appear?*
Answer these questions for each scenario, then match the letter in bold to the answers listed below to find the answer to the riddle.


2 coins are tossed 12 times	**1 die is tossed 12 times**	**Choosing one letter from the word PUZZLES**
Pr(HH) = **D**	Pr(5) = **A**	Pr(Z) = **M**
Pr(HT or TH) = **I**	Pr(greater than 2) = **N**	Pr(consonant) = **O**
Expected number of TT = **R**	Expected number of 6s = **S**	
Expected number of one or more Tails = **T**		

$\frac{1}{2}$ 9 2 , 3 $\frac{1}{6}$ $\frac{2}{3}$ $\frac{1}{4}$ $\frac{5}{7}$ $\frac{2}{7}$


2 Match each description (**1–5**) with the most suitable diagram (**A–E**). Note: There is only one description for each diagram.

1 Don't own an iPhone
2 Own an iPhone only
3 Own both an iPod and an iPhone
4 Own an iPod or an iPhone or both
5 Own an iPod


A




B




C



D



E




3 In a class of 28, each student owns a cat or a dog or both. If 18 students own cats and 16 students own dogs, how many students own both a cat and a dog?

4 Write down the set of five positive integers that has a mean of 5, a mode of 8 and a range of 6.

5 A 6-sided die and a 10-sided die are tossed simultaneously. What total sums have the highest chance of occurring? (A total sum means the sum of the two uppermost faces.)

6 Michael needs to deliver parcels to three places (A, B and C, in order) in the city. This diagram shows the different ways that he can travel.



a Draw a tree diagram showing all the possible combinations of transportation.
b What is the total number of possible outcomes?
c Find the probability that Michael will use a different transport each time.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter summary

Statistics and probability

Data and sampling

A survey can be used to collect information using a sample or population.
A sample needs to be chosen carefully to be representative and to avoid bias.

The type of data can be:

- numerical < discrete, continuous
- categorical < ordinal, nominal

Types of sampling include: simple random, systematic and stratified.

Two-step experiments


These can be represented by tables or tree diagrams. They involve more than one step and can occur with or without replacement.
E.g. a bag contains 2 blue counters and 1 green, 2 are selected at random.
(I) Table with replacement.

Pick 2 \ Pick 1	b	b	g
b	bb	bb	gb
b	bb	bb	gb
g	bg	bg	gg

Sample space of 9 outcomes

$\Pr(2 \text{ of same colour}) = \frac{5}{9}$

(II) Tree without replacement



$\Pr(2 \text{ of same colour}) = \frac{2}{6} = \frac{1}{3}$

Probability review

The sample space is the list of all possible outcomes of an experiment.
For all possible outcomes:

$\Pr(\text{event}) = \frac{\text{number of outcomes where event occurs}}{\text{total number of outcomes}}$

E.g. roll a normal six-sided die


$\Pr(>4) = \frac{2}{6} = \frac{1}{3}$

$0 \leq \Pr(\text{event}) \leq 1$

$\Pr(\text{not A}) = 1 - \Pr(\text{A})$

Venn diagrams and two-way tables

These organise data from two or more categories



Venn diagram

	A	Not A	Totals
B	7	3	10
Not B	5	6	11
Totals	12	9	21

i.e. 6 in neither category, 7 in both categories

$\Pr(\text{A}) = \frac{12}{21} = \frac{4}{7}$

$\Pr(\text{B only}) = \frac{3}{21} = \frac{1}{7}$

Stem-and-leaf plots

These display all the data values using a stem and a leaf.
An ordered back-to-back stem-and-leaf plot.

Leaf	Stem	Leaf
9 8 7 2	1	0 3
7 4 3 3	2	2 2 4
5 2 1	3	3 6 7 8
7	4	4 5 9
0	5	0

skewed; symmetrical

3 | 5 means 35 ← key

Experimental probability/ relative frequency

This is calculated from results of an experiment or survey.

$\text{Experimental probability} = \frac{\text{number of times event occurs}}{\text{total number of trials}}$

$\text{Expected number of occurrences} = \text{probability} \times \text{number of trials}$

Measure of spread

Range = maximum value − minimum value

Summarising data: Measures of centre

Mode is the most common value (there can be more than one). Two modes means data is bimodal.

$\text{Mean} = \frac{\text{sum of all values}}{\text{number of values}}$

Median is the middle value of data that is ordered.

odd data set: 2 4 (7) 10 12 — median

even data set: 2 4 6 10 15 18 — 8

$\text{median} = \frac{6+10}{2}$

An outlier is a value that is not near the rest of the data.

Grouped data


Data values can be grouped into class intervals, e.g. 0–4, 5–9 etc. and recorded in a frequency table.

Frequency table

Class interval	Frequency	Percentage frequency
0–	10	20
5–	20	40
10–	5	10
15–20	15	30
Total	50	100

The frequency or percentage frequency of each interval can be recorded in a histogram.

Frequency histogram



Essential Mathematics Core for the Victorian Curriculum Year 9
ISBN 978-1-009-59315-1
© Greenwood et al. 2025
Photocopying is restricted under law and this material must not be transferred to another party.
Cambridge University Press


Chapter checklist

Chapter checklist

A version of this checklist that you can print out and complete can be downloaded from your Interactive Textbook.



		✔
10A	**1 I can calculate a probability from experiments with equally likely outcomes.** e.g. A letter is chosen from the word COMPLEMENT. Find the probability that the letter is: **a** an E **b** not a vowel **c** an E or a consonant	
10B	**2 I can use a Venn diagram to calculate probabilities.** e.g. From a group of 30 tennis players, 16 do a slice backhand, 18 do a top-spin backhand and 6 do both types of backhand. Illustrate this information in a Venn diagram and **a** state the number of tennis players who do a slice backhand only **b** find the probability that a randomly selected tennis player does a slice or a top-spin backhand.	
10B	**3 I can use a two-way table to calculate probabilities.** e.g. From a group of 20 single car owners, 15 have a car which runs on petrol, 7 have a car which runs on electricity and 4 people have a car which runs on both types of fuel. Illustrate this information using a two-way table and **a** find the number of people who have a car that runs on neither type of fuel **b** find the probability that a randomly selected car owner has a car which runs on petrol only.	
10C	**4 I can construct an array (table) for a two-step experiment with replacement.** e.g. A fair 4-sided die is rolled twice. List all the outcomes (the sum total from the two dice) using a table and find: **a** Pr(a double) **b** Pr(sum of at least 5) **c** Pr(sum not equal to 7).	
10C	**5 I can construct an array (table) for a two-step experiment without replacement.** e.g. Two letters are chosen from the word KEY, without replacement. Construct a table to list the sample space and find the probability of: **a** obtaining the outcome (K, E) **b** selecting an E and a Y in any order.	
10D	**6 I can construct and use a tree diagram to find probabilities for multistage experiments with replacement.** e.g. Two marbles are selected from a container including 2 clear and 2 coloured marbles. Two marbles are randomly chosen with replacement. Represent the selections using a tree diagram that shows all possible outcomes. Find the probability of selecting: **a** one clear marble **b** at least one coloured marble.	
10D	**7 I can construct and use a tree diagram to find probabilities for multistage experiments without replacement.** e.g. A box contains 2 dark and 3 white chocolates. Draw a tree diagram to show the outcomes and probabilities of the selection of two chocolates without replacement. Find the probability of selecting: **a** one white **b** at least one dark chocolate.	
10E	**8 I can find an experimental probability and expected number of occurrences.** e.g. 40 people were surveyed regarding their main mode of transport to work. 25 said *Public transport*, 10 said *Drive* and 5 said *Walk*. Find: **a** the experimental probability that a person drives to work **b** the expected number of people who use public transport from a group of 1000.	

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

		✔
10E	**9 I can classify data.** e.g. Classify the data type from the following surveys. **a** The number of Christmas presents received **b** The breakfast buffet at a hotel: poor, average, good, excellent	
10E	**10 I can choose a sampling method.** e.g. Identify the sampling method used in each of these surveys. **a** An airline chooses every sixth passenger on a flight **b** One hundred students are grouped into year level and then tested on spelling words	
10E	**11 I can identify bias in a data collection method.** e.g. One hundred people at the airport are surveyed about how often they travel. This data is used to make conclusions about that city and how often people travel. Do you think this survey will contain bias? If so, explain why.	
10G	**12 I can find the mean, median, mode and range of a simple data set.** e.g. For the data sets: 3, 4, 5, 2, 3, 3, 1 and 12, 21, 21, 25, 18, 14 find: **a** the mean **b** median **c** mode **d** range	
10G	**13 I can solve problems associated with data means.** e.g. Hilbert received 78, 59, 66 and 69 on his previous four maths tests. What is his current average and what does he need to score in his next test to raise his average to 70?	
10H	**14 I can interpret a common statistical graph.** e.g. This column graph shows how many goals four teams scored in a round of soccer.  **a** How many goals did the Blues score? **b** What was the mean and median score for the four teams? **c** What combined percentage of the goals were scored by the Reds and the Blues?	

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter checklist

		✔
10H	**15 I can decide if a graph may be misleading.** e.g. The following two graphs shows the two-party preferred position of the Labor and Liberal parties after an election. Which one provides a misleading margin of difference and why? a  b  	
10I	**16 I can construct a stem-and-leaf plot from a data set and calculate summary statistics.** e.g. For this data set construct a stem-and-leaf plot and find the mode and median. 26 19 32 42 49 16 35 29 22 25 26 37 15 28	
10I	**17 I can construct a back-to-back stem-and-leaf plot and compare the two data sets.** e.g. The total games scores of two tennis teams for a season of 9 matches is recorded as follows. Team A 31 45 69 40 76 55 49 12 28 Team B 23 34 16 29 27 12 25 29 31 Construct an ordered back-to-back stem-and-leaf plot and compare and comment on the differences between the performance of the two teams.	
10J	**18 I can construct and analyse a frequency table and a histogram.** e.g. The age of 20 tagged sea turtles is recorded as follows. 45 63 79 66 38 97 82 44 59 31 58 62 69 53 85 93 85 77 62 41 Organise the data into a frequency table, using class intervals of 10, and include a percentage frequency column. Construct a frequency histogram and a percentage frequency histogram and determine the percentage of turtles older than 50 years.	

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Short-answer questions

10A **1** Determine the probability of each of the following.
- **a** Rolling more than 2 on a normal six-sided die
- **b** Selecting a vowel from the word EDUCATION
- **c** Selecting a pink or white jelly bean from a packet containing 4 pink, 2 white and 4 black jelly beans

10B **2** From a survey of 50 people, 30 have the newspaper delivered, 25 read it online, 10 do both and 5 do neither.




- **a** Construct a Venn diagram for the survey results.
- **b** How many people only read the newspaper online?
- **c** If one of the 50 people were randomly selected, find:
  - **i** Pr(have paper delivered and read it online)
  - **ii** Pr(don't have it delivered)
  - **iii** Pr(only read it online)

10B **3** **a** Copy and complete this two-way table.

	A	Not A	Total
B		16	
Not B	8		20
Total	17		

- **b** Convert the information into a Venn diagram, as shown.



- **c** Find the following.
  - **i** Pr(not B)
  - **ii** Pr(both A and B)
  - **iii** number of 'A only'
  - **iv** number in either A or B or both A and B

10C **4** A spinner with equal areas of red, green and blue is spun and a four-sided die numbered 1 to 4 is rolled.

		Die			
		1	2	3	4
Spinner	red	(red, 1)	(red, 2)		
	green				
	blue				

- **a** Complete a table like the one shown and state the number of outcomes in the sample space.
- **b** Find the probability that:
  - **i** the outcome is red and an even number
  - **ii** the outcome is blue or green and a 4
  - **iii** the outcome does not involve blue

10D **5** Libby randomly selects two coins from her pocket *without replacement*. Her pocket contains a $1 coin and two 10-cent coins.
- **a** List all the possible combinations using a tree diagram.
- **b** If a chocolate bar costs $1.10, find the probability that she can hand over the two coins to pay for it.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter review

10E **6** A quality controller records the frequency of the types of chocolates from a sample of 120 off its production line.

Centre	Soft	Hard	Nut
Frequency	50	22	48

a What is the experimental probability of randomly selecting a nut centre?
b In a box of 24 chocolates, how many would be expected to have a non-soft centre?

10F **7** Jody is planning to run a survey asking people in her year about their current favourite Netflix show.



a Classify the type of data to be collected.
b Identify the sampling method in each of the following survey methods she is considering.
  i Assigning each student a number and generating some random numbers
  ii Surveying every fifth person in the school canteen queue.
c After planning, Jody surveys everyone in her extension science class, and then will use the data to make conclusions about her year level. Do you think this survey will contain bias? If so, explain why.

10G **8** Claudia records the number of emails she receives each weekday for two weeks as follows.

30 31 33 23 29 31 21 15 24 23

Find:
a the mean **b** the median **c** the mode **d** the range

10I **9** Two mobile phone salespeople are both aiming for a promotion to be the new assistant store manager. The best salesperson over a 15-week period will achieve the promotion. The number of mobile phones they sold each week is recorded below.

Employee 1: 21 34 40 38 46 36 23 51 35 25 39 19 35 53 45
Employee 2: 37 32 29 41 24 17 28 20 37 48 42 38 17 40 45

a Draw an ordered back-to-back stem-and-leaf plot for the data.
b For each employee, find:
  i the median number of sales
  ii the mean number of sales
c By comparing the two sets of data, state, with reasons, who you think should get the promotion.
d Describe each employee's data as approximately symmetrical or skewed.

10J **10** The data below represents the finish times, in minutes, of 25 competitors in a local car rally race.

134 147 162 164 145 159 151 143 136 155 163 157 168
171 152 128 144 161 158 136 178 152 167 154 161

a Record the above data in a frequency table in class intervals of 120−, 130− etc. Include a percentage frequency column.
b Construct a frequency histogram.
c Determine:
  i the number of competitors that finished in less than 140 minutes
  ii the percentage of competitors that finished between 130 and 160 minutes


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10H

11 This column graph shows the value of some of the boats that were badly damaged in cyclone Yasi in North Queensland in 2011.

a Find the mean and median boat value.

b Do you think the mean or the median would better represent the 'average' of these boat values? Give a reason for your answer.



Multiple-choice questions

10A

1 A letter is randomly chosen from the word XYLOPHONE. The probability that it is an O is:

A $\frac{1}{8}$ **B** $\frac{2}{9}$ **C** $\frac{1}{4}$ **D** $\frac{1}{9}$ **E** $\frac{1}{3}$

10B


2 The values of x and y in the two-way table are:


A $x = 12, y = 8$ **B** $x = 12, y = 11$
C $x = 16, y = 4$ **D** $x = 10, y = 1$
E $x = 14, y = 6$


	A	Not A	Total
B		5	9
Not B	8	y	
Total	x		25


10B


3 **i** Which shaded region represents both A and B?
ii Which shaded region represents A only?
iii Which shaded region represents A or B or both?
iv Which shaded region represents B and not A?

A


B


C


D


E


10D

4 A bag contains 2 green balls and 1 red ball. Two balls are randomly selected *without replacement*. Using a tree diagram, the probability of selecting one of each colour is:

A $\frac{1}{2}$ **B** $\frac{2}{3}$ **C** $\frac{5}{6}$ **D** $\frac{1}{3}$ **E** $\frac{3}{4}$

10E

5 From rolling a biased die, a class finds an experimental probability of 0.3 of rolling a 5. From 500 rolls of the die, the expected number of 5s would be:

A 300 **B** 167 **C** 180 **D** 150 **E** 210

10F

6 The type of data generated by the survey question 'How long is your arm length?' is:

A a random number **B** categorical ordinal
C categorical nominal **D** numerical discrete
E numerical continuous

10I

7 **i** The median of the data in this stem-and-leaf plot is:

A 74 **B** 71 **C** 86
D 65 **E** 70

ii The range of the data in the stem-and-leaf plot is:

A 3 **B** 8 **C** 33
D 86 **E** 14

Stem	Leaf
5	3 5 8
6	1 4 7
7	0 2 4 7 9
8	2 6 6

7 | 4 means 74

ISBN 978-1-009-59315-1
© Greenwood et al. 2025
Photocopying is restricted under law and this material must not be transferred to another party.


Chapter review

10G **8** If Jacob achieved scores of 12, 9, 7 and 12 on his last four language vocabulary tests, what score must he get on the fifth test to have a mean of 11?

A 16 **B** 14 **C** 11 **D** 13 **E** 15

10J **9** This frequency histogram shows the times of finishers in a fun run. The percentage of competitors that finished in better than 40 minutes was:

A 55% **B** 85% **C** 50%
D 62.5% **E** 60%



Extended-response question

1 The local Sunday market has a number of fundraising activities.

a For $1 you can spin a spinner numbered 1–5 twice. If you spin two even numbers you receive $2 (your dollar back plus an extra dollar), if you spin two odd numbers you receive your dollar back and for any other result you lose your dollar.

i Complete the table shown to list the sample space.

		First spin				
		1	2	3	4	5
Second spin	1	(1, 1)	(2, 1)			
	2					
	3					
	4					
	5					

ii What is the probability of losing your dollar (i.e. spinning one odd and one even number)?

iii What is the probability of making a dollar profit (i.e. spinning two even numbers)?

iv In 50 attempts, how many times would you expect to lose your dollar?

v If you start with $100 and have 100 attempts, how much money would you expect to end up with?

b Forty-five people were surveyed as they walked through the market as to whether they bought a sausage and/or a drink from the sausage sizzle. Twenty-five people bought a sausage and 30 people bought a drink, with 15 buying both.

i Construct a Venn diagram to represent this information.

ii How many people bought neither a drink nor a sausage?

iii How many people bought a sausage only?

iv If a person was randomly selected from the 45, what is the probability they bought a drink but not a sausage?

v Find Pr(didn't buy a sausage).

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

2 The delay time (in minutes) of the flight departure of the same evening flight of two rival airlines was recorded over 30 consecutive days. The data is shown below.

Airline A	2	11	6	14	18	1	7	4	12	14	9	2	13	4	19
	13	17	3	52	24	19	12	14	0	7	13	18	1	23	8
Airline B	6	12	9	22	2	15	10	5	10	19	5	12	7	11	18
	21	15	10	4	10	7	18	1	18	8	25	4	22	19	26

a Copy and complete this ordered back-to-back stem-and-leaf plot for the data. Use 2 lines for each stem, starting with '0' for leaves $0 - 4$, and then '0*' for leaves $5 - 9$ etc.

Airline A		Airline B
	0	
	0*	
	1	
	1*	
	2	
	2*	
	3	
	3*	
	4	
	4*	
	5	

1 | 2 means 12

1* | 5 means 15

b Does the data for airline A appear to have any outliers (numbers not near the majority of data elements)?

c By removing any outliers listed in part **b**, find the following for each airline, rounding to one decimal place where necessary.

i the median


ii the mean

d Airline A reports that half its flights for that month had a delay time of less than 10 minutes. Is this claim correct? Explain.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

11

Introduction to quadratic equations and graphs



Essential mathematics: why understanding quadratic equations and their graphs is important

Quadratic equations and their graphs, known as parabolas, have practical applications in various fields.

In golf club design, **engineers** use quadratic equations to model the ball's flight path, adjusting the club's angle and weight for better performance. Drivers hit far and low, fairway woods balance distance and control, hybrids are versatile, irons vary in height and distance, wedges are for short, high shots, and putters focus on accuracy. Each club suits different shots!

Meteor showers occur when Earth passes through the debris left by a comet. **Astronomers** use quadratic equations to model the paths of these debris particles. By understanding these parabolic paths, they can predict when and where meteor showers will occur.

Engineers also use quadratic equations to calculate the curves and slopes of roller coasters, ensuring the rides are both thrilling and safe. For example, the path of a roller coaster's hill can be modelled by a parabola, helping designers determine the height and speed of the ride.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.



In this chapter

11A Quadratic equations
11B Solving $ax^2 + bx = 0$ and $x^2 = d$
11C Solving $x^2 + bx + c = 0$
11D Using quadratic equations to solve problems
11E The parabola
11F Sketching $y = ax^2$ with dilations and reflections
11G Sketching translations of $y = x^2$
11H Sketching parabolas using intercept form ★

Victorian Curriculum 2.0

This chapter covers the following content descriptors in the Victorian Curriculum 2.0:

ALGEBRA

VC2M9A02, VC2M9A05, VC2M9A06, VC2M9A07

Please refer to the curriculum support documentation in the teacher resources for a full and comprehensive mapping of this chapter to the related curriculum content descriptors.

© VCAA

Online resources

A host of additional online resources are included as part of your Interactive Textbook, including HOTmaths content, video demonstrations of all worked examples, auto-marked quizzes and much more.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025
Photocopying is restricted under law and this material must not be transferred to another party.
Cambridge University Press

Warm-up quiz

1 Evaluate the following by substituting each of the four values of x into each expression in parts **a**–**d**.

i $x = 0$
ii $x = 2$
iii $x = -1$
iv $x = -3$

a x^2 **b** $2x^2 - 3x$ **c** $-x^2 + 3$ **d** $x^2 + 2x - 1$

2 Does $x = 1$ satisfy the following equations? Substitute to find out.

a $x^2 - 4x + 3 = 0$
b $x^2 - 2x = 0$
c $1 - x^2 = 0$

3 Solve the following for x.

a $2x = 0$ **b** $x - 3 = 0$
c $x + 5 = 0$ **d** $2x + 6 = 0$
e $3x - 12 = 0$ **f** $3x - 4 = 0$

4 Rearrange the following to produce 0 on the right-hand side.

a $a^2 = 2a$ **b** $a^2 = 3a - 1$

5 State the highest common factor of the following pairs of terms.

a $4x$ and $2xy$ **b** x^2 and $2x$
c $3x$ and $9x^2$ **d** $2x^2$ and $4x$
e $4x^2$ and 36 **f** $5x^2$ and 20

6 Expand these expressions.

a $2(x - 3)$ **b** $x(2 - x)$
c $(x + 1)(x - 2)$ **d** $(x - 3)(x - 4)$
e $(x + 2)(x - 2)$ **f** $(x - 5)(x + 5)$

7 Complete these tables of values for y given the x-value and the rule relating x and y.

a $y = x^2 - 4$

x	−3	−2	−1	0	1	2	3
y							

b $y = (x + 2)^2$

x	−3	−2	−1	0	1	2	3
y							

8 Give the x-coordinate of the point halfway between the following pairs of points.

a $(0, 0)$ and $(10, 0)$
b $(2, 0)$ and $(4, 0)$
c $(-1, 0)$ and $(3, 0)$
d $(3, 0)$ and $(8, 0)$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

11A Quadratic equations

Learning intentions

- To be able to identify a quadratic equation
- To know how to write a quadratic equation in standard form
- To understand the Null Factor Law
- To be able to apply the Null Factor Law to solve an equation in factorised form

Key vocabulary: quadratic equation, standard form, Null Factor Law, product

Quadratic equations are commonplace in theoretical and practical applications of mathematics. They are used to solve problems in geometry and measurement as well as in number theory and physics. The path that a projectile takes while flying through the air, for example, can be analysed using quadratic equations.

A quadratic equation can be written in the form $ax^2 + bx + c = 0$ where a, b and c are constants and $a \neq 0$. Examples include $x^2 - 2x + 1 = 0$, $5x^2 - 3 = 0$ and $-0.2x^2 + 4x = 0$. Unlike linear equations which have a single solution, quadratic equations can have zero, one or two solutions. For example, $x = 2$ and $x = -1$ are solutions to the quadratic equation $x^2 - x - 2 = 0$ since $2^2 - 2 - 2 = 0$ and $(-1)^2 - (-1) - 2 = 0$. One method for finding the solutions to quadratic equations involves the use of the Null Factor Law where each factor of a factorised quadratic expression is equated to zero.



Quadratic equations are widely used in many areas of work including: sports scientists analysing the trajectory of a javelin or discus, electronic engineers finding currents and voltages, and financial analysts modelling future revenue and profit.

Lesson starter: Exploring the Null Factor Law

$x = 1$ is not a solution to the quadratic equation $x^2 - x - 12 = 0$ since $1^2 - 1 - 12 \neq 0$.

- Use trial and error to find at least one of the two numbers that are solutions to $x^2 - x - 12 = 0$.
- Rewrite the equation in factorised form.

  $x^2 - x - 12 = 0$ becomes $(\quad\quad)(\quad\quad) = 0$

- Now repeat the first task above to find solutions to the equation using the factorised form.
- Was the factorised form easier to work with? Discuss.

Key ideas

- A **quadratic equation** can be written in the form $ax^2 + bx + c = 0$.
  - This is called **standard form**.
  - a, b and c are constants and $a \neq 0$.
- The **Null Factor Law** states that if the product of two numbers is zero, then either or both of the two numbers is zero.
  - If $p \times q = 0$ then $p = 0$ or $q = 0$.
  - If $(x + 1)(x - 3) = 0$ then $x + 1 = 0$ (so $x = -1$) or $x - 3 = 0$ (so $x = 3$).
  - $x + 1$ and $x - 3$ are the linear factors of $x^2 - 2x - 3$, which factorises to $(x + 1)(x - 3)$.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Exercise 11A

Understanding

1–3 | 2, 3

1 Evaluate these quadratic expressions by substituting the x-value given in the brackets.

a $x^2 - 5 \ (x = 1)$ **b** $x^2 + 2x \ (x = -1)$ **c** $3x^2 - x + 2 \ (x = 5)$

2 Decide if the following equations are quadratics (yes or no).

a $1 - 2x = 0$ **b** $x^2 + 3x + 1 = 0$ **c** $x^2 - 1 = 0$

d $x^3 + x^2 - 2 = 0$ **e** $x^5 + 1 - x = 0$ **f** $x + 2 = 4x^2$

3 Solve these linear equations.

a $x - 1 = 0$ **b** $x + 3 = 0$ **c** $2x + 4 = 0$

Fluency

4–5(½), 6, 8(½) | 4–5(½), 7, 8(½)

Example 1 Writing in standard form

Write these quadratic equations in standard form: $ax^2 + bx + c = 0, a > 0$.

a $x^2 = 2x + 7$ **b** $2(x^2 - 3x) = 5$ **c** $2x - 7 = -3x^2$

Solution

a $x^2 = 2x + 7$

$x^2 - 2x - 7 = 0$

Explanation

We require the form $ax^2 + bx + c = 0$. Subtract $2x$ and 7 from both sides to move all terms to the left-hand side.

b $2(x^2 - 3x) = 5$

$2x^2 - 6x = 5$

$2x^2 - 6x - 5 = 0$

First expand brackets then subtract 5 from both sides.

c $2x - 7 = -3x^2$

$3x^2 + 2x - 7 = 0$

Add $3x^2$ to both sides, to make the x^2 coefficient positive.

Now you try

Write these quadratic equations in standard form: $ax^2 + bx + c = 0, a > 0$.

a $x^2 = 3x + 5$ **b** $2(x^2 + 4x) = 7$ **c** $3x - 4 = -2x^2$

4 Write these quadratic equations in standard form: $ax^2 + bx + c = 0, a > 0$.

a $x^2 = 7x + 2$ **b** $x^2 - 5x = -2$

c $x^2 = 4x - 1$ **d** $3(x^2 - 2x) = 4$

e $2(x^2 + x) + 1 = 0$ **f** $3(x^2 - x) = -4$

g $4 = -3x^2$ **h** $3x = x^2 - 1$



Hint for Q4: Aim to have a positive *a* value on one side

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Example 2 Testing for a solution

Substitute the given x-value into the equation and say whether or not it is a solution.

a $x^2 + x - 6 = 0 \quad (x = 2)$ **b** $2x^2 + 5x - 3 = 0 \quad (x = -4)$

Solution

a $x^2 + x - 6 = 2^2 + 2 - 6$
$= 6 - 6$
$= 0$
$\therefore$ $x = 2$ is a solution

Explanation

Substitute $x = 2$ into the equation to see if the left-hand side equals zero.
$x = 2$ satisfies the equation, so $x = 2$ is a solution.

b $2x^2 + 5x - 3 = 2(-4)^2 + 5(-4) - 3$
$= 2 \times 16 + (-20) - 3$
$= 32 - 20 - 3$
$= 9$
$\therefore$ $x = -4$ is not a solution

Substitute $x = -4$. Recall that $(-4)^2 = 16$ and $5 \times (-4) = -20$.

The equation is not satisfied since it is $\neq 0$, so $x = -4$ is not a solution.

Now you try

Substitute the given x-value into the equation and say whether or not it is a solution.

a $x^2 + x - 3 = 0 \quad (x = 1)$ **b** $3x^2 + 4x - 15 = 0 \quad (x = -3)$

5 Substitute the given x-value into the quadratic equation and say whether or not it is a solution.

a $x^2 - 1 = 0 \ (x = 1)$ **b** $x^2 - 25 = 0 \ (x = 5)$
c $x^2 - 4 = 0 \ (x = 1)$ **d** $2x^2 + 1 = 0 \ (x = 0)$
e $x^2 - 9 = 0 \ (x = -3)$ **f** $x^2 - x - 12 = 0 \ (x = 5)$
g $x^2 + 2x + 1 = 0 \ (x = -1)$ **h** $2x^2 - x + 3 = 0 \ (x = -1)$



Hint for Q5: If when substituted the equation is true then the x value is a solution

6 Substitute $x = -2$ and $x = 5$ into the equation $x^2 - 3x - 10 = 0$. What do you notice?

7 Substitute $x = -3$ and $x = 4$ into the equation $x^2 - x - 12 = 0$. What do you notice?

Example 3 Using the Null Factor Law

Use the Null Factor Law to solve these equations.

a $x(x + 2) = 0$ **b** $(x - 1)(x + 5) = 0$

Solution

a $x(x + 2) = 0$
$x = 0$ or $x + 2 = 0$
$x = 0$ or $x = -2$

Explanation

The factors are x and $(x + 2)$. Solve each linear factor equal to zero.
Check each solution by substituting into the original equation.

Continued on next page

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

11A

b $(x-1)(x+5)=0$	Solve each factor $(x-1)$ and $(x+5)$ equal
$x-1=0$ or $x+5=0$	to zero.
$x=1$ or $x=-5$	Check your solutions using substitution.

Now you try

Use the Null Factor Law to solve these equations.

a $x(x+5)=0$ b $(x+2)(x-3)=0$

8 Use the Null Factor Law to solve these equations.

a $x(x+1)=0$ b $x(x+5)=0$
c $x(x-2)=0$ d $x(x-7)=0$
e $3x(x-1)=0$ f $2x(x+5)=0$
g $(x+1)(x-3)=0$ h $(x-4)(x+2)=0$
i $(x+7)(x-3)=0$ j $(x-5)(x+4)=0$
k $(x+7)(x-6)=0$ l $(x+12)(x-9)=0$



Hint for Q8: If $3x=0$ then $x=0$

Problem-solving and reasoning

9–11(½) | 9–11(½), 12, 13

Example 4 Using the null factor law to find fractional solutions

Use the Null Factor Law to solve.
$(2x-1)(5x+3)=0$

Solution	Explanation
$(2x-1)(5x+3)=0$	The two factors are $(2x-1)$ and $(5x+3)$.
$2x-1=0$ or $5x+3=0$	Each one results in a two-step linear equation.
$2x=1$ or $5x=-3$	
$x=\frac{1}{2}$ or $x=-\frac{3}{5}$	Check your solutions using substitution.

Now you try

Use the Null Factor Law to solve.
$(3x-2)(2x+5)=0$

9 Use the Null Factor Law to solve these equations.

a $(2x-1)(x+2)=0$ b $(x+2)(3x-1)=0$ c $(5x+2)(x+4)=0$
d $(x-1)(3x-1)=0$ e $(x+5)(7x+2)=0$ f $(3x-2)(5x+1)=0$
g $(11x-7)(2x-13)=0$ h $(4x+9)(2x-7)=0$ i $(3x-4)(7x+1)=0$

10 Find the numbers which satisfy the given condition.

a The product of x and a number 3 more than x is zero.
b The product of x and a number 7 less than x is zero.
c The product of a number 1 less than x and a number 4 more than x is zero.
d The product of a number 1 less than twice x and 6 more than x is zero.



Hint for Q10: The product of x and 5 is $x \times 5$ or $5x$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

11 These quadratic equations have two integer solutions between -5 and 5. Use trial and error to find them.

a $x^2 - x - 2 = 0$ **b** $x^2 - 4x + 3 = 0$

c $x^2 - 4x = 0$ **d** $x^2 + 3x = 0$

e $x^2 + 3x - 4 = 0$ **f** $x^2 - 16 = 0$

12 Consider the quadratic equation $(x + 2)^2 = 0$.

a Write the equation in the form (_____)(_____) $= 0$.

b Use the Null Factor Law to find the solutions to the equation. What do you notice?

c Now solve these quadratic equations.

i $(x + 3)^2 = 0$ **ii** $(x - 5)^2 = 0$

13 Consider the equation $3(x - 1)(x + 2) = 0$.

a First divide both sides of the equation by 3. Write down the new equation.

b Solve the equation using the Null Factor Law.

c Compare the given original equation with the equation found in part **a**. Explain why the solutions are the same.

d Solve these equations.

i $7(x + 2)(x - 3) = 0$ **ii** $11x(x + 2) = 0$

Polynomials — 14, 15

14 Polynomials are sums of integer powers of x. They are given names according to the highest power of x in the polynomial expression.

Example	Polynomial name
2	Constant
$2x + 1$	Linear
$x^2 - 2x + 5$	Quadratic
$x^3 - x^2 + 6x - 1$	Cubic
$7x^4 + x^3 + 2x - x + 4$	Quartic
$4x^5 - x + 1$	Quintic

Name these polynomial equations.

a $3x - 1 = 0$ **b** $x^2 + 2 = 0$

c $x^5 - x^4 + 3 = 0$ **d** $5 - 2x + x^3 = 0$

e $3x - 2x^4 + x^2 = 0$ **f** $5 - x^5 = x^4 + x$

15 Solve these polynomial equations using the Null Factor Law, to find the three solutions.

a $(x + 1)(x - 3)(x + 2) = 0$

b $(x - 2)(x - 5)(x + 11) = 0$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

11B Solving $ax^2 + bx = 0$ and $x^2 = d$

Learning intentions

- To know the steps required to solve a quadratic equation
- To be able to factorise a quadratic equation in order to apply the Null Factor Law
- To be able to find the solutions of quadratic equations involving common factors or perfect squares

Key vocabulary: Null Factor Law, factorise, perfect square

When using the Null Factor Law, we notice that equations must first be expressed as a product of two factors. Hence, any equation not in this form must first be factorised. Two types of quadratic equations are studied here. The first is of the form $ax^2 + bx = 0$, where x is a common factor, and the second is of the form $x^2 = d$.



Aerospace engineers use quadratic equations, such as for the trajectory of a rocket launching a communications satellite. An object launched vertically with velocity, u, moving under gravity, with acceleration, g, reaches a maximum height, h, where $u^2 = 2gh$.

Lesson starter: Which factorisation technique?

These two equations may look similar but they are not the same: $x^2 - 9x = 0$ and $x^2 - 9 = 0$.

- Discuss how you could factorise each expression on the left-hand side of the equations.
- How does the factorised form help to solve the equations? What are the solutions? Are the solutions the same for both equations?
- By rearranging $x^2 - 9 = 0$ into $x^2 = 9$, can you explain how to find the two solutions you found above?

Key ideas

- When solving an equation of the form $ax^2 + bx = 0$, factorise by taking out common factors including x.

$$2x^2 - 8x = 0$$
$$2x(x - 4) = 0$$
$$2x = 0 \text{ or } x - 4 = 0$$
$$x = 0 \text{ or } x = 4$$

- When solving an equation of the form $ax^2 = d$, divide both sides by a and then take the square root of each side.

$$5x^2 = 20$$
$$x^2 = 4$$
$$x = 2 \text{ or } x = -2$$

- $x^2 = d^2$ could also be rearranged to $x^2 - d^2 = 0$ and then factorised as a difference of two squares.

$$x^2 - 4 = 0$$
$$(x + 2)(x - 2) = 0$$
$$x = -2 \text{ or } x = 2$$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Exercise 11B

Understanding

1–4 | 1-4

1 State the highest common factor of these pairs of terms.

a $16x$ and 24 **b** x^2 and $2x$ **c** $3x^2$ and $6x$

2 Factorise these expressions fully by first taking out the highest common factor.

a $x^2 - 3x$ **b** $6x^2 + 4x$

c $4x - 16x^2$ **d** $2x^2 - 8$



Hint for Q2: Answer format for part **a** will be $x(_____)$.

3 Use the Null Factor Law to state the solutions to these equations.

a $x(x - 3) = 0$ **b** $4x(x + 1) = 0$

4 Explain why $x = 3$ and $x = -3$ are solutions to $x^2 = 9$.

Fluency

5(½), 7,8(½) | 5–8(½)

Example 5 Solving quadratic equations of the form $ax^2 + bx = 0$

Solve $x^2 + 4x = 0$.

Solution	Explanation
$x^2 + 4x = 0$	Factorise by taking out the common factor x.
$x(x + 4) = 0$	Using the Null Factor Law, set each factor, x and $(x + 4)$, equal to 0.
$x = 0$ or $x + 4 = 0$	Solve for x.
$x = 0$ or $x = -4$	Check your solutions using substitution.

Now you try

Solve $x^2 + 6x = 0$

5 Solve these equations.

a $x^2 + 3x = 0$ **b** $x^2 + 7x = 0$

c $x^2 + 4x = 0$ **d** $x^2 - 5x = 0$

e $x^2 - 8x = 0$ **f** $x^2 - 2x = 0$

g $x^2 + \frac{1}{3}x = 0$ **h** $x^2 - \frac{1}{2}x = 0$



Hint for Q5: First factorise by taking out x.

6 Solve these equations by first taking out the highest common factor.

a $2x^2 - 6x = 0$ **b** $3x^2 - 12x = 0$

c $4x^2 + 20x = 0$ **d** $6x^2 - 18x = 0$

e $-5x^2 + 15x = 0$ **f** $-2x^2 - 8x = 0$



Hint for Q6: Take out the highest common factor.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

11B

Example 6 Solving $ax^2 = bx$

Solve $2x^2 = 8x$.

Solution

$2x^2 = 8x$

$2x^2 - 8x = 0$

$2x(x - 4) = 0$

$2x = 0$ or $x - 4 = 0$

$x = 0$ or $x = 4$

Explanation

Make the right-hand side equal to zero by subtracting $8x$ from both sides.

Factorise by taking out the common factor of $2x$ and apply the Null Factor Law to solve.

For $2 \times x = 0$, x must equal 0.

Now you try

Solve $3x^2 = 15x$.

7 Solve the following equations by first rearranging into standard form.

a $x^2 = 3x$ **b** $5x^2 = 10x$

c $4x^2 = 16x$ **d** $3x^2 = -9x$

e $2x^2 = -8x$ **f** $7x^2 = -21x$



Hint for Q7: First make a zero on the right hand side

Example 7 Solving equations of the form $x^2 = d$

Solve $x^2 = 16$.

Solution

$x^2 = 16$

$x = 4$ or $x = -4$

Alternative method:

$x^2 = 16$

$x^2 - 16 = 0$

$(x + 4)(x - 4) = 0$

$x + 4 = 0$ or $x - 4 = 0$

$x = -4$ or $x = 4$

Explanation

As this is a perfect square, take the square root of both sides.

$x = 4$ or -4 since $4^2 = 16$ and $(-4)^2 = 16$.

Rearrange into standard form.

Factorise using $a^2 - b^2 = (a + b)(a - b)$ then use the Null Factor Law to find the solutions.

Now you try

Solve $x^2 = 49$.

8 Solve the following equations.

a $x^2 = 9$

b $x^2 = 36$

c $x^2 = 25$

d $x^2 - 144 = 0$

e $x^2 - 81 = 0$

f $x^2 - 400 = 0$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Problem-solving and reasoning

9(½), 10, 11 | 9(½), 11, 12

Example 8 Solving $ax^2 = d$

Solve $3x^2 = 18$

Solution

$3x^2 = 18$

$x^2 = 6$

$x = \sqrt{6}$ or $x = -\sqrt{6}$

Alternative method:

$3x^2 = 18$

$3x^2 - 18 = 0$

$3(x^2 - 6) = 0$

$3(x + \sqrt{6})(x - \sqrt{6}) = 0$

$x + \sqrt{6} = 0$ or $x - \sqrt{6} = 0$

$x = -\sqrt{6}$ or $x = \sqrt{6}$

Explanation

Divide both sides by 3 and then take the square root of each side.

As 6 is not a square number leave answers in exact form as $\sqrt{6}$ and $-\sqrt{6}$.

Alternatively, express in standard form and then note the common factor of 3.

Treat $x^2 - 6 = x^2 - \left(\sqrt{6}\right)^2$ as a difference of squares.

Apply the Null Factor Law and solve for x.

Now you try

Solve $2x^2 = 14$

9 Solve each of the following equations by first noting the common factor.

a $7x^2 = 28$ **b** $5x^2 = 45$

c $2x^2 = 50$ **d** $6x^2 = 24$

e $2x^2 = 12$ **f** $3x^2 = 15$

g $5x^2 - 35 = 0$ **h** $8x^2 - 24 = 0$



Hint for Q9: If $x^2 = 5$, $x = \sqrt{5}$ or $-\sqrt{5}$

10 Write an equation and solve it to find the number.

a The square of the number is 7 times the same number.

b The difference between the square of a number and 64 is zero.

c 3 times the square of a number is equal to -12 times the number.

11 Consider the equation $x^2 + 4 = 0$.

a Explain why it cannot be written in the form $(x + 2)(x - 2) = 0$.

b Are there any solutions to the equation $x^2 + 4 = 0$? Why/Why not?

12 Solve these equations by first rearranging where necessary.

a $4 = x^2$

b $-x^2 + 25 = 0$

c $-x^2 = -100$

d $-3x^2 = 21x$

e $-5x^2 + 35x = 0$

f $1 - x^2 = 0$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

11B

More quadratic equation forms — 13–14(½)

13 Remove brackets or fractions to help solve these equations.

a $x - \frac{4}{x} = 0$

b $\frac{3}{x^2} = 3$

c $5(x^2 + 1) = 3x^2 + 7$

d $x(x - 3) = 2x^2 + 4x$

14 Note, for example, that $4x^2 = 9$ becomes $x^2 = \frac{9}{4}$ and then $x = \frac{3}{2}$ or $x = -\frac{3}{2}$, since $\sqrt{\frac{9}{4}} = \frac{3}{2}$.

Now solve these equations.

a $9x^2 = 16$

b $25x^2 = 36$

c $4 = 100x^2$

d $81 - 25x^2 = 0$

Using a CAS calculator 11B: Solving quadratic equations

This activity is in the Interactive Textbook in the form of a printable PDF.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

11C Solving $x^2 + bx + c = 0$

Learning intentions

- To be able to solve quadratic equations involving monic trinomials
- To understand that trinomials that factorise to perfect squares will have only one solution

Key vocabulary: monic quadratic trinomial

Earlier in Chapter 9 we learnt to factorise quadratic trinomials with three terms. For example, $x^2 + 5x + 6$ factorises to $(x + 2)(x + 3)$. This means that the Null Factor Law can be used to solve equations of the form $x^2 + bx + c = 0$.



Mechanical engineers design industrial ventilation systems and can solve quadratic equations to calculate dimensions. For a duct with a rectangular cross-section area 1500 cm^2, a possible equation is: $x(x - 20) = 1500$, i.e. $x^2 - 20x - 1500 = 0$.

Lesson starter: Remembering how to factorise quadratic trinomials

First expand these quadratics using the distributive law.

Distributive law: $(a + b)(c + d) = ac + ad + bc + bd$

- $(x + 1)(x + 2)$
- $(x - 3)(x + 4)$
- $(x - 5)(x - 2)$

Now factorise these expressions.

- $x^2 + 5x + 6$
- $x^2 - x - 12$
- $x^2 - 8x + 7$

Discuss your method for finding the factors of each quadratic above.

Key ideas

- Solve quadratics of the form $x^2 + bx + c = 0$ by factorising the quadratic trinomial.
  - Ask 'What factors of c add to give b?'
  - Then use the Null Factor Law.

$x^2 - 3x - 28 = 0$ $\quad -7 \times 4 = -28$
$(x - 7)(x + 4) = 0$ and $-7 + 4 = -3$
$x - 7 = 0$ or $x + 4 = 0$
$x = 7$ or $x = -4$

- Equations involving perfect squares will give only one solution.

$x^2 - 6x + 9 = 0$
$(x - 3)(x - 3) = 0$
$x - 3 = 0$
$x = 3$

- $x^2 + bx + c$ is called a **monic** quadratic since the coefficient of x^2 is 1.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Exercise 11C

Understanding

1–3 | 1–3

1 Decide which two factors of the first number add to give the second number.

a 6, 5 **b** 10, −7 **c** −5, 4 **d** −12, −1

2 Factorise each of the following.

a $x^2 + 12x + 35$ **b** $x^2 + 4x - 45$ **c** $x^2 - 10x + 16$

3 State the missing components to complete the working to solve each equation.

a $x^2 + 9x + 20 = 0$
$(x + 5)(_____) = 0$
$x + 5 = 0$ or $___ = 0$
$x = ___$ or $x = ___$

b $x^2 - 2x - 24 = 0$
$(x - 6)(_____) = 0$
$x - 6 = 0$ or $___ = 0$
$x = ___$ or $x = ___$

Fluency

4–6(½) | 4–6(½)

Example 9 Solving equations involving quadratic trinomials with $b > 0$ and $c > 0$

Solve $x^2 + 7x + 12 = 0$.

Solution

$x^2 + 7x + 12 = 0$
$(x + 3)(x + 4) = 0$
$x + 3 = 0$ or $x + 4 = 0$
$x = -3$ or $x = -4$

Explanation

Factors of 12 which add to 7 are 3 and 4.
$3 \times 4 = 12, 3 + 4 = 7$
Use the Null Factor Law to solve the equation.

Now you try

Solve $x^2 + 6x + 8 = 0$.

4 Solve these quadratic equations.

a $x^2 + 8x + 12 = 0$ **b** $x^2 + 11x + 24 = 0$
c $x^2 + 7x + 10 = 0$ **d** $x^2 + 10x + 24 = 0$
e $x^2 + 9x + 18 = 0$ **f** $x^2 + 4x + 3 = 0$



Hint for Q4: First factorise by asking what factors of c add to give b

Example 10 Solving $x^2 + bx + c = 0$ if b and/or $c < 0$

Solve these quadratic equations.

a $x^2 - 2x - 8 = 0$ **b** $x^2 - 8x + 15 = 0$

Solution

a $x^2 - 2x - 8 = 0$
$(x - 4)(x + 2) = 0$
$x - 4 = 0$ or $x + 2 = 0$
$x = 4$ or $x = -2$

Explanation

Factors of −8 which add to −2 are −4 and 2.
$-4 \times 2 = -8, -4 + 2 = -2$
Solve using the Null Factor Law.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

b $x^2 - 8x + 15 = 0$

$(x - 5)(x - 3) = 0$

$x - 5 = 0$ or $x - 3 = 0$

$x = 5$ or $x = 3$

The factors of 15 must add to give −8.

$-5 \times (-3) = 15$ and $-5 + (-3) = -8$, so −5 and −3 are the two numbers.

Now you try

Solve the quadratic equations

a $x^2 - 3x - 10 = 0$ **b** $x^2 - 8x + 12 = 0$

5 Solve these quadratic equations.

a $x^2 - 12x + 32 = 0$ **b** $x^2 - 9x + 18 = 0$ **c** $x^2 - 10x + 21 = 0$

d $x^2 - 2x - 15 = 0$ **e** $x^2 - 6x - 16 = 0$ **f** $x^2 - 4x - 45 = 0$

g $x^2 - 10x + 24 = 0$ **h** $x^2 - x - 42 = 0$ **i** $x^2 + 5x - 84 = 0$

Example 11 Solving with perfect squares

Solve $x^2 - 8x + 16 = 0$.

Solution

$x^2 - 8x + 16 = 0$

$(x - 4)(x - 4) = 0$

$x - 4 = 0$

$x = 4$

Explanation

Factors of 16 which add to −8 are −4 and −4.

$(x - 4)(x - 4) = (x - 4)^2$ is a perfect square so there is only one solution.

Now you try

Solve $x^2 - 10x + 25 = 0$.

6 Solve these quadratic equations which include perfect squares.

a $x^2 + 6x + 9 = 0$ **b** $x^2 + 4x + 4 = 0$

c $x^2 + 14x + 49 = 0$ **d** $x^2 + 24x + 144 = 0$

e $x^2 - 10x + 25 = 0$ **f** $x^2 - 16x + 64 = 0$

g $x^2 - 12x + 36 = 0$ **h** $x^2 - 18x + 81 = 0$



Hint for Q6: Both factors should be the same.

Problem-solving and reasoning

7(½), 8, 9 | 7(½), 9–11

Example 12 Solving with trinomials not in standard form

Solve $x^2 = x + 6$.

Solution

$x^2 = x + 6$

$x^2 - x - 6 = 0$

$(x - 3)(x + 2) = 0$

$x - 3 = 0$ or $x + 2 = 0$

$x = 3$ or $x = -2$

Explanation

First make the right-hand side equal zero by subtracting x and 6 from both sides. This is now in standard form. (When rearranging to standard form, keep the x^2 term positive.)

Factors of −6 which add to −1 are −3 and 2.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

11C

Now you try

Solve $x^2 = 3x + 40$

7 Solve these quadratic equations by first rearranging to standard form.

a $x^2 = 3x + 10$ **b** $x^2 = 7x - 10$
c $x^2 = 6x - 9$ **d** $x^2 = 4 - 3x$
e $14 - 5x = x^2$ **f** $x^2 + 16 = 8x$
g $x^2 - 12 = -4x$ **h** $6 - x^2 = 5x$
i $15 = 8x - x^2$ **j** $16 - 6x = x^2$
k $-6x = x^2 + 8$ **l** $-x^2 - 7x = -18$



8 The temperature in °C inside a room at a scientific base in Antarctica after 10 a.m. is given by the expression $t^2 - 9t + 8$ where t is in hours. Find the times in the day when the temperature is 0°C.



9 Write down a quadratic equation in standard form which has the following solutions.

a $x = 1$ and $x = 2$ **b** $x = 3$ and $x = -2$ **c** $x = -4$ and $x = 1$
d $x = -3$ and $x = 10$ **e** $x = 5$ only **f** $x = -11$ only

10 Consider this equation and solution.

$$x^2 - x - 12 = -6$$
$$(x - 4)(x + 3) = -6$$
$$x - 4 = -6 \text{ or } x + 3 = -6$$
$$x = -2 \text{ or } x = -9$$

a Explain what is wrong with the solution.

b Find the correct solutions to the equation.

11 Explain why $x^2 - 2x + 1 = 0$ only has one solution.

Solving non-monics where a is a common a factor — 12(½)

12 Solve these equations by taking out a common factor as the first factorising step.

a $2x^2 - 2x - 12 = 0$ **b** $3x^2 + 24x + 45 = 0$ **c** $4x^2 - 24x - 64 = 0$
d $4x^2 - 20x + 24 = 0$ **e** $2x^2 - 8x + 8 = 0$ **f** $3x^2 + 6x + 3 = 0$
g $7x^2 - 70x + 175 = 0$ **h** $2x^2 + 12x = -18$ **i** $5x^2 = 35 - 30x$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

11D Using quadratic equations to solve problems

Learning intentions

- To be able to form an equation from a word problem using a variable for the unknown
- To be able to recognise a quadratic equation and apply the appropriate steps to solve
- To understand that solutions will need to be checked for validity in the context of the problem

Key vocabulary: variable, product

When using mathematics to solve problems, we often arrive at a quadratic equation. In solving the quadratic equation we obtain the solutions to the problem. Setting up an equation and then interpreting the solution are important parts of the problem-solving process.



When dividing up land for new suburbs, surveyors apply area formulas which form quadratic equations when dimensions are expressed using the same variable. Surveyors re-arrange the quadratic to equal zero and solve it for the unknown variable.

Lesson starter: Solving for the unknown number

The product of a positive number and 6 more than the same number is 16.

- Using x as the unknown number, write an equation describing the given condition.
- Solve your equation for x.
- Are both solutions feasible (allowed)?
- Discuss how this method compares with the method of trial and error.

Key ideas

- When using quadratic equations to solve problems, follow these steps.
  - Define your variable.
    - Write 'Let x be …' e.g. Let x be the width in cm.
  - Use your variable to define any other quantity, e.g. length = $x + 7$.
  - Write an equation relating the facts given in the question.
  - Solve your equation using the Null Factor Law.
  - Check that your solutions are feasible.
    - Some problems may not allow solutions that are negative numbers or fractions.
  - Answer the original question in words and check that your answer seems reasonable.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Exercise 11D


Understanding

1, 2 | 1, 2

1 Solve these equations for x by first expanding and producing a zero on the right-hand side.

a $x(x+3)=18$ **b** $x(x-1)=20$ **c** $(x-1)(x+4)=6$

2 This rectangle has an area of 8 cm^2 and a length that is 2 cm more than its width.



a Using length × width = area, give an equation.

b Solve your equation by expanding and subtracting 8 from both sides. Then use the Null Factor Law.

c Which of your two solutions is feasible for the width of the rectangle?

d State the dimensions (width and length) of the rectangle.

Fluency

3, 4, 6, 7 | 3, 5, 6, 7

Example 13 Solving a problem involving a product

The product of a number and one more than the same number is 56. Write an equation and solve to find the two possible solutions.

Solution	Explanation
Let x be the number	First define your variable.
$x(x+1)=56$	Set up your equation, expand and rearrange.
$x^2+x-56=0$	Factorise your monic quadratic, use the Null Factor Law to solve.
$(x-7)(x+8)=0$	
$x-7=0$ or $x+8=0$	
$x=7$ or $x=-8$	
The number could be 7 or −8	State your answer.

Now you try

The product of a number and 3 more than the same number is 40. Write an equation and solve to find the two possible solutions.

3 The product of a number and 2 more than the same number is 48. Write an equation and solve to find the two possible solutions.



Hint for Q3: The product of x and $x+2$ is $x(x+2)$

4 The product of a number and 7 less than the same number is 60. Write an equation and solve to find the two possible solutions.

5 The product of a number and 13 less than the same number is 30. Write an equation and solve to find the two possible solutions.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Example 14 Solving area problems with quadratic equations

The length of a book is 4 cm more than its width and the area of the face of the book is 320 cm^2. Find the dimensions of the face of the book.

Solution	Explanation
Let x cm be the width of the book face.	Define a variable for width then write the length in terms of the width.
Length $= (x + 4)$ cm	
Area:	
$x(x + 4) = 320$	Write an equation to suit the given situation. Expand, then subtract 320 from both sides.
$x^2 + 4x - 320 = 0$	
$(x + 20)(x - 16) = 0$	Factorise the quadratic trinomial.
$x + 20 = 0$ or $x - 16 = 0$	
$x = -20$ or $x = 16$	Solve using the Null Factor Law, but note that a width of -20 cm is not feasible.
$\therefore x = 16$ since $x > 0$	
$\therefore$ width = 16 cm and length = 20 cm	Finish by writing the dimensions – width and length – as required.
	Length $= x + 4 = 16 + 4 = 20$ cm

Now you try

The length of a rectangular jigsaw puzzle is 12 cm more than its width and the area of the jigsaw is 640 cm^2. Find the dimensions of the jigsaw puzzle.

6 The length of a rectangular magazine is 8 cm more than its width and the area of the magazine is 240 cm^2. Find the dimensions of the magazine.



Hint for Q6: the width is x cm then the length is $(x + 8)$ cm.

7 The length of a rectangular brochure is 5 cm more than its width and the area of the face of the brochure is 36 cm^2. Find the dimensions of the face of the brochure.

Problem-solving and reasoning

8, 9, 11 | 8–10, 12


8 The length of a small rectangular kindergarten play area is 20 metres less than its width and the area is 69 m^2. Find the dimensions of the play area.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


11D

9 An isosceles triangle has height (h) equal to half its base length. Find the value of h if the area is 25 square units.




10 Use Pythagoras' theorem to find the value of x in these right-angled triangles.


a



b




c





Hint for Q10: Recall $c^2 = a^2 + b^2$ so $5^2 = x^2 + (x+1)^2$ in part **a**.

11 The equation for the area of this rectangle is $x^2 + 2x - 48 = 0$, which has solutions $x = -8$ and $x = 6$. Which solution do you ignore and why?




12 The product of an integer and one less than the same integer is 6. The equation for this is $x^2 - x - 6 = 0$. How many different solutions are possible and why?

L-shape problems — 13, 14

13 A square of side length 10 metres has a square of side length x metres removed from one corner.

a Write an expression for the area remaining after the square of side length x metres is removed. (*Hint*: Use subtraction.)


b Find the value of x if the area remaining is 64 m^2.



14 A rectangular farm shed (3 m by 5 m) is to be extended to form an 'L' shape as shown.

a Write an expression for the total area of the extended farm shed.

b Find the value of x if the total area is to be 99 m^2.





ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Progress quiz

11A **1** Write these quadratic equations in standard form.

a $x^2 - 3x = 8$ **b** $3(x^2 - 2) = 4x$ **c** $2(x^2 - 5) = 2x - 7$

11A **2** Substitute the given x-value into the equation and say whether or not it is a solution.

a $x^2 - x - 6 = 0 \quad (x = 2)$ **b** $x^2 - x - 6 = 0 \quad (x = -2)$

11A **3** Use the Null Factor Law to solve these equations.

a $x(x - 7) = 0$ **b** $(x - 5)(x + 2) = 0$

c $(3x - 1)(4x - 3) = 0$ **d** $2(x - 4)(x - 3)$

11B **4** Solve the following equations.

a $x^2 + 11x = 0$ **b** $3x^2 - 18x = 0$

c $-2x^2 - 10x = 0$ **d** $5x^2 = 20x$

11B **5** Solve these equations.

a $x^2 = 49$ **b** $x^2 - 1 = 0$

c $3x^2 = 12$ **d** $7x^2 - 63 = 0$

11C **6** Solve these quadratic equations.

a $x^2 + 11x + 24 = 0$

b $x^2 - 15x + 36 = 0$

c $x^2 - 2x - 35 = 0$

11C **7** Solve these quadratic equations.

a $x^2 + 8x + 16 = 0$

b $x^2 - 14x + 49 = 0$

c $x^2 - 24 = 5x$

11D **8** A rectangular lawn is 3 m longer than it is wide and has an area of 54 m^2. Write an equation and solve it to find the dimensions of this lawn.



11D **9** The product of a number and 5 more than the same number is 84. Write an equation and solve to find the two possible values of the number.

11C **10** Write down a quadratic equation in standard form which has the following solutions.

a $x = 3$ and $x = 2$

b $x = 6$ and $x = -1$

c $x = -3$ only

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


11E The parabola

Learning intentions

- To know that a quadratic relation produces a graph called a parabola
- To know the key features of a parabola including symmetry and the turning point
- To understand that a parabola can have a minimum or maximum turning point
- To be able to identify the key features of a parabola from a graph
- To be able to plot a parabola from a rule by creating a table of values and plotting points

Key vocabulary: Parabola, minimum, maximum, axis of symmetry, turning point

Relations that have rules in which the highest power of x is 2, such as $y = x^2$, $y = 2x^2 - 3$ and $y = 3x^2 + 2x - 4$, are called quadratics and their graphs are called parabolas. Parabolic shapes can be seen in many modern-day objects or situations such as the arches of bridges, the paths of projectiles and the surfaces of reflectors.



When hitting a ball, tennis players, cricketers and golfers can vary the force and launching angle, hence modifying the direction, maximum height and range of the ball's parabolic flight path.


Lesson starter: Finding features

A quadratic is given by the equation $y = x^2 - 2x - 3$. Complete these tasks to discover its graphical features.

- Use the rule to complete this table of values.

x	-2	-1	0	1	2	3	4
y							


- Plot your points on a copy of the axes shown at right and join them to form a smooth curve.
- Describe these features:
  - minimum turning point
  - axis of symmetry
  - coordinates of the y-intercept
  - coordinates of the x-intercepts




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Key ideas


- The graph of a quadratic relation is called a **parabola**. Its basic shape is shown on the right.
  - The basic quadratic rule is $y = x^2$.



- The general equation of a quadratic is $y = ax^2 + bx + c$.
- A parabola is symmetrical about a line called the **axis of symmetry** and it has one **turning point** (a vertex), which may be a **maximum** or a **minimum**.



Here is an example of a parabola with equation $y = x^2 - 4x + 3$, showing all the key features.



Exercise 11E

Understanding

1, 2 | 1, 2

1 Choose a word from this list to complete each sentence.
lowest, parabola, vertex, highest, intercepts, zero

- **a** A maximum turning point is the _____ point on the graph.
- **b** The graph of a quadratic is called a _____.
- **c** The x-_____ are the points where the graph cuts the x-axis.
- **d** The axis of symmetry is a vertical line passing through the _____.
- **e** A minimum turning point is the _____ point on the graph.
- **f** The y-intercept is at x equals _____.

2 State the equation of a vertical line (e.g. $x = 2$) that passes through these points.

- **a** $(3, 0)$
- **b** $(1, 5)$
- **c** $(-2, 4)$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

11E


Fluency

3(½), 4, 5 | 3(½), 4, 6

Example 15 Identifying the features of a parabola

For this graph state the:

a equation of the axis of symmetry
b type of turning point
c coordinates of the turning point
d coordinates of the x-intercepts
e coordinates of the y-intercept.




Solution

a $x = -3$
b Maximum turning point
c Turning point is $(-3, 5)$
d x-intercepts: $(-5, 0)$ and $(-1, 0)$
e y-intercept: $(0, -5)$

Explanation

Graph is symmetrical about the vertical line $x = -3$. Graph has its highest y-coordinate at the turning point, so it is a maximum point.

Highest point: turning point = $(-3, 5)$




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Now you try

For this graph state the:

- **a** equation of the axis of symmetry
- **b** type of turning point
- **c** coordinates of the turning point
- **d** coordinates of the x-intercepts
- **e** coordinates of the y-intercept.




3 For each of the following graphs, state:

- **i** the equation of the axis of symmetry
- **ii** the type of turning point (maximum or minimum)
- **iii** the coordinates of the turning point
- **iv** the coordinates of the x-intercepts
- **v** the coordinates of the y-intercept.


Hint for Q3: The rule for the equation of the axis of symmetry is $x = h$ where h is the x-value of the turning point.




a




b




c




d



e




f




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

11E


g




h




i




j



k



l





The cross-section of a satellite dish is parabolic.

Example 16 Plotting a parabola

Use the quadratic rule $y = x^2 - 4$ to complete these tasks.

a Complete this table of values.

x	−3	−2	−1	0	1	2	3
y							

b Draw a set of axes using a scale that suits the numbers in your table. Then plot the points to form a parabola.

c State these features.

- **i** Type of turning point
- **ii** Axis of symmetry
- **iii** Coordinates of the turning point
- **iv** The coordinates of the y-intercept
- **v** The coordinates of the x-intercepts.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Solution


a

x	-3	-2	-1	0	1	2	3
y	5	0	-3	-4	-3	0	5

Explanation

Substitute each x-value into the rule to find each y-value:
e.g. $x = -3$, $y = (-3)^2 - 4$
$= 9 - 4$
$= 5$

b



Plot each coordinate pair and join to form a smooth curve.

c **i** Minimum turning point
ii $x = 0$ is the axis of symmetry
iii Turning point $(0, -4)$
iv y-intercept: $(0, -4)$
v x-intercepts: $(-2, 0)$ and $(2, 0)$

The turning point at $(0, -4)$ has the lowest y-coordinate for the entire graph. The vertical line $x = 0$ divides the graph like a mirror line. The y-intercept is at $x = 0$. The x-intercepts are at $y = 0$ on the x-axis.

Now you try

Use the quadratic rule $y = x^2 - 9$ to complete these tasks.

a Complete this table of values.

x	-3	-2	-1	0	1	2	3
y							

b Draw a set of axes using a scale that suits the numbers in your table. Then plot the points to form a parabola.

c State these features.
i Type of turning point
ii Axis of symmetry
iii Coordinates of the turning point
iv The coordinates of the y-intercept
v The coordinates of the x-intercepts.

4 Use the quadratic rule $y = x^2 - 1$ to complete these tasks.

a Complete the table of values.

x	-2	-1	0	1	2
y					

b Draw a set of axes using a scale that suits the numbers in your table. Then plot the points to form a parabola.

c State these features.
i Type of turning point
ii Axis of symmetry
iii Coordinates of the turning point
iv The coordinates of the y-intercept
v The coordinates of the x-intercepts

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

11E

5 Use the quadratic rule $y = x^2 + 2x - 3$ to complete these tasks.

a Complete the table of values.

x	-4	-3	-2	-1	0	1	2
y							

b Draw a set of axes using a scale that suits the numbers in your table. Then plot the points to form a parabola.

c State these features.

i Type of turning point
ii Axis of symmetry
iii Coordinates of the turning point
iv The coordinates of the y-intercept
v The coordinates of the x-intercepts

6 Use the quadratic rule $y = 9 - x^2$ to complete these tasks.

a Complete the table of values.

x	-3	-2	-1	0	1	2	3
y							

b Draw a set of axes using a scale that suits the numbers in your table. Then plot the points to form a parabola.


c State these features.

i Type of turning point
ii Axis of symmetry
iii Coordinates of the turning point
iv The coordinates of the y-intercept
v The coordinates of the x-intercepts

Problem-solving and reasoning

7–9 | 8–12


7 This graph shows the height of a cricket ball, y metres, as a function of time t seconds.



a **i** At what times is the ball at a height of 9 m?
ii Why are there two different times?

b **i** At what time is the ball at its greatest height?
ii What is the greatest height the ball reaches?
iii After how many seconds does it hit the ground?

8 The graph gives the height, h m, at time t seconds, of a rocket which is fired up in the air.



a From what height is the rocket launched?

b What is the approximate maximum height that the rocket reaches?

c For how long is the rocket in the air?

d What is the difference in time for when the rocket is going up and when it is going down?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


9 A parabola has x-intercepts at $(-2, 0)$ and $(4, 0)$. The y-coordinate of the turning point is -3.

a What is the equation of its axis of symmetry?


b What are the coordinates of the turning point?

10 Write the rule for these parabolas. Use trial and error to help and check your rule by substituting a known point.


a




b




c




d



e



f



11 Is it possible to draw a parabola with the following properties? If yes, draw an example.

a Two x-intercepts

b One x-intercept

c No x-intercepts

d No y-intercept

12 **a** Mal calculates the y-value for $x = 2$ using $y = -x^2 + 2x$ and gets $y = 8$. Explain his error.

b Mai calculates the y-value for $x = -3$ using $y = x - x^2$ and gets $y = 6$. Explain her error.




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

11E

Solving quadratic equations with parabolas — 13, 14

13 This graph shows the parabola for $y = x^2 - 4$.



- **a** For what values of x is $y = 0$?
- **b** For what value of x is $y = -4$?
- **c** How many values of x give a y-value which is:
  - **i** greater than -4?
  - **ii** equal to -4?
  - **iii** less than -4?

14 This table corresponds to the rule $y = x^2 - 2x$.

x	-1	0	1	2	3	4
y	3	0	-1	0	3	8

- **a** Use this table to solve these equations
  - **i** $0 = x^2 - 2x$ **ii** $3 = x^2 - 2x$
- **b** How many solutions would there be to the equation $8 = x^2 - 2x$? Why?
- **c** How many solutions would there be to the equation $-1 = x^2 - 2x$? Why?
- **d** How many solutions would there be to the equation $-2 = x^2 - 2x$? Why?

Using a CAS calculator 11E: Sketching parabolas
This activity is in the Interactive Textbook in the form of a printable PDF.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

11F Sketching $y = ax^2$ with dilations and reflections

Learning intentions

- To understand the effect of the value and sign of a in the rule $y = ax^2$ on the shape of the graph and key features
- To know which values of a make the parabola appear narrower or wider than $y = x^2$
- To know which values of a cause the parabola to be upright or inverted

Key vocabulary: dilation, reflection, upright (concave up), inverted (concave down).

In geometry we know that shapes can be transformed by applying reflections, rotations, translations and dilations (enlargement). The same types of transformations can also be applied to graphs, including parabolas. Altering the value of a in $y = ax^2$ causes both dilations and reflections.




Suspension bridges have supporting cables that form parabolic curves, such as those on this New York bridge. Engineers model such parabolas with quadratic equations that are then used to calculate the vertical support heights.

Lesson starter: What is the effect of a?

This table and graph show a number of examples of $y = ax^2$ with varying values of a. They could also be produced using technology.

x	-2	-1	0	1	2
$y = x^2$	4	1	0	1	4
$y = 2x^2$	8	2	0	2	8
$y = \frac{1}{2}x^2$	2	$\frac{1}{2}$	0	$\frac{1}{2}$	2
$y = x^2$	-4	-1	0	-1	-4



- Discuss how the different values of a affect the y-values in the table.
- Discuss how the different values of a affect the shape of the graph.
- How would the graphs of the following rules compare to the graphs shown above?


a $y = 3x^2$ **b** $y = \frac{1}{4}x^2$ **c** $y = -\frac{1}{2}x^2$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

11F

Key ideas


- The most basic parabola is the graph of $y = x^2$. It can be graphed quickly using the five key points shown.




  If you know these five points, you can use them to draw many other parabolas.

- The equation $y = ax^2$ describes a family of parabolas including $y = x^2$, $y = -x^2$, $y = 2x^2$, $y = -2x^2$ etc. They contain the following features:
  - the vertex (or turning point) is $(0, 0)$
  - the axis of symmetry is $x = 0$
  - if $a > 0$ the graph is **upright** (or **concave up**) and has a minimum turning point
  - if $a < 0$ the graph is **inverted** (or **concave down**) and has a maximum turning point.

- For $y = ax^2$, if $a > 1$ or $a < -1$:
  the graph appears narrower than the graph of either $y = x^2$ or $y = -x^2$. For example: $y = 2x^2$ or $y = -2x^2$



- For $y = ax^2$, if $-1 < a < 1$:
  the graph appears wider than the graph of either $y = x^2$ or $y = -x^2$. For example: $y = \frac{1}{2}x^2$ or $y = -\frac{1}{2}x^2$



- For $y = 2x^2$ we say that the graph of $y = x^2$ is **dilation** from the x-axis by a factor of 2.
- For $y = -x^2$ we say that the graph of $y = x^2$ is **reflection** in the x-axis.


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Exercise 11F

Understanding

1, 2 | 1, 2

1 Shown here are the graphs of $y = x^2$, $y = 3x^2$, $y = \frac{1}{2}x^2$, $y = -x^2$, $y = -3x^2$ and $y = -\frac{1}{2}x^2$.



- **a** State the rules of the three graphs which have a minimum turning point.
- **b** State the rules of the three graphs which have a maximum turning point.
- **c** What are the coordinates of the turning point for all the graphs?
- **d** What is the equation of the axis of symmetry for all the graphs?
- **e** State the rule of the graph which is:
  - **i** upright (concave up) and narrower than $y = x^2$
  - **ii** upright (concave up) and wider than $y = x^2$
  - **iii** inverted (concave down) and narrower than $y = -x^2$
  - **iv** inverted (concave down) and wider than $y = -x^2$.
- **f** State the rule of the graph which is:
  - **i** a reflection of $y = x^2$ in the x-axis
  - **ii** a reflection of $y = 3x^2$ in the x-axis
  - **iii** a reflection of $y = -\frac{1}{2}x^2$ in the x-axis.

2 Select the word *positive* or *negative* to suit each sentence.

- **a** The graph of $y = ax^2$ will be upright (concave up) with a minimum turning point if a is _____.
- **b** The graph of $y = ax^2$ will be inverted (concave down) with a maximum turning point if a is _____.

Fluency

3–5 | 3–5

Example 17 Comparing graphs of $y = ax^2$, $a > 0$

Complete the following for $y = x^2$, $y = 2x^2$ and $y = \frac{1}{2}x^2$.

- **a** Draw up and complete a table of values for $-2 \leqslant x \leqslant 2$.
- **b** Plot their graphs on the same set of axes.
- **c** Write down the equation of the axis of symmetry and the coordinates of the turning point.
- **d** **i** Does the graph of $y = 2x^2$ appear wider or narrower than the graph of $y = x^2$?
  - **ii** Does the graph of $y = \frac{1}{2}x^2$ appear wider or narrower than the graph of $y = x^2$?

Continued on next page

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

11F

Solution

a $y = x^2$

x	-2	-1	0	1	2
y	4	1	0	1	4

$y = 2x^2$

x	-2	-1	0	1	2
y	8	2	0	2	8

$y = \frac{1}{2}x^2$

x	-2	-1	0	1	2
y	2	$\frac{1}{2}$	0	$\frac{1}{2}$	2

Explanation

Substitute each x-value into $y = x^2$, $y = 2x^2$ and $y = \frac{1}{2}x^2$.


e.g. for $y = 2x^2$, if $x = 2$,

$$\begin{aligned} y &= 2(2)^2 \\ &= 2(4) \\ &= 8 \end{aligned}$$

If $x = -1$,

$$\begin{aligned} y &= 2(-1)^2 \\ &= 2(1) \\ &= 2 \end{aligned}$$

b



Plot the points for each graph using the coordinates from the tables and join them with a smooth curve.

c Axis of symmetry: y-axis $(x = 0)$

Turning point: minimum at $(0, 0)$

Look at graphs to see symmetry about the y-axis and a minimum turning point at the origin.

d i The graph of $y = 2x^2$ appears narrower than the graph of $y = x^2$.

For each value of x, $2x^2$ is twice that of x^2; hence, the graph (y-values) of $y = 2x^2$ rises more quickly.

ii The graph of $y = \frac{1}{2}x^2$ appears wider than the graph of $y = x^2$.

For each value of x, $\frac{1}{2}x^2$ is half that of x^2; hence, the graph of $y = \frac{1}{2}x^2$ rises more slowly.

Now you try

Complete the following for $y = x^2$, $y = 4x^2$ and $y = \frac{1}{4}x^2$.

a Draw up and complete a table of values for $-2 \leqslant x \leqslant 2$.

b Plot their graphs on the same set of axes.

c Write down the equation of the axis of symmetry and the coordinates of the turning point.

d i Does the graph of $y = 4x^2$ appear wider or narrower than the graph of $y = x^2$?

ii Does the graph of $y = \frac{1}{4}x^2$ appear wider or narrower than the graph of $y = x^2$?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3 Complete the following for $y = x^2$, $y = 3x^2$ and $y = \frac{1}{3}x^2$.

- **a** Draw up and complete a table of values for $-2 \leqslant x \leqslant 2$.
- **b** Plot their graphs on the same set of axes.
- **c** Write down the equation of the axis of symmetry and the coordinates of the turning point.
- **d** **i** Does the graph of $y = 3x^2$ appear wider or narrower than the graph of $y = x^2$?
  - **ii** Does the graph of $y = \frac{1}{3}x^2$ appear wider or narrower than the graph of $y = x^2$?

4 For the equations given below, complete these tasks.

- **i** Draw up and complete a table of values for $-2 \leqslant x \leqslant 2$.
- **ii** Plot the graphs of the equations on the same set of axes.
- **iii** Write down the coordinates of the turning point and the equation of the axis of symmetry.
- **iv** Determine whether the graphs of the equations each appear wider or narrower than the graph of $y = x^2$.

a $y = 6x^2$ **b** $y = 5x^2$ **c** $y = \frac{1}{6}x^2$ **d** $y = \frac{1}{5}x^2$

Example 18 Comparing graphs of $y = ax^2$, $a < 0$

Complete the following for $y = -x^2$, $y = -3x^2$ and $y = -\frac{1}{2}x^2$.

- **a** Draw up and complete a table of values for $-2 \leqslant x \leqslant 2$.
- **b** Plot their graphs on the same set of axes.
- **c** Write down the the equation of the axis of symmetry and the coordinates of the turning point.
- **d** **i** Does the graph of $y = -3x^2$ appear wider or narrower than the graph of $y = -x^2$?
  - **ii** Does the graph of $y = -\frac{1}{2}x^2$ appear wider or narrower than the graph of $y = -x^2$?

Solution

a $y = -x^2$

x	-2	-1	0	1	2
y	-4	-1	0	-1	-4

$y = -3x^2$

x	-2	-1	0	1	2
y	-12	-3	0	-3	-12

$y = -\frac{1}{2}x^2$

x	-2	-1	0	1	2
y	-2	$-\frac{1}{2}$	0	$-\frac{1}{2}$	-2

Explanation

Substitute each x-value into $y = -x^2$, $y = -3x^2$ and $y = -\frac{1}{2}x^2$.

e.g. for $y = -3x^2$, if $x = 2$,

$$\begin{aligned} y &= -3(2)^2 \\ &= -3(4) \\ &= -12 \end{aligned}$$

If $x = -1$,


$$\begin{aligned} y &= -3(-1)^2 \\ &= -3(1) \\ &= -3 \end{aligned}$$

Continued on next page

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

11F

b



Plot the coordinates for each graph from the tables and join them with a smooth curve.

c Axis of symmetry: y-axis $(x = 0)$

Turning point: maximum at $(0, 0)$

Graphs are symmetrical about the y-axis with a maximum turning point at the origin.

d **i** The graph of $y = -3x^2$ appears narrower than the graph of $y = -x^2$.

ii The graph of $y = -\frac{1}{2}x^2$ appears wider than the graph of $y = -x^2$.

For each value of x, $-3x^2$ is three times that of $-x^2$; hence, the graph of $y = -3x^2$ gets larger in the negative direction more quickly.

For each value of x, $-\frac{1}{2}x^2$ is half that of $-x^2$.

Now you try

Complete the following for $y = -x^2$, $y = -4x^2$ and $y = -\frac{1}{4}x^2$.

i Draw up and complete a table of values for $-2 \leqslant x \leqslant 2$.

ii Plot their graphs on the same set of axes.

iii Write down the equation of the axis of symmetry and the coordinates of the turning point.

iv **a** Does the graph of $y = -4x^2$ appear wider or narrower than the graph of $y = -x^2$?

b Does the graph of $y = -\frac{1}{4}x^2$ appear wider or narrower than the graph of $y = -x^2$?

5 For the equations given below, complete these tasks.

i Draw up and complete a table of values for $-2 \leqslant x \leqslant 2$.

ii Plot the graphs of the equations on the same set of axes.

iii List the key features for each graph, such as the axis of symmetry, turning point, x-intercept and y-intercept.

iv Determine whether the graphs of the equations each appear wider or narrower than the graph of $y = -x^2$.



a $y = -2x^2$ **b** $y = -3x^2$ **c** $y = -\frac{1}{2}x^2$ **d** $y = -\frac{1}{3}x^2$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Problem-solving and reasoning 6–8 6–9

6 Here are eight quadratics of the form $y = ax^2$.

A $y = 6x^2$ **B** $y = -7x^2$ **C** $y = 4x^2$ **D** $y = \frac{1}{9}x^2$

E $y = \frac{x^2}{7}$ **F** $y = 0.3x^2$ **G** $y = -4.8x^2$ **H** $y = -0.5x^2$


a Which rule would give a graph which is upright (concave up) and the narrowest?

b Which rule would give a graph which is inverted (concave down) and the widest?


7 Match each of the following parabolas with the appropriate equation from the list below. Do a mental check by substituting the coordinates of a known point.

a $y = 3x^2$ **b** $y = -x^2$ **c** $y = 5x^2$ **d** $y = \frac{1}{2}x^2$ **e** $y = -5x^2$ **f** $y = 2x^2$


A




B




C




D



E



F



8 The graph of $y = ax^2$ is reflected in the x-axis and dilated from the x-axis by a given factor. Does it matter which transformation is completed first? Explain.

9 The rule of the form $y = ax^2$ models the shape of the parabolic cables of the Golden Gate Bridge. If the cable is centred at $(0, 0)$ and the top of the right pylon has the coordinates $(492, 67)$, find a possible equation that describes this shape. The numbers given are in metres.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


11F

Describing dilations and reflections — 10, 11

10 The graph of $y = -2x^2$ can be obtained from $y = x^2$ by conducting these transformations:

- reflection in the x-axis
- dilation by a factor of 2 from the x-axis.

In the same way as above, describe the two transformations which take:



a $y = x^2$ to $y = -3x^2$

b $y = x^2$ to $y = -6x^2$

c $y = x^2$ to $y = -\frac{1}{2}x^2$

d $y = -x^2$ to $y = 2x^2$

e $y = -x^2$ to $y = 3x^2$

f $y = -x^2$ to $y = \frac{1}{3}x^2$

11 Write the rule for the graph after each set of transformations.

a The graph of $y = x^2$ is reflected in the x-axis then dilated by a factor of 4 from the x-axis.

b The graph of $y = -x^2$ is reflected in the x-axis then dilated by a factor of $\frac{1}{3}$ from the x-axis.

c The graph of $y = 2x^2$ is reflected in the x-axis then dilated by a factor of 2 from the x-axis.

d The graph of $y = \frac{1}{3}x^2$ is reflected in the x-axis then dilated by a factor of 4 from the x-axis.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

11G Sketching translations of $y = x^2$

Learning intentions

- To understand what it means for a graph to be translated
- To know what causes a graph to be translated vertically and horizontally
- To understand the turning point form of a quadratic equation
- To be able to sketch a graph of a quadratic equation expressed in turning point form, labelling the turning point and y-intercept

Key vocabulary: translation, turning point

As well as reflection and dilation, there is a third type of transformation called translation. Translation involves a shift of every point on the graph horizontally and/or vertically. Unlike reflections and dilations, a translation alters the coordinates of the turning point. The shape of the curve is unchanged, but its position in relation to the origin is altered.




Manufacturers employ mathematicians to develop profit equations, which are quadratics when sales and profit per item are linear relations of the selling price. As the selling price increases, profit increases to the parabola's maximum point and then decreases.

Lesson starter: Which way: Left, right, up or down?

This table and graph shows the quadratics $y = x^2$, $y = (x-2)^2$, $y = (x+1)^2$, $y = x^2 - 4$ and $y = x^2 + 2$. The table could also be produced using technology.

x	-3	-2	-1	0	1	2	3
$y = x^2$	9	4	1	0	1	4	9
$y = (x-2)^2$	25	16	9	4	1	0	1
$y = (x+1)^2$	4	1	0	1	4	9	16
$y = x^2 - 4$	5	0	-3	-4	-3	0	5
$y = x^2 + 2$	11	6	3	2	3	6	11

- Discuss what effect the different numbers in the rules had on the y-values in the table.
- Also discuss what effect the numbers in the rules have on each graph. How are the coordinates of the turning point changed?
- What conclusions could you draw on the effect of h in the rule $y = (x-h)^2$?
- What conclusions could you draw on the effect of k in the rule $y = x^2 + k$?
- What if the rule was $y = -x^2 + 2$ or $y = -(x+1)^2$? Describe how the graphs would look.




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

11G


Key ideas


- A **translation** of a graph involves a shift of every point horizontally and/or vertically.

- Vertical translations: $y = x^2 + k$
  - If $k > 0$, the graph is translated k units up (red curve).
  - If $k < 0$, the graph is translated k units down (green curve).
  - The turning point is $(0, k)$ for all curves.
  - The axis of symmetry is the line $x = 0$ for all curves.
  - The y-intercept is $(0, k)$ for all curves.



- Horizontal translations: $y = (x - h)^2$
  - If $h > 0$, the graph is translated h units to the right.
  - If $h < 0$, the graph is translated h units to the left.





  - The turning point is $(h, 0)$ in both cases.
  - The axis of symmetry is the line $x = h$ in both cases.
  - The y-intercept is $(0, h^2)$ in both cases.

- The **turning point form** of a quadratic is given by:

$y = a(x - h)^2 + k$

$a > 0$ upright (concave up) graph ∪

$a < 0$ graph inverted (concave down) ∩

Translates the graph up or down


$k > 0$ ↑ $k < 0$ ↓

Translates the graph left or right:


$h > 0$ →

$h < 0$ ←

Axis of symmetry: $x = h$



- To sketch a graph of a quadratic equation in turning point form, follow these steps.
  - Draw and label a set of axes.
  - Identify important points including the turning point and y-intercept.
  - Sketch the curve connecting the key points and making the curve symmetrical.




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Exercise 11G

Understanding

1–3 | 1–3

1 This diagram shows the graphs of $y = x^2$, $y = -x^2$, $y = (x+2)^2$ and $y = -(x-3)^2$.




a State the coordinates of the turning point of the graph of:
i $y = (x+2)^2$ **ii** $y = -(x-3)^2$

b Compared to the graph of $y = x^2$, which way has the graph of $y = (x+2)^2$ been translated (left or right)?

c Compared to the graph of $y = -x^2$ which way has the graph of $y = -(x-3)^2$ been translated (left or right)?

2 This diagram shows the graphs of $y = x^2$, $y = -x^2$, $y = x^2 - 3$ and $y = -x^2 + 2$.



a State the coordinates of the turning point of the graph of:
i $y = x^2 - 3$ **ii** $y = -x^2 + 2$

b Compared to the graph of $y = x^2$, which way has the graph of $y = x^2 - 3$ been translated (up or down)?

c Compared to the graph of $y = -x^2$, which way has the graph of $y = -x^2 + 2$ been translated (up or down)?

3 Substitute $x = 0$ to find the coordinates of the y-intercept for these rules.

a $y = x^2 + 3$ **b** $y = -x^2 - 4$

c $y = -(x-2)^2$ **d** $y = (x+5)^2$

Fluency

4–5(½), 6 | 4–5(½), 6

Example 19 Sketching with vertical translations

Sketch the graphs of these rules showing the y-intercept and the coordinates of the turning point.

a $y = x^2 + 2$ **b** $y = -x^2 - 1$


Solution

a $y = x^2 + 2$

Turning point is $(0, 2)$.

y-intercept $(x = 0)$: $y = (0)^2 + 2 = 2$

$\therefore (0, 2)$



Explanation

For $y = x^2 + k$, $k = 2$, so the graph of $y = x^2$ is translated 2 units up.

The point $(0, 0)$ shifts to $(0, 2)$.

Continued on next page

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


11G

b $y = -x^2 - 1$

Turning point is $(0, -1)$.

y-intercept $(x = 0)$: $y = -(0)^2 - 1 = -1$

$\therefore (0, -1)$



The graph of $y = -x^2$ is a reflection of the graph of $y = x^2$ in the x-axis.

For $y = -x^2 + k$, $k = -1$ so the graph of $y = -x^2$ is translated down 1 unit.

Now you try

Sketch the graphs of these rules showing the y-intercept and the coordinates of the turning point.

a $y = x^2 + 4$ **b** $y = -x^2 - 2$

4 Sketch the graphs of these rules showing the y-intercept and the coordinates of the turning point.

a $y = x^2 + 1$
b $y = x^2 + 3$
c $y = x^2 - 2$
d $y = -x^2 + 4$
e $y = -x^2 + 1$
f $y = -x^2 - 5$



Hint for Q4: These involve vertical translations.

Example 20 Sketching with horizontal translations

Sketch the graphs of these rules showing the y-intercept and the coordinates of the turning point.

a $y = (x - 3)^2$
b $y = -(x + 2)^2$

Solution

a $y = (x - 3)^2$


Turning point is $(3, 0)$.

y-intercept $(x = 0)$: $y = (0 - 3)^2$

$= (-3)^2$

$= 9$

$\therefore (0, 9)$



Explanation

For $y = (x - h)^2$, $h = 3$, so the graph of $y = x^2$ is translated 3 units to the right.

The y-intercept is found by substituting $x = 0$ into the rule.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

b $y = -(x+2)^2$


Turning point is $(-2, 0)$.

y-intercept $(x = 0)$: $y = -(0+2)^2$

$= -(2)^2$

$= -4$

$\therefore (0, -4)$



For $y = -(x-h)^2$, $h = -2$ since $-(x-(-2))^2 = -(x+2)^2$. So the graph of $y = -x^2$ is translated 2 units to the left.

The y-intercept is found by substituting $x = 0$.

The negative sign in front means the graph is inverted.

Now you try

Sketch the graphs of these rules showing the y-intercept and the coordinates of the turning point.

a $y = (x-1)^2$


b $y = -(x+3)^2$

5 Sketch the graphs of these rules showing the y-intercept and the coordinates of the turning point.

a $y = (x-2)^2$ **b** $y = (x-4)^2$

c $y = (x+3)^2$ **d** $y = -(x-3)^2$

e $y = -(x+6)^2$ **f** $y = -(x+2)^2$



6 The below diagram shows the graphs of $y = x^2$, $y = -x^2$, $y = -(x-2)^2 + 3$ and $y = (x+1)^2 - 1$.

a State the coordinates of the turning point of the graph of:


i $y = -(x-2)^2 + 3$

ii $y = (x+1)^2 - 1$

b State the missing words and numbers.

i Compared to the graph of $y = x^2$, the graph of $y = (x+1)^2 - 1$ has to be translated _____ unit to the _____ and _____ unit _____.

ii Compared to the graph of $y = -x^2$, the graph of $y = -(x-2)^2 + 3$ has to be translated _____ units to the _____ and _____ units _____.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

11G

Problem-solving and reasoning

7(½), 8 | 7(½), 8–10

Example 21 Sketching with combined translations

Sketch these graphs showing the y-intercept and the coordinates of the turning point.

a $y = (x - 2)^2 - 1$

b $y = -(x + 3)^2 + 2$

Solution

a $y = (x - 2)^2 - 1$


Turning point is $(2, -1)$.

y-intercept $(x = 0)$: $y = (0 - 2)^2 - 1$

$= 4 - 1$

$= 3$

$\therefore (0, 3)$



Explanation

For $y = (x - h)^2 + k$, $h = 2$ and $k = -1$, so the graph of $y = x^2$ is shifted 2 to the right and 1 down.

Substitute $x = 0$ for the y-intercept.

First, position the coordinates of the turning point and y-intercept, then join to form the curve.

b $y = -(x + 3)^2 + 2$


Turning point is $(-3, 2)$.

y-intercept $(x = 0)$: $y = -(0 + 3)^2 + 2$

$= -9 + 2$

$= -7$

$\therefore (0, -7)$



For $y = -(x - h)^2 + k$, $h = -3$ and $k = 2$, so the graph of $y = x^2$ is shifted 3 to the left and 2 up.

First, position the coordinates of the turning point and y-intercept then join to form the curve.

Now you try

Sketch these graphs showing the y-intercept and the coordinates of the turning point.

a $y = (x - 3)^2 - 2$

b $y = -(x + 2)^2 + 1$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

7 Sketch each graph showing the y-intercept and the coordinates of the turning point.

a $y = (x-3)^2 + 2$
b $y = (x-1)^2 - 1$
c $y = (x+2)^2 - 3$
d $y = (x+1)^2 + 7$
e $y = -(x-2)^2 + 1$
f $y = -(x-5)^2 + 3$
g $y = -(x+3)^2 - 4$
h $y = -(x+1)^2 - 5$
i $y = -(x-3)^2 - 6$




Hint for Q7: For $y = (x-h)^2 + k$ then the turning point is (h, k)


8 Match each parabola with the appropriate equation from the list below.

a $y = x^2$
b $y = (x+2)^2$
c $y = x^2 - 4$
d $y = -x^2 - 3$
e $y = (x-2)^2$
f $y = x^2 + 4$
g $y = 4 - x^2$
h $y = (x+3)^2$
i $y = (x-5)^2$
j $y = x^2 + 3$
k $y = -(x+3)^2$
l $y = -x^2 + 3$


A




B




C




D




E




F




G




H




I




J



K



L




ISBN 978-1-009-59315-1
© Greenwood et al. 2025
Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


11G

9 Write the rule for each graph in the turning point form: $y = (x - h)^2 + k$.


a




b




c




d



e



f



10 A quadratic has the rule $y = a(x - h)^2 + k$.

a What are the coordinates of the turning point?

b Write an expression for the y-coordinate of the y-intercept.

The hike track — 11

11 A bike track can be modelled approximately by combining two different quadratic equations.
The first part of the bike path can be modelled by the equation $y = -(x - 2)^2 + 9$ for $-2 \leqslant x \leqslant 5$.
The second part of the bike track can be modelled by the equation $y = (x - 7)^2 - 4$ for $5 \leqslant x \leqslant 10$.

a Find the turning point of the graph of each quadratic equation.

b Sketch each graph on the same set of axes. On your sketch of the bike path you need to show the coordinates of the start and finish of the track and where it crosses the x-axis.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

11H Sketching parabolas using intercept form

Learning Intentions

- To understand how a parabola can have 0, 1 or 2 x-intercepts
- To be able to find the x-intercepts of a quadratic graph by solving an equation using the Null Factor Law
- To know how to use symmetry and the x-intercepts to help locate the turning point
- To be able to sketch a quadratic graph in expanded form using factorisation

Key vocabulary: intercept form, Null Factor Law

So far we have sketched parabolas using rules of the form $y = a(x - h)^2 + k$ where the coordinates of the turning point can be determined directly from the rule. An alternative method for sketching parabolas uses the factorised form of the quadratic rule and the Null Factor Law to find the x-intercepts. The turning point can be found by considering the axis of symmetry, which sits halfway between the two x-intercepts.




Parallel lines reflected from a parabola, and parallel light or radio waves reflected from a parabolic dish, meet at a focal point above the vertex. Scientists use CSIRO's six radio telescopes, with parabolic dishes, to research objects in outer space.

Lesson starter: From x-intercepts to turning point

This graph has the rule $y = x^2 - 2x - 3$ but all its important features are not shown.

Find the features by discussing these questions.

- Can the quadratic rule be factorised?
- What is the value of y at the x-intercepts?
- How can the factorised form of the rule help to find the x-intercepts?
- How does the x-coordinate of the turning point relate to the x-intercepts?
- Discuss how the y-coordinate of the turning point can be found.
- Finish by finding the y-intercept.




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


11H

Key ideas


- All parabolas have one y-intercept and can have zero, one or two x-intercepts.


Zero x-intercepts







One x-intercept






Two x-intercepts





- x-intercepts can be found by substituting $y = 0$ and using the Null Factor Law.
- If the graph has two x-intercepts (at $x = a$ and $x = b$), the turning point can be found by:
  - calculating the x-coordinate of the turning point, which is the midpoint of a and b; that is, $x = \frac{a+b}{2}$
  - calculating the y-coordinate of the turning point by substituting the x-coordinate into the rule for the quadratic.



$$y = (x - a)(x - b)$$

x-intercepts : $0 = (x - a)(x - b)$

$$x - a = 0 \text{ or } x - b = 0$$

$$x = a \text{ or } x = b$$

Exercise 11H

Understanding

1–3 | 1–3

1 Factorise these quadratics.

a $x^2 + 2x$ **b** $x^2 - 3x$ **c** $x^2 - 9$

d $x^2 - 49$ **e** $x^2 - x - 12$ **f** $x^2 - 4x + 4$

2 For these factorised quadratics, use the Null Factor Law to solve for x, then find the x-value halfway between the two values.

a $0 = (x - 2)(x + 2)$ **b** $0 = (x + 1)(x + 5)$

c $0 = (x - 1)(x + 3)$ **d** $0 = (x - 2)(x + 3)$

3 Use substitution to find the y-coordinate of the turning point of these quadratics. The x-coordinate of the turning point is given.

a $y = x^2 - 4, x = 0$ **b** $y = x^2 - 6x, x = 3$ **c** $y = x^2 + 4x + 3, x = -2$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Fluency

4–6(½) 4–6(½)

Example 22 Finding intercepts of quadratic graphs

For each of the following quadratic relations, find:

i the coordinates of the x-intercepts
ii the coordinates of the y-intercept.

a $y = x(x + 1)$

b $y = 2(x + 2)(x - 3)$

Solution

Explanation

a i x-intercepts (let $y = 0$):
$x(x + 1) = 0$
$x = 0$ or $x + 1 = 0$
$x = 0$ or $x = -1$
$\therefore (0, 0)$ and $(-1, 0)$

Let $y = 0$ to find the x-intercepts.

Apply the Null Factor Law to set each factor equal to 0 and solve.

ii y-intercept (let $x = 0$):
$y = 0(0 + 1)$
$y = 0$
$\therefore (0, 0)$

Let $x = 0$ to find the y-intercept.

b i $y = 2(x + 2)(x - 3)$ has two x-intercepts.
x-intercepts (let $y = 0$):
$2(x + 2)(x - 3) = 0$
$x + 2 = 0$ or $x - 3 = 0$
$x = -2$ or $x = 3$
$\therefore (-2, 0)$ and $(3, 0)$

There are two different factors.

Let $y = 0$ to find the x-intercepts. Set each factor equal to 0 and solve.

ii y-intercept (let $x = 0$):
$y = 2(0 + 2)(0 - 3)$
$y = 2(2)(-3)$
$y = -12$
$\therefore (0, -12)$

Let $x = 0$ to find the y-intercept.

Now you try

For each of the following quadratic relations, find:

i the coordinates of the x-intercepts
ii the coordinates of the y-intercept.

a $y = x(x + 5)$

b $y = 3(x + 1)(x - 4)$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

11H

4 For each of the following quadratic relations, find:
- **i** the coordinates of the x-intercepts
- **ii** the coordinates of the y-intercept.

a $y = x(x+7)$
b $y = x(x+3)$
c $y = x(x+4)$
d $y = (x-4)(x+2)$
e $y = (x+2)(x-5)$
f $y = (x-7)(x+3)$
g $y = 2(x+3)(x-1)$
h $y = 3(x+4)(x+1)$
i $y = (2-x)(3-x)$




Hint for Q4: Use the Null Factor Law to find the x-intercepts.

Example 23 Sketching using intercept form

Consider the quadratic relation $y = x^2 - 2x$.

a Find the coordinates of the y-intercept.
b Factorise the relation.
c Find the coordinates of the x-intercepts.
d Find the axis of symmetry.
e Find the turning point.
f Sketch the graph clearly showing all the key features.

Solution	Explanation
a y-intercept (let $x = 0$): $y = 0$ $\therefore (0, 0)$	Let $x = 0$ to find the y-intercept. $y = 0$ since $0^2 - 2(0) = 0$.
b $y = x^2 - 2x$ $= x(x-2)$	Take out the common factor of x.
c x-intercepts (let $y = 0$): $0 = x(x-2)$ $x = 0$ or $x - 2 = 0$ $x = 0$ or $x = 2$ $\therefore (0, 0)$ and $(2, 0)$	Let $y = 0$ to find the x-intercepts. Set each factor equal to 0 and solve.
d Axis of symmetry: $x = \frac{0+2}{2}$ $\therefore x = 1$	The axis of symmetry is halfway between the x-intercepts.
e Turning point occurs when $x = 1$. When $x = 1$, $y = (1)^2 - 2(1)$ $y = -1$ $\therefore$ there is a minimum turning point at $(1, -1)$.	Substitute $x = 1$ into $y = x^2 - 2x$ to find the y-coordinate. The coefficient of x^2 is positive, therefore the basic shape is ∪.
f 	Sketch the graph, labelling the key features found above.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Now you try

Consider the quadratic relation $y = x^2 - 4x$.

a Find the coordinates of the y-intercept.
b Factorise the relation.
c Find the coordinates of the x-intercepts.
d Find the axis of symmetry.
e Find the turning point.
f Sketch the graph clearly showing all the key features.

5 Consider each of the following quadratic relations.

i Find the coordinates of the y-intercept.
ii Factorise the relation.
iii Find the coordinates of the x-intercepts.
iv Find the axis of symmetry.
v Find the turning point.
vi Sketch the graph clearly showing all the key features.



Hint for Q5: First factorise before finding the x-intercepts.

a $y = x^2 - 5x$
b $y = x^2 + x$
c $y = x^2 - 3x$
d $y = 2x + x^2$
e $y = 5x + x^2$
f $y = 3x - x^2$
g $y = -x^2 - 8x$
h $y = -2x - x^2$
i $y = -x^2 + x$

Example 24 Sketching a quadratic trinomial

Consider the quadratic relation $y = x^2 + 2x - 8$.

a Find the coordinates of the y-intercept.
b Factorise the relation.
c Find the coordinates of the x-intercepts.
d Find the axis of symmetry.
e Find the turning point.
f Sketch the graph clearly showing all the key features.

Solution	Explanation
a y-intercept (let $x = 0$): $y = -8$ $\therefore (0, -8)$	Let $x = 0$ to find the y-intercept.
b $y = x^2 + 2x - 8$ $= (x + 4)(x - 2)$	Factorise the quadratic trinomial: $4 \times (-2) = -8$ and $4 + (-2) = 2$.
c x-intercepts (let $y = 0$): $0 = (x + 4)(x - 2)$ $x + 4 = 0$ or $x - 2 = 0$ $x = -4$ or $x = 2$ $\therefore (-4, 0)$ and $(2, 0)$	Let $y = 0$ to find the x-intercepts. Use the factorised form to solve, applying the Null Factor Law.

Continued on next page

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

11H

d Axis of symmetry: $x = \frac{-4+2}{2}$

$\therefore x = -1$

The axis of symmetry is halfway between the x-intercepts.

e Turning point occurs when $x = -1$.

When $x = -1$, $y = (-1)^2 + 2(-1) - 8$

$y = -9$


$\therefore$ there is a minimum turning point at $(-1, -9)$.

Substitute $x = -1$ into $y = x^2 + 2x - 8$ to find the y-coordinate. $y = 1 - 2 - 8$

$x = -1$ and $y = -9$

The coefficient of x^2 is positive, therefore the basic shape is ∪.

f



Sketch the graph showing the key features.

Now you try

Consider the quadratic relation $y = x^2 - 4x - 12$.

a Find the coordinates of the y-intercept.
b Factorise the relation.
c Find the coordinates of the x-intercepts.
d Find the axis of symmetry.
e Find the turning point.
f Sketch the graph clearly showing all the key features.

6 Consider each of the following quadratic relations.

i Find the coordinates of the y-intercept.
ii Factorise the relation.
iii Find the coordinates of the x-intercepts.
iv Find the axis of symmetry.
v Find the turning point.
vi Sketch each graph clearly showing all the key features.


a $y = x^2 - 3x + 2$
b $y = x^2 - 4x + 3$
c $y = x^2 + 2x - 3$
d $y = x^2 + 4x + 4$
e $y = x^2 + 2x + 1$
f $y = x^2 + 2x - 8$

Problem-solving and reasoning


7–9 | 7, 9, 10

7 State the missing number (?) in these quadratic graphs.


a



b




c




Essential Mathematics Core for the Victorian Curriculum Year 9
ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


d



e



f



8 The path of a golf ball is given by the rule $y = 30x - x^2$, where y is the height in metres above the ground and x is the horizontal distance in metres. Find:

a how far the ball travels horizontally

b how high the ball reaches mid-flight.

9 A test rocket is fired and follows a path described by $y = 0.1x(200 - x)$. The height is y metres above ground and x is the horizontal distance in metres.

a How far does the rocket travel horizontally?

b How high does the rocket reach mid-flight?



10 Explain why the coordinates of the x-intercept and the turning point for $y = (x - 2)^2$ are the same.

Mixing it up — 11

11 For each of the following mixed quadratic relations, sketch the graph, clearly showing the x- and y-intercepts and the turning point.

a $y = x^2 - 1$

b $y = 9 - x^2$

c $y = x^2 + 5x$

d $y = 2x^2 - 6x$

e $y = 4x^2 - 8x$

f $y = x^2 - 3x - 4$

g $y = x^2 + x - 12$

h $y = x^2 + 2x + 1$

i $y = x^2 - 6x + 9$



Design structures, such as this building at LAX airport, can be modelled on a set of axes.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


Maths@Work

Fencing Crown land

Land along the side of rivers is usually owned by the government and is sometimes called Crown land. Farmers can often lease this land to graze their sheep or cattle.

A farmer has a permit to fence off a rectangular area of land alongside the river. She has 400 m of fencing available and does not need to fence along the river. She wants to maximise the area of land for her stock using this fixed amount of fencing.





1 **a** Find the length of the rectangular area when the width is:
i 50 m **ii** 120 m **iii** 180 m
b Find the area of the land when the width is:
i 50 m **ii** 120 m **iii** 180 m
c Which width from part **b** above gave the largest area? Explain why the area decreases for small and large values for the width.

2 **a** Using x metres to represent the width, write an expression for the length showing working.
b Write an expression for the area of the land in terms of x.
c Use your area expression from part **b** to find the area of the land when x is:
i 20 **ii** 80 **iii** 160

3 **a** Using your expression from part **b** above, sketch a graph of Area (A) versus x. You should find the following to help complete the graph.
i y-intercept **ii** x-intercepts
iii axis of symmetry **iv** turning point
b What value of x gives the maximum area of land for grazing? Explain your choice and give the dimensions of the rectangular area of land.
c What do you notice about the width and the length when the area is a maximum?
d See if the same is true if the farmer had 600 m of fencing instead. Show your expressions and graph.

Using technology

4 **a** Use graphing software like Desmos to graph your rule from part **2b**.
b Zoom to confirm your answers to some of the above questions.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Bridge support arch

Parabolic shapes are often used as bridge arches due to their self-supporting properties. Using the coordinates $(0, 0)$ for one side of a bridge arch, a model for a bridge parabolic curve is given by $y = kx(a - x)$ where:

- y metres is the vertical distance above water level
- x metres is the horizontal distance from the left-hand side
- a and k are positive numbers.



Present a report for the following tasks and ensure that you show clear mathematical workings and explanations where appropriate.

1 Preliminary task

a If $k = 0.5$ and $a = 30$:

- **i** construct the rule connecting y and x
- **ii** sketch a graph representing the bridge arch by firstly finding the x-intercepts
- **iii** determine the maximum height of the arch.

b Repeat part **a** for $k = 0.2$ and $a = 50$.

2 Modelling task

a The problem is to determine a rule of a new bridge design which clears a river of a certain width and has a maximum arch height of less than or equal to 40 metres. Write down all the relevant information that will help solve this problem with the aid of a rough sketch.

b Construct the rule connecting y and x using the following values for a and k.

- **i** $a = 50$ and $k = 0.1$
- **ii** $a = 70$ and $k = 0.02$

Formulate

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Solve

c If $a = 50$ and $k = 0.1$, sketch a graph of the bridge arch by locating and labelling the following:

i x-intercepts

ii turning point.

d Decide if an arch that is designed using $a = 50$ and $k = 0.1$ meets the requirement that the maximum height is to be less than or equal to 40 metres.

e If $a = 70$ and $k = 0.02$, sketch a graph of the bridge arch by locating and labelling the following:

i x-intercepts **ii** turning point.

f Decide if an arch that is designed using $a = 70$ and $k = 0.02$ meets the requirement that the maximum height is to be less than or equal to 40 metres.

Evaluate and verify

g Choose your own values of a and k so that a bridge that spans a 60 metre-wide river has a maximum arch height of less than 40 metres. Justify your choice by showing your calculations and a graph.

h Choose your own values of a and k so that a bridge that spans a 60 metre-wide river has a maximum arch height of exactly 40 metres. Justify your choice by showing your calculations and a graph.

Communicate

i Summarise your results and describe any key findings.

3 Extension question

The rule for k in terms of a is $k = \frac{160}{a^2}$ if the maximum height of the arch is to be 40 metres.

a Use the rule for k to determine the value of k if the width of the river is 100 m.

b Use the rule for k to determine the width of the river, correct to one decimal place, if $k = 0.3$.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Modelling the bridge

Key technology: Graphing

A parabola is a graph of a quadratic relation and is a shape that appears in many natural settings but also in building and design. For example, the path of an object though the air can be modelled using a parabola. They can also be seen in complex engineering structures including self-supporting arches and bridges.

1 Getting started

We will start by considering the key features of the graphs of the following parabolas.

i $y = x^2$

ii $y = -2x^2$

iii $y = (x+2)^2 - 4$

iv $y = -\frac{1}{2}(x-1)^2 + 3$

For each parabola state the:

a coordinates of the turning point

b equation of the axis of symmetry

c coordinates of the y-intercept.


2 Using technology

We will explore the effect of the pronumerals a, h and k in $y = a(x-h)^2 + k$ by using a graphing package like Desmos.

a Set up a page using the four rules $y = ax^2$, $y = x^2 + k$, $y = (x-h)^2$ and $y = a(x-h)^2 + k$ adding sliders for a, h and k.

b Drag the sliders for the three pronumerals and describe the effect of changing each value on the graph.

c Drag the sliders to find a parabola which has a maximum turning point at $(2, 4)$ and a y-intercept of $(0, -2)$.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Technology and computational thinking

3 Applying an algorithm

The Sydney Harbour Bridge contains two key arches which are central to its design and construction with the lower arch being the one considered here.

For the lower arch use the following approximate measurements.

- Total arch width from ground to ground, 500 m
- Maximum height above ground level, 110 m

a By taking the left point of the arch as $(0, 0)$, state the coordinates of the:
i turning point **ii** x-intercepts.

b Using your graphing package with the sliders, find a parabola which passes through the points found in part **a** above. Follow these steps.
- Step 1: Choose values for h and k to ensure the graph passes through the correct turning point.
- Step 2: Adjust the scale for the slider for a using a smaller increment.
- Step 3: Drag the slider so that the graph passes near the correct x-intercepts
- Step 4: If required repeat from Step 2 until the graph passes through the correct intercepts with a high degree of accuracy.

c Write down your rule which models the lower arch of the Sydney Harbour Bridge.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

1 **a** What do you notice about the sum of these numbers?

i $1+3$

ii $1+3+5$

iii $1+3+5+7$

iv $1+3+5+7+9$

b Find the sum of the first 100 odd integers.

2 Solve these equations.


a $6x^2 = 35 - 11x$

b $\frac{2}{x^2} = 1 - \frac{1}{x}$

c $\sqrt{x-1} = \frac{2}{x-1}$


3 The height of a projectile (in metres) above the ground is given by the expression $t(14-t)$ where time t is in seconds. For how long is the projectile at a height of above 40 m?

4 Find the quadratic rule that relates the number of balls to the term number (n) in the pattern below.




If there are 66 balls in the pattern, what term number is it?

5 Find the value of x in this diagram.



6 **a** A right-angled triangle's shortest side is of length x, the hypotenuse is 9 units longer than the shortest side and the other side is 1 unit longer than the shortest side. Find the side lengths of the triangle.



b The area of a right-angled triangle is 60 square units and the lengths of the two shorter sides differ by 7 units. Find the length of the hypotenuse.

7 Given $a > 0$, for what values of k does $y = a(x-h)^2 + k$ have:

a two x-intercepts? **b** one x-intercept? **c** no x-intercepts?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter summary


Introduction to quadratic equations and graphs

Turning point form
$y = a(x - h)^2 + k$
turning point (h, k)
e.g. $y = 2(x + 1)^2 + 3$
TP: $(-1, 3)$
y-int: $x = 0$, $y = 2(0 + 1)^2 + 3$
$= 2(1)^2 + 3$
$= 5$
$\therefore (0, 5)$


Standard form
A quadratic is of the form
$y = ax^2 + bx + c$, $a \neq 0$

Translation
$y = x^2 + k$ $k > 0$, graph shifts up k units
$k < 0$, graph shifts down k units
$y = (x - h)^2$ $h > 0$, graph shifts h units right
$h < 0$, graph shift h units left


e.g.




Parabola
Basic form $y = x^2$



Turning point form
$y = a(x - h)^2 + k$




Dilation and reflection
$y = ax^2$
$a < 0$, graph is reflected in x-axis




$-1 < a < 1$:
graph appears wider than $y = x^2$ or $y = -x^2$

$a > 1$ or $a < -1$:
graph appears narrower than $y = x^2$ or $y = -x^2$
e.g.



Sketching $y = ax^2 + bx + c$ ⭐
Find x- and y-intercepts and turning point then sketch.
e.g. $y = x^2 - 6x + 5$
y-int: $x = 0$, $y = 0^2 - 6(0) + 5$
$= 5$
x-int: $y = 0$, $0 = x^2 - 6x + 5$
$0 = (x - 1)(x - 5)$
$x - 1 = 0$ or $x - 5 = 0$
$x = 1$ or $x = 5$
Turning point: halfway between the x-intercepts:
$x = \frac{1+5}{2} = 3$
Substitute $x = 3$ into rule:
$y = 3^2 - 6(3) + 5$
$= -4$
$(3, -4)$ minimum turning point




Applications
Define variable.
Set up equation.
Solve using Null Factor Law.
Check answer makes sense.
Answer the question in words.

Solving quadratic equations
- Rewrite equation in standard form: i.e. $ax^2 + bx + c = 0$
- Factorise
- Apply Null Factor Law: If $p \times q = 0$, $p = 0$ or $q = 0$.
- Solve remaining equation:
  e.g. $x^2 + 5x = 0$
  $x(x + 5) = 0$
  $x = 0$ or $x + 5 = 0$
  $x = 0$ or $x = -5$
  e.g. $x^2 - 12 = 4x$
  $x^2 - 4x - 12 = 0$
  $(x - 6)(x + 2) = 0$
  $x - 6 = 0$ or $x + 2 = 0$
  $x = 6$ or $x = -2$
- If $x^2 = 16$, square root both sides so $x = 4$ or $x = -4$.

Essential Mathematics Core for the Victorian Curriculum Year 9 ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter checklist

A version of this checklist that you can print out and complete can be downloaded from your Interactive Textbook.

		✓
11A	**1 I can write a quadratic equation in standard form.** e.g. Write $x(x - 3) = 6$ in standard form.	
11A	**2 I can determine if a value is a solution of an equation.** e.g. Use substitution to determine if $x = -2$ is a solution of $2x^2 + 3x - 2 = 0$.	
11A	**3 I can use the Null Factor Law to solve an equation.** e.g. Use the Null Factor Law to solve $(x + 2)(x - 5) = 0$.	
11B	**4 I can solve a quadratic equation with a common factor.** e.g. Solve $2x^2 - 10x = 0$.	
11B	**5 I can solve a quadratic equation of the form $ax^2 = d$** e.g. Solve $6x^2 = 24$.	
11C	**6 I can solve equations involving quadratic trinomials.** e.g. Solve $x^2 - 3x - 70 = 0$.	
11C	**7 I can solve quadratic equations not in standard form.** e.g. Solve $x^2 = 9x - 20$.	
11D	**8 I can set up and solve a quadratic equation from a word problem.** e.g. The length of a rectangular postcard is 5 cm more than its width and the area of the postcard is 84 cm^2. Find the dimensions of the postcard.	
11E	**9 I can identify the key features of a parabola.** e.g. Consider this graph.  State: i the equation of the axis of symmetry ii the type and coordinates of the turning point iii the x- and y-intercept coordinates.	
11E	**10 I can plot a parabola from a table of values.** e.g. For $y = x^2 - 4x + 4$, fill in the table of values and plot the points on a set of axes to form a parabola. x | −3 | −2 | −1 | 0 | 1 | 2 | 3 y | | | | | | |	

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter checklist

		✔
11E	**11 I can compare graphs of $y = ax^2$, $a > 0$** e.g. Sketch $y = x^2$, $y = 3x^2$ and $y = \frac{1}{3}x^2$ by first completing the table of values below. x: −2, −1, 0, 1, 2 $y = x^2$ $y = 3x^2$ $y = \frac{1}{3}x^2$ State the turning point of each graph and whether the graphs of $y = 3x^2$ and $y = \frac{1}{3}x^2$ are wider or narrower than the graph of $y = x^2$.	
11F	**12 I can compare graphs of $y = ax^2$, $a < 0$** e.g. Sketch $y = -x^2$, $y = -4x^2$ and $y = -\frac{1}{4}x^2$ by first completing the table of values below. x: −2, −1, 0, 1, 2 $y = -x^2$ $y = -4x^2$ $y = -\frac{1}{4}x^2$ State the turning point of each graph and whether the graphs of $y = -4x^2$ and $y = -\frac{1}{4}x^2$ are wider or narrower than the graph of $y = -x^2$.	
11G	**13 I can sketch parabolas with vertical translations.** e.g. Sketch $y = x^2 + 3$ showing the y-intercept and turning point.	
11G	**14 I can sketch parabolas with horizontal translations.** e.g. Sketch $y = -(x + 1)^2$ showing the y-intercept and turning point.	
11G	**15 I can sketch parabolas with combined translations.** e.g. Sketch $y = (x - 1)^2 - 3$ showing the y-intercept and turning point.	
11H ★	**16 I can sketch parabolas using intercept form.** e.g. For the quadratic relation $y = x^2 - 6x$, find the x- and y-intercepts and the turning point and sketch its graph, labelling key features.	
11H ★	**17 I can sketch a quadratic trinomial.** e.g. For the quadratic relation $y = x^2 - 4x - 12$, find the x- and y-intercepts and the turning point and sketch its graph, labelling key features.	

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Short-answer questions

11E **1** Consider the quadratic $y = x^2 - 2x - 3$.

a Complete this table of values for the equation.

x	-3	-2	-1	0	1	2	3
y							

b Plot the points in part **a** on a Cartesian plane and join in a smooth curve.

11A **2** Use the Null Factor Law to solve the following equations.

a $x(x + 2) = 0$

b $3x(x - 4) = 0$

c $(x + 3)(x - 7) = 0$

d $(x - 2)(2x + 4) = 0$

e $(x + 1)(5x - 2) = 0$

f $(2x - 1)(3x - 4) = 0$

11B/C **3** Solve the following quadratic equations by first factorising.

a $x^2 + 3x = 0$

b $2x^2 - 8x = 0$

c $x^2 = 25$

d $x^2 = 81$

e $5x^2 = 20$

f $3x^2 - 30 = 0$

g $x^2 + 10x + 21 = 0$

h $x^2 - 3x - 40 = 0$

i $x^2 - 8x + 16 = 0$

11C **4** Write the following quadratic equations in standard form and solve for x.

a $x^2 = 5x$

b $3x^2 = 18x$

c $x^2 + 12 = -8x$


d $2x + 15 = x^2$

e $x^2 + 15 = 8x$

f $4 - x^2 = 3x$


11D **5** Set up and solve a quadratic equation to determine the value of x that gives the specified area of the shapes below.

a



$A = 80$ units^2

b



$A = 30$ units^2

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


Chapter review

11E **6** For the following graphs state:


i the axis of symmetry

ii the turning point and its type.

a




b




11F/G **7** Match the following quadratic rules with their graphs.

$y = x^2 - 4$, $y = 2x^2$, $y = (x+1)^2$, $y = (x-3)^2$, $y = -x^2 + 2$


a




b




c



d



e



11G **8** Sketch the following graphs labelling the turning point and y-intercept.

a $y = x^2 + 2$

b $y = -x^2 - 5$

c $y = (x+2)^2$

d $y = -(x-3)^2$

e $y = (x-1)^2 + 3$

f $y = 2(x+2)^2 - 4$

11H **9** Sketch the following graphs labelling the y-intercept, turning point and x-intercepts.

a $y = x^2 - 8x + 12$

b $y = x^2 + 10x + 16$

c $y = x^2 + 2x - 15$

d $y = x^2 + 4x - 5$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Multiple-choice questions

11A **1** $(1, 3)$ is a point on a curve with which equation?

A $y = x^2$ **B** $y = (x - 3)^2$ **C** $y = x^2 + 2x - 3$
D $y = x^2 + 2$ **E** $y = 6 - x^2$

11A **2** The solution(s) to $2x(x - 3) = 0$ is/are:

A $x = -3$ **B** $x = 0$ or $x = 3$ **C** $x = 2$ or $x = 3$
D $x = 0$ or $x = -3$ **E** $x = 0$

11A **3** The quadratic equation $x^2 = 7x - 12$ in standard form is:

A $x^2 - 7x - 12 = 0$ **B** $-x^2 + 7x + 12 = 0$ **C** $x^2 - 7x + 12 = 0$
D $x^2 + 7x - 12 = 0$ **E** $x^2 + 7x + 12 = 0$

11B **4** The solution(s) to $x(x + 2) = 2x + 9$ is/are:

A $x = -3$ or $x = 3$ **B** $x = 0$ or $x = -2$ **C** $x = 3$
D $x = 9$ or $x = -1$ **E** $x = 9$ or $x = 0$

The following applies to Questions **5** and **6**.

The height, h metres, of a toy rocket above the ground t seconds after launch is given by $h = 6t - t^2$.

11D **5** The rocket returns to ground level after:

A 5 seconds **B** 3 seconds **C** 12 seconds
D 6 seconds **E** 8 seconds

11H **6** The rocket reaches its maximum height after:

A 6 seconds
B 3 seconds
C 10 seconds
D 4 seconds
E 9 seconds

11G **7** The turning point of $y = (x - 2)^2 - 4$ is:

A a maximum at $(2, 4)$
B a minimum at $(-2, 4)$
C a maximum at $(-2, 4)$
D a minimum at $(-2, -4)$
E a minimum at $(2, -4)$

11G **8** The transformation of the graph of $y = x^2$ to $y = x^2 - 2$ is described by:

A a translation of 2 units to the left
B a translation of 2 units to the right
C a translation of 2 units down
D a translation of 2 units up
E a translation of 2 units right and 2 units down

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter review


11F **9** Compared to $y = x^2$, the narrowest graph is:

A $y = 5x^2$ **B** $y = 0.2x^2$ **C** $y = 2x^2$


D $y = \frac{1}{2}x^2$ **E** $y = 3.5x^2$

11G **10** The graph of $y = -(x - 3)^2$ is:


A




B




C



D



E



Extended-response questions

1 A sail of a yacht is in the shape of a right-angled triangle. It has a base length of $2x$ metres and its height is 5 metres more than half its base.

a Write an expression for the height of the sail.

b Give an expression for the area of the sail in expanded form.

c If the area of the sail is 14 m^2, find the value of x.

d Hence, state the dimensions of the sail.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

2 Connor and Sam are playing in the park with toy rockets they have made. They launch their rockets at the same time to see whose is better.

a The path of Sam's rocket is modelled by the equation $h = 12t - 2t^2$, where h is the height of the rocket in metres after t seconds.

i Find the coordinates of the axis intercepts.

ii Find the turning point.

iii Sketch a graph of the height of Sam's rocket over time.

b The path of Connor's rocket is modelled by the equation $h = -(t - 4)^2 + 16$, where h is the height of the rocket in metres after t seconds.

i Find the h-intercept coordinates (i.e. $t = 0$).

ii State the turning point.

iii Use the answers from parts **b i** and **ii** to state the coordinates of the two t-intercepts.

iv Sketch a graph of the height of Connor's rocket over time.

c **i** Whose rocket was in the air for longer?

ii Whose rocket reached the greater height and by how much?

iii How high was Sam's rocket when Connor's rocket was at its maximum height?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Semester review 2

Measurement

Short-answer questions


1 A digital scale records the weight of a parcel to be 6.4 kg.

a If the reading is mistakenly recorded as 6.0 kg, what is the percentage error?


b What is the lower and upper limit of accuracy of the true recording?

2 Find the area of these basic shapes.


a



b



c




3 Find the circumference and area of these circles, correct to two decimal places.


a



b




c




4 Find the area of each of the figures below. Round to two decimal places where necessary.


a




b



c




d




5 Find the surface area of these solid objects. Round to two decimal places where necessary.


a



b




c




6 Find the volume of these solids. Round to two decimal places for part **c**.


a



b




c



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


Multiple-choice questions

1 The perimeter and area of this triangle, respectively, are:



A 48 cm, 24 m^2 **B** 68 cm, 80 cm^2
C 24 cm, 24 cm^2 **D** 24 cm, 12 cm^3
E 24 cm, 48 cm^2

2 The circumference and area for this circle, correct to two decimal places, respectively, are:




A 37.70 m, 18.85 m^2 **B** 9.42 m, 56.55 m^2
C 18.85 m, 28.27 m^2 **D** 18.85 m, 18.85 m^2
E 9.42 m, 28.27 m^2

3 420 cm^2 is equal to:


A 4.2 m^2 **B** 0.42 m^2 **C** 42 000 m^2 **D** 0.042 m^2 **E** 0.0042 m^2

4 The perimeter and area of the figure shown, respectively, are:



A 20.2 m, 22.42 m^2 **B** 24.8 m, 22.42 m^2
C 24.8 m, 25.15 m^2 **D** 20.2 m, 25.15 m^2
E 21.6 m, 24.63 m^2


5 The volume of the cylinder shown is closest to:



A 703.7 cm^3 **B** 351.9 cm^3
C 2814.9 cm^3 **D** 452.4 cm^3
E 1105.8 cm^3

Extended-response questions

A pond is to be built inside a rectangular garden, as shown.



a Find the area of the pond, correct to two decimal places.
b What is the area of the garden, not including the pond?
c The pond is to be surrounded by a low wall that costs $50 per metre. What will it cost to build this wall? Round to the nearest dollar.



Essential Mathematics Core for the Victorian Curriculum Year 9 ISBN 978-1-009-59315-1 © Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Semester review 2

Indices

Short-answer questions

1 Use index laws to simplify the following.

a $5p^2q \times 3pq$ **b** $\dfrac{9a^6b^3}{18a^4b^2}$ **c** $(-3x^4y^2)^2 \times 6xy^2$ **d** $-3x^0 + (5x)^0$

2 Write each of the following using positive indices and then write as a basic numeral.

a $\dfrac{3}{4^{-2}}$ **b** 4×10^{-3}

3 Write these numbers as basic numerals.

a 2.4×10^3 **b** 1.08×10^6 **c** 7.1×10^{-3} **d** 2.06×10^{-5}

4 Write these numbers using scientific notation.

a 60 300 **b** 2 700 000 **c** 0.004 **d** 0.000703

5 Convert these numbers to the units given in brackets. Write your answer using scientific notation using three significant figures.

a 30.71 g (kg) **b** 4236 tonnes (kg)
c 3.4 hours (seconds) **d** 235 seconds (years)

Multiple-choice questions

1 $3a^2b^3 \times 4ab^2$ is equivalent to:

A $12a^2b^6$ **B** $7a^3b^5$ **C** $12a^3b^5$ **D** $12a^4b^5$ **E** $7a^2b^6$

2 $\left(\dfrac{2x}{5}\right)^3$ is equivalent to:

A $\dfrac{6x^3}{5}$ **B** $\dfrac{8x^3}{125}$ **C** $\dfrac{2x^3}{5}$ **D** $\dfrac{2x^4}{15}$ **E** $\dfrac{2x^3}{125}$

3 4^{-2} can be expressed as:

A $\dfrac{1}{4^{-2}}$ **B** $\dfrac{1}{8}$ **C** -16 **D** $\dfrac{1}{16}$ **E** -8

4 3×10^{-4} written with positive indices is:


A -3×10^4 **B** $\dfrac{1}{3 \times 10^4}$ **C** $\dfrac{-3}{10^4}$ **D** $\dfrac{1}{3 \times 10^{-4}}$ **E** $\dfrac{3}{10^4}$

5 0.00371 in scientific notation is:

A 0.371×10^{-3} **B** 3.7×10^{-2} **C** 3.71×10^{-3} **D** 3.71×10^3 **E** 371×10^3

Extended-response question

The average human body contains about 74 billion cells.
(Note: 1 billion = 1 thousand million)



a Write this number of cells:
i as a basic numeral
ii using scientific notation

b If the population of a particular city is 2.521×10^6, how many human cells are there in the city? Give your answer using scientific notation correct to three significant figures.

c If the average human weighs 64.5 kilograms, what is the average mass of one cell in grams? Give your answer using scientific notation correct to three significant figures.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


Geometry

Short-answer questions


1 State the angle sums for these shapes.
- **a** Triangle
- **b** Quadrilateral
- **c** Pentagon
- **d** Heptagon
- **e** Octagon
- **f** Decagon

2 Find the value of each pronumeral in the following.


a




b




c




d




e




f




g




h




3 The two given triangles are similar.
- **a** Are they congruent? Give a reason.
- **b** Write a congruence statement.
- **c** Which side on $\triangle ABC$ corresponds to side DE on $\triangle DEF$?



4 For the given pair of triangles (not to scale):
- **a** give a reason (SSS, SAS, AAA or RHS) why they are similar
- **b** find the scale factor



5 For this pair of triangles:
- **a** give a reason why the two triangles are similar
- **b** find the value of x



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Semester review 2

Multiple-choice questions

1 The supplementary angle to 55° is:

A 55° **B** 35° **C** 125° **D** 135° **E** 70°

2 A quadrilateral with all four sides equal and opposite sides parallel is best described by a:


A parallelogram **B** rhombus **C** rectangle
D trapezium **E** kite

3 The size of the interior angle in a regular pentagon with angle sum 540° is:

A 108° **B** 120° **C** 96° **D** 28° **E** 115°


4 The test that proves congruence in these two triangles is:

A SAS **B** RHS **C** AAA
D SSS **E** AAS




5 What is the scale factor that enlarges shape 1 to shape 2 in these similar figures, and what is the value of x?

A 2 and $x = 8$ **B** 2.5 and $x = 7.5$
C 3.33 and $x = 13.33$ **D** 2.5 and $x = 12.5$
E 2 and $x = 6$



Extended-response question

A vertical wall is being supported by a piece of timber that touches the ground 10 metres from the base of the wall. A vertical metal support 4.5 m high is placed under the timber support 4 m from the wall.



a Give a reason why $\triangle ABD \parallel\!\parallel \triangle ECD$.
b What is the length DC?
c What is the scale factor for the two triangles formed by the timber and support?
d Find how far the timber reaches up the wall.
e How far above the ground is the point halfway along the timber support?

Algebraic techniques

Short-answer questions

1 Expand these expressions.


a $-3(x+2)$ **b** $x(1-x)$ **c** $(x+1)(x-1)$
d $(x-4)^2$ **e** $(5x-2)(5x+2)$ **f** $(3x-4)^2$
g $(4x-1)(2x+3)$ **h** $5x-3(x-1)$ **i** $(2-x)^2$

2 Simplify these expressions.


a $-3x \times 2yx$ **b** $7a \times 6ab^2$ **c** $4ab \div (8b)$
d $-6x^2y \div (xy^2)$ **e** $a^2b \div (ab^2)$ **f** $2xy \div (4xy^2)$

3 Find the area of the following shapes in expanded form.


a



b



c



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

4 Factorise the following.

a $3x - 12$ **b** $-7x - 14$ **c** $5x^2 + 2x$ **d** $14x^2 - 21x$

5 Factorise the following.

a $x^2 - 9$ **b** $25 - 4y^2$

c $x^2 - 6x + 3x - 18$ **d** $3x^2 - 10 + 6x - 5x$

6 Factorise the following quadratic trinomials.

a $x^2 + 9x + 20$ **b** $x^2 + 2x - 8$ **c** $x^2 - x - 56$ **d** $2x^2 - 4x - 48$

7 Simplify these expressions.

a $\frac{5x^2}{3} \times \frac{9}{10x}$ **b** $\frac{x-1}{2} \times \frac{3}{2(x-1)}$ **c** $\frac{4x-8}{x-2}$

d $\frac{-6x^2 - x}{6x + 1}$ **e** $\frac{4x}{3} \div \frac{8x}{9}$ **f** $\frac{3(x-2)}{5a} \div \frac{x(x-2)}{15a^2}$

8 Simplify these algebraic fractions.

a $\frac{x}{3} - \frac{x}{4}$ **b** $\frac{5x}{4} + \frac{7x}{5}$ **c** $\frac{x+1}{2} + \frac{x}{5}$ **d** $\frac{2x-1}{5} + \frac{x+2}{3}$

Multiple-choice questions

1 $-3(x - 4)$ is equal to:

A $-3x + 7$ **B** $-3x + 4$ **C** $-3x - 4$ **D** $-3x - 12$ **E** $-3x + 12$

2 The expanded form of $(x + 4)(3x - 2)$ is:

A $3x^2 + 12x - 8$ **B** $4x^2 + 14x - 8$ **C** $3x^2 + 12x - 10$

D $3x^2 - 8$ **E** $3x^2 + 10x - 8$

3 $(2x - 1)^2$ expands to:

A $4x^2 - 4x + 1$ **B** $2x^2 - 2x - 1$ **C** $2x^2 - 4x + 1$ **D** $4x^2 - 2x + 1$ **E** $4x^2 - 1$

4 $\frac{2x - 6}{2}$ simplifies to:


A $2x - 3$ **B** $x - 3$ **C** $2x - 4$ **D** $x - 6$ **E** $x - 4$

5 $\frac{x+1}{3} + \frac{x}{4}$ simplifies to:

A $\frac{7x+1}{12}$ **B** $\frac{7x+4}{7}$ **C** $\frac{5x+4}{7}$ **D** $\frac{7x+4}{12}$ **E** $\frac{5x-3}{8}$

Extended-response question

A rectangular room that is 10 metres long and 8 metres wide has a rectangular rug in the middle of it that leaves a border, x metres wide, all the way around it as shown.



a Write expressions for the length and the width of the rug.

b Write an expression for the area of the rug in expanded form.

c What is the area of the rug when $x = 1$?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Statistics and probability

Short-answer questions

1 Identify the sampling method used in each of these surveys.

a Employees are selected from each department in a workplace and surveyed on their job satisfaction.

b Every tenth customer on a store is surveyed regarding their thoughts on customer service in the store.

2 In a survey of 30 people, 18 people drink coffee during the day, 14 people drink tea and 8 people drink both. Let C be the set of people who drink coffee and T the set of people who drink tea.

a Construct a Venn diagram for the survey results.

b Find:

i the number who drink either coffee or tea or both

ii the number who do not drink tea

c If one of the 30 people was randomly selected, find:

i Pr(drinks neither coffee nor tea)

ii Pr(C only)

3 A coin and a four-sided die are tossed.

a Draw a table to show the sample space.

b How many outcomes are in the sample space?

c What is the probability of tossing T4?

d What is the probability of tossing tails and an odd number?

4 A spinner has equal areas for the colours blue, orange and yellow. It is spun twice.

a Draw a tree diagram and list the outcomes.

b Find the probability of orange then blue.

c Find the probability that both outcomes are the same colour.

d Find the probability that blue and yellow show in any order.

5 Two ice creams are randomly selected without replacement from a box containing one vanilla (V), two strawberry (S) and one chocolate (C) flavoured ice creams.

a Draw a tree diagram to show each of the possible outcomes.

b What is the probability of selecting:

i a vanilla and a strawberry flavoured ice cream?

ii two strawberry flavoured ice creams?

iii no vanilla-flavoured ice creams?

6 The data below shows the number of aces served by a player in each of their grand slam tennis matches for the year.

15 22 11 17 25 25 12 31 26 18 32 11 25 32 13 10

a Classify the data as either numerical continuous or numerical discrete.

b Construct a stem-and-leaf plot for the data.

c From the stem-and-leaf plot, find the mode and median number of aces.

d Is the data symmetrical or skewed?

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

7 The frequency table shows the number of visitors, in intervals of 50, to a theme park each day in April.

a Complete the frequency table shown. Round to one decimal place where necessary.

b Construct a frequency histogram.

c **i** On how many days were there fewer than 100 visitors?

ii What percentage of days had between 50 and 199 visitors?

Class interval	Frequency	Percentage frequency
0–	2	
50–	4	
100–	5	
150–	9	
200–		
250–	3	
Total	30	

Multiple-choice questions

1 The probability of rolling a number less than five on a normal six-sided die is:

A $\frac{1}{3}$ **B** 4 **C** $\frac{1}{2}$ **D** $\frac{2}{3}$ **E** 3

2 From the two-way table, Pr(both A and B) is:

A $\frac{1}{5}$

B 4

C $\frac{9}{20}$

D $\frac{1}{4}$

E 16

	A	Not A	Total
B		7	
Not B	5		
Total		11	20

3 In a bag of 40 marbles, 28 are blue.

a The probability of selecting a blue marble is:

A 0.28 **B** 0.4 **C** 0.7 **D** 0.54 **E** 0.75

b If a marble is selected 50 times *with replacement*, the number of blue marbles you would expect to select is:

A 28 **B** 38 **C** 70 **D** 35 **E** 40

4 The median, mean and range of the data set

12 3 1 6 10 1 5 18 11 15

are respectively:

A 5.5, 8.2, 1 – 18 **B** 8, 8.2, 17 **C** 5.5, 8, 17 **D** 8, 8.2, 1 – 18 **E** 8, 74.5, 18

5 The median of this stem-and-leaf plot is:

A 55 **B** 67 **C** 7

D 69 **E** 4

Stem	Leaf
5	1 3 5 5 5 5
6	2 4 5 6 7 7 9
7	1 3 4 8 9
8	3 4
9	1 2 6

5 | 1 means 51

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Semester review 2

Extended-response question

A game at the school fair involves randomly selecting a green ball and a red ball, each numbered 1, 2 or 3.



a List the outcomes in a table.

b What is the probability of getting an odd and an even number?

c Participants win $1 when they draw each ball showing the same number.

i What is the probability of winning $1?

ii If someone wins six times, how many games are they likely to have played?

d The ages of those playing the game in the first hour are recorded and are shown below.

12 16 7 24 28 9 11 17 18 18 37 9 40 16 32 42 14

i Approximately 50% of the participants are below what age?

ii If this data is used as a model for the 120 participants throughout the day, how many would be expected to be aged less than 30?

Introduction to quadratic equations and graphs

Short-answer questions

1 Solve the following quadratic equations.

a $(x+5)(x-3)=0$ **b** $(2x-1)(3x+5)=0$

c $4x^2+8x=0$ **d** $5x^2=45$

e $x^2-9x+14=0$ **f** $8x=-x^2-16$

2 The length of a rectangular swimming pool is x m and its width is 7 m less than its length. If the area occupied by the pool is 120 m^2, solve an appropriate equation to find the dimensions of the pool.

3 Sketch the following graphs showing the y-intercept and turning point, and state the transformations that have taken place from the graph of $y=x^2$.

a $y=x^2+3$ **b** $y=-(x+4)^2$ **c** $y=(x-2)^2+5$

4 Consider the quadratic relation $y=x^2-4x-12$.

a Find the coordinates of the y-intercept.

b Factorise the relation and find the coordinates of the x-intercepts.

c Find the coordinates of the turning point.

d Sketch the graph.

Multiple-choice questions

1 The solution(s) to $3x(x+5)=0$ is/are:

A $x=-5$ **B** $x=0$ or $x=-5$ **C** $x=3$ or $x=5$

D $x=5$ **E** $x=0$ or $x=5$

2 $x^2=3x-2$ is the same as the equation:

A $x^2+3x-2=0$ **B** $x^2-3x+2=0$ **C** $x^2-3x-2=0$

D $-x^2+3x+2=0$ **E** $x^2+3x+2=0$

3 The *incorrect* statement about the graph of $y=2x^2$ is:

A the graph is the shape of a parabola

B the point $(-2,\ 8)$ is on the graph

C its turning point is at $(0,\ 0)$

D it has a minimum turning point


E the graph is wider than the graph of $y=x^2$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

4 The type and coordinates of the turning point of the graph of $y = -(x+3)^2 + 2$ are:

A a minimum at $(3, 2)$
B a maximum at $(3, 2)$
C a maximum at $(-3, 2)$
D a minimum at $(-3, 2)$
E a minimum at $(3, -2)$

5 The graph shown has the equation:



A $y = 4x^2$
B $y = x^2 + 4$
C $y = (x+2)^2$
D $y = x^2 + 2$
E $y = (x-2)^2$

Extended-response question

The flight path of a soccer ball kicked upwards from the ground is given by the equation $y = 120x - 20x^2$, where y is the height of the ball above the ground in centimetres at any time x seconds.

a Find the x-intercepts to determine when the ball lands on the ground.
b Find the coordinates of the turning point and state:
  i the maximum height reached by the ball
  ii after how many seconds the ball reaches this maximum height.
c At what times was the ball at a height of 160 cm?
d Sketch a graph of the path of the ball until it returns to the ground.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Glossary

A

Absolute value A positive value ignoring a negative sign

Acute angle Between 0 and 90 degrees

Adjacent (side) In a right-angled triangle, the side adjacent to (next to) the unknown angle

Algebraic fraction A fraction including pronumerals

Algorithm A sequence of steps that, when followed, lead to the solution of a problem

Alternate angles A pair of angles lying on opposite sides of a transversal but inside two other lines

Angle The difference in direction between two lines

Angle of depression The angle of your line of sight from the horizontal when looking down at an object

Angle of elevation The angle of your line of sight from the horizontal when looking up at an object

Angle sum The total measure of the angles in a plane figure

Area A measure of surface in square units

Array A rectangular table for listing outcomes

Average The central value of a data set

Axis of symmetry A vertical line that divides a parabola into two symmetrical halves

B

Back-to-back stem-and-leaf plot A visual representation of two sets of data that groups the scores and lists them in order horizontally on either side of the stem

Base A number or pronumeral that is being raised to a power

Best buy A cost per unit which represents the best value

Bias A tendancy which influences the outcome

Bimodal A data set with two modes

Binomial An expression with two terms

Binomial product The product of two binomial expressions

C

Capacity The amount of liquid a container can hold


Cartesian plane A plane on which every point is related to a pair of numbers called coordinates

Categorical data Data which can be sorted into groups/categories

Chance The probability of an event occurring

Circle A simple round shape with a centre and radius

Circumference The curved boundary of a circle



Class interval A numerical interval for grouped data

Coefficient A numeral multiplied by a pronumeral

Cointerior angles A pair of angles lying between two lines on the same side of a transversal

Column graph A graph that uses columns to compare data in categories

Commission The proportion of a sales amount earned by an employee

Common factor A number/expression which divides into two or more expressions

Complement A set of outcomes containing the elements that are not in another given set

Complementary angles Angles that sum to 90 degrees

Composite shape A complex shape made up of two or more basic shapes

Congruence statement A statement linking the vertices of congruent figures

Congruent figures Figures that are exactly the same size and shape

Constant term The part of an equation or expression without any pronumerals

Convex (polygon) A polygon with all interior angles less than 180 degrees

Coordinate pair (coordinates) An ordered pair written in the form (x, y) that states the location of a point on the Cartesian plane

Corresponding angles A pair of angles sitting in similar positions after a transversal cuts two or more lines

Corresponding sides Sides that are in the same position in two or more shapes

Cosine The ratio of the length of the adjacent side to the length of the hypotenuse

Cost price The price at which goods have been bought by a retailer

Critical digit The digit that determines whether you round the previous digit up or down

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Cross-section The plane figure formed when you slice a solid figure parallel to one of its surfaces

Cube (number) The product of a number multiplied by itself twice

Cylinder A solid with a uniform circular cross-section

D

Decimal A number expressed using a system of counting based on the number ten: Three-fifths as a decimal is 0.6

Decimal places The number of digits to the right of the decimal point

Deductions Amounts of money taken from gross income

Denominator Parts in the whole. The part of a fraction that sits below the dividing line.

Dependent variable The value that changes in response to changes made in the independent variable


Diagonal A line segment across a shape joining two vertices

Diameter A line passing through the centre of a circle with its end points on the circumference

Difference of two squares When one square term is subtracted from another

Dilation A stretch in a given direction

Direct proportion The relationship between two quantities that increase or decrease at the same rate



Discount An amount subtracted from a price

Distributive law Adding numbers in brackets, *then* multiplying the total, gives the same answer as multiplying each number in the brackets separately first, *then* adding the products

Dot plot A statistical graph which uses dots piled vertically to illustrate frequency

Double time A pay rate of overtime that is 2 times the normal hourly rate

E

Endpoints The coordinates of the points at the end of an interval

Enlargement A transformation that changes the size of a figure without changing its shape

Equation A statement that two expressions have the same value

Equilateral triangle A triangle with three equal sides and all angles 60 degrees

Equivalent equations Two equations that produce the same solution

Equivalent fractions Fractions with the same value

Error The difference between the measured value and the actual value

Estimate An Approximation of the actual value

Evaluate To work out the numerical value

Event Outcomes resulting from an experiment

Expand Remove grouping symbols (such as brackets)

Expanded form An expression without brackets

Expected number of occurrences The expected number of favourable outcomes from an experiment

Experiment A situation involving chance or probability trials

Experimental probability Probability based on measuring the outcomes of trials

Exponent A number that shows how many times another number (the base) is to be multiplied by itself. Also called the Index or power.

Expression A collection of mathematical terms containing no equals sign

Exterior angle The angle formed between the extended side and the adjacent side of a polygon

F

Factor A number or expression that divides in without a remainder

Factorise To write an expression as a product

Favourable outcome An outcome that is desired from an experiment

Formula A rule for finding the value of one quantity given the values of others

Fraction A number that results from dividing one whole number by another


Frequency table A table summarising data by showing all possible scores from lowest to highest in one column, and the frequency of each score in another column

G

Gradient The steepness of a straight line or interval

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Gradient–intercept form The equation of a straight line, written with y as the subject of the equation



Gross income Total income before any deductions are made

Grouping in pairs A strategy for factorising by first grouping in pairs and factorising each pair


H

Highest common factor (HCF) The largest number/expression which divides into two or more expressions

Histogram A special type of column graph with no gaps between the columns

Horizontal Parallel to the ground, at right angles to the vertical

Hypotenuse The longest side of a right-angled triangle



I

Image The resulting figure after a transformation

Improper fraction A fraction where the numerator is larger than or equal to the denominator

Income tax An amount paid to the government by people earning an income

Independent variable The value that is deliberately controlled when collecting data

Index The number of times a factor is repeated under multiplication

Index form A method of writing numbers that are multiplied by themselves

Inequality An equation that includes a $<, \leqslant, >$ or $\geqslant$

Integer A number in the infinite set $\{\ldots, -3, -2, -1, 0, 1, 2, 3, \ldots\}$

Intercept form A rule in factorised form

Inverse Opposite in relation to something. The inverse of addition is subtraction.

Inverse cosine The angle of which the cosine ratio is given

Inverse sine The angle of which the sine ratio is given

Inverse tangent The angle of which the tangent ratio is given

Invented (concave down) Curving downwards

Irrational number A real number that cannot be expressed as a fraction

Isosceles triangle A triangle with a pair of equal sides and a pair of equal angles

K

Kite A quadrilateral with two pairs of adjacent equal sides

L

Length A measure of distance

Like terms Terms with the exact same pronumerals including powers

Limits of accuracy The smallest and largest possible values that a measurement could be

Line A continuous set of points without breadth or thickness

Line segment A section of a straight line

Linear relation A set of ordered pairs that give a straight line

Litres (L) A unit of capacity where $1\text{ L} = 1000\text{ mL} = 1000$ cubic centimetres

Long-run proportion The experimental probability for a sufficiently large number of trails

Loss The amount of money lost by selling something for less than its cost

Lowest common denominator (LCD) The lowest common multiple of the denominators of a set of fractions

M

Mark-up An amount added by the retailer to the cost price of goods

Maximum The largest value

Mean ($\bar{x}$) The average value of the scores in a set of data

Median The middle score (or mean of two middle scores) when a set of data is arranged in order

Midpoint The point on a line segment that is an equal distance from each of the end points of the segment

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Millilitres (mL) A unit of capacity where $1\text{ L} = 1000\text{ mL}$

Minimum The smallest value

Mixed number A combination of a whole number and a fraction

Mode The most frequently occurring value in a set of data

Monic quadratic trinomial A quadratic expression with the coefficient of the squared term equal to 1

N

Negative To be less than zero

Net A diagram showing how the plane faces of a solid are joined to each other

Net income Income that remains after deductions have been made

Non-convex (polygon) A polygon with at least one reflex angle

Null Factor Law If $p \times q = 0$ then $p = 0$ or $q = 0$


Numerator Parts of the whole. The part of a fraction that sits above the dividing line.

Numerical data Data which are numbers

O

Obtuse angles Between 90 and 180 degrees

Opposite (side) In a right-angled triangle, the side opposite the unknown angle



Origin The point $(0, 0)$ where the x- and y-axes of the Cartesian plane intersect

Outcome One of the possibilities resulting from a chance experiment

Outlier Any value that is much larger or much smaller than the rest of the data in a set

Overtime Time worked in addition to normal working hours

P

Parabola A smooth U-shaped curve which is the graph of a quadratic relation

Parallel Lines that are side by side and always have the same distance between them

Parallelogram A quadrilateral with two pairs of parallel sides

Per annum (p.a.) Annually; that is, per year

Percentage A number expressed as a part of 100

Percentage error An error expressed as a percentage

Percentage frequency histogram A special type of histogram that shows the frequencies of the data as percentages of the total

Perfect square An algebraic expression that can be written as a single square

Perimeter The total distance (length) around the outside of a figure

Perpendicular To be at right angles (90 degrees)

Pi A special number ($3.14159\ldots$) which connects a circle's radius with its circumference and area

Pie chart A circular graph showing sectors which represent statistical categories

Polygon A two-dimensional shape where three or more straight lines are joined together to form a closed figure

Population The complete group to be investigated

Positive To be greater than zero

Power An expression that includes an index

Prime A number with two factors

Principal An amount of money invested in or loaned to a person or organisation

Prism A solid with a uniform cross-section and the remaining sides parallelograms

Probability A measure of the likelihood that an event will occur

Product Involving multiplication

Profit The amount of money made by selling something for more than its cost

Pronumeral A letter or symbol used to represent a number (also called a variable)

Proper fraction A fraction where the numerator is smaller than the denominator

Pythagoras' theorem In a right-angled triangle, the square of the hypotenuse is equal to the sum of the squares on the other two sides

Q

Quadrant A quarter circle or one quarter of a Cartesian plane

Quadratic equation An equation involving a quadratic expression

Quadratic relation An equation or rule with a squared term but no term with a higher power

Quadrilateral A four-sided plane figure with straight sides

ISBN 978-1-009-59315-1
© Greenwood et al. 2025
Photocopying is restricted under law and this material must not be transferred to another party.

R

Radius The distance from the centre of a circle to its outside edge

Random sampling Where each member of the population has an equal chance of being selected

Range The difference between the highest score and the lowest score in a set of data

Rate The number of units of one quantity for each single unit of another quantity

Ratio A way of comparing quantities of the same unit, separated by a colon

Rational number A real number that can be expressed as a fraction

Real numbers Any positive or negative number or zero

Reciprocal An inverted fraction

Rectangle A parallelogram with two pairs of equal sides and all angles 90 degrees

Recurring decimal A decimal with an infinite repeating pattern

Reflection A mirror image over a given reflection line

Reflex angle Between 180 and 360 degrees

Regular polygon A polygon with all sides equal and all angles equal

Repayment An amount paid to a financial institution at regular intervals to repay a loan, with interest included

Retainer A set weekly or monthly fee paid to an employee

Revolution 360 degrees

Rhombus A parallelogram with four equal sides

Right-angled Containing an angle of 90°

Right prism A solid with a uniform cross-section, and remaining sides are rectangles

Rise The change in value in the vertical direction

Run The change in value in the horizontal direction from left to right

S

Salary A fixed agreed yearly amount that an employee earns

Sample A selection from the population

Sample space The list of all the possible outcomes of an event

Scale factor The number by which you multiply each side length to enlarge or reduce the size of a shape

Scalene triangle A triangle with three different side lengths

Scientific notation A method used to express very large and very small numbers

Sector A portion of a circle formed by an arc and two radii

Selling price The price for which a retailer sells goods to a buyer

Semicircle Half a circle

Significant figure A digit that indicates how accurate a number is

Similar figures Figures of the same shape but not the same size

Simple interest The percentage rate per year that is paid on a loan or investment amount

Sine The ratio of the length of the opposite side to the length of the hypotenuse

Skewed data Scores in a set of data that are unevenly distributed around either side of the mean and the median

Solid A three-dimensional shape with volume

Solution The answer

Solution set A group of numbers which are the solution to an equation or inequality

Solve To find the answer

Square (geometry) A parallelogram with four equal sides and all angles 90 degrees

Square (number) A number multiplied by itself

Square root The number that when squared produces the number under the square root sign

Standard form (equation) An equation with a zero on one side

Straight angle 180 degrees

Stem-and-leaf plot A visual representation of data that groups the scores in a set of data and lists them in order horizontally

Stratified sampling Where members of the population are divided into subgroups before being surveyed

Subject The pronumeral that is alone on the left hand side (LHS) in an equation or formula

Substitute To replace pronumerals with numerical values

Substitution To replace a pronumeral with a number

Supplementary angles Angles that sum to 180 degrees

Surd An irrational number expressed using a radical sign (root)

Survey A set of questions or other activity designed to collect data

Symmetrical data A distribution of data that is balanced on either side of the mean and the median

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Systematic sampling Where members of the population are selected at regular time intervals

T

Tangent The ratio of the length of the opposite side to the length of the adjacent side

Taxable income Gross income minus tax deductions

Term A combination of numbers and pronumerals connected with only multiplication and division

Time and a half A pay rate of overtime that is 1.5 times the normal hourly rate

Total surface area The number of square units needed to cover the outside of a solid

Translation A shift in a particular direction

Transpose To rearrange a formula to make a different variable the subject

Transversal A line that cuts two or more lines

Trapezium A quadrilateral with at least one pair of parallel sides

Tree diagram A diagram used to show all the possibilities when several different options are available

Triangle A three-sided shape

Trigonometric ratio The ratios that relate the sides and angles of right-angled triangles

Turning point The point at which a curve changes direction

Two-step experiment A multi-stage experiment involving two steps

Two-way table A table used to organise, display and compare two sets of outcomes

U

Uniform With a constant cross-section

Unitary method A way of solving a problem by reducing one of the units to 1

Upright (concave up) Curving upwards

V

Variable An unknown, which can take on any value

Venn diagram A diagram using overlapping circles to organise and show two or more sets of outcomes

Vertical Such that the top is directly above the bottom, at right angles to the horizontal

Vertically opposite Equal angles opposite at a point formed by intersecting lines

Volume The amount of three-dimensional space inside an object

W

Wage Earnings paid to an employee based on an hourly rate

With replacement Selecting items and replacing them before the next selection

Without replacement Selecting items and not replacing them before the next selection

X

***x*-axis** The horizontal axis of the Cartesian plane

***x*-coordinate** The first coordinate of an ordered pair, describing the horizontal position from the origin

***x*-intercept** The point at which a line or curve cuts the x-axis

Y

***y*-axis** The vertical axis of the Cartesian plane

***y*-coordinate** The second coordinate of an ordered pair, describing the vertical position from the origin

***y*-intercept** The point at which a line or curve cuts the y-axis

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Answers

Chapter 1

Warm-up quiz

1 Addition: **a** Sum, **b** Total, **i** Add, **l** More than
Subtraction: **c** Less than, **g** Take away, **h** Difference, **k** Minus
Multiplication: **d** Lots of, **e** Product, **j** Times
Division: **f** Into, **m** Quotient

2 **a** 4 **b** 32 **c** 4 **d** 52
e 74 **f** 5 **g** 9 **h** 40

3 **a** 56 **b** 100 **c** 0 **d** 133
e 63 **f** 80 **g** 1020 **h** 53

4 **a** −12 **b** −3 **c** 3 **d** 0
e −1 **f** −1 **g** 0 **h** −12

5 **a** 3 **b** 3 **c** − **d** +

6 **a** 26 073 260 **b** 26 073 300
c 26 073 000 **d** 26 000 000

7 **a** 50 **b** 5 **c** 5000
d $\frac{5}{10}$ **e** $\frac{5}{100}$ **f** $\frac{5}{1000}$

8 **a** 2.654, 2.645, 2.564, 2.465
b 0.654, 0.564, 0.456, 0.0456

9 **a** 7.99 **b** 10.11 **c** 7.11

10 **a** 1.4 **b** 0.06 **c** 3.68
d 16.38 **e** 3.7 **f** 180

11 **a** 34.5 **b** 374 000
c 0.03754 **d** 0.000037754

12 **a** 6 **b** 12 **c** 30 **d** 1

13 **a** 15 **b** 12 **c** 10

14 **a** $\frac{5}{7}$ **b** $1\frac{1}{4}$ **c** 8 **d** $\frac{3}{8}$

1A

Now you try

Example 1

a −3 **b** −10

Example 2

4

Example 3

−8

Exercise 1A

1 **a** iii **b** v **c** iv **d** i **e** ii

2 **a** iv **b** ii **c** v **d** i **e** iii

3 **a** T **b** T **c** F

4 **a** −2 **b** 2 **c** −3 **d** 4 **e** −4
f −5 **g** −2 **h** 0 **i** −6

5 **a** 3 **b** 2 **c** 1 **d** 0 **e** −1
f −9 **g** −1 **h** −17 **i** −10 **j** 9
k −2 **l** 11 **m** −7 **n** −18 **o** −1

6 **a** 5 **b** 20 **c** −20 **d** 0 **e** −7
f 6 **g** −24 **h** 5 **i** −15

7 **a** 4 **b** −4 **c** −4 **d** −9
e −32 **f** −38 **g** −189 **h** −24
i −18 **j** 0 **k** −805 **l** −57
m −167 **n** 142 **o** −804

8 **a** −2 **b** 4 **c** 13 **d** 9 **e** 24
f 0 **g** −28 **h** −81 **i** 180 **j** 2
k −3 **l** 240 **m** 755 **n** 94 **o** 10

9 **a** −27 **b** −3 **c** 10 **d** 26
e 0 **f** −24 **g** −91 **h** 125

10 **a** 2 **b** 11 **c** −3
d −1 **e** −7 **f** 11

11 **a** −8 **b** 4 **c** 13
d 124 **e** −19 **f** 0
g −22 **h** 98 **i** −26

12 −12°C

13 **a**

−8	6	−4
2	−2	−6
0	−10	4

b Example answer (other answers are possible):

2	−13	8
5	−1	−7
−10	11	−4

c

5	−2	7	3
0	10	−3	6
−1	4	2	8
9	1	7	−4

1B

Now you try

Example 4

a 32 **b** −35

Example 5

a 64 **b** −27

Example 6

a −5 **b** 4

Example 7

12

Essential Mathematics Core for the Victorian Curriculum Year 9 ISBN 978-1-009-59315-1 © Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Exercise 1B

1 **a** Positive **b** Negative
c Positive **d** Negative
e Negative
2 **a** Negative **b** Negative
c Negative **d** Positive
e Positive **f** Positive
3 **a** T **b** F **c** T
4 **a** −48 **b** −20 **c** −72 **d** 72
e 45 **f** −90 **g** −600 **h** 81
i −144 **j** −143 **k** −143 **l** −34
m 72 **n** 90 **o** −108
5 **a** 36 **b** 64 **c** 144 **d** 9 **e** 100
f 169 **g** −8 **h** −1 **i** −64
6 **a** −9 **b** 6 **c** 10 **d** −10 **e** 3
f 5 **g** 2 **h** −1 **i** −8 **j** 25
k −104 **l** 5 **m** −37 **n** −214 **o** 13
7 **a** 16 **b** −26 **c** −12 **d** −14 **e** −20
f −28 **g** −72 **h** −8 **i** −90 **j** 10
k −24 **l** −18 **m** 9 **n** −26
8 **a** 7 **b** −32 **c** −20 **d** −4 **e** −1
f 6 **g** −9 **h** −5 **i** −15
9 −5 and 2
10 **a** 12 and −12 **b** 8 and −8 **c** 100 and −100
11 Two negatives with × and ÷ produce a positive.
12 **a** Positive **b** Positive **c** Negative
d Positive **e** Negative **f** Positive
13 **a** −2 **b** −38 **c** −8 **d** 27
e 1 **f** 24 **g** 21 **h** 0

1C

Now you try

Example 8

a 107.39 **b** 0.03 **c** 2.80

Example 9

a 3500 **b** 1800 **c** 0.0023

Example 10

10, reasonable as real answer is 9.5

Exercise 1C

1 **a** 3 **b** 1 **c** 2 **d** 0
2 **a** 2 **b** 4 **c** 1 **d** 4
3 **a** Down **b** Up **c** Up
4 **a** 32 100 **b** 432 **c** 5.89
d 0.443 **e** 0.00197
5 **a** 17.96 **b** 11.08 **c** 72.99
d 47.86 **e** 63.93 **f** 23.81
g 804.53 **h** 500.57 **i** 821.27
j 5810.25 **k** 1005.00 **l** 2650.00
6 **a** 2400 **b** 35 000 **c** 0.060
d 34 **e** 110 000 **f** 0.0025
g 2.1 **h** 0.71 **i** 4700
j 59 000 **k** 0.46 **l** 1.1
7 **a** 1.46, 1.5 **b** 0.09, 0.094
c 23.71, 24 **d** 0.01, 0.0078
e 100.47, 100
8 **a** 7 **b** 73 **c** 130 **d** 36 200
9 **a** 30 000 **b** 200 **c** 0.05 **d** 0.0006
e 5000 **f** 900 **g** 0.9 **h** 0.0003 **i** 1
10 **a** 3600, 3693 **b** 760, 759.4
c 4000, 4127.16 **d** 3000, 3523.78
e 0, 0.72216 **f** 4, 0.716245
g 0.12, 0.1186 **h** 0.02, 0.02254
i 0.2, 0.2
11 **a** A: 54.3, B: 53.8, 0.5 **b** A: 54.28, B: 53.79, 0.49
c A: 54, B: 54, 0 **d** A: 50, B: 50, 0
12 0.143 tonnes
13 As magnesium in this case would be zero if rounded to two decimal places rather than two significant figures
14 2.14999 is closer to 2.1 correct to one decimal place, ∴ round down
15 **a** **i** 8 **ii** 1 **iii** 8
b 0.1428571428571
c **i** 4 **ii** 2 **iii** 8
16 Teacher to check

1D

Now you try

Example 11

$\frac{5}{8}$, 0.32 and $-\sqrt{25}$ are rational numbers

Example 12

a $3\frac{2}{5}$ **b** $\frac{9}{4}$

Example 13

2.3125

Example 14

$0.41\dot{6}$

Example 15

a $\frac{9}{20}$ **b** $3\frac{9}{40}$

Example 16

$\frac{4}{9} > \frac{5}{12}$

Exercise 1D

1 **a** T **b** F **c** T **d** T
2 **a** Proper **b** Mixed number
c Improper **d** Proper
3 **a** $\frac{7}{10}$ **b** $\frac{3}{10}$ **c** $\frac{23}{100}$ **d** $\frac{9}{100}$
4 a, b, c, d, e, f, g, i, k
5 **a** **i** $2\frac{2}{5}$ **ii** $2\frac{5}{6}$ **iii** $5\frac{3}{4}$ **iv** $1\frac{1}{8}$
b **i** $\frac{16}{5}$ **ii** $\frac{44}{7}$ **iii** $\frac{24}{5}$ **iv** $\frac{83}{8}$
6 **a** 2.75 **b** 0.35 **c** 3.4 **d** 1.875
e 2.625 **f** 3.8 **g** $2.3\overline{125}$ **h** 0.25
7 **a** $0.\overline{27}$ **b** $0.\dot{7}$ **c** $1.\overline{285714}$
d $0.58\dot{3}$ **e** $1.\dot{1}$ **f** $3.8\dot{3}$
g $7.2\dot{6}$ **h** $2.\overline{63}$
8 **a** $\frac{7}{20}$ **b** $\frac{3}{50}$ **c** $3\frac{7}{10}$ **d** $\frac{14}{25}$
e $1\frac{7}{100}$ **f** $\frac{3}{40}$ **g** $3\frac{8}{25}$ **h** $7\frac{3}{8}$
i $2\frac{1}{200}$ **j** $10\frac{11}{250}$ **k** $6\frac{9}{20}$ **l** $2\frac{101}{1000}$
9 **a** $\frac{5}{6}$ **b** $\frac{13}{20}$ **c** $\frac{7}{10}$ **d** $\frac{5}{12}$
e $\frac{7}{16}$ **f** $\frac{11}{14}$ **g** $\frac{19}{30}$ **h** $\frac{11}{27}$
10 **a** $\frac{9}{20}$ **b** $\frac{3}{20}$ **c** $\frac{32}{45}$ **d** $\frac{23}{75}$
11 **a** $\frac{11}{6}, \frac{7}{3}$ **b** $\frac{2}{5}, \frac{2}{15}$ **c** $\frac{11}{12}, 1$ **d** $\frac{5}{7}, \frac{11}{14}$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

12 Weather forecast

13 **a** $\frac{3}{5}$ **b** $\frac{5}{9}$ **c** $\frac{8}{13}$ **d** $\frac{23}{31}$

14 **a** 31, 32 **b** 36, 37, 38, …, 55 **c** 4, 5

15 **a** $\frac{8}{9}$ **b** $1\frac{2}{9}$

c $\frac{81}{99} = \frac{9}{11}$ **d** $3\frac{43}{99}$

1E

Now you try

Example 17

a $1\frac{1}{7}$ **b** $\frac{2}{5}$

Example 18

$\frac{19}{21}$

Example 19

a $4\frac{3}{8}$ **b** $\frac{8}{15}$

Exercise 1E

1 **a** 12 **b** 6 **c** 15

d 49 **e** 15 **f** 35

2 **a** T **b** F

3 **a** $\frac{3}{10} + \frac{4}{10} = \frac{7}{10}$ **b** $\frac{4}{8} - \frac{3}{8} = \frac{1}{8}$

c $1\frac{2}{4} + \frac{1}{4} = 1\frac{3}{4}$

4 **a** 6 **b** 63 **c** 30

d 8 **e** 33 **f** 60

5 **a** $\frac{3}{5}$ **b** $\frac{5}{8}$ **c** $\frac{7}{10}$ **d** $\frac{4}{7}$

e $\frac{5}{17}$ **f** $\frac{1}{5}$ **g** $\frac{10}{6} = 1\frac{2}{3}$ **h** $\frac{10}{8} = 1\frac{1}{4}$

i $\frac{7}{5} = 1\frac{2}{5}$ **j** $1\frac{4}{5}$ **k** $2\frac{1}{3}$ **l** $\frac{1}{2}$

m $\frac{2}{5}$ **n** $4\frac{1}{2}$ **o** $1\frac{2}{5}$ **p** $6\frac{1}{2}$

6 **a** $\frac{5}{8}$ **b** $\frac{1}{3}$ **c** $\frac{1}{2}$ **d** $\frac{1}{2}$

e $\frac{9}{10}$ **f** $\frac{7}{12}$ **g** $\frac{7}{10}$ **h** $\frac{3}{20}$

i $\frac{11}{24}$ **j** $1\frac{1}{6}$ **k** $1\frac{8}{21}$ **l** $\frac{17}{24}$

m $\frac{11}{36}$ **n** $\frac{4}{7}$ **o** $\frac{19}{100}$ **p** $\frac{19}{90}$

7 **a** $3\frac{5}{6}$ **b** $\frac{7}{12}$ **c** $2\frac{4}{5}$ **d** $2\frac{7}{12}$

e $6\frac{3}{4}$ **f** $3\frac{5}{6}$ **g** $\frac{1}{8}$ **h** $1\frac{5}{12}$

i $1\frac{7}{10}$ **j** $\frac{11}{12}$ **k** $4\frac{7}{10}$ **l** $1\frac{7}{12}$

8 $\frac{7}{8}$ tonnes

9 $5\frac{29}{56}$ tonnes

10 **a** $4\frac{19}{20}$ litres **b** 33

11 $\frac{1}{20}$

12 $\frac{5}{6}$, problem is the use of negatives in the method, since $\frac{1}{3} < \frac{1}{2}$

13 **a** **i** $\frac{1}{8} + \frac{3}{4}$ **ii** $\frac{2}{3} - \frac{5}{12}$ **iii** $\frac{5}{6} - \frac{1}{2}$

b $\frac{1}{2} + \frac{2}{3} + \frac{5}{6} = 2$

Progress quiz

1 **a** -8 **b** -14 **c** 25

d 8 **e** 4 **f** -3

2 **a** 42 **b** -60 **c** -8

d 8 **e** -64 **f** 81

3 **a** -9 **b** 19 **c** -7

4 **a** **i** 24.19 **ii** -6.14 **iii** 7.60

b **i** 3.3 **ii** 240 **iii** 7900 **iv** 0.0079

5 **a** 100, 121 **b** 89.8, 93.17302

6 **a** 0.625 **b** 3.32 **c** $0.58\dot{3}$ **d** $2.\overline{63}$

7 **a** $\frac{4}{5}$ **b** $\frac{9}{25}$ **c** $\frac{1}{250}$ **d** $4\frac{41}{200}$

8 $\frac{3}{8}, \frac{5}{12}, \frac{11}{24}$

9 **a** $\frac{5}{8}$ **b** $\frac{2}{21}$ **c** $3\frac{11}{20}$ **d** $1\frac{9}{20}$

10 $3\frac{7}{12}$

1F

Now you try

Example 20

a $\frac{12}{35}$ **b** $\frac{5}{18}$

Example 21

$4\frac{1}{5}$

Example 22

a $\frac{3}{2}$ or $1\frac{1}{2}$ **b** $\frac{9}{5}$ or $1\frac{4}{5}$

Exercise 1F

1 **a** T **b** F **c** T

2 **a** 12 **b** 4 **c** 12 **d** 180

3 **a** $\frac{4}{3}$ **b** 7 **c** $\frac{1}{6}$ **d** $\frac{3}{5}$

4 **a** $\frac{3}{20}$ **b** $\frac{4}{15}$ **c** $\frac{15}{28}$ **d** $\frac{1}{6}$

e $\frac{1}{20}$ **f** $\frac{2}{27}$ **g** $\frac{10}{17}$ **h** $\frac{3}{7}$

i $\frac{2}{5}$ **j** $\frac{2}{3}$ **k** $\frac{2}{5}$ **l** $\frac{4}{5}$

m $\frac{3}{4}$ **n** $\frac{2}{3}$ **o** $\frac{2}{9}$ **p** $\frac{3}{16}$

5 **a** $\frac{1}{2}$ **b** $1\frac{1}{4}$ **c** $2\frac{1}{10}$ **d** 2 **e** $4\frac{1}{5}$

f $\frac{6}{7}$ **g** $6\frac{1}{4}$ **h** 4 **i** $2\frac{11}{12}$ **j** 6

6 **a** $\frac{20}{21}$ **b** $1\frac{1}{8}$ **c** $\frac{45}{56}$ **d** $\frac{27}{28}$

e $1\frac{1}{3}$ **f** $1\frac{1}{2}$ **g** 6 **h** $\frac{7}{8}$

i 18 **j** 9 **k** $\frac{1}{10}$ **l** $\frac{4}{27}$

7 **a** $3\frac{1}{3}$ **b** 4 **c** $\frac{1}{6}$

d $1\frac{1}{2}$ **e** $\frac{7}{8}$ **f** 27

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

8 **a** \$10 **b** \$3.50 **c** \$6
d \$18 **e** \$6400 **f** 90c

9 **a** $\frac{2}{7}$ **b** $\frac{8}{15}$ **c** 16

10 $\frac{3}{12}$ hour = 15 minutes

11 160 km

12 90 minutes each

13 **a** **i** 0.5 **ii** 0.2 **iii** 0.16129…
iv $0.\dot{6}$ **v** -1
b 3.39
c 25
d Numbers greater than zero but less than one
e 1
f 0

1G

Now you try

Example 23

5 : 3

Example 24

a 25 : 16 **b** 26 : 15

Example 25

7 : 40

Example 26

\$350

Exercise 1G

1 **a** No **b** Yes **c** Yes **d** No

2

a	1 : 3	1 : 4
b	2 : 3	2 : 5
c	1 : 2	1 : 3

3 **a** 5 : 1 **b** 1 : 2 **c** 2 : 1
d 10 : 1 **e** 5 : 16

4 **a** 1 : 5 **b** 2 : 5 **c** 4 : 5 **d** 1 : 10
e 5 : 6 **f** 1 : 3 **g** 3 : 1 **h** 2 : 3
i 4 : 3 **j** 4 : 1 **k** 1 : 2 **l** 3 : 1
m 5 : 14 **n** 1 : 2 : 4 **o** 3 : 6 : 2

5 **a** 1 : 2 **b** 3 : 41 **c** 5 : 32
d 1 : 3 **e** 1 : 5 **f** 10 : 3
g 40 : 1 **h** 1500 : 1 **i** 20 : 7

6 **a** 3 : 4 **b** 8 : 3 **c** 9 : 20
d 45 : 28 **e** 3 : 14 **f** 22 : 39

7 **a** 1 : 10 **b** 1 : 5 **c** 2 : 3 **d** 7 : 8
e 25 : 4 **f** 1 : 4 **g** 1 : 4 **h** 24 : 5
i 4 : 20 : 5 **j** 4 : 3 : 10 **k** 5 : 72 **l** 3 : 10 : 40

8 **a** \$100, \$400 **b** \$16, \$20
c 24 kg, 64 kg **d** \$56, \$40
e \$200, \$300 **f** 750 g, 1250 g
g \$70, \$30 **h** \$300, \$300
i \$14, \$49, \$7 **j** 336 g, 84 g

9 126 g

10 120

11 \$3000, \$1200, \$1800 respectively

12 108 L

13 Amy : \$2400, Belinda : \$3600, Candice : \$1200, Diane : \$4800

14 **a** **i** 100 mL **ii** 200 mL
b **i** 250 mL **ii** 270 mL
c **i** 300 mL **ii** 1 : 4
d **i** 1 : 3 **ii** 7 : 19 **iii** 26 : 97 **iv** 21 : 52
e Jug 3 and 4

1H

Now you try

Example 27

a \$30/h **b** 1200 revs/min

Example 28

a 60 km/h **b** 4 hours 10 minutes

Example 29

The 3 L is cheaper

Exercise 1H

1 **a** 60 km/h **b** \$84/h **c** 60 km/h

2 **a** \$4 **b** \$3 **c** \$2.50

3 **a** 10 m/s **b** 60 km/h **c** 14 km/h

4 B

5 **a** \$28/h **b** \$50/h **c** 6 kg/min
d \$7/kg **e** 4 runs/over **f** 89 pts/game
g 19 cm/y **h** 14 m/s **i** 92 beats/min
j 8 mL/s

6 **a** 70 km/h **b** 15 km/h **c** 4 km/h
d 54 km/h **e** 80 km/h

7 **a** 13 hours **b** 16 hours 15 minutes

8 **a** 3 kg deal **b** Red delicious
c 2.4 L **d** 0.7 GB

9 C

10 **a** 55 km **b** $16\frac{1}{2}$ km **c** $5\frac{1}{2}$ km

11 **a** Coffee A: \$3.60, coffee B: \$3.90. Therefore, coffee A is the best buy.
b Pasta A: \$1.25, pasta B: \$0.94. Therefore, pasta B is the best buy.
c Cereal A: \$0.37, cereal B: \$0.40. Therefore, cereal A is the best buy.

12 **a** 40 sheep/h **b** 220 sheep
c 25 hours


13 **a** 1 080 000 000 km/h **b** 36 000 000 km
c $8\frac{5}{18}$ min

14 **a**

Time in hours	0	1	2	3	4	5	6	7	8
Distance in km	0	20	40	60	80	100	120	140	160

b 20 km/h

c, e



d Straight line, speed was constant

Maths@Work: Cooks and chefs

1 **a** 3 : 4 **b** 9 egg whites
c 5 egg whites **d** 20 servings
e 216 g **f** 45 g

ISBN 978-1-009-59315-1
© Greenwood et al. 2025
Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

2 **a** 3 loaves **b** 6 cartons **c** 3 : 5
d 1350 grams **e** 6 : 5
f Recipe for 1 large loaf of banana bread:
3 cups of plain flour
1½ cups sugar
1 teaspoon of bicarb of soda
375 grams of melted butter
3 eggs
$3\frac{1}{3}$ mashed bananas

3 **a** **i** 2 : 1 **ii** 1 : 25 **iii** 2 : 1
b 8 eggs
c $2\frac{1}{2}$ cups

4 3.5

5 **a** **i** 150 mL oil **ii** 2.5 cups flour
iii 250 g chocolate
b **i** 4 eggs **ii** 8 tablespoons apple sauce
iii 640 g brown sugar

Puzzles and games


1 $1 + 2 + 3 + 4 + 5 + 6 + 7 + (8 \times 9) = 100$

2 48 ways

3

1. **9**	2. **2**	3. **5**	**3**
4. **3**	**5**	5. **5**	6. **6**
7. **9**	8. **9**	9. **2**	**4**
10. **1**	**2**	**8**	**0**

4 **a** **i** $1 + 2 + 3 = 6$ **ii** $28\,(1 + 2 + 4 + 7 + 14 = 28)$
b **i**

1	2	3	4	5	6
					
1	3	6	10	15	21

ii 28, 36
c **i** 0, 1, 1, 2, 3, 5, 8, 13, 21, 34
ii −21, 13, −8, 5, −3, 2, −1, 1, 0, 1, …, ∴ −21, −8, −3, −1

Checklist Answers

1 **a** −10 **b** −10
2 −5
3 10
4 **a** 28 **b** −8
5 **a** 16 **b** −125
6 −11
7 **a** 32.24 **b** 3.71 **c** 1.50
8 **a** 7800 **b** 0.038
9 $1000 - 500 \times 0.05 = 975$ calculator = 1040.688
10 rational are: $1.4, \sqrt{36}, -\frac{3}{5}$
11 **a** $3\frac{3}{5}$ **b** $\frac{17}{7}$
12 **a** 2.875 **b** $0.\dot{3}\hat{6}$
13 **a** $\frac{9}{25}$ **b** $\frac{443}{200}$
14 $\frac{7}{9} > \frac{11}{15}$
15 **a** $\frac{5}{17}$ **b** $2\frac{3}{4}$
16 $\frac{43}{30} = 1\frac{13}{30}$
17 **a** $\frac{55}{12} = 4\frac{7}{12}$ **b** $\frac{19}{21}$
18 **a** $\frac{5}{24}$ **b** $\frac{4}{9}$
19 **a** $\frac{7}{6}$
20 $1\frac{5}{7}$
21 **a** 5 : 4 **b** 10 : 7 **c** 25 : 18
22 4 : 25
23 $180
24 **a** 60 L/h **b** 72 strokes/min
25 60 km/h there and 2 hours 24 minutes return trip
26 7 Litre is best buy

Short-answer questions

1 **a** 8 **b** −45 **c** 270
d −64 **e** 6 **f** 8
2 **a** 21.5 **b** 29 100 **c** 0.153 **d** 0.00241
3 **a** 14.98 **b** 0.71 **c** 2.00
4 **a** 2.125 **b** $0.8\dot{3}$ **c** $1.\overline{857142}$
5 **a** $\frac{3}{4}$ **b** $1\frac{3}{5}$ **c** $2\frac{11}{20}$
6 **a** $\frac{4}{3}$ **b** $\frac{5}{1} = 5$ **c** $\frac{1}{8}$ **d** $\frac{7}{9}$
7 **a** $\frac{1}{2}$ **b** $2\frac{1}{6}$ **c** $\frac{7}{24}$
d 2 **e** $3\frac{3}{4}$ **f** $2\frac{19}{28}$
8 **a** 5 : 2 **b** 16 : 9 **c** 75 : 14
9 **a** 50, 30 **b** 25, 55 **c** 10, 20, 50
10 **a** Store A: $2.25/kg; store B: $2.58/kg. Store A is the best buy.
b Store A: 444 g/$; store B: 388 g/$
11 $132, $198, $330
12 **a** 32 **b** 7 **c** 192
d $14.80 **e** 82.5
13 **a** 156 km/h **b** 36 km/h
14 $6000

Multiple-choice questions

1 D **2** B **3** C **4** A **5** E
6 E **7** D **8** E **9** C **10** B
11 C **12** C

Extended-response questions

1 **a** 12 **b** 10 adults, 18 children **c** 5 : 9
2 **a** 85.6 km/h **b** 14 hours
c 61.1 km/h **d** 611 km

Chapter 2

Warm-up quiz

1 **a** $\frac{3}{25}$ **b** $\frac{1}{5}$ **c** $\frac{7}{20}$
d $\frac{3}{4}$ **e** $\frac{3}{5}$ **f** $\frac{1}{2}$
2 **a** 99 **b** 58 **c** 90
d 122 **e** 8 **f** 150
3 **a** 55% **b** $\frac{3}{4}$ **c** 98%
4 **a** 61 **b** 9 **c** 37
d 121 **e** 1 **f** 75, 4

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5

Fraction	Decimal	Percentage
$\frac{1}{100}$	0.01	1%
$\frac{1}{10}$	0.1	10%
$\frac{1}{4}$	0.25	25%
$\frac{1}{2}$	0.5	50%
$\frac{3}{4}$	0.75	75%

6 **a** 10 g **b** 7 km **c** \$45
d \$800 **e** 50 c **f** 9 c
7 **a** \$100 **b** 60 m **c** \$1.50
d \$67.80 **e** 6 days **f** 3 days
8 **a** 100 **b** 10 **c** 20 **d** 3 **e** 4
9 2100 km
10 25% of 100
11 **a** 7 **b** 365 **c** 52 **d** 12
12 **a** 14 **b** 6 **c** 12.5

2A

Now you try

Example 1

a 55% **b** $\frac{1}{8}$

Example 2

a 23% **b** 0.48

Example 3

25%

Exercise 2A

1 **a** multiply **b** divide
c multiply **d** divide

2

%	10%	20%	25%	50%	75%	$33\frac{1}{3}\%$	$66\frac{2}{3}\%$
Fraction	$\frac{1}{10}$	$\frac{1}{5}$	$\frac{1}{4}$	$\frac{1}{2}$	$\frac{3}{4}$	$\frac{1}{3}$	$\frac{2}{3}$
Decimal	0.1	0.2	0.25	0.5	0.75	$0.\dot{3}$	$0.\dot{6}$

3 **a** 25% **b** 50% **c** 100% **d** 40%
4 Sarah, $79\% > \frac{38}{50} = 76\%$
5 **a** 20% **b** 80% **c** 80% **d** 30%
e 25% **f** $12\frac{1}{2}\%$ **g** 75% **h** 60%
i 56% **j** 35% **k** 9% **l** $7\frac{1}{2}\%$
6 **a** $\frac{19}{100}$ **b** $\frac{23}{100}$ **c** $\frac{99}{100}$ **d** $\frac{1}{20}$
e $\frac{11}{50}$ **f** $\frac{9}{20}$ **g** $\frac{37}{50}$ **h** $\frac{3}{4}$
i $\frac{1}{40}$ **j** $\frac{69}{400}$ **k** $\frac{1}{100}$ **l** $1\frac{1}{4}$
7 **a** 78% **b** 95% **c** 65% **d** 48%
e 75% **f** 142% **g** 7% **h** 30%
i 3% **j** 104% **k** 12% **l** 12.25%
8 **a** 0.12 **b** 0.83 **c** 0.57 **d** 0.88
e 0.99 **f** 1.0 **g** 1.2 **h** 0.05
9 **a** 2.5% **b** 10% **c** 5% **d** 5%
e 20% **f** 20% **g** 200% **h** 75%

10

Sport	Number of students who chose sport	Fraction of the total	Percentage of the total
Swimming	44	$\frac{22}{75}$	$29\frac{1}{3}\%$
Golf	12	$\frac{2}{25}$	8%
Volleyball	58	$\frac{29}{75}$	$38\frac{2}{3}\%$
Cricket	36	$\frac{6}{25}$	24%
Total	150	1	100%

11 31.5%
12 $16\frac{2}{3}\%$
13 $6\frac{1}{4}\%$
14 Yes, 2.8% fat content
15 Answers will vary.

2B

Now you try

Example 4

\$10.80

Example 5

\$500

Exercise 2B

1 **a** True **b** False **c** False **d** True
2 **a** 12 **b** 9 **c** 6
3 **a** \$400 **b** \$750
4 **a** 2 **b** 10 **c** 8 **d** 7.5
e 57.6 **f** 34 **g** 1230 **h** 42
i 450 **j** 33 **k** 198 **l** 1.5
5 **a** \$36 **b** \$210 **c** 48 kg
d 30 km **e** 15 apples **f** 350 m
g 250 people **h** 200 cars **i** \$49
6 **a** \$120 **b** \$700 **c** \$300 **d** \$7
e \$0.20 **f** \$450 **g** \$800 **h** \$360
7 **a** \$540 **b** \$600 **c** \$508
d \$1250 **e** \$120 **f** \$40
8 **a** \$5.80 **b** \$4.20 **c** \$0.46
d \$0.50 **e** \$44 **f** \$40
9 64
10 **a** Divide by 10. **b** Divide by 2.
c Divide by 2 and then 2 again (or just 4).
11 48 kg
12 15 students
13 $\frac{10}{100} \times 24 = \frac{24}{100} \times 10$; can multiply in any order
14 $x = 2, y = 24$
15 **a** 72 **b** $\frac{10}{11}$ **c** 280%
d $3\frac{1}{4}$ **e** 150%

2C

Now you try

Example 6

\$156

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Example 7

$54.40

Example 8

a 40% **b** 16.0%

Example 9

$8.85

Exercise 2C

1 **a** 1.4 **b** 1.26 **c** 60% **d** 21%
2 **a** 0.8 **b** 0.27 **c** 6% **d** 69%
3 **a** 25% **b** 200%
4 **a** 61.6 **b** 1176 **c** 112 **d** 934.5
e 207 **f** 540 **g** 12 **h** 196
5 **a** 76 **b** 540 **c** 22.5 **d** 616
e 7360 **f** 337.5 **g** 17
h 8910 **i** 0 **j** 9850
6 20%
7 **a**

Original amount	New amount	Increase	Percentage change
40	60	20	50%
12	16	4	$33\frac{1}{3}$%
100	125	25	25%
24	30	6	25%
88	100	12	13.6%

b

Original amount	New amount	Decrease	Percentage change
90	81	9	10%
100	78	22	22%
20	15	5	25%
24	18	6	25%
150	50	100	$66\frac{2}{3}$%

8 **a** $3 **b** $80 **c** $200
d $500 **e** $12 500
9 $1785
10 $84 634.25
11 24 534 cars
12 50%
13 28%
14 $21.50
15 $10.91
16 **a** $900 **b** $990
c As 10% of 1000 = 100 but 10% of 900 = 90
17 **a** $594 **b** $3235.65
c $189 **d** $61.48
18 **a** 79.86 g **b** $97 240.50
c $336 199.68 **d** 7.10 cm

2D

Now you try

Example 10

$1488

Example 11

$52

Example 12

a $13.50 **b** 37.5%

Example 13

$220

Exercise 2D

1 **a** C **b** D **c** A **d** B
2 $3 profit, $2.50 loss, $65.95 loss, $1180 profit
3 40.95, 179.95, 799.95, 8995
4 28, 7.25, 199, 2037
5

Item	Cost price	% mark-up	Selling price
Jeans	$60	28%	$76.80
Toaster	$40	80%	$72
Car	$22 000	45%	$31 900
Can of drink	$1.20	140%	$2.88
Loaf of bread	$1.80	85%	$3.33
Handbag	$80	70%	$136
Tablet	$320	35%	$432

6

Item	Cost price	% discount	Selling price
Camera	$900	15%	$765
Car	$24 000	20%	$19 200
Bike	$600	25%	$450
Shoes	$195	30%	$136.50
Blu-ray player	$245	50%	$122.50
Electric razor	$129	20%	$103.20
Lawn mower	$880	5%	$836

7 **a**

Cost price ($)	Selling price ($)	Profit ($)	Profit (%)
10	15	5	50%
24	30	6	25%
100	150	50	50%
250	255	5	2%
17.50	20	2.50	14.29%

b

Cost price ($)	Selling price ($)	Loss ($)	Loss (%)
10	8	2	20%
16	12	4	25%
100	80	20	20%
34	19	15	44.12%
95	80.75	14.25	15%

8 **a** $5.50 **b** $200 **c** $1500 **d** $125
9 **a** **i** $2 **ii** 20%
b **i** $5 **ii** 25%
c **i** $16.80 **ii** 14%
d **i** $2450 **ii** 175%
10 $1001.25
11 42.3%
12 $613.33
13 **a** $400 **b** $900 **c** $765 **d** No
14 **a** $350 000
b **i** 71.4% **ii** 157.1% **iii** 114.3%
c 25%

2E

Now you try

Example 14

a $1484.62 **b** Ben **c** $58 500

Example 15

a $36 **b** $48 **c** $720

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Example 16

$8500

Exercise 2E

1 **a** C **b** D **c** B **d** A
2 **a** $25.40 **b** $101.60 **c** $482.60
3 **a** $36 **b** $48
4 **a** $11 406.25 **b** $2632.21 **c** $375
5 **a** $1115.38 **b** $29.35/h, therefore less **c** $17 264
6 **a** $41 080 **b** $3423.33 **c** $19.75
7

Employee	Hourly rate	Hours worked	Income
Adam	$20.40	8	$163.20
Betty	$15.50	$8\frac{1}{2}$	$131.75
Ceanna	$19.70	15	$295.50
David	$24.30	38	$923.40
Edward	$57.85	42	$2429.70
Francis	$30	27	$810
George	$35.20	7.25	$255.20

8 **a** $262.80 **b** $379.60
c $584 **d** $876
e $817.60 **f** $1284.80
9

Person	Weekly retainer	Rate of commission	Commission earned	Weekly wage
Andrew	$0	12% on $7000	$840	$840
Byron	$160	8% on $600	$48	$208
Cindy	$300	5% on $680	$34	$334
Deanne	$260	5% on $40 000	$2000	$2260
Elizabeth	$500	8% on $5600	$448	$948
Faruq	$900	2% on $110 000	$2200	$3100
Gary	$1000	1.5% on $45 000	$675	$1675

10 **a** 7 **b** 18 **c** 33
11 $29 per hour
12 $24.40 per hour
13 $490
14 $5010
15 $1678.10
16 **a** $751.64 **b** $151.64 **c** 7 hours

Progress quiz

1 **a** $\frac{8}{25}$ **b** 0.07 **c** 23%
d 12.5% **e** 35% **f** $\frac{9}{200}$
2 **a** 10% **b** 80% **c** 12.5%
3 **a** 16 km **b** $42
c 60 g **d** $21
4 **a** $400 **b** $300 **c** $750
5 **a** 90.2 **b** 237.6
c 93.5 **d** 235
6 **a** 20% **b** 14.9%
7 775
8 **a** $1392 **b** $100.80
9 61.3%
10 **a** $220 **b** $65
11 **a** $1425
b $400.40
c $188.75
d $620

2F

Now you try

Example 17

a $4712 **b** 24% **c** $64 712

Example 18

$13 288

Exercise 2F

1

Gross income	Deductions	Net income
$5600	$450	$5150
$87 000	$28 000	$59 000
$50 000	$6700	$43 300

2 $1405
3 $25 000
4 **a** Nil **b** $3008 **c** $14 788
5 **a** **i** $40 035 **ii** 17.0%
b **i** $53 905.80 **ii** 20.1%
c **i** $41 218.20 **ii** 15%
d **i** $30 052.56 **ii** 22.2%
6 **a** $1830 **b** $8043
c $12 617.50 **d** $23 772.80
7 **a** $19 656 **b** $53 144
8 **a** $0 **b** $1888 **c** $14 788
d $39 058 **e** $56 138 **f** $416 138
9 $67 400
10 **a** $51 200 **b** $6148
c $45 052 **d** $1732.77
11 $302.85
12 **a** $45 001 – $135 000 **b** $47 373
13 **a** $81 120 **b** $15 124
c $65 996 **d** $2538.31
14 **a** $166 960 **b** $43 113.20
c $123 846.80 **d** $4763.34
15 $1112.52

2G

Now you try

Example 19

$400

Example 20

$618.75

Example 21

Interest = $570

Total = $5570

Exercise 2G

1 **a** $12 000 **b** 6% p.a. **c** 3.5 years
2 **a** $1120 **b** $2800 **c** $5600
3 **a** $3000 **b** $3600 **c** $416
4 **a** $300 **b** $900 **c** $1600
d $3150 **e** $1196.25
5 $2700, $17 700
6 **a** $52.50 **b** $100 **c** $40
d $264.38 **e** $1027.40
7 $1980, $23 980

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

8

a	$840	$7840
b	$840	$2340
c	$1500	$41 500
d	$4550	$74 550
e	$200	$2200

9 $2560
10 **a** $3600 **b** $18 000 **c** $38 000
11 Choice 2
12 **a** $645 **b** $194 **c** $129
d $19 **e** $55 148 **f** $12 320
g $2580 **h** $58 050 **i** $129 000
13 **a** **i** $12 **ii** $31 **iii** $10
b **i** $335 **ii** $500 **iii** $235

2H

Now you try

Example 22

2.5 years or 2 years 6 months

Example 23

5% p.a.

Example 24

a $16 800 **b** $51 800 **c** $719.44

Exercise 2H

1 **a** 2 years **b** 4 years **c** 50 years **d** $2\frac{1}{2}$ years
2 5 years
3 $2400
4 $450
5 3 years
6 $4\frac{1}{2}$ years
7 18 months
8 **a** 6.25% **b** 2% **c** 6.5%
d 5.5% **e** 21% **f** 3.5%
9

Interest	Total amount to be repaid	Monthly repayment
$5250	$10 250	$170.83
$10 500	$24 500	$408.33
$2400	$12 400	$258.33
$44 000	$99 000	$825
$525 000	$775 000	$2152.78

10 $2083.33
11 **a** $14 400 **b** $240
12 10%
13 **a** $\$P$ **b** 12.5%
c **i** 20 years **ii** 40 years **iii** Double
14 **a** Final amount = $732.05 **b** $232.05
15 **a** $5743.27
b $5743.27, same as final answer in part **a**.
c Increases the amount by 5% five times, giving the same result
d 5.5%

Maths@Work: Facebook cake-decorating business

1 $294
2 **a** 8 cm **b** 10 cm **c** 23 cm **d** 25 cm
3 21%
4 **a** White: 5 kg at $40; Red: 2.5 kg at $36.95; Blue: 500g at $7.95
b $1.55/100g
c When a specific quantity was required and any extra would be wasted
5 **a** $231 **b** 49% **c** 290%, 480%, 350%, 330%
d $26/h; $44/h; $32/h; $30/h
6 **a** $427
b Birthday – choc fudge mud; Birthday – caramel mud; Birthday – choc Jaffa
c $47/h wedding cake 3 tiers; $46/h wedding cake 2 tiers; $43/h child's theme birthday cake
d The higher rate of pay for these cakes is due to their complexity requiring a higher level of skill for baking and decorating.

Puzzles and games

1

S	A	G	R	O	S	S	W	H	K	L	O	M
A	P	E	R	C	E	N	T	A	G	E	C	O
L	A	R	C	O	M	M	I	S	S	I	O	N
A	S	E	R	V	I	N	T	E	R	E	S	T
R	T	Y	H	E	T	S	L	O	S	S	T	H
Y	E	F	O	R	T	N	I	G	H	T	I	L
W	S	M	O	T	A	X	A	T	I	O	N	Y
E	I	O	L	I	D	I	D	N	P	Y	I	E
K	M	N	Q	M	O	N	T	R	N	I	S	C
E	P	Y	U	E	D	I	S	C	O	U	N	T
M	L	R	E	P	A	Y	M	E	N	T	A	H
D	E	D	U	C	T	I	O	N	S	I	X	L

2 EARTH, AIR, FIRE, WATER

Checklist Answers

1 **a** 17.5% **b** 37.5%
2 **a** 30% **b** 0.72
3 15%
4 $26.25
5 $600
6 **a** $89.60 **b** $36.80
7 15.6%
8 1300
9 $1320
10 62.5%
11 $125
12 **a** $44.53 **b** $82 368
13 $672
14 $3032
15 $4104
16 tax = $15 548 net = $62 452
17 $17 788
18 interest = $480, Amount = $4480
19 $945
20 18 months (1.5 years)
21 $880

Short-answers questions

1

Decimal	Fraction	Percentage
0.6	$\frac{3}{5}$	60%
$0.\dot{3}$	$\frac{1}{3}$	$33\frac{1}{3}$%
0.0325	$\frac{13}{400}$	$3\frac{1}{4}$%
0.75	$\frac{3}{4}$	75%
1.2	$1\frac{1}{5}$	120%
2	2	200%

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

2 **a** \$77.50 **b** 1.65
3 **a** 150 **b** 25
4 **a** 72 **b** 1.17 **c** 20%
5 12.5 kg
6 \$1800
7 \$4375
8 **a** \$25 **b** $16\frac{2}{3}\%$
9 **a** \$38.50 **b** \$14.30
10 \$50 592
11 \$525
12 3 years
13 \$33 138

Multiple-choice questions

1 C **2** C **3** B **4** D **5** C
6 C **7** C **8** A **9** C **10** E

Extended-response questions

1 **a** \$231
b \$651
c **i** \$63 **ii** \$34.65
2 **a** \$25.68 **b** \$38.52 **c** \$154.08
d 6.5 **e** \$46 547.31 **f** \$558.57

Chapter 3

Warm-up quiz

1 **a** $3x$ **b** $4ab$ **c** $10x$ **d** $21b$
2 **a** 8 **b** 7 **c** 12
d 6 **e** 9 **f** 14
3 **a** $x+3$ **b** ab **c** $2y-3$ **d** $\frac{x+2}{3}$
4 **a** -18 **b** -28 **c** 40 **d** 3
e 3 **f** -5 **g** -4 **h** -8
5 **a** $3x$ **b** $6y+5$ **c** $8x$
d $17y$ **e** $10a+9a^2$ **f** $5xy-7y$
6 **a** $6a$ **b** $-21xy$ **c** $4b$ **d** $\frac{3m}{2}$
7 **a** $2x+6$ **b** $3a-15$ **c** $8x+4$
8 **a** 1 **b** -2 **c** 5 **d** -6
9 b, c
10 **a** 9 **b** 10 **c** 4 **d** 21

3A

Now you try

Example 1

a 4
b **i** -2 **ii** 6
c 5

Example 2

a $g+5$ **b** $\$4x$ **c** $\frac{600}{y}$ mL

Example 3

a $y+7$ **b** $2x-6$ **c** $\frac{m+n}{3}$ **d** $\sqrt{a+b}$

Example 4

a 24 **b** -13 **c** 33

Exercise 3A

1 **a** expression **b** constant term
c pronumeral **d** coefficient
e terms
2 **a** $7y$ **b** $-2x$ **c** ab
d $\frac{y}{2}$ **e** $\frac{x}{y}$ **f** $\frac{2}{a}$
3 **a** 5 **b** 8 **c** 14 **d** 4
4 **a** Yes **b** No **c** Yes **d** No
5 **a** **i** 3 **ii** 2 **iii** 3
b **i** 2 **ii** -3 **iii** 0
c **i** 3 **ii** 7 **iii** -4
d **i** 4 **ii** $\frac{1}{2}$ **iii** -1
6 **a** **i** $4+r$ **ii** $t+2$ **iii** $x+y+z$
b **i** $\$6P$ **ii** $\$10n$ **iii** $\$2D$ **iv** $\$(5P+2D)$
c $\frac{500}{C}$
7 **a** $2+x$ **b** $ab+y$ **c** $x-5$ **d** $2y-7$
e $3x$ **f** $3p$ **g** $2x+4$ **h** $\frac{x+y}{5}$
i $4x-10$ **j** $1-3x$ **k** $\frac{3+y}{2}$ **l** $\frac{1}{2}(x+1)$
m $(m+n)^2$ **n** m^2+n^2
8 **a** 20 **b** 25 **c** 15 **d** 0 **e** 6
f 5 **g** 28 **h** -96 **i** 5
9 **a** 60 m^2 **b** Length $= 12+x$, width $= 5-y$
c $17+x-y$
10 **a** \$26 **b** \$50 **c** No **d** 12 hours
11 **a** 18 square units **b** 1, 2, 3, 4, 5
12 **a** $\frac{A}{20}$ **b** $\frac{3A}{20}$ **c** $\frac{nA}{20}$
13 **a** **i** $P=2x+2y$ **ii** $A=xy$
b **i** $P=4p$ **ii** $A=p^2$
c **i** $P=6+x+y$ **ii** $A=3x$
14 **a** **i** No **ii** No
b **i** $3x+1$ or $3(x+1)$
ii $5+\frac{x}{3}$ or $\frac{5+x}{3}$
iii $\frac{1}{2}(x+y)$ or $\frac{1}{2}x+y$
c Answers may vary.
i The sum of 2 and x is multiplied by 4.
ii 3 is added to a half of x.
iii A third of the sum of m and n
iv 7 more than 5 lots of x
v The sum of x and y is divided by 2.
vi The sum of b and a half of a

3B

Now you try

Example 5

a $5m$ and $2m$ are like terms
b $7y$ and $-y$ are like terms
$4xz$ and zx are like terms

Example 6

a $8t$ **b** $4y+3$

Example 7

a $6a+13b$ **b** $3st+4s$ **c** $5x^2y-6xy$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Exercise 3B

1 **a** like **b** coefficients
c pronumeral/variable **d** 1
2 **a** Y **b** N **c** N
d Y **e** Y **f** N
3 a, c, d, f
4 **a** $4y$ and $2y$, $3xy$ and $7xy$ **b** $3x$ and $7x$
c $7ab$ and $-3ab$, $2a$ and $5a$ **d** $2a^2$ and $-3a^2$, $4a$ and a
e $3x^2y$ and $7yx^2$ **f** $5ab^2$ and $4ab^2$, $3ab$ and $7ba$
5 **a** $10a$ **b** $7n$ **c** $8y$ **d** $11x$
e $4ab$ **f** $8mn$ **g** $7y + 8$ **h** $3x + 5$
i $4m + 2$ **j** $12ab + 3a$ **k** $7xy + 4y$ **l** $3bc - 4$
6 **a** $5a + 9b$ **b** $6x + 5y$ **c** $5xy + 4x$
d $4t + 6$ **e** $11x + 4$ **f** $7mn - 1$
g $5ab - a$ **h** 0 **i** $10ac - 9c$
j $3xy^2$ **k** $3m^2n$ **l** $7x^2y - 4xy$
m $4x^2 - 3x$ **n** $4a^2b - 3ab$ **o** $7pq^2 - 8pq$
7 **a** $x + y$ **b** $4x + 2y$
8 **a** $4x + 2y$ **b** $12s$ **c** $30x$
d $5x + 4$ **e** $6x + 16$ **f** $12a + 4b + 4$
9 $8x$ metres
10 **a** $4t$ **b** $9x$ **c** $2x$ **d** $7a$
e $5n$ **f** $4pq$ **g** $2xy$ **h** $3b$
11 **a** $3a + 7b$ **b** 4 units each
c a **d** $2a - b$

3C

Now you try

Example 8

a $20x$ **b** $-18mp$

Example 9

a $\frac{x}{3}$ **b** $\frac{9q}{2}$

Example 10

a $42a^2b$ **b** $5y$

Exercise 3C

1 **a** $7xy$ **b** 6 **c** $20x$ **d** 2
2 **a** T **b** T **c** F
d F **e** F **f** T
3 **a** $10m$ **b** $12b$ **c** $15p$ **d** $6x$
e $18pr$ **f** $16mn$ **g** $-14xy$ **h** $-15mn$
i $-12cd$ **j** $30ab$ **k** $-24rs$ **l** $-40jk$
4 **a** $4b$ **b** $\frac{a}{3}$ **c** $\frac{2ab}{3}$
d $\frac{m}{2}$ **e** $\frac{x}{4}$ **f** $\frac{5s}{3}$
g 3 **h** $\frac{9q}{2}$ **i** $\frac{x}{2}$
j $6b$ **k** $\frac{7m}{3}$ **l** $\frac{2}{5x}$
5 **a** $24n^2$ **b** $-3q^2$ **c** $10s^2$
d $21a^2b$ **e** $-15mn^2$ **f** $18gh^2$
g $3b$ **h** $5x$ **i** $\frac{m}{2}$
j $\frac{1}{4y}$ **k** $\frac{3ab}{5}$ **l** $3pq$
6 **a** $\frac{4x}{y}$ **b** $\frac{m}{3}$ **c** $\frac{2a}{b}$ **d** $\frac{10x}{y}$ **e** $8n$
f $5b$ **g** 5 **h** $6xy$ **i** $\frac{p^2}{2}$ **j** $2ab$
7 **a** $12ab$ **b** $4x^2$ **c** $10xy$ **d** $6x^2$
8 $2x$
9 **a** $4n$ **b** $7y$ **c** $9y$
d $5ab$ **e** $14x$ **f** $10y$
g $12a$ **h** $8x^2y$ **i** $10x^2$
10 **a** **i** $3x^2$ m^2 **ii** $5x$ metres
b **i** 7500 m^2 **ii** 250 m
11 **a** $4b$ **b** $2ab$ **c** b

3D

Now you try

Example 11

a $4x + 8$ **b** $6x - 60$

Example 12

a $-3x - 21$ **b** $-5x + 45$

Example 13

a $5a + 20b$ **b** $15y^2 - 12y$

Example 14

a $7x + 6$ **b** $18 - 2y$

Exercise 3D

1 **a** **i** $5x$ **ii** 10
b $5x + 10$
c $x + 2$
d $5(x + 2)$
e $5x + 10$
2 term, multiplied, inside
3 C
4 **a** 3, 3, 15 **b** -4, -4, $-4x$, 8
5 **a** $2x + 6$ **b** $5x + 60$ **c** $2x + 14$
d $7x + 63$ **e** $3x - 21$ **f** $6x - 66$
g $5x - 45$ **h** $10x - 10$ **i** $6 + 3x$
j $21 + 7x$ **k** $28 - 4x$ **l** $40 - 5x$
6 **a** $-3x - 6$ **b** $-2x - 22$
c $-5x + 15$ **d** $-6x + 36$
e $-8 + 4x$ **f** $-39 - 13x$
g $-72 - 8x$ **h** $-300 + 300x$
7 **a** $2a + 2b$ **b** $5a + 10b$
c $6m + 3y$ **d** $16x - 40$
e $-12x - 15$ **f** $4x^2 - 8xy$
g $2t^2 - 3t$ **h** $3a^2 + 4a$
i $2d^2 - 5d$ **j** $6b^2 - 10b$
k $8x^2 + 2x$ **l** $5y - 15y^2$
8 **a** $2x + 11$ **b** $6x + 22$ **c** $5x + 6$
d $3x + 10$ **e** $4x - 5$ **f** $2x + 1$
g $8 - 3x$ **h** $21 - 5x$ **i** $-1 - 2x$
j $-3x$ **k** $3 - x$ **l** $6 - 3x$
9 **a** $4x + 4y$ **b** $8x + 4$ **c** $2x - 2$
10 **a** $2x + 2$ **b** $8x - 12$
c $6x^2 + 2x$ **d** $6x + 15$
11 $2x + 120$
12 **a** $2x + 12$ **b** $x^2 - 4x$
c $-3x - 12$ **d** $-7x + 49$
e $5x + 10 + 4 = 5x + 14$ **f** $5 - 2x + 14 = 19 - 2x$
13 **a** $x - 10\,000$ **b** $0.2(x - 10\,000) = 0.2x - 2000$
14 **a** $5x + 12$ **b** $4x + 6$ **c** $11x + 7$
d $17x - 7$ **e** $x - 7$ **f** $2x - 4$
g $3x^2 + 10x$ **h** $5x - 3 + x^2$
15 **a** $2x^2 + 4x$ **b** $x^2 + 2x + 6$
c $20 + 8x - x^2$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3E

Now you try

Example 15

a $x = 5$ **b** $x = 3$

Example 16

a $x = -2$ **b** $x = -\frac{4}{5}$

Exercise 3E

1 c, d

2 **a** 6 **b** 8 **c** 4

3 **a** $2x = 6$ **b** $3x = 21$ **c** $x = 9$

4 **a** Subtract 9 from both sides.
b Add 1 to both sides.
c Divide both sides by 3.

5 **a** $x = 4$ **b** $a = 1$ **c** $m = 4$ **d** $x = 4$
e $n = 2$ **f** $x = -4$ **g** $b = 3$ **h** $y = 7$
i $a = 4$ **j** $b = 8$ **k** $x = -1$ **l** $y = 2$

6 **a** $x = -3$ **b** $x = -1$ **c** $x = -2$ **d** $x = -8$
e $x = 2$ **f** $x = 2$ **g** $x = 2$ **h** $x = 6$

7 **a** $x = \frac{3}{2}$ **b** $x = \frac{3}{5}$ **c** $x = \frac{10}{11}$
d $x = \frac{1}{2}$ **e** $x = -\frac{1}{3}$ **f** $x = \frac{4}{5}$

8 **a** $x + 8 = 34$, $x = 26$ **b** $x - 7 = 21$, $x = 28$
c $2x + 4 = 10$, $x = 3$ **d** $3x - 4 = 11$, $x = 5$
e $3x - 4 = 20$, $x = 8$ **f** $25 - 2x = 7$, $x = 9$

9 **a** Added 2 on the right hand side rather than subtracting 2.
b Multiplied the right hand side by 2 in the last step rather than dividing by 2.
c The negative was dropped, $-x = 7$ so $x = -7$.
d Needed to subtract 3 from both sides before dividing both sides by 2.

10 **a** $x = 5$ **b** $x = -9$ **c** $x = -2$

11 **a** 10 s **b** 8 s

3F

Now you try

Example 17

a $x = 8$ **b** $x = 30$

Example 18

$x = 17$

Exercise 3F

1 **a** 5 **b** $\frac{3}{4}$ **c** $2x$

2 **a** 20 **b** 24 **c** 11 **d** 15

3 C

4 **a** iv, add 1 to both sides
b iii, multiply both sides by 4
c i, multiply both sides by 3
d ii, multiply both sides by 2

5 **a** $x = 15$ **b** $y = 14$ **c** $y = 10$ **d** $x = 4$
e $x = 8$ **f** $x = 2$ **g** $b = 12$ **h** $t = -6$
i $a = -6$ **j** $y = 30$ **k** $x = 15$ **l** $s = 4$
m $x = 12$ **n** $m = -4$ **o** $y = -35$

6 **a** $x = 11$ **b** $x = 6$ **c** $y = 2$ **d** $b = 0$
e $y = 20$ **f** $t = 39$ **g** $k = 57$ **h** $x = 70$
i $x = -10$ **j** $b = -24$ **k** $y = -22$ **l** $y = 1$

7 **a** $t = 3$ **b** $m = 40$ **c** $y = -2$
d $x = 19$ **e** $y = -6$ **f** $t = -\frac{3}{4}$

8 **a** $\frac{x}{3} = 12$, $x = 36$ **b** $\frac{2x}{5} = 4$, $x = 10$
c $\frac{x}{2} - 4 = 10$, $x = 28$ **d** $\frac{x+3}{4} = 6$, $x = 21$
e $\frac{7x}{3} = 8$, $x = \frac{24}{7}$ **f** $\frac{x}{3} + 5 = 16$, $x = 33$

9 21

10 **a** Needed to $\times$ 3 before $-$ 2, $x + 2 = 21$, $x = 19$
b Need to + 4 before $\times$ 3, $\frac{x}{3} = 6$, $x = 18$

11 **a** $x = 12$ **b** $x = 12$ **c** $x = -4$
d $b = 6$ **e** $x = 5$ **f** $m = 5$
g $y = -1$ **h** $t = -7$ **i** $x = -3$

12 **a** $x = 9$ **b** $x = 36$ **c** $x = -7$
d $x = 8$ **e** $x = 5$ **f** $x = -10$

13 No, order does not matter for $\times$ or $\div$. By dividing by 7 first you deal with smaller numbers, since 7 is a factor of 14.

Progress quiz

1 **a** 4 **b** -3 **c** 8

2 **a** $\$3d$ **b** $\frac{\$500}{x}$ **c** $3y - 4$ **d** $2(m + n)$

3 **a** -1 **b** -8 **c** 12

4 **a** $12x$ **b** $7b$ **c** $9a + 5$
d $8xy - 4x$ **e** $3a + 10b$ **f** $6mn - 3n$
g $2xy^2 - 2xy$ **h** $2a^2b + 7ab^2$ **i** $-2m^2n - 2mn^2$

5 **a** $24b$ **b** $24ac$ **c** $-20xy$
d $28ab^2$ **e** $\frac{x}{4}$ **f** $\frac{3y}{4}$
g $\frac{3xy}{2}$ **h** $8x$ **i** $\frac{5ab}{2}$

6 **a** $3x + 15$ **b** $4y - 4$ **c** $-2x - 10$
d $-5a + 10$ **e** $12x - 30y$ **f** $12m^2 + 3mn$

7 **a** $6x - 8$ **b** $-4x - 4$ **c** $2 - 2x$

8 **a** $x = 3$ **b** $x = 4$ **c** $x = -9$ **d** $x = -\frac{3}{5}$

9 **a** $x = 28$ **b** $x = 48$ **c** $x = 27$ **d** $x = -22$

10 **a** $3x + 5 = 32$, number is 9 **b** $\frac{x-3}{4} = 5$, number is 23

3G

Now you try

Example 19

$x = \frac{11}{3}$ or $3\frac{2}{3}$

Example 20

$x = 5$

Exercise 3G

1 **a** $3 \times \underline{x} + \underline{3} \times 7 = 25$
$3x + \underline{21} = 25$
b $\underline{4} \times 2x + 4 \times (\underline{-1}) = 15$
$\underline{8x} - 4 = 15$

2 **a** $x + 2 = 3$ **b** $x - 3 = 5$ **c** $x + 6 = -4$

3 **a** $4x - 12$ **b** $2x - 15$ **c** $-2x + 8$

4 **a** $x = \frac{5}{2}$ or $2\frac{1}{2}$ **b** $a = \frac{3}{5}$ **c** $m = \frac{19}{3}$ or $6\frac{1}{3}$
d $y = \frac{47}{5}$ or $9\frac{2}{5}$ **e** $p = \frac{35}{4}$ or $8\frac{3}{4}$ **f** $k = \frac{19}{2}$ or $9\frac{1}{2}$
g $b = -\frac{1}{4}$ **h** $m = -\frac{11}{2}$ or $-5\frac{1}{2}$
i $x = -\frac{4}{5}$ **j** $a = \frac{1}{14}$ **k** $x = \frac{19}{6}$ or $3\frac{1}{6}$
l $n = \frac{2}{3}$ **m** $x = -\frac{1}{10}$ **n** $y = -\frac{1}{6}$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5 **a** $x = 1$ **b** $x = 2$ **c** $x = 2$ **d** $x = 12$
e $x = 3$ **f** $x = 1\frac{2}{3}$ **g** $x = -5$ **h** $x = -1$
6 **a** $x = 1$ **b** $x = 2$ **c** $x = 1$ **d** $x = 9$
e $x = -4$ **f** $x = 1$
7 **a** $3(x + 1) = 4, x = \frac{1}{3}$ **b** $2(x - 2) = 19, x = 11\frac{1}{2}$
c $2(x + 3) = 7, x = \frac{1}{2}$ **d** $3(x - 4) = 8, x = 6\frac{2}{3}$
e $2(3x - 2) = 5, x = 1\frac{1}{2}$ **f** $3(2x + 5) = 10, x = -\frac{5}{6}$
8 \$5/hour
9 **a** $x = 5$ **b** $x = 5$
c Dividing both sides by 3 is faster because $9 \div 3$ is a whole number
10 **a** $x = 4\frac{1}{3}$ **b** $x = 4\frac{1}{3}$
c Expanding the brackets is easier because $7 \div 3$ gives a fraction answer
11 **a** No **b** No **c** Yes, $4(x + 1) = 20$
d Yes, $2(x - 4) + 3x + 1 = 13$
e Yes, $3x + 2 + 2(x - 5) = 12$
f Yes, $2x - 3 - 3(x - 1) = -4$

3H

Now you try

Example 21

a $x = 4$ **b** $x = 3$

Example 22

a $x = 4$ **b** $x = 2$

Example 23

a $x = -4$ **b** $x = 8$

Exercise 3H

1 **a** Left **b** Left **c** Right **d** Right
2 **a** Subtract $3x$ from both sides.
b Subtract $5x$ from both sides.
c Add x to both sides.
d Add $2x$ to both sides.
3 **a** $2x = 6$ **b** $2x - 3 = 1$ **c** $5x = 1$
d $4 = 8 + 6x$
4 **a** $x = 4$ **b** $x = 3$ **c** $x = 3$ **d** $x = 7$
e $x = -3$ **f** $x = -2$ **g** $x = 2$ **h** $x = 4$
i $x = 6$ **j** $x = 2$
5 **a** $x = 8$ **b** $a = 3$ **c** $m = 8$ **d** $x = 4$
e $x = 5$ **f** $x = 3$ **g** $x = 1$ **h** $y = 2$
i $m = 3$ **j** $x = 3$ **k** $b = -1$ **l** $m = -3$
6 **a** $x = 7$ **b** $a = 5$ **c** $y = 2$ **d** $x = -14$
e $b = -16$ **f** $m = 3$ **g** $a = 19$ **h** $x = -3$
i $x = -13$ **j** $n = 14$ **k** $a = 2$ **l** $x = \frac{5}{2}$
7 **a** $2x = 3x - 4, x = 4$ **b** $2x + 4 = 5x, x = 1\frac{1}{3}$
c $3x + 10 = 5x, x = 5$ **d** $3x + 2 = 5x - 6, x = 4$
e $2x - 1 = 3x + 5, x = -6$ **f** $2x + 4 = 13 - x, x = 3$
8 Mardy is 8 years old.
9 **a** $x = 4$ **b** $x = 4$
c Method a: don't have to deal with negatives
10 **a** 48 **b** 38 **c** 26 **d** 48

3I

Now you try

Example 24

5

Example 25

5 hours

Example 26

Max kicked 4 goals and Tim 7 goals


Exercise 3I


1 **a** iii **b** iv **c** i **d** ii
2 ii, v, iv, i, iii
3 age, $x + 4$, $x + 4$, $2x$, 30, 15, Bernard is 15 and Sam is 19.
4 **a** 7 **b** 5 **c** 21 **d** 8 **e** 2
5 **a** Let d be the number of days the car is rented for.
b $50 + 40d = 290$ **c** $d = 6$
d The car was rented for 6 days
6 48 items
7 **a** Let e be the number of goals for Emma. **b** $e + 8$
c $e + e + 8 = 28$ **d** $e = 10$
e Emma scored 10 goals, Leonie scored 18 goals
8 **a** Let w be the width in centimetres. **b** Length $= 4w$
c $2w + 2(4w) = 560$ **d** $w = 56$
e Length = 224 cm, width = 56 cm
9 Benita \$360, Adele \$640
10 Andrew \$102.50, Brenda \$175, Cammi \$122.50
11 Walked 10 km, ran 20 km
12 15 and 45
13 Rectangle $L = 55$ m, $W = 50$ m. Triangle side = 70 m
14 **a** 27, 28, 29
b **i** $x, x + 2, x + 4$ **ii** 4, 6, 8
15 **a** $T = 8x + 7200$ **b** 300 **c** $R = 24x$
d $x = 350$ **e** 3825


3J

Now you try

Example 27

a


b


c



Example 28


a $x < 11$
b $x < -1$


Example 29


a $a \geqslant -9$
b $a \geqslant -\frac{7}{2}$

Exercise 3J

1 **a** $3 > 2$ **b** $-1 < 4$ **c** $-7 < -3$ **d** $5 > -50$
2 **a** C **b** A **c** B
3 **a**



b


c



d


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

e



f



4 **a** $x < 3$ **b** $b > 5$ **c** $y > 6$ **d** $x \geqslant 3$
e $t > -5$ **f** $y \geqslant -10$ **g** $m < 6$ **h** $a \geqslant 1\frac{1}{2}$
i $x < 1$ **j** $x < \frac{1}{3}$

5 **a** $x < 4$ **b** $n \leqslant -1$ **c** $x \geqslant \frac{3}{5}$
d $a \geqslant 4$ **e** $x \geqslant -6$ **f** $x \geqslant 10$
g $x < -6$ **h** $t \leqslant -2$ **i** $m > -4\frac{5}{6}$

6 **a** $x \leqslant 16$ **b** $x \leqslant -9$ **c** $x \leqslant 20$
d $x < 11$ **e** $x > -2\frac{2}{3}$ **f** $x \geqslant -2$

7 **a** $x < 1$ **b** $a < -8$ **c** $x \leqslant -2$
d $x < 2\frac{1}{2}$ **e** $y < -3\frac{1}{5}$ **f** $x < -\frac{4}{7}$

8 **a** $x \geqslant 2\frac{1}{2}$ **b** $t > -\frac{3}{5}$ **c** $y \leqslant \frac{3}{8}$
d $a < 1\frac{1}{5}$ **e** $m \geqslant 6$ **f** $b < 5\frac{1}{2}$

9 less than 18
10 less than 13 cm
11 399 km
12 **a** $x < 3$ **b** $x < 3$
c Reverse the inequality sign when dividing by a negative number.

13 **a** $x < -13$ **b** $x \geqslant -3$ **c** $x > \frac{4}{7}$
d $x \leq \frac{13}{5}$ **e** $x > \frac{10}{17}$ **f** $x \geqslant \frac{3}{4}$
g $x > \frac{b-c}{a}$ **h** $x \leqslant \frac{ac}{b}$ **i** $x < \frac{c}{a} - b$

3K

Now you try

Example 30

a 7 **b** 8

Example 31

a $A = 3$ **b** $r = 3.2$

Example 32

a $a = \frac{v-u}{t}$ **b** $v = \sqrt{\frac{2E}{m}}$

Exercise 3K

1 Answers will vary, e.g. $A = lw$, $A = \frac{1}{2}bh$, $A = \pi r^2 \ldots$

2 **a** A **b** D **c** M **d** A

3 **a** $x = \frac{b}{a}$ **b** $x = cd$
c $x = ab + d$ **d** $x = c^2$

4 **a** 21 **b** 24 **c** 2 **d** 6
e 452.39 **f** 33.51 **g** 25.06 **h** 14.95
i 249.86 **j** 80

5 **a** $F = 36$ **b** $w = 5$ **c** $t = 7.5$
d $v = 25$ **e** $r = 3.91$ **f** $a = 20$

6 **a** $d = \frac{C}{\pi}$ **b** $x = \frac{d-a}{b}$
c $n = \frac{p}{m} - x$ or $\frac{p - mx}{m}$ **d** $r = \frac{100I}{Pt}$
e $v = \sqrt{PR}$ **f** $h = \frac{A}{2\pi r}$
g $r = \sqrt{\frac{V}{\pi h}}$ **h** $A = (4C - B)^2$
i $a = \sqrt{c^2 - b^2}$ **j** $b = a^2 - c$

7 **a** 88.89 km/h
b **i** $d = st$ **ii** 285 km

8 **a** 35 m/s **b** 2 s

9 **a** Decrease **b** 3988 L **c** 417 min

10 **a** **i** 212°F **ii** 100.4°F
b **i** −10°C **ii** 36.7°C

11 **a** 73 **b** 7 **c** 40.025

Maths@Work: Plumber

1 **a** 6.3 L **b** 7.88 L **c** 18.75 L
d 15 L

2 **a** **i** 98.1 kPa **ii** 196.2 kPa
iii 294.3 kPa **iv** 392.4 kPa
b $h = \frac{P}{9.81}$
i 2.51 m **ii** 2.46 m **iii** 1.38 m **iv** 2.84 m

3 **a** 150 mm **b** 357 mm
c 150 mm **d** 333 mm

4

The number of drain pipes for a house			
House	Total roof catchment area m^2	Decimal number of downpipes	Whole number of downpipes
i	131	2.79	3
ii	147	3.27	4
iii	180.5	3.92	4
iv	257	5.47	6

Puzzles and games

1 $x = 3$

4	9	2
3	5	7
8	1	6


2 $5x^2 - xy$
3 WORD
4 Eric is 18 years old now.
5 1st leg = 54 km, 2nd leg = 27 km, 3rd leg = 18 km, 4th leg = 54 km
6 $140
7 **a** $x = \frac{12}{5}$ **b** $x = \frac{23}{2}$

Chapter checklist

1 **a** 3 **b** −5 **c** 6
2 **a** $\$15n$ **b** $3x - 4$ **c** $\frac{y^2}{4}$
3 **a** 19 **b** −23
4 $4x, -5x$ and $3xy, yx$
5 **a** $8a - 4$ **b** $3xy + 10y$
6 **a** $20st$ **b** $-24xy^2$
7 **a** $\frac{5x}{4}$ **b** $\frac{2b}{3}$
8 **a** $4x + 20$ **b** $-3x + 18$ **c** $6x^2 - 2xy$
9 **a** $3x + 2$ **b** $3 - 2x$
10 **a** $x = 6$ **b** $x = \frac{-7}{3}$
11 **a** $x = 12$ **b** $x = 30$
12 $x = 23$
13 $x = \frac{7}{9}$
14 $x = \frac{5}{2}$
15 $x = 2$
16 **a** $x = 5$ **b** $x = 2$
17 $x = -2$

Essential Mathematics Core for the Victorian Curriculum Year 9 ISBN 978-1-009-59315-1 © Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


18 $2x + 8 = 30,\ x = 11$
19 $15 + 12n = 51,\ n = 3$
20 $x + (x + 140) = 370$ Trey earns \$115 and Toni earns \$255
21




22 $x > -4$
23 a $A = 35$ **b** $h = 4.2$
24 $u = \sqrt{v^2 - 2as}$

Short-answer questions


1 a $x + y$ **b** $7m$ **c** $\$3m$ **d** $\frac{n-3}{4}$
2 a 11 **b** 7 **c** 12 **d** 8 **e** −11
3 a $8n$ **b** $6xy$ **c** $4a$ **d** $\frac{xy}{3}$
4 a $7b$ **b** $4x + 3$ **c** $3p + 2q$ **d** $2m - 3mn + n$
5 a $2x + 14$ **b** $-6x - 15$ **c** $6x^2 - 8x$ **d** $-10a + 8a^2$ **e** $4x + 13$ **f** $3x - 7$
6 a $x = 9$ **b** $x = 2$ **c** $x = -3$ **d** $x = -9$
7 a $x = 10$ **b** $x = 8$ **c** $x = 26$ **d** $x = 6$
8 a $2n + 3 = 21, n = 9$ **b** $\frac{l-5}{3} = 7, l = 26$ **c** $\frac{x}{4} - 5 = 0, x = 20$
9 a $x = 5$ **b** $x = \frac{11}{6}$ or $1\frac{5}{6}$
10 a $x = 4$ **b** $x = 2$ **c** $x = 1$ **d** $x = -5$
11 \$260
12 a




b




c




d



e



f



13 a $x < 12$ **b** $m > -1$ **c** $y \geq -6$ **d** $x < 17$ **e** $a > 4$ **f** $x \geq 2$
14 a $E = 60$ **b** $a = 5$ **c** $h = 10$
15 a $x = \frac{v^2 - u^2}{2a}$ **b** $I = \sqrt{\frac{P}{R}}$

Multiple-choice questions

1 C **2** C **3** D **4** C **5** D **6** A
7 B **8** B **9** B **10** B **11** A **12** D

Extended-response questions

1 a i \$85 **ii** \$110
b i Let h be the number of hours of hire. **ii** $60 + 25h = 210$
iii 6 hours
2 a $h = \frac{A}{10}$
b 10 m
c 17 m

Chapter 4

Warm-up quiz

1 a 5 cm **b** 13 m **c** 41 mm
2 a 9 **b** 49 **c** 20 **d** 45
3 a 0.4568 **b** 0.3457 **c** 0.0456 **d** 0.2800
4 a 4.23 **b** 5.68 **c** 76.90 **d** 23.90
5 a 5.92 **b** 15.36 **c** 4.86 **d** 8.09
6 a $x = 64$ **b** $x = 28$ **c** $x = 108$ **d** $x = 60$ **e** $x = 50$ **f** $x = 55$
7 a $x = 2$ **b** $x = 3$ **c** $x = 12$ **d** $x = 6$ **e** $x = 12$ **f** $x = 30$ **g** $x = 28$ **h** $x = 182$
8 a $x = 0.6$ **b** $x = 0.6$ **c** $x = 2.1$ **d** $x = 0.5$
9 a $x = 2$ **b** $x = 4$ **c** $x = 5$ **d** $x = 13$

4A

Now you try

Example 1

$x^2 + z^2 = y^2$ or $y^2 = x^2 + z^2$

Example 2

$10^2 + 24^2 = 26^2$

Exercise 4A

1 hypotenuse, triangle
2 a c **b** x **c** u
3 a False **b** True **c** True **d** False
4 a $a^2 + b^2 = c^2$ **b** $d^2 + e^2 = f^2$ **c** $x^2 + y^2 = z^2$ **d** $s^2 + t^2 = v^2$ **e** $f^2 + g^2 = e^2$ **f** $a^2 + b^2 = x^2$
5 a $3^2 + 4^2 = 5^2$ **b** $8^2 + 15^2 = 17^2$ **c** $9^2 + 12^2 = 15^2$ **d** $5^2 + 12^2 = 13^2$ **e** $9^2 + 40^2 = 41^2$ **f** $2.5^2 + 6^2 = 6.5^2$
6 a $a^2 + b^2 = x^2$ **b** $a^2 + b^2 = d^2$ **c** $d^2 + h^2 = x^2$
7

a	b	c	a^2	b^2	$a^2 + b^2$	c^2
3	4	5	9	16	25	25
6	8	10	36	64	100	100
8	15	17	64	225	289	289

a The last two columns
b i 13 **ii** 20
c i 25 **ii** 110
8 a No
b No, $a^2 + b^2 = c^2$ should be true for a right-angled triangle.
9 a Yes **b** No **c** No **d** Yes **e** No **f** Yes
10/11 {(3, 4, 5), (6, 8, 10), (9, 12, 15), (12, 16, 20), (15, 20, 25), (18, 24, 30), (21, 28, 35), (24, 32, 40), (27, 36, 45), (30, 40, 50), (33, 44, 55), (36, 48, 60), (39, 52, 65), (42, 56, 70), (45, 60, 75), (48, 64, 80), (51, 68, 85), (54, 72, 90), (57, 76, 95)}, {(5, 12, 13), (10, 24, 26), (15, 36, 39), (20, 48, 52), (25, 60, 65), (30, 72, 78), (35, 84, 91)}, {(7, 24, 25), (14, 48, 50), (21, 72, 75)}, {(8, 15, 17), (16, 30, 34), (24, 45, 51), (32, 60, 68), (40, 75, 85)}, {(9, 40, 41), (18, 80, 82)}, {(11, 60, 61)}, {(20, 21, 29), (40, 42, 58), (60, 63, 87)}, {(12, 35, 37), (24, 70, 74)}, {(28, 45, 53)}, {(33, 56, 65)}, {(16, 63, 65)}, {(48, 55, 73)}, {(13, 84, 85)}, {(36, 77, 85)}, {(39, 80, 89)}, {(65, 72, 97)};
50 Pythagorean triples

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

4B

Now you try

Example 3

$c = 5$

Example 4

6.61

Example 5

$\sqrt{53}$

Exercise 4B

1 **a** 17 **b** 50 **c** $\sqrt{8}$
2 **a** $c^2 = 3^2 + 5^2$ **b** $c^2 = 2^2 + 7^2$
3 a, f
4 **a** $c = 6.7$ **b** $c = 6.3$ **c** $c = \sqrt{22}$ **d** $c = \sqrt{34}$
5 **a** $c = 10$ **b** $c = 13$ **c** $c = 17$ **d** $c = 15$
e $c = 25$ **f** $c = 41$ **g** $c = 50$ **h** $c = 30$
i $c = 25$
6 **a** 4.47 **b** 3.16 **c** 15.62 **d** 11.35
e 7.07 **f** 0.15
7 **a** $\sqrt{5}$ **b** $\sqrt{58}$ **c** $\sqrt{34}$ **d** $\sqrt{37}$
e $\sqrt{109}$ **f** $\sqrt{353}$
8 4.4 m
9 495 m
10 2.4 m
11 **a** 8.61 m **b** 5.24 m **c** 13.21 cm
d 0.19 m **e** 17.07 mm **f** 10.93 cm
12 250 m
13 42 units
14 5.83 m
15 **a** 77.78 cm **b** 1.39 m **c** Reduce by 7.5 cm

4C

Now you try

Example 6

$a = 7$

Example 7

$x = 6.08$

Example 8

$x = \sqrt{\frac{25}{2}}$ or $\frac{5}{\sqrt{2}}$

Exercise 4C


1 **a** subtract **b** 16 **c** 11
2 **a** T **b** F **c** T **d** F
3 **a** 4 **b** 24 **c** 8
4 **a** 16 **b** 24 **c** 6 **d** 21
e 60 **f** 27
5 **a** 8.66 **b** 11.31 **c** 5.11 **d** 17.55
e 7.19 **f** 0.74
6 5.3 m
7 14.2 m
8 49 cm
9 1.86 m
10 **a** $\sqrt{2}$ **b** $\sqrt{8}$ **c** $\sqrt{\frac{1}{2}}$
11 **a** $\sqrt{187}$ **b** $\sqrt{567}$ **c** 40

4D

Now you try

Example 9

a




b 22.25 m

Example 10

The cable is 32.02 m long

Exercise 4D

1 B
2 **a** B, III **b** C, I **c** A, II
3 **a**




b 31.22 m
4 7.07 cm
5 **a** 6.4 m **b** 5.7 cm **c** 6.3 m **d** 6.0 m
6 3.0 m
7 142.9 m
8 3823 mm
9 1060 m
10 466.18 m
11 **a** 27 m **b** 118.3 m

4E

Now you try

Example 11




Example 12


a $\cos\theta = \frac{7}{9}$ **b** $\sin\theta = \frac{6}{7}$ **c** $\tan\theta = \frac{8}{5}$

Exercise 4E

1 **a** 13 m **b** 5 m **c** 12 m
2 **a** hypotenuse **b** opposite **c** adjacent **d** opposite
e hypotenuse **f** adjacent
3 **a**




b




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


c




d




e




f



g



h



4 **a** $\sin\theta = \frac{4}{7}$ **b** $\tan\theta = \frac{5}{4}$ **c** $\cos\theta = \frac{3}{5}$

d $\sin\theta = \frac{2}{3}$ **e** $\tan\theta = 1$ **f** $\cos\theta = \frac{2}{3}$

g $\tan\theta = \frac{11}{12}$ **h** $\cos\theta = \frac{1}{2}$ **i** $\tan\theta = \frac{5}{3}$

5 **a** $\sin\theta = \frac{3}{5}$ $\cos\theta = \frac{4}{5}$ $\tan\theta = \frac{3}{4}$

b $\sin\theta = \frac{12}{13}$ $\cos\theta = \frac{5}{13}$ $\tan\theta = \frac{12}{5}$

c $\sin\theta = \frac{12}{13}$ $\cos\theta = \frac{5}{13}$ $\tan\theta = \frac{12}{5}$

6 **a** **i** $\frac{5}{13}$ **ii** $\frac{5}{13}$ **iii** The same

b **i** $\frac{12}{13}$ **ii** $\frac{12}{13}$ **iii** The same

c **i** $\frac{5}{12}$ **ii** $\frac{5}{12}$ **iii** The same


7 **a** $\frac{3}{5}$ **b** $\frac{4}{5}$ **c** $\frac{3}{4}$

8 $\frac{3}{4}$

9 **a** 5 **b** 4 **c** 3 **d** 3 **e** 4

10 **a** $\frac{4}{5}$ **b** $\frac{3}{5}$ **c** $\frac{3}{5}$ **d** $\frac{3}{4}$ **e** $\frac{4}{5}$ **f** $\frac{4}{3}$

11 **a**



b 8

c $\sin\theta = \frac{15}{17}$, $\cos\theta = \frac{8}{17}$, $\tan\theta = \frac{15}{8}$

12 **a** **i** $\frac{1}{2}$ **ii** $\frac{\sqrt{3}}{2}$ **iii** $\frac{1}{\sqrt{3}}$ **iv** $\frac{\sqrt{3}}{2}$

v $\frac{1}{2}$ **vi** $\sqrt{3}$

b **i** They are equal. **ii** They are equal.

13 **b** **i** 0.766 **ii** 0.643 **iii** 0.839 **iv** 0.766
v 1.192 **vi** 0.643

c $\sin 40° = \cos 50°$, $\sin 50° = \cos 40°$, $\tan 50° = \frac{1}{\tan 40°}$

$\tan 40° = \frac{1}{\tan 50°}$

4F

Now you try

Example 13

a 0.34 **b** 0.79 **c** 2.36

Example 14

3.83

Example 15

a 1.69 **b** 5.03 **c** 8.48

Exercise 4F

1 **a** A **b** O **c** H
2 **a** sin **b** tan **c** cos
3 B
4 **a** 0.34 **b** 0.80 **c** 2.05 **d** 0.73
e 0.10 **f** 0.25 **g** 0.46 **h** 0.24
5 **a** 3.06 **b** 18.94 **c** 5.03
d 0.91 **e** 1.71 **f** 9.00
g 2.36 **h** 4.79 **i** 7.60
6 **a** 5.95 **b** 0.39 **c** 13.38 **d** 3.83
e 8.40 **f** 1.36 **g** 29.00 **h** 1.62
i 40.10 **j** 4.23 **k** 14.72 **l** 13.42
m 17.62 **n** 5.48
7 10.39 m
8 1.12 m
9 44.99 m
10 10.11 m
11 **a** 65° **b** 2.113 **c** 2.113
12 **a** 20.95 m **b** 10 cm

Progress quiz

1 **a** $x^2 + y^2 = z^2$ **b** $r^2 + s^2 = t^2$
c $5^2 + 12^2 = 13^2$
2 **a** 15 **b** $\sqrt{73}$ **c** 7.2
3 **a** $a = 10$ **b** $b = 6.32$
4 2.2 m
5 3.46 m
6 3.3 m
7 **a** $\sin\theta = \frac{4}{7}$ **b** $\tan\theta = \frac{5}{9}$ **c** $\cos\theta = \frac{6}{11}$
8 **a** 8.19 **b** 5.96
9 2.6 m

4G

Now you try

Example 16

$x = 3.58$

Example 17

$x = 5.85$

Example 18

$x = 4.45$, $y = 10.95$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Exercise 4G

1 **a** $\cos\theta = \frac{7}{x}$ **b** $\sin\theta = \frac{10}{x}$ **c** $\tan\theta = \frac{2.3}{x}$

2 C

3 **a** $\cos 25°$ **b** $\tan 65°$ **c** $\frac{6}{\sin 72°}$

4 **a** 4.10 **b** 6.81 **c** 37.88
d 0.98 **e** 12.80 **f** 14.43
g 9.52 **h** 114.83 **i** 22.05

5 **a** 13.45 **b** 16.50 **c** 57.90 **d** 26.33
e 15.53 **f** 38.12 **g** 9.15 **h** 32.56
i 21.75 **j** 49.81 **k** 47.02 **l** 28.70

6 40 m

7 3848 m

8 **a** $x = 7.5, y = 6.4$ **b** $a = 7.5, b = 10.3$
c $a = 6.7, b = 7.8$ **d** $x = 9.5, y = 12.4$
e $x = 12.4, y = 9.2$ **f** $x = 21.1, y = 18.8$

9 **a** 27 m **b** 104 m

10 **a** B as student B did not use an approximation in their working out.
b Use your calculator and do not round sin 31° during working.

11 **a** 23.7 m **b** 124.9 m

4H

Now you try

Example 19

28°

Example 20

48.19°

Example 21

25°

Exercise 4H

1 **a** $\sin\theta = \frac{8}{9}$ **b** $\sin\theta = \frac{5}{7}$
c $\cos\theta = \frac{14}{23}$ **d** $\tan\theta = \frac{15}{26}$

2 **a** 11.54 **b** 64.53 **c** 68.20

3 **a** $\frac{1}{2}$ **b** 0.6 **c** $\tan^{-1}\left(\frac{5}{4}\right)$

4 **a** 30° **b** 60° **c** 45° **d** 30°
e 45° **f** 30° **g** 90° **h** 50°
i 90° **j** 55° **k** 0° **l** 70°

5 **a** 34.85° **b** 19.47° **c** 64.16°
d 75.52° **e** 36.87° **f** 38.94°
g 30.96° **h** 57.99° **i** 85.24°

6 **a** 43° **b** 31° **c** 41° **d** 16° **e** 55°
f 50° **g** 49° **h** 41° **i** 34°

7 17°

8 23.13°

9 **a** $\frac{1}{2}$
b 50°
c $45° = \tan^{-1}(1)$
d 0.707

10 25.4°

11 26.6°

12 45°

13 **a** 18° **b** 27° **c** 45°
d 5.67 m **e** up to 90°

4I

Now you try

Example 22


29 m

Example 23


16°

Exercise 4I

1 **a** **i**



ii



b Yes, $\theta = \alpha$, alternate angles are equal on parallel lines.

2 C

3 **a** 22° **b** 22°

4 $a = 65, b = 25$

5 16 m

6 157 m

7 38 m

8 90 m

9 37°

10 6°

11 10°

12 **a** 6° **b** 210 m

13 4634 mm


14 Yes, by 244.8 m

15 **a** **i** 47.5 km **ii** 16.25 km
b No
c Yes

Maths@Work: Carpenter

1 **a** 450 mm **b** 2400 mm **c** 9010 mm
d 2700 mm **e** 46 000 mm


2




Essential Mathematics Core for the Victorian Curriculum Year 9 ISBN 978-1-009-59315-1 © Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3 Diagrams are not to scale.

a 770 mm by 1040 mm



b 2430 mm by 1540 mm



4 **a** $BC = 72$ mm **b** $BC = 106$ mm
c $AB = 187$ mm **d** $AB = 364$ mm

5 1370 tiles

6 170 m^2

7

Location	Volume of rain in litres
Strahan, TAS	232 713
Melbourne, VIC	94 860
Port Lincoln, SA	77 724
Perth, WA	111 384
Newcastle, NSW	183 600
Tully, QLD	628 065
Alice Springs, NT	37 179

Puzzles and games

1 44

2 **a** Answer will vary **b** a^2, b^2, c^2
c $a^2 + b^2 = c^2$

3 **a** A diagonal of the square **b** $\sqrt{2}$ cm **c** 63.7%

4 Round peg square hole

5 171 cm

Chapter checklist

1 **a** $y^2 = x^2 + w^2$ **b** $5^2 = 3^2 + 4^2$

2 **a** $c = 15$ **b** $c = 7.16$

3 $c^2 = 9^2 + 2^2$, $c = \sqrt{85}$


4 $a = 6.63$

5 $\frac{7}{\sqrt{2}}$

6 $\sqrt{8} = 2.83$ m (2 decimal places)

7 $\sqrt{116} = 10.77$ m (2 decimal places)

8



9 **a** $\sin\theta = \frac{4}{9}$ **b** $\tan\theta = \frac{6}{7}$

10 0.70

11 8.48

12 9.43


13 $y = 17.01$, $x = 13.76$

14 **a** $\theta = 51°$ **b** 25°

15 66°

16 6.88 m ≈ 7 m (nearest metre)

17



Short-answer questions

1 **a** $x^2 + y^2 = z^2$ **b** $s^2 + t^2 = u^2$ **c** $e^2 + f^2 = d^2$

2 **a** 10 **b** 25 **c** 13 **d** 37
e 7.21 **f** 7.07

3 **a** 3.32 **b** 7.55 **c** 9.95

4 4.49 m

5 **a** 13 cm **b** 13.93 cm

6 19 m

7 **a** 0.64 **b** 2.25 **c** 0.72

8 **a** $\sin\theta$ **b** $\tan\theta$ **c** $\cos\theta$

9 **a** 11.33 **b** 4.88 **c** 48.02
d 10.31 **e** 50.71° **f** 60°

10 28.01 m

11 25 m

12 177.91 m


13 63.2 m

Multiple-choice questions

1 A **2** B **3** A **4** C **5** B **6** C
7 D **8** C **9** A **10** B

Extended-response questions

1 **a**



b 470 m **c** 3° **d** 1530 m


2 **a** 2.15 m **b** 0.95 m **c** 3.05 m

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Chapter 5

Warm-up quiz

1



2 **a** 1 **b** −1 **c** −3 **d** −4

3 **a** $x = 5$ **b** $x = -2$ **c** $x = -3$ **d** $x = 4$
e $x = -2$ **f** $x = 4$

4 **a** 3 **b** −4 **c** 4 **d** −3

5

	x	−2	−1	0	1	2
a	y	1	2	3	4	5
b	y	−5	−3	−1	1	3

6 **a** False **b** True **c** True **d** False

7 **a** 7 **b** 6.5 **c** 3 **d** −5

8 **a** 5 **b** 4 **c** 5 **d** 3

9 **a** 4 **b** 5 **c** 7 **d** 3

5A

Now you try

Example 1

a

x	−2	−1	0	1	2
y	−8	−5	−2	1	4


b

x	−2	−1	0	1	2
y	8	6	4	2	0

Example 2


a

x	−3	−2	−1	0	1	2	3
y	−5	−4	−3	−2	−1	0	1



b

x	−3	−2	−1	0	1	2	3
y	13	10	7	4	1	−2	−5




Example 3

a Yes **b** No


Exercise 5A

1 **a**



b **i** x-coordinate **ii** y-coordinate
iii right, down

2 **a**



b $A(4, 1)$, $B(2, 0)$, $C(0, -3)$, $D(-2, 2)$, $E(-2, -4)$, $F(3, -4)$

3 C

4 A, C

5 **a**

x	−2	−1	0	1	2	3
y	0	1	2	3	4	5

b

x	−2	−1	0	1	2
y	−1	1	3	5	7

c

x	−2	−1	0	1	2
y	4	2	0	−2	−4

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

d

x	−2	−1	0	1	2	3
y	5	2	−1	−4	−7	−10

6 a

x	−3	−2	−1	0	1	2	3
y	−4	−3	−2	−1	0	1	2


b

x	−3	−2	−1	0	1	2	3
y	0	1	2	3	4	5	6

c

x	−3	−2	−1	0	1	2	3
y	−9	−7	−5	−3	−1	1	3

a $y = x - 1$ **b** $y = x + 3$ **c** $y = 2x - 3$



d

x	−3	−2	−1	0	1	2	3
y	9	6	3	0	−3	−6	−9


e

y	−3	−2	−1	0	1	2	3
y	5	3	1	−1	−3	−5	−7

f

x	−3	−2	−1	0	1	2	3
y	7	6	5	4	3	2	1

d $y = -3x$ **e** $y = -2x - 1$ **f** $y = -x + 4$



7 a No **b** Yes **c** No
8 a Yes **b** Yes **c** No
9 a $y = x + 2$ **b** $y = 2x$ **c** $y = 2x + 1$
d $y = 3x - 1$ **e** $y = x - 2$
10 $y = 3x$
11 a iv **b** iii **c** i **d** ii
12 a $y = 2x + 7$ **b** $y = 20 - x$ **c** $y = 2x - 1$
d $y = \frac{1}{2}x + 1$ **e** $y = -3x + 4$ **f** $y = 5x + 4$

5B

Now you try

Example 4

a x-intercept is $(-1, 0)$, y-intercept is $(0, 2)$
b x-intercept is $(3, 0)$, y-intercept is $(0, 2)$

Example 5


a $(0, 5)$ **b** $(0, 3)$

Example 6

a $(2, 0)$ **b** $(-3, 0)$

Exercise 5B

1 a x-axis **b** x-coordinate **c** x-intercept
2 a A(1, 0), D(−2, 0), H(3, 0) **b** B(0, 3), E(0, −1)
3 a $y = 0$ **b** $x = 0$
4 a (1, 0), (0, 1) **b** (−2, 0), (0, 2)
c (4, 0), (0, 8) **d** (−5, 0), (0, 10)
e (2, 0), (0, 3) **f** (7, 0), (0, −3)
g (−11, 0), (0, 5) **h** (−2, 0), (0, −5)
5 a (0, 5) **b** (0, 1) **c** (0, −7) **d** (0, −3)
e (0, 2) **f** (0, −4) **g** (0, −2) **h** (0, 10)
i (0, 11) **j** (0, 3) **k** (0, 2) **l** (0, 6)
m (0, −3) **n** (0, −1) **o** (0, −3) **p** (0, 3)
6 a (−6, 0) **b** (4, 0) **c** (4, 0) **d** (−2, 0)
e (−4, 0) **f** (3, 0) **g** (7, 0) **h** (2, 0)
i (4, 0) **j** (6, 0) **k** (1, 0) **l** (−3, 0)
7 a $\left(\frac{1}{2}, 0\right)$, $(0, -1)$ **b** (4, 0), (0, −2)
c (12, 0), (0, 4) **d** $(2\frac{1}{3}, 0)$, $(0, 3\frac{1}{2})$
8 a 100 m **b** 20 s
9 a 20 minutes **b** 80 litres
10 When you substitute $x = 0$ into the rule, $y = 3 \times (0) = 0$. It means the graph passes through the origin.
11 a $A = -60$ **b** She makes a loss of $60.
c 30 badges
d She then starts to make a profit from her stall.
e




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


5C

Now you try

Example 7



Example 8




Exercise 5C

1 **a** 2 **b** 0 **c** 0


2 **a** $(0, 3)$ **b** $(0, -4)$

3 **a** $(-2, 0)$ **b** $(2, 0)$


4 **a**




b




c




d




5 **a**




b




c




d




e




f




g




h



i




j




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


k




l




6 a




b




c




d




e




f




g




h



i




j




7 a 8 m
b 4 seconds


c




8 a



b



c



9 a For 0.5 across, graph moves 1 down.
b For 1.5 across, graph moves one up.


10 a $x + y = 4$ **b** $x + y = 2$
c $x - y = 3$ **d** $x - y = -1$

5D


Now you try

Example 9


a



b




c



ISBN 978-1-009-59315-1
© Greenwood et al. 2025
Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Example 10



Example 11

a $x = 1$ **b** $y = 2$

Exercise 5D


1 **a** Horizontal **b** Horizontal **c** Vertical
d Horizontal **e** Vertical **f** Vertical

2 **a** ii, iv **b** i, iii


3 **a** **i** 2 **ii** −3
b **i** 3 **ii** −2

4 b, e, f


5 **a**




b




c




d




e




f




g




h




6 **a**




b




c




d




e




f




g



h




i



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

7



8 **a** $y = -2$ **b** $y = 4$ **c** $x = -2$
d $x = 5$ **e** $y = 1.5$ **f** $x = -6.7$
9 **a** $y = 3$ **b** $x = 5$ **c** $x = -2$ **d** $y = 0$
10 **a** $y = 250$ **b** $y = -45$
11 **a** $y = 2x$ **b** $y = 3x$
c $y = -x$ **d** $y = -3x$
12 **a** $A = 15$ square units **b** $A = 68$ square units
13 **a** **i** $y = 1$ or $y = -5$ **ii** $y = 0$ or $y = -4$
iii $y = 3\frac{1}{2}$ or $y = -7\frac{1}{2}$
b **i** $y = 1$ or $y = -5$ **ii** $y = 7$ or $y = -11$
iii $y = 9\frac{1}{2}$ or $y = -13\frac{1}{2}$

5E

Now you try

Example 12

a Zero **b** Positive, 2 **c** Undefined
d Negative, $-\frac{2}{5}$

Example 13

a $m = 2$ **b** $m = -\frac{8}{5}$

Exercise 5E

1 **a** **i** 2 **ii** 4
b **i** 2 **ii** 1
c **i** 1 **ii** 3
d **i** 3 **ii** -3
e **i** 1 **ii** -3
f **i** 4 **ii** -1
2 **a** Zero **b** Negative
c Positive **d** Undefined
3 **a** Positive, 1 **b** Positive, 2 **c** Zero
d Zero **e** Negative, $-\frac{2}{3}$ **f** Negative $-\frac{3}{4}$
g Undefined **h** Undefined **i** Positive, $\frac{1}{2}$
j Positive, 3 **k** Positive, 2 **l** Negative, -4
4 A 1 B -2 C 1
D $\frac{1}{2}$ E -2 F 3
5 **a** 2 **b** -1 **c** -2 **d** 1
e 2 **f** -2 **g** -1 **h** $\frac{1}{2}$
i $\frac{3}{2}$ **j** $-\frac{5}{2}$ **k** $\frac{1}{3}$ **l** $\frac{5}{2}$
6 **a** $-\frac{1}{20}$ **b** 17.5 or $\frac{35}{2}$ **c** $-\frac{1}{5}$ **d** 7.5 or $\frac{15}{2}$
7 40 metres
8 90 metres
9 **a** 6 **b** 1 **c** 0 **d** -4
10 **a** **i** -1 **ii** -3
b (4.5, 0)
c **i** (6, 0) **ii** (1.5, 0) **iii** (2.4, 0)

5F


Now you try

Example 14

$V = 2t$

Example 15


a 50 L/min
b




c **i** 50 **ii** $V = 50t$
d **i** 1000 L **ii** 40 mins

Exercise 5F

1 **a** No **b** Yes **c** Yes **d** No
2 **a** 60 km/h **b** 9 m/s **c** 5 L/min
d \$1.50/L **e** \$30/h **f** 3°C/min
3 **a** **i** 10 km **ii** 20 km **iii** 30 km
b 10 km/h
c 10
d They are the same.
4 **a** $d = 60t$ **b** $g = 2t$ **c** $C = 1.5n$
d $P = 20t$
5 **a** 100 L/h
b



c **i** 100 **ii** $V = 100t$
d **i** 150 L **ii** 2 h
6 **a** 25 km/h
b




c **i** 25 **ii** $d = 25t$
d **i** 62.5 km **ii** 1.6 h
7 **a** 100 km/h **b** 7 cm/s
c 2.5 cm/min **d** 49 mm/min
8 Sally
9 The panther
10 **a** 10.2 L **b** 72.25 L **c** 800 km

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

11 a $A = 12w$

b



c Yes, the rule for the graph is in the form $y = mx$.

12 a $\frac{1}{2}$ **b** $\frac{1}{4}$ **c** $\frac{3}{4}$

d 1 min 20 s or $1\frac{1}{3}$ min

13 1.2 h or 1 h 12 min

5G

Now you try

Example 16


a Gradient = 3, y-intercept = -2
b Gradient = -1, y-intercept = 0

Example 17


a $y = -3x + 5$ **b** $y = -2x + 3$

Example 18

a Gradient = 3, y-intercept = -2



b Gradient = $\frac{-5}{4}$, y-intercept = 1




Example 19

$y = -\frac{2}{3}x + 4$

Gradient = $-\frac{2}{3} = \frac{-2}{3}$

y-intercept = 4



Exercise 5G

1 a gradient **b** y-intercept **c** subject
2 a iii **b** i **c** ii **d** iv


3 a **i** (2, 6) **ii** (3, 1)
b **i** (1, 1) **ii** (3, −5)

4 a 3, 5 **b** 1, 4 **c** 2, −1


5 a Gradient = 3, y-intercept = (0, −4)
b Gradient = −5, y-intercept = (0, −2)
c Gradient = −2, y-intercept = (0, 3)
d Gradient = $\frac{1}{3}$, y-intercept = (0, 4)
e Gradient = −4, y-intercept = (0, 0)
f Gradient = 2, y-intercept = (0, 0)
g Gradient = 2.3, y-intercept = (0, 0)
h Gradient = −0.7, y-intercept = (0, 0)
i Gradient = 1, y-intercept = (0, 0)

6 a $y = -3x + 4$ **b** $y = -2x + 7$
c $y = 3x + 2$ **d** $y = 2x + 7$
e $y = 5x - 3$ **f** $y = -2x + 4$
g $y = -\frac{3}{2}x + 3$ **h** $y = 2x - \frac{5}{2}$
i $y = \frac{5}{3}x + 2$


7 a $m = 1$, y-int = (0, −2)




b $m = 2$, y-int = (0, 1)



c $m = 3$, y-int = (0, 2)




d $m = \frac{1}{2}$, y-int = (0, 2)




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


e $m = -3$, y-int $= (0, 3)$




f $m = \frac{3}{2}$, y-int $= (0, 1)$




g $m = 2$, y-int $= (0, 0)$




h $m = \frac{-4}{3}$, y-int $= (0, 0)$




i $m = \frac{-2}{3}$, y-int $= (0, 0)$




8 a $m = -1$, y-int $= (0, 4)$




b $m = 1$, y-int $= (0, 6)$




c $m = -2$, y-int $= (0, 3)$




d $m = \frac{-1}{2}$, y-int $= (0, 4)$



e $m = \frac{-2}{3}$, y-int $= (0, 2)$




f $m = \frac{-4}{3}$, y-int $= (0, 4)$




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


g $m = \frac{1}{2}$, y-int = $(0, -2)$




h $m = \frac{2}{3}$, y-int = $(0, -2)$




i $m = \frac{3}{4}$, y-int = $(0, -3)$




j $m = \frac{-1}{4}$, y-int = $(0, 0)$




k $m = \frac{1}{5}$, y-int = $(0, 0)$



l $m = 4$, y-int = $(0, 0)$



9 a Gradient = 4, V-intercept = $(0, 2)$



b Gradient is the rate the water level is increasing (the rate of rain fall in mL/h); the V-intercept is the amount of water in the gauge before the storm.

10 Yes, she loses the bet. 6 needs to be divided by 2 also to give $2x + 3$.

11 c, d, f

12 a $y = 2x + 2$, y-intercept is $(0, 2)$ **b** Expand brackets

13 a $y = 2x + 3$ **b** $y = -x + 2$

c $y = 2x + 3$ **d** $y = \frac{1}{2}x + 3$


e $y = \frac{1}{2}x + 3\frac{1}{2}$

Progress quiz

1 a i

x	−3	−2	−1	0	1	2	3
y	−7	−4	−1	2	5	8	11

ii




ISBN 978-1-009-59315-1
© Greenwood et al. 2025
Photocopying is restricted under law and this material must not be transferred to another party.

b i

x	−3	−2	−1	0	1	2	3
y	10	8	6	4	2	0	−2

ii




2 a No **b** Yes

3 a (1, 0) and (0, −2) **b** (3, 0) and (0, −5)


4 a i y-int = (0, 6) **ii** x-int = (−2, 0)

iii




b i y-int = (0, −8) **ii** x-int = (−4, 0)

iii




c i y-int = (0, −2) **ii** x-int = (5, 0)

iii




5 a 10 km **b** 20 min


c




6 a




b



c



7



8 a $m = 1$

b $m = 0$

c $m = \frac{-3}{2}$

d Undefined

e i $m = 2$ **ii** $m = \frac{7}{3}$


iii $m = -5$ **iv** $m = 0$

9 a $d = 50t$ **b** $E = 16t$ **c** $r = 8o$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

10 a 60 L/min

b



c **i** 60 **ii** $V = 60t$

d **i** 120 L **ii** 3.5 min

11 a Gradient = 2, y-intercept = $(0, -3)$


b Gradient = -4, y-intercept = $(0, 1)$

c Gradient = $\frac{1}{2}$, y-intercept = $(0, 0)$


d Gradient = 2, y-intercept = $(0, 3)$

e Gradient = 2, y-intercept = $(0, 4)$


12 a Gradient = 3, y-intercept = $(0, -2)$



b Gradient = $\frac{-1}{2}$, y-intercept = $(0, 3)$



c Gradient = $\frac{-3}{2}$, y-intercept = $(0, 4)$



5H

Now you try

Example 20

$y = 2x + 6$

Example 21

$y = 2x - 5$

Exercise 5H

1 gradient, y-intercept, point

2 **a** $y = 2x + 5$ **b** $y = 4x - 1$

c $y = -2x + 5$ **d** $y = -x - \frac{1}{2}$

3 C

4 **a** 1 **b** 3 **c** 5

5 **a** $y = 2x + 6$ **b** $y = x + 2$

c $y = -4x - 4$ **d** $y = 2x - 2$

e $y = 8x + 8$ **f** $y = -x + 4$

g $y = -2x + 6$

6 **a** $y = \frac{3}{4}x + 3$ **b** $y = -\frac{3}{4}x + 3$

c $y = -\frac{5}{4}x + 3$ **d** $y = \frac{3}{2}x + 4$

e $y = \frac{3}{5}x$ **f** $y = -\frac{1}{3}x - 1$

7 **a** $y = 3x + 5$ **b** $y = -2x - 1$

c $y = -3x + 8$ **d** $y = x - 3$

e $y = -3x + 3$ **f** $y = 5x - 1$

g $y = -x + 8$ **h** $y = -3x + 6$

i $y = -2x + 2$ **j** $y = -4x - 9$

8 $y = 5x + 1$

9 $(2.5, 0)$

10 **a** $y = 35x + 80$ **b** After 12 weeks

11 **a** **i** 2 **ii** $y = 2x + 2$

b **i** -4 **ii** $y = -4x + 11$

c **i** -1 **ii** $y = -x + 3$

d **i** 1 **ii** $y = x - 4$

12 **a** $y = -20x + 120$ **b** 120 L

13 **a** $y = 2x - 3$

b $(1.5, 0)$, $(0, -3)$

c The x-intercept is when the temperature reaches 0°C and the y-intercept represents the temperature at midnight.

5I

Now you try

Example 22

a $(4, 5)$ **b** $(-2, 2.5)$

Example 23

6.71

Exercise 5I

1 **a** 8 **b** 1 **c** -3 **d** 5.5

2 **a** 5 **b** 10 **c** 5.1 **d** 6.7

3 **a** **i** 3 **ii** 4

b **i** 2 **ii** 7

4 **a** $(3, 3)$ **b** $(2, 2)$ **c** $(1, 5)$

d $(4, 1)$ **e** $(-1, 3)$ **f** $(-1, -1)$

g $(1.5, 1.5)$ **h** $(2.5, 2)$ **i** $(0.5, 3)$

j $(-1.5, -2.5)$ **k** $(-3, -8.5)$ **l** $(0.5, -2.5)$

5 **a** 5.10 **b** 2.83 **c** 5.39 **d** 4.47

e 3.61 **f** 2.83 **g** 8.94 **h** 7.21

i 6.71

6 $(-1, 5.5)$ and $(5, 0.5)$

7 $B(8, 0)$, $A(-6, 5)$, $A(-6, 9)$

8 **a** $(-3, 1)$ **b** $(1, -4)$ **c** $(8, 3)$

9 **a** 12.8 **b** 24.2

10 **a** $(3, 5)$, $(3, -1)$, $(0, 2)$, $(6, 2)$ are the obvious points

b A circle

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

11 a $\frac{1}{3}$

b $\frac{1}{3}$


c i $(1, 1)$ **ii** $(-2, -0.5)$ **iii** $(2, 1.5)$

5J

Now you try

Example 24

a $C = 60 + 30n$ **b**



c \$150 **d** 5 hours

Exercise 5J

1 B


2 a i \$1200 **ii** \$1500 **iii** \$2200

b $A = 1000 + 100n$

3 a i dependent **ii** independent

b i Volume **ii** Height **iii** Amount of money

4 a $W = 20x + 400$ **b**




c \$640 **d** 30

5 a $C = 50n + 40$ **b** \$240 **c** \$640

6 a $C = 25n + 500$


b



c \$1500 **d** 70

7 a $C = 120t + 300$

b



c \$1020 **d** 3 h

8 a $F = 18t + 12$ **b** 3 min

9 a $V = 4000 - 20t$ **b** 2200 L

c 200 min **d** 175 min

10 a 80 km/h


b Rate changes; i.e. new gradient = 70, $d = 50 + 70t$

11 a i 20 m/s **ii** 350 m **iii** 17.5 s

b Increasing altitude (i.e. rising) at rate of 20 m/s

12 a $d = 65t$

b and **e**



c 350 km

d 4 h and 40 min ($4\frac{2}{3}$ hours)


f i 2.5 h **ii** 162.5 km

Maths@Work: Trading in foreign currencies

1


	x (input)	y (output)	Linear equation	Points
i	EUR	AUD	$y = 1.5687x$	$(0, 0)(10, 15.687)$
ii	GBP	AUD	$y = 2.0306x$	$(0, 0)(10, 20.306)$
iii	USD	CNY	$y = 6.5181x$	$(0, 0)(10, 65.181)$
iv	EUR	GBP	$y = 0.7725x$	$(0, 0)(10, 7.725)$
v	AUD	EUR	$y = 0.6375x$	$(0, 0)(10, 6.375)$
vi	AUD	USD	$y = 0.7158x$	$(0, 0)(10, 7.185)$
vii	AUD	SGD	$y = 1.0038x$	$(0, 0)(10, 10.038)$
viii	AUD	NZD	$y = 1.0753x$	$(0, 0)(10, 10.753)$

2 a i




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


ii



iii




iv



b The exchange rate is the gradient of each graph

c



d The slope of the line increases as the exchange rate increases.

3 a *See figure at bottom of page.*

b $y = 0.6235x + 79.691$; x is the date in May; y is the Japanese exchange rate for 1 Australian dollar on that date.

c The gradient tells us that the exchange rate is increasing by 0.6235 Japanese yen per day in May.

d 85.926 Japanese yen

e 133565.25 Japanese yen

f Exchange rates can vary at any time and don't necessarily follow a trend line.

Puzzles and games

1 101
2 602
3 20 days
4 (0.5, 5.5), diagonals intersect at their midpoint
5 Length AB = length $AC = \sqrt{17}$
6 31 hours

Chapter checklist

1

x	−2	−1	0	1	2
y	5	3	1	−1	−3


2 a no **b** yes

3 a x-intercept (1, 0) y-intercept (0, 4)

b x-intercept (−2, 0) y-intercept (0, 4)

4 a y-intercept (0, −4) **b** y-intercept (0, −4)


3a




ISBN 978-1-009-59315-1
© Greenwood et al. 2025
Photocopying is restricted under law and this material must not be transferred to another party.

5 **a** x-intercept $\left(\frac{4}{3}, 0\right)$ **b** x-intercept $(5, 0)$


6




7




8 **a**




b



c



9




10 **a** $y = 2$ **b** $x = -1$

11 **a** undefined **b** -3

12 gradient $m = \frac{4}{3}$

13 $d = 60t$

14 **a**




b 10 L/min and $V = 10t$ **c** 25 minutes

15 $m = 3$ y-int $(0, -2)$


16 **a** $y = -3x + 4$ **b** $y = -3x + 6$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

17 a $y = 4x - 3 \quad m = 4 \quad y$-int $(0, -3)$



b $y = \dfrac{10 - 3x}{2} \quad m = -\dfrac{3}{2}$ y-int $(0, 5)$



18 $y = 2x - 6$


19 $y = 2x - 8$

20 $M\left(3, \dfrac{5}{2}\right)$

21 distance = 6.40

22 a $C = 200 + 80t$

b




c 2.5 hours

Short-answer questions


1

x	−2	−1	0	1	2	3
y	−7	−5	−3	−1	1	3




2 a $(-2, 0)$ and $(0, 4)$ **b** $(3, 0)$ and $(0, -2)$

3 a




b




ISBN 978-1-009-59315-1
© Greenwood et al. 2025
Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


c




d




e




f




g




h



4 **a** Zero **b** Zero **c** Undefined



d Undefined **e** Positive **f** Negative




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5 **a** 2 **b** -3


6 **a** 2 **b** -1 **c** 2 **d** $\frac{-10}{3}$

7 **a** 2000 L/h **b**




c $V = 2000t$ **d** 2.5 hours

8 **a** Gradient = 2, y-intercept = (0, 3)



b Gradient = -3, y-intercept = (0, 7)



9 **a** $y = -2x + 6$, gradient = -2, y-intercept = (0, 6)

b $y = -\frac{1}{2}x + 2$, gradient = $\frac{-1}{2}$, y-intercept = (0, 2)

10 **a** $y = 2x - 2$ **b** $y = -3x + 6$

c $y = 3x - 2$

11 **a** $y = 3x + 1$ **b** $y = -2x + 3$

12 **a** $M(4, 6)$, 5.66 **b** $M(7.5,\ 4.5)$, 7.07

c $M(0, 4)$, 7.21 **d** $M(-3,\ 2.5)$, 9.85


Multiple-choice questions

1 D **2** C **3** D **4** D **5** A

6 D **7** B **8** B **9** A **10** C


Extended-response questions

1 **a** $C = 50t + 100$ **b**



c \$300 **d** $2\frac{1}{2}$ hours

2 **a**



b 42 km **c** 3 hours **d** -14 **e** 14 km/h

Semester review 1

Reviewing number

Short-answer questions

1 **a** $\frac{8}{2} = 2$ **b** $\frac{10}{10} = 1$ **c** $2\frac{3}{4}$ **d** $\frac{7}{9}$

2 **a** $\frac{3}{8}$ **b** $1\frac{4}{5}$

3 **a** 4.126 **b** 21.002 **c** 0.010

4 **a** \$160, \$640 **b** \$320, \$480 **c** \$700, \$100

5 **a** 5 : 3 **b** 55 km/h **c** 24 mL/h

6 181.3 km

7 5 kg

Multiple-choice questions

1 B **2** D **3** D **4** C **5** C

Extended-response question

a 24 km

b **i** 14 km **ii** 2 p.m.

Financial mathematics

Short-answer questions

1 **a** 60% **b** 31.25% **c** 10% **d** 25%

2 **a** \$9.60 **b** \$765 **c** \$11

3 **a** 84 m **b** \$50.76

4 4 h 48 min

5 \$892

6 \$67 484

7 **a** \$72 **b** \$872

Multiple-choice questions

1 B **2** D **3** A **4** D **5** A

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Extended-response question

a \$17 500 **b** \$23 520
c 9 years **d** 27%

Expressions and equations

Short-answer questions

1 **a** $-2x+7y$ **b** $-15mn$ **c** $\frac{y}{3}$

2 **a** $x=6$ **b** $x=9$ **c** $m=\frac{3}{8}$
d $a=3$ **e** $x>-2$ **f** $x\leqslant -3$

3 **a** $2m+1=15$ **b** \$7

4 **a** 155 **b** 18

Multiple-choice questions

1 D **2** E **3** A **4** C **5** C

Extended-response question

a $p+12$
b $p+(p+12)=38$; i.e. $2p+12=38$
c Michael scored 13 points and Chris 25 points

Pythagoras' theorem and trigonometry

Short-answer questions

1 **a** 6.3 **b** 15.1 **c** 9.9 **d** 6.3

2 **a** 8.1 **b** 11.2 **c** 4.2
d 1.5 **e** 53.1° **f** 29.5°

3 **a** 19.21 m **b** 38.7°

4 **a** $x=13$ **b** $y=14.72$

Multiple-choice questions

1 B **2** A **3** A **4** C **5** A


Extended-response question

a 17.75 m **b** 14.3° **c** 18.8 m


Linear relationships

Short-answer questions


1 **a**




b



c



d



2 **a** $\frac{2}{3}$ **b** -3 **c** -2 **d** $\frac{4}{3}$

3 **a** $y=-3x+6$ **b** $y=3x-1$


4 **a** (3, 6) and 10 units
b $\left(\frac{1}{2},3\right)$ and 5 units

Multiple-choice questions

1 C **2** D **3** A **4** B **5** A

Extended-response question

a 12 kg/h
b $S=180-12t$
c



d 15 hours
e **i** $P=40+25h$ **ii** \$415

Chapter 6

Warm-up quiz

1 **a** 4 cm^2 **b** 6 m^2 **c** 8 cm^2
2 **a** 8 cm **b** 10 m **c** 18 cm
3 **a** 3 **b** 6 **c** 27
4 **a** Cylinder **b** Circle
5 **a** 23 **b** 48 **c** 2.7
d 5.2134 **e** 50 **f** 72.16
6 7
7 **a** 30 mm **b** 2000 cm **c** 1600 m
d 2.3 cm **e** 3.167 km **f** 0.72 m
g 20 km **h** 30 m **i** 0.0756 km
8 **a** 30 cm^2 **b** 4 m^2 **c** 49 km^2
d 3 m^2 **e** 24 cm^2 **f** 66 m^2
9 $C=31.42$ m, $A=78.54$ m^2

6A

Now you try

Example 1

a 10% **b** 6%

Example 2

a 57.5 g and 58.5 g **b** 575 g and 585 g

Exercise 6A

1 **a** 1 cm **b** 0.5° C **c** 0.25 L
2 **a** 23 g **b** 1.78 L **c** 41.1 s
3 **a** \$20 **b** 25%
4 **a** yes **b** yes **c** no
d yes **e** no
5 **a** 10% **b** 5% **c** 4% **d** 12.5%
6 **a** 5% **b** 6% **c** 6.25% **d** 3.75%
7 **a** 25.45 g and 25.55 g **b** 509 g and 511 g
8 **a** 14.5 g and 15.5 g **b** 435 g and 465 g

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

9 **a** 35.5 cm, 36.5 cm
b 102.5 m, 103.5 m
c 3.15 mm, 3.25 mm
d 6.75 min, 6.85 min
e 45.235 hours, 45.245 hours
f 83.505 g, 83.515 g
g 3.1055 kg, 3.1065 kg
h 0.03565 L, 0.03575 L
i 0.015 mL, 0.025 mL
10 **a** 1.505 L and 1.515 L **b** 15.05 L and 15.15 L
11 13.9%
12 15 g
13 172.5 cm or 174.3 cm
14 **a** 22.975 km **b** 0.54%
15 **a** 60.6% **b** 2.9% **c** Answers will vary

6B

Now you try

Example 3

a 240 cm **b** 42 cm

Example 4


a 9 cm **b** 6 mm

Example 5

33.71 m

Exercise 6B

1



2 add
3 **a** 7 **b** 26 **c** 2.1
4 **a** 50 mm **b** 410 mm **c** 280 cm
d 40 cm **e** 4600 m **f** 900 m
g 52.1 cm **h** 3.6 cm **i** 2.4 m
j 0.837 m **k** 7 km **l** 2.17 km
5 825 cm, 2.25 cm
6 **a** 15 mm **b** 31 m **c** 86 cm
d 12 m **e** 27 cm **f** 24 mm
g 18 km **h** 10 m **i** 42 cm
7 **a** $x = 4$ **b** $x = 2.2$ **c** $x = 14$
8 **a** $a = 3, b = 6$ **b** $a = 12, b = 4$ **c** $a = 6.2, b = 2$
9 **a** 90 cm **b** 80 cm **c** 170 cm
d 30.57 m **e** 25.5 cm **f** 15.4 km
10 108 m
11 **a** 8000 mm **b** 110 m **c** 1 cm
d 20 mm **e** 0.284 km **f** 62.743 km
12 **a** 86 cm **b** 13.6 m **c** 40.4 cm
13 88 cm
14 **a** **i** 96 cm **ii** 104 cm **iii** 120 cm
b $P = 4(20 + 2x)$ or $P = 80 + 8x$
c 109.6 cm

6C

Now you try

Example 6

23.88 m

Example 7

20.11 km

Example 8

10.71 cm

Exercise 6C

1 **a** Radius **b** Diameter **c** Circumference
2 **a** 2.8 cm **b** 96 mm
3 **a** $C = 2\pi r$ **b** $C = \pi d$
4 **a** $\frac{1}{4}$ **b** $\frac{1}{2}$ **c** $\frac{3}{4}$
5 **a** 50.27 m **b** 87.96 cm **c** 31.42 m
d 6.91 m **e** 45.24 cm **f** 101.79 mm
6 **a** 31.42 m **b** 78.54 cm **c** 9.42 mm
d 12.57 km **e** 5.65 mm **f** 23.25 m
7 **a** 12.56 m **b** 62.8 cm **c** 22 mm **d** 44 m
8 **a** 10.28 m **b** 51.42 cm **c** 14.28 m
d 10.00 cm **e** 20.05 m **f** 106.73 km
9 28.27 m
10 4.1 m
11 **a** 12.57 cm **b** 102.83 mm **c** 41.06 m
12 **a** 188.50 cm
b **i** 376.99 cm **ii** 1979.20 cm
c 531

6D

Now you try

Example 9

a 3.2 m^2 **b** 400 000 m^2

Example 10

a 36 m^2 **b** 6.25 m^2 **c** 60 cm^2

Example 11

a 56 cm^2 **b** 15 m^2

Exercise 6D

1 **a** 100 **b** 10 000 **c** 1 000 000
2 **a** 6 **b** 16 **c** 12
d 1 **e** 12 **f** 153
3 **a** Rectangle **b** Rhombus/kite **c** Triangle
d Trapezium **e** Parallelogram **f** Square
4 **a** 200 mm^2 **b** 40 mm^2 **c** 5 cm^2
d 3.1 cm^2 **e** 21 000 cm^2 **f** 2000 cm^2
g 21 m^2 **h** 0.37 m^2 **i** 1000 m^2
j 4 300 000 m^2 **k** 3.2 km^2 **l** 0.0394 km^2
5 **a** 24 m^2 **b** 10.5 cm^2 **c** 20 km^2
d 25.2 m^2 **e** 15 m^2 **f** 36.8 m^2
6 **a** 21 mm^2 **b** 12 cm^2 **c** 17 cm^2
d 63 m^2 **e** 6.205 m^2 **f** 15.19 km^2
7 500 000 m^2
8 0.175 km^2
9 **a** 12.25 cm^2 **b** 3.04 m^2 **c** 0.09 cm^2
d 6.5 mm^2 **e** 18 cm^2 **f** 2.4613 cm^2
10 0.51 m^2
11 **a** 1.5×10^{10}(15 000 000 000) cm^2 **b** 5 mm^2
c 0.075 m^2
12 **a** **i** 1.5 m **ii** 1.5 m
b 78 m^2
c Yes

6E

Now you try

Example 12

0.50 m^2

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Example 13

25.13 cm^2

Exercise 6E

1 **a** 4.1 m **b** 7.5 m **c** 3.8 cm
2 C
3 **a** $\frac{1}{4}$ **b** $\frac{1}{2}$ **c** $\frac{3}{4}$
4 **a** 12.57 m^2 **b** 113.10 cm^2 **c** 254.47 km^2
d 60.82 mm^2 **e** 415.48 km^2 **f** 45 238.93 cm^2
5 **a** 314.16 cm^2 **b** 60.82 m^2 **c** 17.35 cm^2
6 **a** 21.23 m^2 **b** 216.51 km^2 **c** 196.07 cm^2
7 **a** 12.57 m^2 **b** 157.08 cm^2 **c** 84.82 m^2
8 177 cm^2
9 **a** 2 m **b** 12.57 m^2 **c** 3.43 m^2 **d** 21.5%
10 12.89%
11 **a** 1.8 cm **b** 6.1 m **c** 2 km
12 31%

6F

Now you try

Example 14

$P = 32$ m
$A = 64$ m^2

Example 15

$P = 142.83$ cm
$A = 171.68$ cm^2

Exercise 6F

1 **a** Semicircle and rectangle
b Triangle and semicircle **c** Rhombus and parallelogram
2 **a** $\frac{1}{2}$ **b** $\frac{1}{4}$ **c** $\frac{3}{4}$ **d** $\frac{1}{4}$
3 **a** 4 cm^2 **b** 3.14 cm^2 **c** 7.14 cm^2
4 **a** 46 m, 97 m^2 **b** 34 m, 76 m^2
c 40 m, 90 m^2 **d** 18.28 m, 22.28 m^2
e 19.42 m, 26.14 m^2 **f** 85.42 mm, 326.37 mm^2
5 **a** $P = 14.28$ m, $A = 3.43$ m^2
b $P = 35.71$ m, $A = 10.73$ m^2
c $P = 41.13$ km, $A = 27.47$ km^2
6 **a** 108 m^2 **b** 33 cm^2 **c** 98 m^2
d 300 m^2 **e** 16 cm^2 **f** 22.5 m^2
7 189.27 m^2
8 68.67 cm^2
9 **a** 90 cm^2 **b** 15 m^2 **c** 9 m^2
d 7.51 cm^2 **e** 7.95 m^2 **f** 180.03 cm^2
g 8.74 mm^2 **h** 21.99 cm^2 **i** 23.83 mm^2
10 **a** 17 cm^2 **b** 3.5 cm^2 **c** 21.74 cm^2
11 **a** 37.70 m, 92.55 m^2 **b** 20.57 mm, 16 mm^2
c 18.00 cm, 11.61 cm^2 **d** 12.57 m, 6.28 m^2
e 25.71 cm, 23.14 cm^2 **f** 33.56 m, 83.90 m^2

6G

Now you try

Example 16


A = 90 cm^2

Example 17

A = 336 mm^2


Exercise 6G

1 **a** 6 **b** 6 **c** 5
2 **a**




Cube

b



Rectangular prism

c



Triangular prism

3 **a** $A = 2 \times (8 \times 7) + 2 \times (8 \times 3) + 2 \times (7 \times 3)$
$= 112 + 48 + 42$
$= 202$ m^2
b $A = 2 \times \frac{1}{2} \times 4 \times 3 + 5 \times 7 + 4 \times 7 + 3 \times 7$
$= 12 + 35 + 28 + 21$
$= 96$ cm^2
4 **a** 52 m^2 **b** 242 cm^2 **c** 76 m^2
d 192 cm^2 **e** 68.16 m^2 **f** 85.76 m^2
5 **a** 96 cm^2 **b** 240 m^2 **c** 199.8 cm^2 **d** 238 cm^2
6 6 m^2
7 14.54 m^2
8 34 000 cm^2
9 **a** 44.4 m^2 **b** 4.44 L
10 **a** [6, 10, 14, 18, 22, 26, 30, 34, 38] **b** $S = 4n + 2$
c 402
11 **a** 39 mm^2 **b** 224 cm^2 **c** 9.01 m^2

Progress quiz

1 20%
2 25.35 cm and 25.45 cm
3 **a** 21 cm **b** 42 m
4 **a** 25.13 cm **b** 31.42 m
5 **a** 8.24 m **b** 300 mm **c** 1240 m
d 36 000 cm^2 **e** 45 000 mm^2 **f** 0.0832 m^2
6 **a** 45 cm^2 **b** 40 m^2 **c** 40 mm^2 **d** 20 cm^2

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

7 **a** 153.9 mm^2 **b** 176.7 cm^2
8 **a** 46.3 cm **b** 127.2 cm^2
9 **a** **i** 33.4 cm **ii** 64.3 cm^2
b **i** 26 m **ii** 30 m^2
10 **a** 392 cm^2 **b** 360 cm^2

6H

Now you try

Example 18


175.93 m^2

Example 19


56.27 m^2

Exercise 6H

1 **a** $2\pi rh$ **b** $2\pi r^2$
2 **a** 25.13 m
b 18.85 mm
c 26.39 cm
3 **a**



b



4 **a** 22 cm by 10 cm
b 12.57 cm by 8 cm
c 50.27 m by 5 m
5 **a** 25.13 m^2 **b** 471.24 cm^2 **c** 50.27 m^2
6 **a** 44.0 cm^2 **b** 603.2 cm^2 **c** 113.1 m^2
7 **a** 251.33 cm^2 **b** 207.35 mm^2 **c** 24.13 m^2
8 395.84 cm^2
9 **a** 54.56 m^2 **b** 218.23 m^2 **c** 63.98 cm^2
d 71.91 cm^2 **e** 270.80 m^2 **f** 313.65 km^2
10 7539.82 cm^2
11 Half cylinder is more than half surface area as it includes new rectangular surface.
12 **a** 18 849.556 cm^2
b **i** 1.88 m^2 **ii** 37.70 m^2
c 239

6I

Now you try

Example 20

a 600 mm^3 **b** 4.52 m^3

Example 21


40 m^3

Example 22


15 mm^3

Exercise 6I

1 **a** Square **b** Triangle **c** Rectangle
d Trapezium **e** Hexagon **f** Triangle
2 **a**




b




c




d



e



f



3 **a** 10 **b** 100 **c** 1000 **d** 1 000 000
e 1 000 000 000 **f** 1000 **g** 1000 **h** 1
4 **a** 3000 mm^3 **b** 300 mm^3 **c** 2 cm^3 **d** 1000 cm^3
e 8 700 000 cm^3 **f** 0.0059 m^3 **g** 10 000 m^3
h 0.0000217 km^3 **i** 0.43 m^3
5 **a** 3000 mL **b** 200 L **c** 3.5 L
d 21 mL **e** 37 kL **f** 42.9 mL
g 2 mL **h** 2000 cm^3 **i** 1000 L
6 **a** 8 cm^3 **b** 84 m^3 **c** 21 mm^3
7 **a** 16 cm^3 **b** 42.875 m^3 **c** 15 cm^3
8 **a** 4 cm^3 **b** 10.5 m^3 **c** 11.96 cm^3
d 29 cm^3 **e** 14.88 m^3 **f** 8.1351 cm^3
9 **a** 75 m^3 **b** 30 cm^3 **c** 1.25 cm^3
10 1.6 L
11 Yes, the tank only holds 20 L
12 **a** 8 L **b** 0.36 L **c** 0.48 L
13 **a** 55 m^2 **b** 825 000 L
14 **a** Answers will vary **b** $\frac{1}{3}$ **c** Answers will vary

6J

Now you try

Example 23

a 1696.46 cm^3 **b** 22 167.08 mm^3

Example 24

14 litres

Exercise 6J

1 **a** 12.57 cm^2 **b** 8.04 m^2
c 78.54 cm^2 **d** 2.54 km^2

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

2 **a** 2 L **b** 4.3 mL **c** 3700 cm^3
d 1000 L **e** 38 m^3 **f** 200 mL
3 **a** $r = 4$ m, $h = 10$ m **b** $r = 2.6$ cm, $h = 11.1$ cm
c $r = 2.9$ m, $h = 12.8$ m **d** $r = 9$ m, $h = 23$ m
e $r = 5.8$ cm, $h = 15.1$ cm
f $r = 10.65$ cm, $h = 10.4$ cm
4 **a** 226.19 cm^3 **b** 18.85 m^3 **c** 137.44 m^3
d 100.53 cm^3 **e** 8.48 m^3 **f** 68.05 m^3
5 **a** 18 L **b** 503 L **c** 20 L **d** 2 L
6 **a** 1.571 m^3 **b** 1571 L
7 Cylinder by 131 cm^3
8 **a** 25.133 m^3 **b** 25 133 L
9 37 699 L
10 Answers will vary. Require $h = 2\pi r$.
11 **a** 502.65 cm^3 **b** 1.02 m^3
c 294.52 m^3 **d** 35 342.92 m^3
e 47.12 cm^3 **f** 1017.88 cm^3

Maths@Work: Vegetable and fruit growers

1 **a** **i** 0.89 ML **ii** 60 ML **iii** 800 ML **iv** 380 ML
v 190 000 mL
b **i** 9000 m^3 **ii** 1200 m^3 **iii** 120 000 m^3
iv 980 m^3 **v** 12 000 000 m^3
2 **a** 720 m^3 **b** 400 m^3
c 1280 m^3 **d** 10 000 m^3
e 38 400 m^3
3 **a** 95 058 L; 95 kL **b** 29 452 L; 29 kL
4

Plant	Percentage moisture used by plant	Plant watering depth in mm
Onions	4%	16
Carrots	5%	30
Tomatoes	4%	48
Apple trees	6%	90
Banana trees	11%	66

5

Plant	Area of irrigation in m^2	Litres of water per plant per irrigation	Litres of water per 100 trees
Apple trees	0.1385	12	1247
Banana trees	0.3217	21	2123

Puzzles and games

1 **a** 35 **b** 19
2 8
3 314 m
4 163.4 m^2
5 7.1 cm
6 11
7 100 L

Chapter checklist

1 23.1%
2 6.65 kg to 6.75 kg
3 **a** 92 mm **b** 610 m
4 **a** 21 m **b** 36 cm
5 15.71 m
6 20.57 cm
7 **a** 532 mm^2 **b** 72.8 mm^2
8 56 m^2
9 46 cm^2
10 201.06 cm^2
11 38.48 m^2
12 $P = 35.42$ cm $A = 74.14$ cm^2
13 180 cm^2
14 108 m^2
15 439.82 cm^2
16 135.96 m^2
17 **a** 3000 cm^3 **b** 640 cm^3
18 120 m^3
19 180 cm^3
20 282.74 cm^3
21 9 L

Short-answer questions

1 **a** 380 cm **b** 1270 m **c** 4.8 cm **d** 2.73 cm^2
e 52 000 cm^2 **f** 10 000 cm^3 **g** 53.1 cm^3 **h** 3.1 L
i 43 mL **j** 2830 L **k** 4 L **l** 1000 L
2 **a** 4% **b** 40%
3 **a** 14 m **b** 51 mm **c** 16.2 cm
4 **a** 4 cm^2 **b** 1122 mm^2 **c** 30.34 mm^2
d 7.5 m^2 **e** 15 cm^2 **f** 3 cm^2
5 **a** 2.5 m^2 **b** 37.4 m^2
6 **a** $A = 28.27$ cm^2, $P = 18.85$ cm
b $A = 1.57$ m^2, $P = 5.14$ m
c $A = 2.36$ cm^2, $P = 6.71$ cm
7 **a** $P = 22.28$ m, $A = 30.28$ m^2
b $P = 45.56$ cm, $A = 128.54$ cm^2
8 **a** 46 cm^2 **b** 114 m^2
9 **a** 659.73 mm^2 **b** 207.35 m^2
10 **a** 30 cm^3 **b** 54 m^3
c 31.42 mm^3 **d** 60 m^3

Multiple-choice questions

1 E **2** B **3** C **4** E **5** B **6** A
7 D **8** B **9** E **10** C **11** E


Extended-response questions

1 **a** 135 cm^3
b 174 cm^2
c \$43.50
d **i** Yes, by ~119 cm^3 **ii** Yes, by ~52 cm^2
2 **a** 5.17 m **b** \$65 **c** 1.59 m^2, claim is correct

Chapter 7

Warm-up quiz

1



$36 = 2 \times 2 \times 3 \times 3$
2 **a** 24, 1, 12, 2, 8, 3, 6, 4 **b** 45, 1, 15, 3, 9, 5
c 2, 3 **d** 3, 5
3 **a** 7^3 **b** 5^2 2^3 **c** 3^2 **d** 4^1 or 4
4 **a** 2^3 **b** 2^6 **c** 2^5
5 **a** 25 **b** 100 **c** 1 **d** 5 **e** 1
6 **a** $3 \times 2 \times 2 = 12$ **b** $4 \times 3 \times 3 = 36$
c $5 \times 2 \times 2 \times 2 = 40$ **d** $3 \times 10 \times 10 \times 10 = 30\,000$
7 **a** 3^4 **b** 2^{12} **c** 3^2
8 **a** 0.003 **b** 0.04 **c** 0.000 002 **d** 0.0008
9 **a** 3.73 **b** 24.62 **c** 18.37 **d** 4.40
10 **a** Tens **b** Thousandths **c** Millions **d** Tenths
11 **a** 5 **b** 4 **c** 2 **d** 4

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

12 a 38 **b** 2310 **c** 0.172
d 0.0018 **e** 1000 **f** 10 000

7A

Now you try

Example 1

a $6 \times 6 \times 6$ **b** $b \times b \times b \times b$
c $mn \times mn$ **d** $3 \times x \times x \times y$

Example 2

a 8 **b** -27 **c** $\frac{9}{25}$

Example 3

a 9^3 **b** $2a^2$ **c** 11^2b^3

Example 4

a $\left(\frac{5}{7}\right)^4$ **b** $\left(\frac{2}{5}\right)^2 \times \left(\frac{4}{7}\right)^3$

Example 5

a $7^3x^3y^2$ **b** $2^37^2m^3s^2$ **c** $3^4a^4b^4$

Example 6

$2^2 \times 3 \times 5$

Example 7

a 36 **b** $-\frac{8}{27}$ **c** 33

Exercise 7A

1 a expanded **b** index **c** power
d base **e** index, power or exponent
f prime **g** prime factors
2 a 25 **b** 8 **c** 27 **d** 16
3 a 3 **b** -7 **c** $\frac{2}{3}$ **d** w
4 a 3 **b** 7 **c** 4 **d** 13
5 a 2, 3 **b** 3, 5 **c** 2, 3, 5 **d** 7, 11
6 a $4 \times 4 \times 4$ **b** $7 \times 7 \times 7 \times 7$
c $3 \times 3 \times 3 \times 3 \times 3$ **d** $5 \times 5 \times 5$
e $a \times a \times a \times a$ **f** $b \times b \times b$
g $x \times x \times x$ **h** $xp \times xp \times xp \times xp \times xp \times xp$
i $5a \times 5a \times 5a \times 5a$ **j** $3y \times 3y \times 3y$
k $4 \times x \times x \times x \times y \times y \times y \times y \times y$
l $pq \times pq$
m $-3 \times s \times s \times s \times t \times t$
n $6 \times x \times x \times x \times x \times y \times y \times y \times y \times y$
o $5 \times yz \times yz \times yz \times yz \times yz \times yz$
p $4 \times ab \times ab \times ab$
7 a 36 **b** 16 **c** 243 **d** 12
e -8 **f** -1 **g** 81 **h** 25
i $\frac{8}{27}$ **j** $\frac{9}{16}$ **k** $\frac{1}{216}$ **l** $\frac{25}{4}$
m $-\frac{8}{27}$ **n** $\frac{81}{256}$ **o** $\frac{1}{16}$ **p** $-\frac{3125}{32}$
8 a 3^3 **b** 8^6 **c** y^2 **d** $3x^3$
e $4c^5$ **f** 5^3d^2 **g** x^2y^3 **h** 7^3b^2
9 a $\left(\frac{2}{3}\right)^4$ **b** $\left(\frac{3}{5}\right)^5$ **c** $\left(\frac{4}{7}\right)^2 \times \left(\frac{1}{5}\right)^4$
d $\left(\frac{7x}{9}\right)^2 \times \left(\frac{y}{4}\right)^3$
10 a $3^3x^3y^2$ **b** $(3x)^2(2y)^2$ or $3^22^2x^2y^2$
c $(4d)^2(2e)^2$ or $4^22^2d^2e^2$ **d** $(6by)^3$ or $6^3b^33y^3$
e $(3pq)^4$ or $3^4p^4q^4$ **f** $(7mn)^3$ or $7^3m^3n^3$
11 a 2×5 **b** 2^3 **c** $2^4 \times 3^2$
d 3×5^2 **e** 3×7^2 **f** $2^2 \times 5^3$
12 a $3 \times 3 \times a \times a \times a = 3^2a^3$
b $5 \times 5 \times k \times k \times k = 5^2k^3$
c $\frac{2}{7} \times \frac{2}{7} \times \frac{2}{7} = \left(\frac{2}{7}\right)^3$
d $3p^2q = 3 \times p \times p \times q$
e $(abc)^1 = a \times b \times c$
f $4ab^2 = -2 \times (-2) \times a \times b \times b$
13 a 36 **b** -216 **c** 1 **d** $-\frac{8}{27}$
e -18 **f** 15 **g** -36 **h** 216
14 a 4 **b** 8 **c** 5
d 2 **e** -4 **f** -2
g $\frac{1}{2}$ **h** 4 **i** 1
15 a

Time in minutes	Number of bacteria	Number in index form
0	1	2^0
1	$1 \times 2 = 2$	2^1
2	$2 \times 2 = 4$	2^2
3	$2 \times 2 \times 2 = 8$	2^3
4	$2 \times 2 \times 2 \times 2 = 16$	2^4
5	$2 \times 2 \times 2 \times 2 \times 2 = 32$	2^5
6	$2 \times 2 \times 2 \times 2 \times 2 \times 2 = 64$	2^6
7	$2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2 = 128$	2^7
8	$2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2 = 256$	2^8
9	$2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2 = 512$	2^9
10	$2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2 = 1024$	2^{10}

b i 2 min **ii** 4 min **iii** 6 min
c $2^{24} = 16\,777\,216$ cells

7B

Now you try

Example 8

a 2^{11} **b** 7^6 **c** 11^8 **d** 5^9

Example 9

a a^8 **b** m^3n^7

Example 10

y^6

Example 11

a $20a^5$ **b** $4b^5$ **c** $\frac{3}{4}x^2y = \frac{3x^2y}{4}$

Example 12

a a **b** $6m^2n^2$

Exercise 7B

1 a multiply, base, add
b divide, base, subtract

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

2 **a** $3 \times 3 \times 3 \times 3 \times 3 \times 3 \times 3 = 3^6$
b $k \times k \times k \times k \times k \times k = k^5$

3 **a** $\dfrac{5 \times 5 \times \cancel{5}^{1} \times \cancel{5}^{1} \times \cancel{5}^{1}}{\cancel{5}^{1} \times \cancel{5}^{1} \times \cancel{5}^{1}} = 5^2$

b $\dfrac{a \times a \times a \times a \times \cancel{a}^{1} \times \cancel{a}^{1}}{\cancel{a}^{1} \times \cancel{a}^{1}} = a^4$

4 **a** $6^{5+7} = 6^{12}$ **b** $a^{13+2} = a^{15}$
c $5^{12-4} = 5^8$ **d** $m^{16-2} = m^{14}$

5 **a** 2^7 **b** 5^9 **c** 7^6 **d** 8^{10}
e 3^8 **f** 6^{14} **g** 3^3 **h** 6^5
i 5^3 **j** 10 **k** 9^3 **l** $(-2)^2$

6 **a** x^7 **b** a^9 **c** t^8 **d** y^5
e d^3 **f** y^7 **g** b^8 **h** q^{11}
i a^5m^4 **j** k^5p^3 **k** x^6y^8 **l** m^7e^4

7 **a** a^2 **b** x^3 **c** q^{10} **d** d
e b^5 **f** d^5 **g** a^7 **h** y^8

8 **a** $7x^7y^5$ **b** $3x^9y^4$ **c** $5x^4y^9$
d $4x^2y^5z$ **e** $15m^5$ **f** $8e^6f^4$
g $20c^7d^2$ **h** $18y^2z^7$ **i** $3m$
j $7x^3$ **k** $5y^2$ **l** $3a$
m $3m^5$ **n** $\dfrac{w}{5}$ **o** $\dfrac{a}{5}$
p $\dfrac{x^4}{9}$ **q** $\dfrac{4x^6y^3}{3}$ **r** $\dfrac{3st^2}{7}$
s $\dfrac{4mn}{3}$ **t** $5x$

9 **a** b^6 **b** y^6 **c** c^7 **d** x
e t **f** p^6 **g** d^6 **h** x^{10}
i $4x^2y^3$ **j** $6b^2g$

10 **a** $\dfrac{m^5}{n^5}$ **b** $\dfrac{x^4}{y^2}$ **c** a^3b^3 **d** $\dfrac{6a^5}{c^7}$
e $6f^6$ **f** $12x^4b^2$

11 **a** 12 **b** 8 **c** 3 **d** 3
e 1 **f** 18 **g** 12 **h** 11
i 4 **j** 15 **k** 2 **l** 39

12 **a** $7^2 = 49$ **b** 10 **c** $13^2 = 169$
d $2^3 = 8$ **e** 101 **f** $200^2 = 40\,000$
g $7 \times 31 = 217$ **h** $3 \times 50^2 = 7500$
i $-4 \times 10^3 = -4000$

13 **a** **i** $10a^{10}$ **ii** $9m^8$ **iii** $150m^5$ **iv** $60x^6y^5$
b **i** $4am$ **ii** $6xy^2$

7C

Now you try

Example 13

a a^{21} **b** $2b^8$

Example 14

a 1 **b** 2 **c** 3

Example 15

a^{17}

Example 16

a^2

Example 17

$\dfrac{3}{2}$

Exercise 7C

1 **a** multiply **b** 1
2 **a** 16, 8, 4, 2, 1 **b** 64, 16, 4, 1

3 **a** $4^2 \times 4^2 \times 4^2 = (4 \times 4) \times (4 \times 4) \times (4 \times 4) = 4^6$
b $12^3 \times 12^3 = (12 \times 12 \times 12) \times (12 \times 12 \times 12) = 12^6$
c $x^4 \times x^4 = (x \times x \times x \times x) \times (x \times x \times x \times x) = x^8$
d $a^2 \times a^2 \times a^2 \times a^2 \times a^2$
$= (a \times a) \times (a \times a) \times (a \times a) \times (a \times a) \times (a \times a) = a^{10}$

4 **a** 1 **b** 1 **c** 1 **d** 1
e 1 **f** 1 **g** 1 **h** 1

5 **a** y^{12} **b** m^{18} **c** x^{10} **d** b^{12}
e 3^6 **f** 4^{15} **g** 3^{30} **h** 7^{10}
i $5m^{16}$ **j** $4q^{28}$ **k** $-3c^{10}$ **l** $2j^{24}$

6 **a** 1 **b** 1 **c** 1 **d** 1
e 1 **f** 1 **g** 1 **h** 1
i 5 **j** -3 **k** 4 **l** -6
m 1 **n** 3 **o** 4 **p** 0

7 **a** 4^7 **b** 3^9 **c** x **d** y^{13} **e** b^{14}
f a^{10} **g** d^{24} **h** y^{16} **i** z^{25}

8 **a** b^6 **b** x^5 **c** y^6 **d** 7^2 **e** 4
f 3^8 **g** 1 **h** y^3 **i** h^2

9 **a** 2 **b** $10x$ **c** $3x^8$
d $\dfrac{d^2e}{2}$ **e** $\dfrac{2m^6n}{5}$ **f** $\dfrac{a^{12}}{8}$

10 5 ways: $(a^{16})^1 = a^{16}$, $(a^1)^{16} = a^{16}$, $(a^2)^8 = a^{16}$, $(a^8)^2 = a^{16}$, $(a^4)^4 = a^{16}$

11 **a** 4×5 not $4 + 5$, a^{20}
b Power of 2 only applies to x^4 and 4×2 not $4 + 2$, $3x^8$
c Power zero applies to whole bracket, 1

12 **a** **i** 400 **ii** 6400 **iii** 100
b **i** 800 **ii** 12 800 **iii** 102 400
c 13 years

7D

Now you try

Example 18

a $64x^3$ **b** $16y^{16}$

Example 19

a $\dfrac{a^2}{b^2}$ **b** $\dfrac{27a^3}{b^6}$

Example 20

a $16a^4b^2$ **b** $6a^{12}b^8$

Example 21

$-\dfrac{27x^6}{8y^3z^9}$

Example 22

a $18a^2b^4$ **b** x

Exercise 7D

1 **a** $a^m \times b^m$ **b** $\dfrac{a^m}{b^m}$

2 **a** $ab \times ab \times ab \times ab$
$= a \times a \times a \times a \times b \times b \times b \times b$
$= a^4 \times b^4$

b $\dfrac{x}{6} \times \dfrac{x}{6} \times \dfrac{x}{6}$
$= \dfrac{x \times x \times x}{6 \times 6 \times 6}$
$= \dfrac{x^3}{6^3}$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3 **a** x^5y^5 **b** 2^3m^3 **c** $a^4b^4c^4$ **d** $\frac{5^3}{k^3}$

e $\frac{a^5}{m^5}$ **f** $\frac{2^4}{3^4}=\frac{16}{81}$

4 **a** $6a^2b^2$

b $4x^3y^3$

c $7\times 2^2m^2=7\times 4\times m^2=28m^2$

d $4\times 3^2a^2b^2=4\times 9a^2b^2=36a^2b^2$

5 **a** $8x^3$ **b** $25y^2$ **c** $64a^3$ **d** $9r^2$

e $81b^4$ **f** $343r^3$ **g** $16h^8$ **h** $625c^8$

i $27a^6$ **j** $49p^8$ **k** $25m^6$ **l** $9y^{20}$

6 **a** $\frac{p^3}{q^3}$ **b** $\frac{x^4}{y^4}$ **c** $\frac{64}{y^3}$ **d** $\frac{625}{p^8}$

e $\frac{4}{r^6}$ **f** $\frac{s^6}{49}$ **g** $\frac{49}{x^6}$ **h** $\frac{w^8}{100}$

i $\frac{a^6}{125}$ **j** $\frac{m^8}{16}$ **k** $\frac{32m^5}{n^5}$ **l** $\frac{8a^6}{27}$

7 **a** $32x^{15}y^{10}$ **b** $9p^6q^{12}$ **c** $2x^6y^2$

d 1 **e** $-27w^9y^3$ **f** $-4p^8q^2r^2$

g $25s^{14}t^2$ **h** $8x^{12}y^3z^9$ **i** $24a^3b^6$

8 **a** $\frac{27n^9}{8m^{12}}$ **b** $\frac{16r^4}{n^4}$ **c** $\frac{9f^2}{64g^{10}}$ **d** $\frac{25w^8y^2}{4x^6}$

e $\frac{9x^2}{4y^6g^{10}}$ **f** $\frac{27k^3m^9}{64n^{21}}$ **g** $-\frac{25w^8y^2}{4z^2x^6}$ **h** $\frac{9x^4y^6}{4a^{10}b^6}$

9 **a** $9ab^2$ **b** $27ab^6$ **c** $-12a^8b^8$

d $54x^6y^9$ **e** $-64b^6c^{15}d^3$ **f** $8a^4$

g $9a^5$ **h** $-40a^{15}b^3$ **i** $160m^{15}p^5t^{10}$

j $\frac{-27x^6}{125a^{15}b^9}$ **k** $a^{11}bc^5$ **l** $x^{11}y^2z$

10 **a** **i** 8 **ii** 125

b $N=\frac{t^3}{8}$

c **i** 27 **ii** 8

d **i** 8 **ii** 2

7E

Now you try

Example 23

a $\frac{1}{6^4}$ **b** $\frac{7}{2^4}$

Example 24

a 10^3 **b** 5×9^4

Example 25

$\frac{7}{5^2}=\frac{7}{25}$

Example 26

$\frac{3}{100}=0.03$

Example 27

a $\frac{1}{32}$ **b** 54 **c** $\frac{25}{16}$

Exercise 7E

1 **a** $\frac{1}{2^2}$ **b** $\frac{1}{3^2}$ **c** $\frac{1}{5^3}$ **d** $\frac{1}{3^3}$

2 **a**

Index form	3^4	3^3	3^2	3^1
Whole number or fraction	81	27	9	3

Index form	3^0	3^{-1}	3^{-2}	3^{-3}
Whole number or fraction	1	$\frac{1}{3}$	$\frac{1}{9}=\frac{1}{3^2}$	$\frac{1}{27}=\frac{1}{3^3}$

b

Index form	10^4	10^3	10^2	10^1
Whole number or fraction	10 000	1000	100	10

Index form	10^0	10^{-1}	10^{-2}	10^{-3}
Whole number or fraction	1	$\frac{1}{10}$	$\frac{1}{100}=\frac{1}{10^2}$	$\frac{1}{1000}=\frac{1}{10^3}$

3 **a** $10^{-4}=\frac{1}{10^4}$ **b** $3^{-2}=\frac{1}{3^2}$ **c** $7^{-3}=\frac{1}{7^3}$

4 **a** $\frac{1}{3^{-4}}=3^4$ **b** $\frac{1}{7^{-6}}=7^6$ **c** $\frac{1}{8^{-3}}=8^3$

5 **a** $\frac{1}{5^2}$ **b** $\frac{1}{7^4}$ **c** $\frac{1}{8^3}$ **d** $\frac{1}{3^5}$

e $\frac{1}{9^2}$ **f** $\frac{1}{10^3}$

6 **a** $\frac{3}{2^4}$ **b** $\frac{5}{4^3}$ **c** $\frac{7}{5^6}$ **d** $\frac{2}{3^4}$ **e** $\frac{4}{3^5}$

f $\frac{9}{5^2}$ **g** $\frac{8}{7^3}$ **h** $\frac{6}{5^6}$ **i** $\frac{1}{4^2}$

7 **a** 2^4 **b** 3^2 **c** 4^3 **d** 6^5

e 5^3 **f** 8^5 **g** 7^3 **h** 9^4

8 **a** 6×4^3 **b** 5×8^2 **c** 4×7^5 **d** 3×2^5

e 12×5^4 **f** 6×10^3 **g** 7×9^4 **h** 7×10^6

9 **a** $\frac{6}{25}$ **b** $\frac{2}{9}$ **c** $\frac{4}{125}$ **d** $\frac{6}{7}$

e $\frac{1}{250}$ **f** $\frac{1}{5}$ **g** $\frac{5}{16}$ **h** $\frac{4}{5}$ **i** $\frac{1}{9}$

10 **a** $\frac{2}{1000}=0.002$ **b** $\frac{5}{100}=0.05$

c $\frac{7}{10}=0.7$ **d** $\frac{3}{10\,000}=0.0003$

e $\frac{5}{10\,000}=0.0005$ **f** $\frac{8}{100\,000}=0.00008$

g $\frac{2}{1\,000\,000}=0.000002$

h $\frac{4}{100\,000\,000}=0.00000004$

11 **a** $\frac{1}{5}$ **b** $\frac{1}{9}$ **c** $\frac{1}{25}$ **d** $-\frac{1}{200}$

e $\frac{1}{2}$ **f** $\frac{1}{36}$ **g** $\frac{1}{8}$ **h** 8

i 16 **j** 64 **k** $\frac{64}{9}$ **l** $\frac{27}{64}$

12 **a** Negative power only applies to x, $\frac{2}{x^2}$

b $5=5^1$ has a positive power, $5a^{-4}$

c $\frac{2}{3^{-2}b^{-2}}=2\times 3^2\times b^2=18b^2$

13 **a** 1.95 g **b** $3\,486\,784.401t$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Progress quiz

1 **a** $b \times b \times b \times b$ **b** $2 \times x \times x \times y \times y \times y$
c $3x \times 3x \times 3x$ **d** $4 \times 4 \times 4 = 64$
e $-5 \times (-5) \times (-5) \times (-5) = 625$
f $\frac{3}{5} \times \frac{3}{5} \times \frac{3}{5} = \frac{27}{125}$

2 **a** 5^2y^4 **b** $\left(\frac{2}{3}\right)^3 \times \left(\frac{5}{7}\right)^2$ **c** $4^3a^2b^4$

3 **a** $45 = 3^2 \times 5$ **b** $120 = 2^3 \times 3 \times 5$

4 **a** -27 **b** $\frac{1}{16}$ **c** 48

5 **a** 4^7 **b** 9^4 **c** x^8
d a^5b^5 **e** y^6 **f** $3a^3$
g $12m^3n^7$ **h** $\frac{x^3y^2}{2}$ **i** $-2ab$

6 **a** x^8 **b** $5y^9$ **c** 1 **d** 1
e -1 **f** 5

7 **a** $12mn^3$ **b** $\frac{x^2}{6}$ **c** $\frac{2a^2}{3}$

8 **a** $4x^2$ **b** $16m^{12}$ **c** $\frac{x^8}{81}$
d $-32a^3b^6$ **e** $\frac{27a^6}{125}$ **f** $\frac{4a^4c^2}{49b^6}$

9 **a** $\frac{2y^2}{x^3}$ **b** $\frac{27x^2}{32}$

10 **a** $\frac{1}{5^2}$ **b** $\frac{3}{4^4}$
c 3^2 **d** 5×2^3

11 **a** $\frac{1}{81}$ **b** $\frac{5}{4}$ **c** 96
d 0.0006 **e** $\frac{125}{64}$

7F

Now you try

Example 28

7.1×10^4

Example 29

2.75×10^{-5}

Example 30

24 000

Example 31

0.0094

Exercise 7F

1

Scientific notation	Power of 10 expanded	Basic numeral
5×10^3	5×1000	5000
3×10^4	$3 \times 10\,000$	30 000
2×10^5	$2 \times 100\,000$	200 000
7×10^2	7×100	700
7×10^4	$7 \times 10\,000$	70 000
4×10^5	$4 \times 100\,000$	400 000

2

Scientific notation	Positive power	Fraction	Basic numeral
2×10^{-4}	$\frac{2}{10^4}$	$\frac{2}{10\,000}$	0.0002
3×10^{-2}	$\frac{3}{10^2}$	$\frac{3}{100}$	0.03
5×10^{-3}	$\frac{5}{10^3}$	$\frac{5}{1000}$	0.005
7×10^{-6}	$\frac{7}{10^6}$	$\frac{7}{1\,000\,000}$	0.000007
9×10^{-3}	$\frac{9}{10^3}$	$\frac{9}{1000}$	0.009
8×10^{-2}	$\frac{8}{10^2}$	$\frac{8}{100}$	0.08

3 **a** 10 000 **b** 1000 **c** 100 000 **d** 1000
e 100 000 **f** 10 000

4 **a** Positive **b** Negative **c** Positive **d** Negative

5 **a** 4×10^4 **b** 2.3×10^{12} **c** 1.6×10^{10} **d** 7.2×10^6
e 3.5×10^3 **f** 8.8×10^6 **g** 5.2×10^3 **h** 3×10^6
i 2.1×10^4

6 **a** 3×10^{-6} **b** 4×10^{-4} **c** 8.76×10^{-3}
d 7.3×10^{-10} **e** 3×10^{-5} **f** 1.25×10^{-10}
g 8.09×10^{-9} **h** 2.4×10^{-8} **i** 3.45×10^{-5}

7 **a** 6×10^3 **b** 7.2×10^5 **c** 3.245×10^2
d 7.86903×10^3 **e** 8.45912×10^3 **f** 2×10^{-1}
g 3.28×10^{-4} **h** 9.87×10^{-3} **i** 1×10^{-5}
j 4.601×10^8 **k** 1.7467×10^4 **l** 1.28×10^2

8 **a** 57 000 **b** 3 600 000 **c** 430 000 000
d 32 100 000 **e** 423 000 **f** 90 400 000 000
g 197 000 000 **h** 709 **i** 635 700

9 **a** 0.00012 **b** 0.0000046 **c** 0.0000000008
d 0.0000352 **e** 0.3678 **f** 0.000000123
g 0.00009 **h** 0.05 **i** 0.4

10 **a** 6×10^{24} kg **b** 4×10^7 m
c 1×10^{-10} m **d** 1.5×10^8 km
e 6.67×10^{-11} Nm^2/kg^2 **f** 1.5×10^{-4} s
g 4.5×10^9 years

11 **a** 4 600 000 000 km **b** 8 000 000 000 000 organisms
c 384 000 km **d** 0.0038 m
e 0.00000000000001 m **f** 720 000 people

12 **a** 1.62×10^9 km **b** 2.126×10^{-2} g

13 **a** 3.2×10^4 **b** 4.1×10^6 **c** 1.3×10^4 **d** 9.2×10^1
e 6.1×10^{-2} **f** 4.24 **g** 2×10^{-5} **h** 4×10^{-6}

7G

Now you try

Example 32

a 4 **b** 2 **c** 3

Example 33

a 4.72×10^5 **b** 3.25×10^{-4}

Example 34

9.942×10^7

Example 35

1.653×10^{-2}

Exercise 7G

1 **a** **i** 300 **ii** 32 700 **iii** 18 400
b **i** 0.1 **ii** 0.2 **iii** 21.0
c **i** 268 000 **ii** 38 000 **iii** 4 061 000

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

2 42, 0.0071
3 **a** 57 260, 57 300, 57 000, 60 000
b 0.003661, 0.00366, 0.0037, 0.004
4 **a** Yes **b** No **c** No
d No **e** Yes **f** Yes
5 **a** 3 **b** 4 **c** 5 **d** 2
e 3 **f** 2 **g** 3 **h** 3
i 3 **j** 4 **k** 3 **l** 3
6 **a** 2.42×10^5 **b** 1.71×10^5 **c** 2.83×10^3 **d** 3.25×10^6
e 3.43×10^{-4} **f** 6.86×10^{-3} **g** 1.46×10^{-2} **h** 1.03×10^{-3}
i 2.34×10^1 **j** 3.26×10^2 **k** 1.96×10^1 **l** 1.72×10^{-1}
7 **a** 4.78×10^4 **b** 2.2×10^4 **c** 4.833×10^6
d 3.7×10^1 **e** 9.95×10^1 **f** 1.443×10^{-2}
g 2×10^{-3} **h** 9×10^{-2} **i** 1×10^{-4}
8 **a** 2.441×10^{-4} **b** 2.107×10^{-6}
c -4.824×10^{15} **d** 4.550×10^{-5}
e 1.917×10^{12} **f** 1.995×10^8
g 3.843×10^2 **h** 1.804×10^{-11}
i 1.524×10^8 **j** 3.325×10^{15}
k 4.067×10^3 **l** -9.077×10^{-1}
9 **a** 9.3574×10^1 **b** 2.1893×10^5 **c** 8.6000×10^5
d 8.6288×10^{-2} **e** 2.2985×10^{15} **f** 3.5741×10^{28}
g 6.4000×10^7 **h** 1.2333×10^9 **i** 1.8293
j 5.4459×10^{-1}
10 1.98×10^{30} kg
11 1.39×10^6 km
12 $2421 \times 10^3, 24.2 \times 10^5, 2.41 \times 10^6, 0.239 \times 10^7, 0.02 \times 10^8$
13 **a** 4.26×10^6 **b** 9.1×10^{-3} **c** 5.04×10^{11}
d 1.931×10^{-1} **e** 2.1×10^6 **f** 6.14×10^{-11}
14 **a** $5.400\,46 \times 10^{12}$
b **i** 1.08×10^{13} **ii** 4.32×10^{13}

Maths@Work: Lab technician

1 **a** $1 \times 10^2 \mu m^3$ **b** $4 \times 10^3 \mu m^3$
c $1.3 \times 10^2 \mu m^3$ **d** $1 \times 10^3 \mu m^3$ **e** $6 \times 10^5 \mu m^3$
2 **a** 0.00012 m **b** 1.2×10^{-4} m
3 **a** 2×10^{-9} g **b** 1×10^{-6} g **c** 3 g
4 **a** 45 000 000 000 000 cells **b** 29 000 000 000 000 cells
5 **a** 5.7×10^3 years **b** 4.5×10^9 years
c 1.57×10^7 years **d** 1.5×10^{-4} seconds
6 **a** 2.08×10^6 days **b** 1.64×10^{12} days **c** 5.73×10^9 days
7 Just under 15.7 hours
8 **a** 4.00×10^{-3} g **b** 1.05×10^3 g
9 $\frac{1}{2}, \frac{1}{2^2}, \frac{1}{2^3}, \frac{1}{2^4}, \frac{1}{2^5}, \frac{1}{2^6}, \frac{1}{2^7}, \frac{1}{2^8}, \frac{1}{2^9}, \frac{1}{2^{10}}$ g

$2^{-1}, 2^{-2}, 2^{-3}, 2^{-4}, 2^{-5}, 2^{-6}, 2^{-7}, 2^{-8}, 2^{-9}, 2^{-10}$ g

10 **a**

	Quantity remaining in mg			
Radioactive isotope	After 1 half-life	After 2 half-lives	After 4 half-lives	After 10 half-lives
Hydrogen-3	4.0000	2.0000	0.5000	0.0078
Carbon-11	32.0640	16.0320	4.0080	0.0626
Sodium-24	12.3650	6.1825	1.5456	0.0242
Iron-59	3.9300	1.9650	0.4913	0.0077
Radium-226	2.1250	1.0625	0.2656	0.0042

b

	Total time			
Radioactive isotope	For 2 half-lives	For 4 half-lives	For 6 half-lives	For 10 half-lives
Hydrogen-3	24.64	49.28	73.92	123.2
Carbon-11	40.6	81.2	121.8	203
Sodium-24	29.902	59.804	89.706	149.51
Iron-59	88.99	177.98	266.97	444.95
Radium-226	3200	6400	9600	16000

Puzzles and games

1 EXPONENTIAL
2 **a** $10^8 = 100\,000\,000$ **b** $100^5 = 10\,000\,000\,000$
4 $4a \times (3m)^3 \times \left(\frac{a}{6m}\right)^2 = 3a^3m$
5

Pile	Height	Value
1	2 mm	$0.10
5	3.2 cm	$1.60
10	1.024 m	$51.20
15	32.77 m	$1638.40
20	1.05 km	$52 428.80
25	33.55 km	$1 677 721.60
30	1073.74 km	$53 687 091.20

6 14th day
7 **a** 1.84×10^{11} tonnes
b $50 729 000 000 000 or more than 50 trillion dollars

Chapter checklist

1 **a** $8a^3b^3$ **b** $\frac{4}{49}$
2 **a** $\left(\frac{7}{9}\right)^4$ **b** $2a^3b^2$
3 $2 \times 2 \times 23 = 2^2 \times 23$
4 **a** -216 **b** $\frac{9}{121}$
5 **a** 4^7 **b** 7^4
6 **a** a^{10} **b** $27x^5$
7 y^4
8 $\frac{3ab}{2}$
9 4
10 $5x^{18}$
11 **a** 1 **b** $\frac{7a^2}{2}$
12 **a** $8a^{12}$ **b** $\frac{16b^8}{a^{12}}$
13 **a** $-x^4y^6$ **b** a^2b^4
14 **a** $\frac{4}{x^2}$ **b** $5 \times 2^3 = 40$
15 **a** $\frac{5}{8}$ **b** $\frac{1}{25}$
16 **a** 4900 **b** 0.00000301
17 **a** 2.7×10^4 **b** 3.75×10^{-5}
18 **a** 9.14×10^6 **b** 3.20×10^{-4}
19 8.750×10^{-1} or 0.8750

Short-answer questions

1 **a** 3^4 **b** $2x^3y^2$ **c** $3a^2b^2$ **d** $\left(\frac{3}{5}\right)^3 \times \left(\frac{1}{7}\right)^2$
2 **a** $3^2 \times 5$ **b** $2^2 \times 3 \times 5^2$
3 **a** x^{10} **b** $12a^5b^6c$ **c** $24m^7n^4$
d a^9 **e** x^3y^2 **f** $\frac{a^2b^2}{2}$
4 **a** m^6 **b** $9a^8$ **c** $-32a^{10}b^5$
d $3b$ **e** 2 **f** $\frac{a^6}{27}$
5 **a** $\frac{1}{2^3}$ **b** $\frac{3}{4^5}$ **c** $\frac{6}{10^3}$
d 4^2 **e** 6×5^3 **f** 2×10^5
6 **a** k^6 **b** $6a^2m$ **c** $4a^{10}$
d h^{16} **e** $\frac{a^6b^2}{5}$ **f** $45p^4q^2$
7 $0.0012, 35.4 \times 10^{-3}, 3.22 \times 10^{-1}, 0.4, 0.007 \times 10^2, 2.35$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

8 **a** 324 **b** 172 500 **c** 0.2753
d 0.00149
9 **a** 2.25×10^7 people **b** 9.63×10^6 km^2
c 3.34×10^{-9} s **d** 2.94×10^{-7} m
10 **a** 2.19×10^5 h **b** 3.6×10^7 months
c 4.3×10^5 kg **d** 5×10^6 g

Multiple-choice questions

1 D **2** B **3** E **4** C **5** D **6** B
7 D **8** C **9** C **10** E

Extended-response questions

1 **a** 1.2×10^{55} kg^2 **b** 4.3×10^{-5} m/s
2 **a** **i** 5.79×10^7 km **ii** 1.50×10^8 km
iii 7.78×10^8 km **iv** 1.43×10^9 km
b **i** 2.41×10^{-1} **ii** 1.88 **iii** 2.95×10^1
c **i** 4.87×10^{24} kg **ii** 6.42×10^{23} kg
iii 1.90×10^{30} kg

Chapter 8

Warm-up quiz

1 **a** Obtuse **b** Acute **c** Reflex
d Right **e** Straight **f** Revolution
2 **a** 35 **b** 142 **c** 260
3 **a** Isosceles **b** Equilateral **c** Scalene
4 **a** Obtuse **b** Acute **c** Right
5 **a** 60° **b** 75° each **c** 50°
6 **a** Parallelogram, rectangle, kite
b Parallelogram, rectangle, square, rhombus
c Square, rectangle
7 **a** $a = 60$ **b** $b = 110$ **c** $a = 60, b = 120$
8 **a** Pentagon **b** Parallelogram **c** Trapezium

8A

Now you try

Example 1

a 106° **b** 16°

Example 2

a Obtuse scalene, $a = 35$ **b** Right isosceles, $b = 45$
c Acute scalene, $c = 65$

Example 3

a $b = 55$ **b** $a = 40$

Exercise 8A

1 **a** right **b** 180° **c** revolution
d Obtuse **e** Acute **f** 180° **g** 90°
h Supplementary **i** 180° **j** equal
2 **a** Isosceles triangle **b** Obtuse-angled triangle
c Equilateral triangle **d** Isosceles triangle
e Acute-angled triangle **f** Scalene triangle
g Right-angled triangle **h** Right-angled isosceles triangle
3 **a** 50° **b** 90° **c** 101°
d 202° **e** 180° **f** 360°
4 **a** **i** 125° **ii** 35°
b **i** 149° **ii** 59°
c **i** 106° **ii** 16°
d **i** 170° **ii** 80°
e **i** 91° **ii** 1°
f **i** 158° **ii** 68°
g **i** 142° **ii** 52°
h **i** 115° **ii** 25°
i **i** 133° **ii** 43°
j **i** 103° **ii** 13°
5 **a** $a = 63$ **b** $a = 71$ **c** $a = 38$
d $a = 147$ **e** $a = 233$ **f** $a = 33$
6 **a** Obtuse isosceles, $a = 40$
b Acute scalene, $b = 30$ **c** Right-angled scalene, $c = 90$
d Equilateral, $d = 60$ **e** Obtuse isosceles, $e = 100$
f Right-angled isosceles, $f = 45$
g Obtuse scalene, $g = 100$ **h** Equilateral, $h = 60$
i Obtuse isosceles, $i = 120$ **j** Obtuse isosceles, $j = 35$
k Right-angled scalene, $k = 90$
l Acute scalene, $l = 70$
7 **a** **i** $\angle BAC$ **ii** Obtuse **iii** 120°
b **i** $\angle PRQ$ **ii** Acute **iii** 30°
c **i** $\angle XYZ$ **ii** Reflex **iii** 315°
d **i** $\angle SRT$ **ii** Straight angle **iii** 180°
e **i** $\angle ROB$ **ii** Obtuse **iii** 103°
f **i** $\angle AOB$ **ii** Right **iii** 90°
8 **a** $s = 120$ **b** $t = 20$
c $r = 70$ **d** $a = 60, x = 120$
e $a = 100, b = 140$ **f** $c = 115, d = 65$
g $x = 56$ **h** $x = 155$
9 $AO = BO$ (radii)
ΔAOB is isosceles, 2 sides equal, $\angle AOB = 116°$
$\angle OAB = 32°$, base angles of isosceles triangle
10 **a** 360° **b** 90° **c** 60°
d 90° **e** 432° **f** 6°
g 720° **h** 8640°
11 **a** 90° **b** 150° **c** 15°
d 165° **e** 157.5° **f** 80°
g 177.5° **h** 171°

8B

Now you try

Example 4

a No. The two cointerior angles are not supplementary.
b Yes. The alternate angles are equal.
c Yes. The corresponding angles are equal.

Example 5

a $a = 59$ (supplementary) $b = 59$ (alternate)
b $a = 128$ (cointerior) $b = 128$ (vertically opposite)

Exercise 8B

1 **a** Corresponding **b** Alternate
c Cointerior **d** Alternate
e Cointerior **f** Corresponding
2 **a** equal **b** equal **c** supplementary
3 **a** No, alternate angles are not equal.
b Yes, corresponding angles are equal.
c Yes, alternate angles are equal.
d No, cointerior angles don't add to 180°.
e Yes, cointerior angles add to 180°.
f Yes, corresponding angles are equal.
g No, corresponding angles are not equal.
h No, alternate angles are not equal.
i No, cointerior angles do not add to 180°.
4 **a** $x = 125$, alternate angles in $\parallel$ lines
b $y = 110$, cointerior angles in $\parallel$ lines
c $r = 80$, corresponding angles in $\parallel$ lines
d $s = 66$, alternate angles in $\parallel$ lines
e $v = 106$, corresponding angles in $\parallel$ lines
f $q = 116$, cointerior angles in $\parallel$ lines

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5 **a** $a = 60, b = 120$ **b** $c = 95, d = 95$
c $e = 100, f = 100, g = 100$ **d** $a = 110, b = 70$
e $a = 100, b = 80, c = 80$
f $e = 140, f = 140, d = 140$
6 $a = 40, b = 140, c = 40, d = 40, e = 140$
7 **a** $x = 70, y = 40$ **b** $t = 58, z = 122$
c $u = 110, v = 50, w = 50$ **d** $x = 118$
e $x = 295$ **f** $x = 79$
8 **a** 56 **b** 120 **c** 265
9 **a** 105° **b** 105° **c** 56°
d 105° **e** 90° **f** 85°

8C

Now you try

Example 6

a Parallelograms including rhombus, rectangle and square
b Kite

Example 7

$x = 80$

Example 8

a $m = 40$ **b** $n = 104$

Exercise 8C

1 Parallelogram, rhombus, rectangle, square, kite, trapezium
2 **a** Yes **b** No **c** Yes **d** Yes **e** No
3 **a** parallel **b** right **c** trapezium **d** equal
4 **a** Parallelogram, rectangle, kite **b** Rectangle, square
c Square, rectangle **d** Square, rhombus, kite
e Trapezium
f Parallelogram, rhombus, kite, trapezium
5 **a** 115 **b** 159 **c** 30
d 121 **e** 140 **f** 220
6 **a** $a = 110$ **b** $x = 70$ **c** $b = 54$
d $a = 33$ **e** $y = 63$ **f** $a = 109$
7 **a** Yes **b** No **c** Yes **d** No
8 60°
9 A parallelogram has opposite sides parallel and equal and rectangles, squares and rhombi have these properties (and more) and are therefore all parallelograms.
10 **a** 110 **b** 55 **c** 120
11 **a** 255 **b** 80 **c** 115
d 37 **e** 28 **f** 111

8D

Now you try

Example 9

$S = 900°$
$a = 250$

Example 10

$a = 144$

Exercise 8D

1 **a** hexagon **b** regular
c non-convex **d** $S = (n - 2) \times 180°$
2 **a** 5 **b** 7 **c** 4 **d** 11 **e** 9 **f** 12
3 **a** 720° **b** 1080° **c** 1620° **d** 900°
4 **a** Convex quadrilateral **b** Non-convex hexagon
c Non-convex heptagon **d** Non-convex decagon
5 **a** 140 **b** 100 **c** 100 **d** 120
6 **a** 110 **b** 150 **c** 210
d 130 **e** 25 **f** 285
7 **a** 90° **b** 108° **c** 140°
d 120° **e** 128.57° **f** 135°
8 **a** 16 **b** 25 **c** 102
9 **a** Yes **b** Yes
10 **a** 105 **b** 240 **c** 85 **d** 150
11 **a** **i** No **ii** Yes **iii** No
b **i** One **ii** Two **iii** Five
c $(n - 3)$

8E

Now you try

Example 11

a RHS **b** SSS **c** SAS **d** AAS

Example 12

a $x = 11, y = 5$ **b** $a = 71, b = 38$

Exercise 8E

1 **a** **i** Y **ii** X **iii** Z
b **i** XY **ii** XZ **iii** YZ
c **i** $\angle A$ **ii** $\angle B$ **iii** $\angle C$
2 **a** size **b** ΔSTU **c** SAS, RHS, AAS
3 **a** $\Delta ABC \equiv \Delta FGH$ **b** $\Delta DEF \equiv \Delta STU$
4 **a** SAS **b** AAS **c** RHS **d** SAS
e SSS **f** RHS **g** AAS **h** SSS
5 **a** $x = 3, y = 4$ **b** $x = 2, y = 6$
c $a = 105, b = 40$ **d** $a = 65, b = 85$
e $x = 2.5, b = 29$ **f** $a = 142, x = 9.21, b = 7$
g $y = 4.2, a = 28$ **h** $a = 6.5, b = 60$
6 **a** A and C **b** RHS
7 **a** $\Delta ABC \equiv \Delta STU$, RHS **b** $\Delta DEF \equiv \Delta GHI$, SSS
c $\Delta ABC \equiv \Delta DEF$, SAS **d** $\Delta ABC \equiv \Delta GHI$, AAS
8 It is SAS; the hypotenuse is not given.
9 No – they can all be different sizes, one might have all sides 2 cm and another all sides 5 cm.
10 **a** SSS **b** Equal
11 $\Delta PBR \equiv \Delta FDE$
$\Delta LMN \equiv \Delta KIJ$
$\Delta FGH \equiv \Delta BCD$
$\Delta MNO \equiv \Delta RQP$

Progress quiz

1 **a** 27° **b** 117°
2 **a** Right-angled scalene, $a = 55$
b Acute-angled isosceles, $b = 40$
c Obtuse-angled scalene, $c = 154$
3 **a** Alternate angles are not equal.
b Cointerior angles are not supplementary.
4 **a** $a = 55$ (supplementary angles), $b = 55$ (corresponding angles in parallel lines)
b $a = 65$ (cointerior angles in parallel lines), $b = 65$ (vertically opposite)
c $a = 40$ (alternate angles in parallel lines and angle sum of a triangle)
5 **a** $a = 100$ **b** Parallelogram, $b = 58$
c Kite, $c = 80$
6 **a** $S = 720, a = 109$ **b** $S = 540, a = 108$
7 **a** SAS **b** SSS
8 $x = 6, y = 40$

8F

Now you try


Example 13

a 2.5 **b** 40 **c** 13


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Example 14

a



b



Exercise 8F

1 **a** image **b** shape **c** $|||$ or $\sim$
2 **a** $\angle F$ **b** $\angle D$ **c** GH **d** AE **e** 2
3 **a** OA' is double OA **b** OB' is double OB
c OC' is double OC **d** 2 **e** Yes
4 **a** OA' is a quarter of OA **b** OD' is a quarter of OD
c $\frac{1}{4}$ **d** Yes
5 **a** **i** 2 **ii** 14 **iii** 10
b **i** 1.5 **ii** 10.5 **iii** 8
c **i** 1.5 **ii** 45 **iii** 24
d **i** 2.5 **ii** 1 **iii** 1.4
e **i** 2.5 **ii** 0.6 **iii** 2
f **i** 1.75 **ii** 3.5 **iii** 3
6 **a** $A'B'C'$ should have sides $\frac{1}{3}$ that of ABC.
b $A'B'C'$ should have sides double that of ABC.
7 **b** **i** $A'B'C'D'$ should have sides lengths $\frac{1}{2}$ that of $ABCD$
ii $A'B'C'D'$ should have side lengths 1.5 times that of $ABCD$
8 **a** Yes **b** 8 cm **c** 25 cm
9 3.7 cm
10 **a** **i** 2 **ii** $(0, 0)$
b **i** $\frac{1}{2}$ **ii** $(0, 0)$
c **i** $\frac{1}{2}$ **ii** $(3, 0)$
d **i** 3 **ii** $(1, 0)$
11 **a** All angles of any square equal 90°, with only 1 side length.
b All angles in any equilateral triangle equal 60°, with only 1 side length.
c The ratio of the length to the width may vary between different rectangles.
d 2 isosceles triangles do not have to have equal angles.
12 **a** 12.7 cm **b** 3 cm **c** 3 m
13 **a** 100 000 cm = 1 km **b** 24 cm

8G

Now you try

Example 15

a SAS **b** AAA **c** SSS **d** RHS

Example 16

a AAA **b** 2.5 **c** 4.8

Exercise 8G

1 **a** SAS, AAA, and RHS **b** shape, size
c triangle **d** similar
2 **a** E **b** C **c** DF
d BC **e** $\angle A$ **f** $\angle E$
3 **a** AAA **b** RHS **c** SSS **d** SAS
e RHS **f** AAA **g** SAS **h** SSS
4 **a** $\Delta ABC \,|||\, \Delta GHI$ **b** $\Delta ABC \,|||\, \Delta MNO$
c $\Delta ABC \,|||\, \Delta ADE$ **d** $\Delta HFG \,|||\, \Delta HJI$
e $\Delta ADC \,|||\, \Delta AEB$ **f** $\Delta ABD \,|||\, \Delta ECD$
5 2.5
6 **a** AAA **b** 1.5 **c** 12
7 **a** RHS **b** 2.5 **c** 8
8 2 m
9 **a** ΔDEF **b** ΔDEF **c** ΔABC **d** ΔDEF
10 **a** **i** AAA **ii** 6.5
b **i** AAA **ii** 10
c **i** AAA **ii** 24
11 $\angle ACB = 25°$, AAA
12 **a**

Triangle	Original	Image 1	2	3
Length scale factor	1	2	3	4
Area	4	16	36	64
Area scale factor	1	4	9	16

b Area scale factor = (length scale factor)^2
c **i** 100 **ii** 400 **iii** 10 000

8H

Now you try

Example 17


a AAA **b** 10 m

Exercise 8H

1 **a** Two **b** AAA **c** 2 **d** 12
2 $\angle C$
3 **a** $\angle ACB$ and $\angle ECD$
b $\angle BAC = \angle DEC$ and $\angle CBA = \angle CDE$
4 **a** 3 **b** 3.6 m
5 **a** AAA **b** 40 m
6 6 m
7 **a** AAA **b** 7.5 m
8 **a** To create two similar triangles
b AAA **c** 24 m
9 7.2 m
10 20 m
11 **a** **i** 3.6 m **ii** 9 m **iii** 2.7 m
b **i** 5.4 m **ii** 6.3 m
c **i** 4 m **ii** 6 m
12 $\frac{55}{6}$
13 **a** 3
b **i** $BD = \frac{25}{3}$ **ii** $AC = 4$ **iii** $AB = \frac{20}{3}$


Maths@Work: Animator

2 **a**



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


b



3 12 pairs of congruent triangles; 2 pairs of similar triangles
4 30° clockwise; 60° clockwise; 20° clockwise
5 Answers will vary.

Puzzles and games


1



2 110
3 10
4 11
5 Base of altitudes (an altitude is a line through a vertex that meets the opposite side at right angles)
6 **b** **i** $\frac{l}{2}$ **ii** $\frac{l}{4}$ **iii** $\frac{l}{128}$
c **i** $\frac{3}{4}$ **ii** $\frac{9}{16}$ **iii** $\frac{243}{1024}$
d Zero

Chapter checklist

1 Complement $= 90° - 25° = 65°$ Supplement $= 180° - 25° = 155°$
2 Right angled isosceles triangle $a = 45, b = 135$
3 $a = 113$, supplementary angles
$b = 113$, corresponding angles in 11 lines
$c = 11$, vertically opposite angles
$d = 67$, corresponding angles in 11 lines
4 The co-interior angles are not supplementary (they do not add to 180 degrees) so the lines are not parallel.
5 Rhombus, square and kite have diagonals that intersect at 90 degrees.
6 $a = 50$
7 $a = 30$
8 $S = 900, a = 285$
9 135° $(6 \times 180° \div 8)$
10 $\Delta ABC \equiv \Delta MON$ (SAS)
11 $a = 9, x = 20$
12 $SF = 1.5, x = 6, y = \frac{4}{3}$
13



14 2 sides in proportion and the included angle equal (SAS similarity)
15 $x = 8.75$
16 **a** All angles equal **b** 16 m

Short-answer questions

1 **a** $a = 142$ **b** $b = 40$ **c** $c = 120$
2 **a** Acute-angled isosceles, $x = 50, y = 80$
b Right-angled, $x = 25$
c Obtuse-angled scalene, $x = 30, y = 110$
3 **a** No – corresponding angles are not equal.
b Yes – cointerior angles are supplementary.
c Yes – alternate angles are equal.
4 **a** $a = 60, b = 120, c = 120$ **b** $a = 40, b = 80$
c $a = 99, b = 20$
5 **a** Square, rectangle **b** Square, rhombus, kite
c Parallelogram, rectangle, kite
6 **a** $a = 70, b = 110$ **b** $x = 115$
c $x = 15$
7 **a** $x = 95$ **b** $a = 120$ **c** $x = 135$
8 **a** SSS, $x = 60$ **b** Not congruent
c RHS, $x = 12$ **d** AAS, $x = 9$
9 For image $\Delta A'B'C'$, $OA' = 3OA, OB' = 3OB, OC' = 3OC$
10 **a** Yes, SAS **b** Yes, AAA
c Not similar **d** Yes, RHS
11 **a** 3.5 **b** 4 **c** 18
12 **a** 3 **b** 4.8 m

Multiple-choice questions

1 D **2** A **3** B **4** D **5** C **6** D
7 E **8** B **9** D **10** A

Extended-response questions

1 **a** Isosceles, right-angled
b $a = 40$, angle sum in isosceles triangle
$b = 70$, angles in parallel lines
$c = 120$, angle sum of a quadrilateral
$d = 25$, supplementary angles
$e = 65$, angle sum of a triangle
2 **a** $\angle ECD$
b $\angle ABC = \angle EDC$ (given 90°)
$\angle ACB = \angle ECD$ (vertically opposite)
$\therefore \Delta ABC \,|||\, \Delta EDC$ (AAA)
c 19.8 m

Chapter 9

Warm-up quiz

1 **a** 4 **b** 6 **c** −2 **d** −7
2 **a** 5 **b** 2 **c** 1 **d** −6
3 **a** −10 **b** 16 **c** 11 **d** −100
e 2 **f** 48 **g** −20 **h** −70
4 **a** $2x + 6$ **b** $3a - 15$ **c** $12x - 8xy$
d $3 - 6b$
5 **a** 2 **b** 6 **c** $2x$ **d** $3y$
e $5x$ **f** ab
6 **a** $2(a + 3)$ **b** $3(x + 4y)$
c $5x(x - 3)$ **d** $2m(2 - 3n)$
7 **a** $\frac{5}{7}$ **b** $-\frac{1}{9}$ **c** $\frac{7}{6}$ **d** $-\frac{1}{8}$
8 **a** $\frac{8}{15}$ **b** $\frac{1}{2}$ **c** $\frac{1}{3}$ **d** 1
e 2 **f** $\frac{3}{2}$ **g** $\frac{9}{8}$ **h** $\frac{7}{12}$
9 **a** $3x + 2$ **b** $x + 4$ **c** $-10 - 3x$
10 $x(x + 2), x^2 + 2x$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

9A

Now you try

Example 1

a $2m - n$ **b** $-2x^2y - xy$

Example 2

a $-15xy^2$ **b** $\frac{a}{2}$

Example 3

a $-3x - 6$ **b** $6x^2 - 4x$

Exercise 9A

1 **a** **i** $\$11x$ **ii** $50n$ cents
b **i** $x + y$ **ii** $b + g$
c $\$(50 + 60n)$

2 **a** 5 **b** -2 **c** -7 **d** No
e -10 **f** 6

3 **a** $5 + 2x$ **b** $4a - 7$ **c** $y^2 - 1$
d xz **e** $a^2 + b^2$ **f** $\sqrt{x + 8}$

4 **a** $-a$ **b** $11x$ **c** $8ab$ **d** $-4xy$
e $a - 1$ **f** $2x - 4y$ **g** $6x - 4y$ **h** $2ab$
i $-4a^2b - 3$ **j** $-3xy$ **k** $12ar^2$ **l** $st^2 - st$

5 **a** $15x$ **b** $14a$ **c** $-8xy$ **d** $-42a^2$
e $-3ab^2$ **f** $-12a^2b^2$ **g** $2x$ **h** 7
i 3 **j** $2a$ **k** $\frac{10}{a}$ **l** $4x$
m ab **n** $\frac{3a}{c}$ **o** $\frac{1}{3y}$

6 **a** $3x + 6$ **b** $8 + 2x$ **c** $-3x - 12$
d $-6x - 6$ **e** $-2x + 6$ **f** $-x^2 - x$
g $-6y + 9y^2$ **h** $-7a^2 + 7ab$ **i** $2x + 1$
j $4 - x$ **k** $3 - 4x$ **l** $2x + 6$
m $4x - 9$ **n** $-17x - 3$ **o** $13x - 40$ **p** $3x - 11$

7 **a** **i** $2a + 2b$ or $2(a + b)$ **ii** ab
b **i** $4x$ **ii** x^2
c **i** $a + b + c$ **ii** $\frac{1}{2}ab$

8 **a** 4 **b** 24 **c** -1 **d** 10
e 13 **f** -8 **g** $\frac{25}{2}$ **h** 1
i -2 **j** 35 **k** -12 **l** $\sqrt{13}$

9 **a** **i** \$260 **ii** \$820
b $100 + 80n$

10 **a** $2x^2 - x$ **b** $x^2 + 6x$ **c** $x^2 + x + 2$

11 **a** $-3x + 3$ **b** $\frac{a}{2}$ **c** $\frac{2}{a}$ **d** 9

12 **a** **i** \$120 **ii** \$600
b **i** \$140 **ii** \$540
c $\$60n$.
d $\$(40 + 50n)$
e 4 hours

9B

Now you try

Example 4

$x^2 + 11x + 18$

Example 5

a $x^2 - 3x - 18$ **b** $3x^2 - 14x + 8$ **c** $18x^2 - 9x - 20$

Exercise 9B

1 **a** $x^2, 2x, 3x, 6$ **b** $x^2 + 3x + 2x + 6 = x^2 + 5x + 6$

2 **a** $2x^2, 2x, 3x, 3$
b $(2x + 3)(x + 1) = 2x^2 + 2x + 3x + 3$
$= 2x^2 + 5x + 3$

3 **a** $x^2 + 5x + x + 5 = x^2 + 6x + 5$
b $x^2 + 2x - 3x - 6 = x^2 - x - 6$
c $21x^2 + 6x - 14x - 4 = 21x^2 - 8x - 4$
d $12x^2 - 16x - 3x + 4 = 12x^2 - 19x + 4$

4 **a** $x^2 + 7x + 10$ **b** $b^2 + 7b + 12$
c $t^2 + 15t + 56$ **d** $p^2 + 12p + 36$
e $x^2 + 15x + 54$ **f** $d^2 + 19d + 60$
g $a^2 + 8a + 7$ **h** $y^2 + 12y + 20$
i $m^2 + 16m + 48$

5 **a** $x^2 - x - 12$ **b** $x^2 + 3x - 10$
c $x^2 - 4x - 32$ **d** $x^2 - 4x - 12$
e $x^2 + 9x - 10$ **f** $x^2 + 2x - 63$
g $x^2 + 5x - 14$ **h** $x^2 - 3x + 2$
i $x^2 - 9x + 20$ **j** $8x^2 + 26x + 15$
k $6x^2 + 7x + 2$ **l** $15x^2 + 17x + 4$
m $6x^2 + x - 15$ **n** $24x^2 + 23x - 12$
o $6x^2 - x - 2$ **p** $10x^2 - 31x - 14$
q $6x^2 + 5x - 6$ **r** $16x^2 - 16x - 5$
s $18x^2 - 27x + 10$ **t** $15x^2 - 11x + 2$
u $21x^2 - 37x + 12$

6 **a** **i** $x + 10$ **ii** $x + 7$
b $x^2 + 17x + 70$
c 108 m^2

7 **a** $x^2 + 9x + 20$
b **i** 56 m^2 **ii** 36 m^2

8 **a** **i** $2x + 3$ **ii** $2x + 2$
b $4x^2 + 10x + 6$
c 20 m^2

9 **a** 3 **b** 2 **c** 6, 6 **d** 2, 18
e 2, 6 **f** 3, 15

10 **a** $a^2 + ac + ab + bc$ **b** $a^2 + ac - ab - bc$
c $ab + bc - a^2 - ac$ **d** $2x^2 - 3xy - 2y^2$
e $2a^2 - ab - b^2$ **f** $6x^2 + xy - y^2$

11 **a** $2x^2$ **b** $2x^2 - 30x + 100$

12 **a** **i** $15 - 2x$ **ii** $10 - 2x$
b $150 - 50x + 4x^2$
c 66 m^2
d $x = 2.5$

9C

Now you try

Example 6

a $x^2 + 16x + 64$ **b** $x^2 - 10x + 25$

Example 7

$25x^2 - 30x + 9$

Exercise 9C

1 **a** $x^2, 3x, 3x, 9$ **b** $x^2 + 6x + 9$
c $(x + 3)(x + 3) = x^2 + 3x + 3x + 9$
$= x^2 + 6x + 9$

2 **a** $+4x + 16 = x^2 + 8x + 16$ **b** $+5x + 25 = x^2 + 10x + 25$
c $-2x + 4 = x^2 - 4x + 4$ **d** $-7x + 49 = x^2 - 14x + 49$

3 **a** **i** $x^2 + 6x + 9$ **ii** $x^2 + 22x + 121$
b **i** $x^2 - 4x + 4$ **ii** $x^2 - 18x + 81$

4 **a** $x^2 + 2x + 1$ **b** $x^2 + 6x + 9$
c $x^2 + 4x + 4$ **d** $x^2 + 10x + 25$
e $x^2 + 8x + 16$ **f** $x^2 + 18x + 81$
g $x^2 + 14x + 49$ **h** $x^2 + 20x + 100$
i $x^2 - 4x + 4$ **j** $x^2 - 12x + 36$
k $x^2 - 2x + 1$ **l** $x^2 - 6x + 9$
m $x^2 - 18x + 81$ **n** $x^2 - 14x + 49$
o $x^2 - 8x + 16$ **p** $x^2 - 24x + 144$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

5 **a** $4x^2 + 4x + 1$ **b** $4x^2 + 20x + 25$
c $9x^2 + 12x + 4$ **d** $9x^2 + 6x + 1$
e $25x^2 + 20x + 4$ **f** $16x^2 + 24x + 9$
g $49 + 28x + 4x^2$ **h** $25 + 30x + 9x^2$
i $4x^2 - 12x + 9$ **j** $9x^2 - 6x + 1$
k $16x^2 - 40x + 25$ **l** $4x^2 - 36x + 81$

6 **a** $9 - 6x + x^2$ **b** $25 - 10x + x^2$
c $1 - 2x + x^2$ **d** $36 - 12x + x^2$
e $121 - 22x + x^2$ **f** $16 - 8x + x^2$
g $49 - 14x + x^2$ **h** $144 - 24x + x^2$
i $64 - 32x + 4x^2$ **j** $4 - 12x + 9x^2$
k $81 - 36x + 4x^2$ **l** $100 - 80x + 16x^2$

7 **a** $x + 12$
b $x^2 + 24x + 144$
c **i** 196 m^2 **ii** 289 m^2

8 **a** $10 - x$
b $x^2 - 20x + 100$
c **i** 64 cm^2 **ii** 16 cm^2
d $x = 5$

9 **a** $20 - 2x$
b $(20 - 2x)(20 - 2x) = 400 - 80x + 4x^2$
c 196 cm^2
d 588 cm^3

10 **a** $a^2 - b^2$
b **i** $(a - b)^2 = a^2 - 2ab + b^2$
ii $b(a - b) = ab - b^2$ **iii** $b(a - b) = ab - b^2$
c Yes, $a^2 - 2ab + b^2 + ab - b^2 + ab - b^2 = a^2 - b^2$

11 **a** $a + b$
b $(a + b)(a + b) = a^2 + 2ab + b^2$
c $a - b$
d $(a - b)(a - b) = a^2 - 2ab + b^2$
e $4ab$
f ab so, yes, four courts' area is $4ab$

9D

Now you try

Example 8

a $x^2 - 25$ **b** $x^2 - 225$

Example 9

$144x^2 - 25y^2$

Exercise 9D

1 Because $-ax + ax = 0$

2 **a** Single **b** Difference **c** Difference
d Single

3 **a** $+4x - 16$, $x^2 - 16$
b $+2x - 2x - 1$, $4x^2 - 1$

4 **a** $x^2 - 1$ **b** $x^2 - 9$ **c** $x^2 - 64$
d $x^2 - 16$ **e** $x^2 - 144$ **f** $x^2 - 121$
g $x^2 - 81$ **h** $x^2 - 25$ **i** $x^2 - 36$
j $25 - x^2$ **k** $4 - x^2$ **l** $49 - x^2$

5 **a** $9x^2 - 4$ **b** $25x^2 - 16$
c $16x^2 - 9$ **d** $49x^2 - 9y^2$
e $81x^2 - 25y^2$ **f** $121x^2 - y^2$
g $64x^2 - 4y^2$ **h** $100x^2 - 81y^2$
i $49x^2 - 25y^2$ **j** $36x^2 - 121y^2$
k $64x^2 - 9y^2$ **l** $81x^2 - 16y^2$

6 **a** 899 **b** 1599 **c** 624 **d** 2499
e 396 **f** 391 **g** 875 **h** 2484

7 **a** **i** x^2 **ii** $x^2 - 4$
b No, they differ by 4.

8 **a** $x^2 - 1$
b No, area of rectangle is 1 square unit less.

9 **a** **i** x^2 **ii** $x + 3$ **iii** $x - 3$
b $x^2 - 9$
c No, 9 m^2 less

9E

Now you try

Example 10

a $5b$ **b** $7xy$

Example 11

a $5(3a + 4)$ **b** $2x(x - 3)$

Example 12

$-11a(1 + 2a)$

Exercise 9E

1 **a** 4 **b** 10 **c** 5 **d** 6

2 **a** x **b** x **c** a **d** $2a$
e $-2y$ **f** $-3x$

3 **a** **i** 6 **ii** $3x$ **iii** $6x$
b iii

4 **a** $2x$ **b** $6a$ **c** 2 **d** 4
e 3 **f** 1 **g** $3x$ **h** $3n$
i $2y$ **j** $2x$ **k** $2xy$ **l** $5ab$

5 **a** $7(x + 1)$ **b** $3(x + 1)$ **c** $4(x - 1)$
d $5(x - 1)$ **e** $4(1 + 2y)$ **f** $5(2 + a)$
g $3(1 - 3b)$ **h** $2(3 - x)$ **i** $3(4a + b)$
j $6(m + n)$ **k** $2(5x - 4y)$ **l** $4(a - 5b)$
m $x(x + 2)$ **n** $a(a - 4)$ **o** $y(y - 7)$
p $x(1 - x)$ **q** $3p(p + 1)$ **r** $8x(1 - x)$
s $4b(b + 3)$ **t** $2y(3 - 5y)$

6 **a** $-4(2x + 1)$ **b** $-2(2x + 1)$
c $-5(2x + y)$ **d** $-7(a + 2b)$
e $-3(3x + 4)$ **f** $-2(3y + 4)$
g $-5(2x + 3y)$ **h** $-4(m + 5n)$
i $-3x(x + 6)$ **j** $-4x(2x + 3)$
k $-2y(8y + 3)$ **l** $-5a(a + 2)$

7 **a** 3 **b** y **c** a
d $5x$ **e** -7 **f** $-12a$

8 **a** $4(x + 2)$ **b** $2(x + 3)$
c $10(x + 2)$ **d** $2(x + 7)$
e $2(2x + 3)$ **f** $2(x + 7)$

9 $4x$

10 **a** $t(5 - t)$
b **i** 0 m **ii** 6 m **iii** 4 m
c 5 s

11 **a** 63 **b** 72 **c** -20
d -70 **e** 69 **f** 189

12 **a** $3(a^2 + 3a + 4)$ **b** $z(5z - 10 + y)$
c $x(x - 2y + xy)$ **d** $2b(2y - 1 + 3b)$
e $-4y(3x + 2z + 5xz)$ **f** $ab(3 + 4b + 6a)$

13 **a** $(x + 3)(4 + x)$ **b** $(x + 1)(3 + x)$
c $(m - 3)(7 + m)$ **d** $(x - 7)(x + 2)$
e $(a + 4)(8 - a)$ **f** $(x + 1)(5 - x)$
g $(y + 3)(y - 2)$ **h** $(x + 2)(a - x)$
i $(2t + 5)(t + 3)$ **j** $(5m - 2)(m + 4)$
k $(4y - 1)(y - 1)$ **l** $(7 - 3x)(1 + x)$

Progress quiz

1 **a** $8x - 6y$ **b** $7ab^2 - ab$ **c** $20a^2b$ **d** $\frac{y}{3}$

2 **a** $3x - 12$ **b** $6y^2 + 10y$ **c** $3 - 6x$

3 **a** $x^2 + 10x + 24$ **b** $x^2 + x - 6$
c $2x^2 + 5x - 12$ **d** $6x^2 - 19x + 10$

4 **a** $x^2 + 12x + 36$ **b** $x^2 - 16x + 64$
c $9x^2 + 42x + 49$ **d** $25 - 10x + x^2$

5 **a** $(2x^2 - 5x - 3)$ units^2 **b** $(9x^2 - 6x + 1)$ m^2

6 **a** $x^2 - 100$ **b** $x^2 - 1$ **c** $9x^2 - 4$ **d** $16x^2 - 25y^2$

7 $2x + 2$

Essential Mathematics Core for the Victorian Curriculum Year 9 ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

8 **a** 4 **b** $3a$
c $5ab$ **d** $3x$
9 **a** $6(x+2)$ **b** $5(3y-4)$
c $4a(2b+5)$ **d** $2x(2x-1)$
e $-5(2x+1)$ **f** $-2x(2x+5)$
10 **a** $2(4x+3)$ **b** $6(x+2)$

9F

Now you try

Example 13

a $(x+4)(x-4)$ **b** $(7+y)(7-y)$

Example 14

a $(2a+3)(2a-3)$ **b** $(5a+b)(5a-b)$

Example 15

$3(a+3)(a-3)$

Exercise 9F

1 **a** x^2-4 **b** x^2-49 **c** $9x^2-y^2$
2 **a** 3 **b** 11 **c** $5b$ **d** $7y$
3 **a** $(x+4)(x-4)$
b $x^2-(12)^2=(x+12)(x-12)$
c $(4x)^2-(1)^2=(4x+1)(4x-1)$
d $(3a)^2-(2b)^2=(3a+2b)(3a-2b)$
4 **a** $(x+3)(x-3)$ **b** $(y+5)(y-5)$
c $(y+1)(y-1)$ **d** $(x+8)(x-8)$
e $(x+4)(x-4)$ **f** $(b+7)(b-7)$
g $(a+9)(a-9)$ **h** $(x+y)(x-y)$
i $(a+b)(a-b)$ **j** $(4+a)(4-a)$
k $(5+x)(5-x)$ **l** $(1+b)(1-b)$
m $(6+y)(6-y)$ **n** $(11+b)(11-b)$
o $(x+20)(x-20)$ **p** $(30+y)(30-y)$
5 **a** $(2x+5)(2x-5)$ **b** $(3x+7)(3x-7)$
c $(5b+2)(5b-2)$ **d** $(2m+11)(2m-11)$
e $(10y+3)(10y-3)$ **f** $(9a+2)(9a-2)$
g $(1+2x)(1-2x)$ **h** $(5+8b)(5-8b)$
i $(4+3y)(4-3y)$ **j** $(6x+y)(6x-y)$
k $(2x+5y)(2x-5y)$ **l** $(8a+7b)(8a-7b)$
m $(2p+5q)(2p-5q)$ **n** $(9m+2n)(9m-2n)$
o $(5a+7b)(5a-7b)$ **p** $(10a+3b)(10a-3b)$
6 **a** $3(x+6)(x-6)$ **b** $10(a+1)(a-1)$
c $6(x+2)(x-2)$ **d** $4(y+4)(y-4)$
e $2(7+x)(7-x)$ **f** $8(2+m)(2-m)$
g $5(xy+1)(xy-1)$ **h** $3(1+xy)(1-xy)$
i $7(3+ab)(3-ab)$
7 **a** $4(3+t)(3-t)$
b **i** 36 m **ii** 20 m
c 3 seconds
8 **a** $(x+3)(x-3)$ **b** $(4x+11)(4x-11)$
c $(2+5a)(2-5a)$ **d** $(x+y)(x-y)$
9 **a** Factorise each binomial:
$(4x+2)(4x-2)=2(2x+1)2(2x-1)$
$=4(2x+1)(2x-1)$
b Take out common factor of 4.
10 **a** $(x+8)(x+2)$ **b** $(x+5)(x+1)$
c $(x+14)(x+6)$ **d** $(x+2)(x-8)$
e $(x-6)(x-8)$ **f** $(x+3)(x-9)$
11 **a** **i** x^2 **ii** $(30+x)(30-x)$
b **i** 500 cm^2 **ii** 675 cm^2

9G

Now you try

Example 16

$(x+4)(x+2)$

Example 17

$(x+2)(x-3)$

Example 18

$(x-5)(2x+3)$

Exercise 9G

1 **a** $2x-2$ **b** $-5+5a$
c a^2+5a **d** $4y-y^2$
e $ax+x+2a+2$ **f** $bx-2b-3x+6$
2 **a** $2+x$ **b** $5-x$ **c** $a+1$ **d** $1-a$
3 **a** $(x-3)(x-2)$ **b** $(x+4)(x+3)$
c $(2x+1)(3-x)$ **d** $(3x-2)(4-x)$
4 **a** $(x+3)(x+2)$ **b** $(x+4)(x+3)$
c $(x+7)(x+2)$ **d** $(x-6)(x+4)$
e $(x-4)(x+6)$ **f** $(x-3)(x+10)$
5 **a** $(x+4)(x-3)$ **b** $(x+2)(x-6)$
c $(x+2)(x-18)$ **d** $(x+3)(x-14)$
e $(x+4)(x-18)$ **f** $(x-2)(x-a)$
6 **a** $(x-b)(x+1)$ **b** $(x-c)(x+1)$
c $(x+b)(x+1)$ **d** $(x+c)(x-1)$
e $(x+a)(x-1)$ **f** $(x-b)(x-1)$
7 **a** $(x-7)(2x+1)$ **b** $(x+5)(x+2)$
c $(x+3)(2x-1)$ **d** $(1-2x)(3x+4)$
e $(x-5)(11+a)$ **f** $(3-2x)(4y-1)$
g $(n+2)(3m-1)$ **h** $(3-r)(5p+8)$
8 **a** $(x+2)(x+5)$ **b** $(x+3)(x+5)$
c $(x+4)(x+6)$ **d** $(x-3)(x+2)$
e $(x+6)(x-2)$ **f** $(x-9)(x-2)$
9 **a** $(3a+5c)(b+d)$ **b** $(4b-7c)(a+d)$
c $(y-4z)(2x+3w)$ **d** $(s-2)(5r+t)$
e $(x+3y)(4x-3)$ **f** $(2b-a)(a-c)$
10 Answers may vary.
11 **a** Method 1: $a(x+7)-3(x+7)=(x+7)(a-3)$
Method 2: $x(a-3)+7(a-3)=(a-3)(x+7)$
b **i** $(x-3)(b+2)$ **ii** $(x+2)(y-4)$
iii $(2m+3)(2m-5n)$ **iv** $(2-n)(m-3)$
v $(1-2b)(4a+3b)$ **vi** $(3a-1)(b+4c)$

9H

Now you try

Example 19

$(x+2)(x+7)$

Example 20

$(x+6)(x-2)$

Example 21

$(x-3)(x-4)$

Example 22

$3(x+3)(x-5)$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Exercise 9H

1 **a** $x^2+9x+14$ **b** $x^2+14x+33$
c $x^2+7x-60$ **d** $x^2-8x+12$
2 **a** 3, 2 **b** 12, 1 **c** 4, 5
d 5, −3 **e** −6, 5 **f** −9, −2
3 **a** $(x+6)(x+2)$ **b** $(x+9)(x+1)$
c $(x+7)(x+1)$ **d** $(x+14)(x+1)$
e $(x+3)(x+4)$ **f** $(x+8)(x+2)$
g $(x+5)(x+3)$ **h** $(x+4)(x+5)$
i $(x+8)(x+3)$
4 **a** $(x+4)(x-1)$ **b** $(x+2)(x-1)$
c $(x+5)(x-1)$ **d** $(x+7)(x-2)$
e $(x+5)(x-3)$ **f** $(x+10)(x-2)$
g $(x-6)(x+3)$ **h** $(x-9)(x+2)$
i $(x-4)(x+3)$
5 **a** $(x-5)(x-1)$ **b** $(x-1)(x-1)$
c $(x-1)(x-4)$ **d** $(x-8)(x-1)$
e $(x-2)(x-2)$ **f** $(x-6)(x-2)$
g $(x-9)(x-2)$ **h** $(x-7)(x-3)$
i $(x-3)(x-2)$
6 **a** $(x-8)(x+1)$ **b** $(x-4)(x+1)$
c $(x-6)(x+1)$ **d** $(x-8)(x+2)$
e $(x-6)(x+4)$ **f** $(x-5)(x+3)$
g $(x-4)(x+3)$ **h** $(x-12)(x+1)$
i $(x-6)(x+2)$
7 **a** $2(x+4)(x+1)$ **b** $2(x+10)(x+1)$
c $3(x+2)(x+4)$ **d** $2(x+10)(x-3)$
e $2(x-9)(x+2)$ **f** $4(x-1)(x-1)$
g $2(x-2)(x+3)$ **h** $6(x-6)(x+1)$
i $5(x-4)(x-2)$ **j** $3(x-5)(x-6)$
k $2(x-5)(x+2)$ **l** $3(x-4)(x+3)$
8 **a** $6x$ **b** $6x$ or $10x$ **c** x, $4x$, $11x$
d x, $4x$, $11x$ **e** $9x$, $11x$, $19x$ **f** $9x$, $11x$, $19x$
g $0x$, $6x$, $15x$ **h** $0x$, $24x$
9 **a** $(x+4)^2$ **b** $(x+5)^2$
c $(x+15)^2$ **d** $(x-1)^2$
e $(x-7)^2$ **f** $(x-13)^2$
g $2(x+1)^2$ **h** $5(x-3)^2$
i $-3(x-6)^2$
10 **a** **i** $x(x+2)$ **ii** $x^2+2x-15$
iii $(x+5)(x-3)$
b **i** 105 m^2 **ii** 48 m^2

9I

Now try

Example 23

a $\frac{-ab}{2}$ **b** $-(x+7)$ or $-x-7$

Example 24

a $a-6$ **b** -7

Example 25

a $6a$ **b** 2

Example 26

a $\frac{-3x}{2}$ **b** $\frac{2}{15}$

Exercise 9I

1 **a** 2 **b** $2x$ **c** $2(x+2)$ **d** $2(x-1)$
2 **a** $\frac{3}{2}$ **b** $\frac{3}{4}$ **c** $\frac{2}{7x}$
d $\frac{4a^2}{5}$ **e** $\frac{1}{7}$ **f** $\frac{1}{6x}$
3 **a** $\frac{8}{9}$ **b** $\frac{7}{6}$ **c** $\frac{3}{2}$
d $\frac{1}{4}$ **e** 6 **f** 1
4 **a** $\frac{x}{2}$ **b** $\frac{x}{2}$ **c** $\frac{-ab}{3}$ **d** $\frac{a}{2b}$
e $x-1$ **f** $-2(2x+1)$ **g** $\frac{-1}{3}$ **h** $\frac{a}{10}$
i 3 **j** -2 **k** $\frac{1}{2}$ **l** $\frac{-2}{3}$
m $x-4$ **n** $2(x-1)$ **o** $-2(x+2)$ **p** -1
5 **a** $x+2$ **b** $x-3$ **c** $x-4$ **d** $\frac{1}{x+1}$
e $\frac{-1}{2x+3}$ **f** $\frac{a}{3a-2}$ **g** 2 **h** 7
i -5 **j** -2 **k** 6 **l** -5
6 **a** 1 **b** $\frac{1}{2}$ **c** $\frac{6a}{5}$ **d** $\frac{3}{2a}$
e $\frac{7xy}{3}$ **f** $\frac{5x}{6}$ **g** 4 **h** 5
i $\frac{9}{2}$ **j** $\frac{x+1}{4}$ **k** 1 **l** $\frac{x+6}{2(x+7)}$
7 **a** $\frac{7x}{6}$ **b** $\frac{2x}{3}$ **c** $\frac{-21}{8}$ **d** $\frac{-a^2}{2}$
e 6 **f** $\frac{4}{3}$ **g** 11 **h** $\frac{-3}{2}$
8 **a** $x+1$ **b** $-(x+2)$ **c** $\frac{3}{2}$ **d** $x(x+2)$
9 **a** 4 **b** 5 **c** -18 **d** 15
e -34 **f** 14
10 **a** $x+3$ **b** $2x-1$ **c** $-5(x-3)$ **d** $\frac{-1}{3(x-4)}$
11 **a** -1 **b** -1 **c** $\frac{-4}{3}$ **d** 2
e $\frac{-1}{x-2}$ **f** $-4(x-6)$

9J

Now try

Example 27

a $\frac{7a}{10}$ **b** $\frac{5x}{24}$

Example 28

$\frac{7x-17}{12}$

Exercise 9J

1 **a** 24 **b** 15 **c** 143 **d** 36
2 **a** $\frac{5}{6}$ **b** $\frac{17}{20}$ **c** $\frac{13}{8}$ **d** $\frac{7}{10}$
e $-\frac{2}{15}$ **f** $\frac{1}{12}$
3 **a** $4x$ **b** $21x$ **c** 3
d 2 **e** 8 **f** 90
4 **a** $\frac{3x}{12}+\frac{8x}{12}=\frac{11x}{12}$
b $\frac{25x}{35}-\frac{14x}{35}=\frac{11x}{35}$
5 **a** 15 **b** 14 **c** 8
d 6 **e** 10

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

6 **a** $\frac{9x}{14}$ **b** $\frac{2x}{5}$ **c** $\frac{x}{8}$ **d** $\frac{14x}{45}$
e $\frac{y}{56}$ **f** $\frac{13a}{22}$ **g** $\frac{2b}{9}$ **h** $\frac{m}{6}$
i $\frac{11m}{12}$ **j** $\frac{15a}{28}$ **k** $\frac{x}{2}$ **l** $-\frac{20p}{63}$
m $-\frac{5b}{18}$ **n** $\frac{61y}{40}$ **o** $\frac{13x}{35}$

7 **a** $\frac{7x+11}{10}$ **b** $\frac{7x}{12}$ **c** $\frac{15a-51}{56}$
d $\frac{11y+9}{30}$ **e** $\frac{13m+28}{40}$ **f** $\frac{5x-13}{24}$
g $\frac{11b-6}{24}$ **h** $\frac{7x}{6}$ **i** $\frac{7y-8}{14}$
j $\frac{5t-4}{16}$ **k** $\frac{34-10x}{21}$ **l** $\frac{8m-9}{12}$

8 **a** $\frac{4x}{8}+\frac{6x}{8}=\frac{10x}{8}=\frac{5x}{4}$
b $\frac{2x}{4}+\frac{3x}{4}=\frac{5x}{4}$
c Using denominator 8 does not give answer in simplified form and requires extra steps. Preferable to use actual LCD.

9 **a** Didn't make a common denominator and subtracted denominators, $\frac{7x}{15}$
b Didn't use brackets: $2(x+1)=2x+2, \frac{7x+2}{10}$

10 **a** $\frac{11}{2x}$ **b** $\frac{1}{3x}$ **c** $-\frac{3}{4x}$ **d** $\frac{14}{9x}$

11 **a** $-\frac{43x}{30}$ **b** $\frac{5x}{12}$ **c** $\frac{13x}{24}$
d $\frac{43x-5}{60}$ **e** $\frac{23x-35}{42}$ **f** $\frac{29x+28}{40}$

12 **a** $\frac{x}{3}$ **b** $\frac{x}{8}$ **c** $\frac{x}{2}$
d $\frac{x}{5}$ **e** $\frac{8x}{9}$ **f** $\frac{x}{4}$

Maths@Work: Automotive technology

1 **a** TSA $= 2lw+2wh+2hl$
$= 2(lw+wh+hl)$
b TSA $= ah+bh+ch+ab$
c TSA $= 2\pi r^2+2\pi rh$
TSA $= 2\pi r(r+h)$
d TSA $= \frac{\pi d^2}{2}+\pi dh$
TSA $= \pi d\left(\frac{d}{2}+h\right)$

2 **i** c **ii** a
iii b **iv** d

3 **a** Area of cross-section $= \pi R^2-\pi r^2$
$= \pi(R^2-r^2)$
$= \pi\left(\frac{D^2}{2^2}-\frac{d^2}{2^2}\right)$
$= \frac{\pi}{4}(D^2-d^2)$
$= \frac{\pi}{4}(D+d)(D-d)$
b **i** 76π mm^2 **ii** 31π mm^2
iii $\frac{\pi}{2}$ cm^2 **iv** 3π cm^2

4

Pipe material	Area of cross-section in cm^2	Length in cm	Weight in g	Weight in kg
Poly pipe a	3.77	100	360.03	0.36
Poly pipe b	11.50	100	1098.08	1.10
Poly pipe c	141.37	100	13500.99	13.50
Copper pipe a	0.35	100	308.94	0.31
Copper pipe b	0.31	100	273.84	0.27
Copper pipe c	1.54	100	1376.21	1.38
Steel pipe a	2.89	100	2268.86	2.27
Steel pipe b	8.64	100	6781.91	6.78
Steel pipe c	25.13	100	19729.20	19.73

a Copper pipe c = 1.38 kg/m; steel pipe b = 6.78 kg/m
Steel pipe b is 5.4 kg/m heavier than copper pipe c.
b 3.5 m of poly pipe a = 1.26 kg; 3.5 m of steel pipe a = 7.94 kg
3.5 m of steel pipe a is 6.68 kg heavier than 3.5 m of poly pipe a.
c 2.75 m of poly pipe c = 37.13 kg; 5 m of steel pipe c = 98.65 kg.
5 m of steel pipe c is 61.52 kg heavier than 2.75 m of poly pipe c.

Puzzles and games

1 RECIPROCAL
2 **a** $4x^2-80x+400$ **b** $x=5$
3 **a** n^2 **b** $n(n+1)$ **c** $\frac{1}{2}n(n+1)$
4 **a** $\frac{x+10}{12}$ **b** $\frac{-11x-2}{15}$
5 **a** 48, 49 **b** 33, 35 **c** 12, 15
6 Ryan

Chapter checklist

1 **a** $2a^2-ab+1$ **b** $-x^2y-xy$
2 35
3 **a** $14a^2b$ **b** $\frac{y}{3}$
4 **a** $-3x+6$ **b** $-8a^2+10a$
5 **a** $2x^2-x-15$ **b** $6x^2-29x+35$
6 **a** $x^2+14x+49$ **b** $16x^2-24x+9$
7 **a** x^2-16 **b** $9x^2-25$
8 HCF $= 6xy$
9 **a** $5(a+3)$ **b** $11x(x-3y)$ **c** $-7m(2n+1)$
10 **a** $(x+10)(x-10)$ **b** $(2a-3)(2a+3)$
11 $(x+6)(x-3)$
12 $(x-3)(2x+3)$
13 $(x+7)(x-3)$
14 $2(x+3)(x-3)$ and $3(x-4)(x-3)$
15 $\frac{2(x-2)}{x-2}=2$
16 $\frac{3}{2}$
17 $\frac{1}{4}$
18 **a** $\frac{x}{10}$ **b** $\frac{11x-29}{12}$

Short-answer questions

1 **a** $6x^2$ **b** $-18ab^2$ **c** $\frac{b}{2}$ **d** $-4x$
2 **a** $3x-12$ **b** $-2x-12$ **c** $-x^2+x$ **d** $-8x^2+12x$
3 **a** $7x-1$ **b** $5ab$ **c** $-3a^2b$ **d** $8x^2-xy$
4 **a** x^2+x-12 **b** $x^2-9x+14$
c $6x^2-5x-6$ **d** $3x^2+x-4$
5 **a** x^2+6x+9 **b** $x^2-8x+16$
c $9x^2-12x+4$ **d** x^2-25
e $49-x^2$ **f** $121x^2-16$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

6 **a** $4(a+3b)$ **b** $x(x+1)$
c $3x(2-3x)$ **d** $-3(1+3x)$
7 **a** $(x+10)(x-10)$ **b** $(5x+y)(5x-y)$
c $(7+3x)(7-3x)$ **d** $3(x+4)(x-4)$
8 **a** $(x-3)(x+6)$ **b** $(2x+5)(2x-1)$
c $(x-4)(3+2b)$
9 **a** $(x+3)(x+5)$ **b** $(x-6)(x+3)$
c $(x-6)(x-1)$ **d** $3(x+7)(x-2)$
10 **a** $\frac{x-1}{2}$ **b** $x+1$ **c** $x+4$
d $\frac{-(x+3)}{2}$ **e** $\frac{2}{3}$ **f** $2(x-1)$
11 **a** $\frac{1}{4}$ **b** $\frac{a}{2}$ **c** $\frac{x-4}{2}$
d $\frac{2a}{3}$ **e** 4 **f** $\frac{2}{5}$
12 **a** $\frac{11x}{12}$ **b** $-\frac{x}{24}$ **c** $\frac{a}{4}$
d $\frac{5x-2}{6}$ **e** $\frac{10x-5}{12}$ **f** $\frac{7x+5}{24}$

Multiple-choice questions

1 E **2** C **3** B **4** B **5** C
6 A **7** E **8** D **9** C **10** B
11 A **12** E **13** E

Extended-response questions

1 **a** $(x+3)$ m
b **i** No change
ii (x^2-1) m^2, 1 square metre less in area
c **i** 3 m^2 **ii** $x=4$
2 **a** 400 m^2
b **i** $L=W=(20+2x)$m **ii** $(4x^2+80x+400)$ m^2
c $4x(x+20)$ m^2
d $x=5$

Chapter 10

Warm-up quiz

1 **a** 0.1 **b** 0.25 **c** 0.3 **d** 0.85 **e** 0.237
2 **a** $\frac{1}{2}$ **b** $\frac{1}{3}$ **c** $\frac{2}{3}$ **d** 1 **e** 0
f $\frac{1}{12}$ **g** $\frac{1}{2}$ **h** $\frac{18}{29}$ **i** $\frac{2}{3}$ **j** $\frac{2}{7}$
3 **a** 1, 2, 3, 4, 5 or 6
b **i** 3 **ii** 2 **iii** 3 **iv** 5 **v** 5 **vi** 3
4 **a** 5 **b** 7 **c** 6 **d** 7 **e** 4 **f** 6
5 **a** 8 **b** 20 **c** $\frac{3}{20}$
6 **a** 9
b 6
c 5
d 5
e 12
f **i** $\frac{2}{9}$ **ii** $\frac{7}{9}$ **iii** $\frac{5}{9}$

10A

Now you try

Example 1


a $\{1,2,3\}$ **b** $\frac{1}{3}$ **c** $\frac{1}{2}$ **d** $\frac{1}{2}$ **e** $\frac{5}{6}$

Example 2





a $\frac{1}{10}$ **b** $\frac{7}{10}$ **c** $\frac{2}{5}$
d $\frac{3}{5}$ **e** $\frac{9}{10}$ **f** $\frac{1}{2}$

Exercise 10A

1 **a** sample space **b** event **c** probability
d impossible, certain **e** even **f** complement
2 **a** **i** $\frac{1}{4}$ **ii** 0.25 **iii** 25%
b



3

	Percentage	Decimal	Fraction	Number line
a	50%	0.5	$\frac{1}{2}$	
b	25%	0.25	$\frac{1}{4}$	
c	75%	0.75	$\frac{3}{4}$	
d	20%	0.2	$\frac{1}{5}$	

4 0.15, $\frac{2}{9}$, 1 in 4, 0.28, $\frac{1}{3}$, $\frac{2}{5}$, $\frac{3}{7}$, 2 in 3, 0.7, 0.9
5 **a** **i** $\{1,2,3,4,5,6,7\}$ **ii** $\frac{1}{7}$ **iii** $\frac{6}{7}$ **iv** $\frac{2}{7}$ **v** $\frac{6}{7}$
b **i** $\{2,6,7\}$ **ii** $\frac{1}{2}$ **iii** $\frac{1}{2}$ **iv** $\frac{1}{2}$ **v** 1
c **i** $\{1,2,3\}$ **ii** $\frac{2}{3}$ **iii** $\frac{1}{3}$ **iv** $\frac{5}{6}$ **v** $\frac{5}{6}$
d **i** $\{1,2,3\}$ **ii** $\frac{1}{3}$ **iii** $\frac{2}{3}$ **iv** $\frac{2}{3}$ **v** $\frac{2}{3}$
e **i** $\{1,2,3,4\}$ **ii** $\frac{1}{7}$ **iii** $\frac{6}{7}$ **iv** $\frac{3}{7}$ **v** $\frac{5}{7}$
f **i** $\{2\}$ **ii** 1 **iii** 0 **iv** 1 **v** 1
6 **a** $\frac{1}{2}$ **b** $\frac{3}{8}$ **c** $\frac{1}{6}$ **d** $\frac{1}{4}$ **e** 1 **f** 0
7 **a** $\frac{1}{2}$ **b** $\frac{5}{8}$ **c** $\frac{5}{6}$ **d** $\frac{3}{4}$ **e** 0 **f** 1
8 **a** $\frac{1}{8}$ **b** $\frac{7}{8}$
9 **a** $\frac{1}{8}$ **b** $\frac{1}{4}$ **c** $\frac{3}{8}$ **d** $\frac{3}{8}$ **e** $\frac{5}{8}$
f 1 **g** 0 **h** $\frac{1}{4}$ **i** $\frac{3}{4}$ **j** $\frac{3}{4}$
10 **a** {Hayley, Alisa, Rocco, Stuart}
b **i** $\frac{1}{4}$ **ii** $\frac{1}{2}$ **iii** $\frac{1}{2}$
11 **a** $\frac{1}{52}$ **b** $\frac{1}{13}$ **c** $\frac{1}{26}$ **d** $\frac{1}{2}$
e $\frac{2}{13}$ **f** $\frac{12}{13}$ **g** $\frac{23}{26}$ **h** $\frac{25}{26}$
12 **a** $\frac{1}{6}$ **b** $\frac{1}{6}$ **c** $\frac{5}{6}$ **d** $\frac{1}{3}$ **e** $\frac{2}{3}$
f 1 **g** $\frac{1}{3}$ **h** $\frac{1}{2}$ **i** $\frac{5}{6}$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

13 a $\frac{2}{11}$ **b** $\frac{9}{11}$ **c** $\frac{4}{11}$ **d** $\frac{7}{11}$
e $\frac{7}{11}$ **f** $\frac{3}{11}$ **g** $\frac{7}{11}$ **h** $\frac{4}{11}$


14 a 31 min
b $\frac{3}{31}$
c i $\frac{4}{31}$ **ii** $\frac{4}{31}$ **iii** $\frac{20}{31}$
iv $\frac{5}{31}$ **v** $\frac{8}{31}$ **vi** $\frac{11}{31}$

10B

Now you try

Example 3

a



b i 43 **ii** 25 **iii** 18
c i $\frac{7}{25}$ **ii** $\frac{7}{50}$ **iii** $\frac{11}{50}$

Example 4

a

	Natural	Not Natural	Total
Artificial	20	25	45
Not artificial	45	10	55
Total	65	35	100

b i 55 **ii** 10
c i $\frac{13}{20}$ **ii** $\frac{9}{20}$ **iii** $\frac{9}{10}$

Exercise 10B


1 a 26
b i 10 **ii** 14 **iii** 5 **iv** 9
v 4 **vi** 7 **vii** 19
c i 12 **ii** 17
2 a 22 **b** 12 **c** 10
d 39 **e** 13 **f** 1
3 a B **b** D **c** A **d** C
4 a

	A	Not A	Total
B	7	8	15
Not B	3	1	4
Total	10	9	19

b

	A	Not A	Total
B	2	5	7
Not B	9	4	13
Total	11	9	20

5 a



b i 28 **ii** 21 **iii** 6
c i $\frac{1}{10}$ **ii** $\frac{1}{15}$ **iii** $\frac{19}{30}$

6 i Not A: diagram T **ii** A or B: diagram R
iii A and B: diagram U **iv** B: diagram P
v B only: diagram Q **vi** Neither A nor B: diagram S

7 a i $\frac{2}{5}$ **ii** $\frac{1}{3}$ **iii** $\frac{7}{15}$
iv $\frac{1}{15}$ **v** $\frac{13}{15}$ **vi** $\frac{2}{15}$
b i $\frac{3}{7}$ **ii** $\frac{12}{35}$ **iii** $\frac{13}{35}$
iv $\frac{3}{35}$ **v** $\frac{34}{35}$ **vi** $\frac{1}{35}$

8 a

	Cream	Not cream	Total
Ice cream	5	20	25
Not ice cream	16	9	25
Total	21	29	50

b i 29 **ii** 9
c i $\frac{21}{50}$ **ii** $\frac{8}{25}$ **iii** $\frac{41}{50}$

9 a

	A	Not A	Total
B	3	1	4
Not B	2	2	4
Total	5	3	8

i $\frac{5}{8}$ **ii** $\frac{3}{8}$ **iii** $\frac{3}{8}$ **iv** $\frac{3}{4}$ **v** $\frac{1}{8}$ **vi** $\frac{1}{4}$


b

	A	Not A	Total
B	11	4	15
Not B	6	5	11
Total	17	9	26

i $\frac{17}{26}$ **ii** $\frac{9}{26}$ **iii** $\frac{11}{26}$ **iv** $\frac{21}{26}$ **v** $\frac{2}{13}$ **vi** $\frac{5}{26}$


10 a

	A	Not A	Total
B	2	6	8
Not B	3	1	4
Total	5	7	12




b

	A	Not A	Total
B	3	4	7
Not B	9	4	13
Total	12	8	20



11 a



b 5 **c** $\frac{1}{17}$


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

12 a

	Tank water	No tank water	Total
Tap water	12	36	48
No tap water	11	41	52
Total	23	77	100

b 12 **c** $\frac{9}{25}$ **d** $\frac{59}{100}$


13 a



b 23 **c** $\frac{11}{23}$ **d** $\frac{4}{16} = \frac{1}{4}$ **e** $\frac{4}{11}$

14 a $\frac{3}{10}$ **b** $\frac{1}{3}$

15



a 32 **b** 45 **c** 70 **d** 38 **e** 13
f 13 **g** 62

10C

Now you try

Example 5

a

	1	2	3	4	5	6	7	8
H	H1	H2	H3	H4	H5	H6	H7	H8
T	T1	T2	T3	T4	T5	T6	T7	T8

b 16 **c** $\frac{1}{16}$ **d** $\frac{3}{16}$

Example 6

a

	3	5	6
3	×	(3, 5)	(3, 6)
5	(5, 3)	×	(5, 6)
6	(6, 3)	(6, 5)	×

b $\frac{1}{6}$ **c** $\frac{1}{3}$

Exercise 10C

1

	B	Y	E
H	HB	HY	HE
I	IB	IY	IE

2 a

	1	2	3	4
H	H1	H2	H3	H4
T	T1	T2	T3	T4

b 8 outcomes in the sample space **c** $\frac{1}{8}$ **d** $\frac{1}{8}$

3 a 10 **b** H2, H4, T2, T4 **c** $\frac{2}{5}$

d T1, T3, T5 **e** $\frac{3}{10}$

4 a $\frac{1}{4}$ **b** HH, TT **c** $\frac{1}{2}$

5 a

	1	2	3	4	5	6
H	H1	H2	H3	H4	H5	H6
T	T1	T2	T3	T4	T5	T6

b 12 **c** $\frac{1}{12}$ **d** $\frac{1}{4}$

6 a

	R	I	D	E
L	LR	LI	LD	LE
I	IR	II	ID	IE
N	NR	NI	ND	NE
E	ER	EI	ED	EE

b 16 **c** $\frac{1}{16}$ **d** $\frac{1}{16}$

e $\frac{1}{4}$ **f** $\frac{1}{4}$ **g** $\frac{1}{8}$

7 a

	R	P	B
R	RR	RP	RB
P	PR	PP	PB
G	GR	GP	GB
B	BR	BP	BB

b $\frac{1}{12}$ **c** $\frac{1}{12}$ **d** $\frac{1}{6}$ **e** $\frac{1}{4}$

8 a

Sum	1	2	3	4	5	6
1	2	3	4	5	6	7
2	3	4	5	6	7	8
3	4	5	6	7	8	9
4	5	6	7	8	9	10
5	6	7	8	9	10	11
6	7	8	9	10	11	12

36 outcomes **b** $\frac{1}{9}$

c $\frac{1}{2}$ **d** 7 **e** 2 and 12

9 a

	Y	W	B	B	B
W	WY	WW	WB	WB	WB
O	OY	OW	OB	OB	OB
O	OY	OW	OB	OB	OB

b $\frac{2}{15}$ **c** $\frac{1}{15}$ **d** $\frac{2}{15}$

e $\frac{1}{5}$ **f** $\frac{1}{15}$

10 a

	D	O	G
D	X	(O, D)	(G, D)
O	(D, O)	X	(G, O)
G	(D, G)	(O, G)	X

b $\frac{1}{6}$ **c** $\frac{2}{3}$

11 a

		1st			
		1	**2**	**3**	**4**
2nd	**1**	X	(2, 1)	(3, 1)	(4, 1)
	2	(1, 2)	X	(3, 2)	(4, 2)
	3	(1, 3)	(2, 3)	X	(4, 3)
	4	(1, 4)	(2, 4)	(3, 4)	X

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

b i $\frac{1}{12}$ **ii** $\frac{1}{12}$

c i $\frac{1}{6}$ **ii** $\frac{5}{6}$ **iii** $\frac{1}{6}$ **iv** $\frac{1}{2}$


12 a $\frac{1}{8}$ **b** $\frac{1}{3}$

10D

Now you try

Example 7

a




b 16

c i $\frac{1}{16}$ **ii** $\frac{3}{8}$ **iii** $\frac{15}{16}$

Example 8

a




b i $\frac{1}{3}$ **ii** $\frac{2}{3}$ **iii** 1

Exercise 10D

1 a A **b** B


2 a i replacement **ii** 3

b




3 a i Without replacement **ii** 2

b




4 a



b 4

c i $\frac{1}{4}$ **ii** $\frac{1}{2}$ **iii** $\frac{3}{4}$ **iv** $\frac{3}{4}$


5 a



b $\frac{1}{9}$ **c** $\frac{1}{9}$ **d** $\frac{2}{9}$

e $\frac{1}{3}$ **f** $\frac{1}{3}$


6 a



b i $\frac{1}{3}$ **ii** $\frac{2}{3}$ **iii** 1


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

7 a



b i $\frac{1}{3}$ **ii** $\frac{1}{6}$ **iii** $\frac{1}{6}$ **iv** $\frac{5}{6}$

8 a



b $\frac{1}{8}$ **c** $\frac{1}{8}$ **d** $\frac{3}{8}$

e Getting exactly 2 tails

9 a $\frac{3}{5}$ **b** $\frac{2}{5}$ **c** $\frac{7}{10}$

10 a $\frac{1}{9}$ **b** $\frac{7}{18}$ **c** $\frac{1}{2}$ **d** $\frac{1}{6}$ **e** $\frac{1}{9}$ **f** $\frac{1}{9}$

g $\frac{7}{18}$ **h** 0 **i** $\frac{2}{3}$ **j** $\frac{17}{18}$

10E

Now you try

Example 9

a $\frac{4}{5}$ or 0.8 **b** 40

Example 10

a $\frac{5}{24}$ **b** 200

Exercise 10E

1 a Experimental **b** Theoretical **c** Theoretical
d Experimental **e** Expected

2 a $\frac{9}{10}$ **b** $\frac{4}{5}$ **c** $\frac{5}{9}$

3 a $\frac{2}{5}$ **b** 40

4 a 5 **b** 10 **c** 50 **d** 4

5 a $\frac{35}{90} = \frac{7}{18}$ **b** 1050

6 a 0.6
b i 60 **ii** 120 **iii** 360

7 a $\frac{7}{8}$
b i 350 **ii** 4375 **iii** 35

8 a $\frac{1}{15}$ **b** $\frac{2}{5}$ **c** $\frac{11}{60}$


9 a B: $\frac{5}{20} = 0.25$ C: $\frac{30}{100} = 0.3$
b C as it is a larger sample size

10 0.41, from the 100 throws as the more times an experiment is carried out the closer the experimental probability becomes to the actual/theoretical probability

11 a 20 **b** 40 **c** 60 **d** 40

12 a 20
b i $\frac{1}{4}$ **ii** $\frac{7}{20}$
c i 25 **ii** 20 **iii** 45

13 a i 0.52 **ii** 0.48 **iii** 0.78
b 78
c $\frac{1}{2}, \frac{1}{4}, \frac{1}{4}$



14 a $\frac{\text{Shaded area}}{\text{Total area}} = 0.225$, $\therefore$ 100 shots ≈ 23

b $\frac{1}{10} \times 100 = 10$

c $\frac{150 - 32}{150} \times 100 \approx 79$

d $\frac{225\pi - 25\pi}{225\pi} \times 100 \approx 89$

Progress quiz

1 a $\{1, 2, 4, 5, 7\}$
b i $\frac{1}{3}$ **ii** $\frac{2}{3}$ **iii** $\frac{1}{2}$ **iv** $\frac{5}{6}$


2 a $\frac{1}{8}$ **b** $\frac{1}{4}$ **c** $\frac{3}{4}$ **d** $\frac{1}{4}$ **e** $\frac{1}{2}$

3 a

	A	Not A	Total
B	4	5	9
Not B	4	7	11
Total	8	12	20

b i $\frac{2}{5}$ **ii** $\frac{1}{5}$ **iii** $\frac{1}{4}$ **iv** $\frac{13}{20}$

4 a



b i 8 **ii** 15 **iii** 3
c i $\frac{1}{4}$ **ii** $\frac{1}{4}$ **iii** $\frac{7}{20}$

d

	Run	Not run	Total
Class	5	3	8
Not class	7	5	12
Total	12	8	20

5 a

	1	2	3	4
H	H1	H2	H3	H4
T	T1	T2	T3	T4

b 8
c i $\frac{1}{8}$ **ii** $\frac{3}{8}$


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

6 a

		1st choice		
		B	**E**	**E**
2nd choice	**B**	×	(E, B)	(E, B)
	E	(B, E)	×	(E, E)
	E	(B, E)	(E, E)	×

b $\frac{1}{3}$ **c** $\frac{2}{3}$

7 a



b i $\frac{1}{3}$ **ii** $\frac{2}{3}$

8 a $\frac{3}{5}$ **b** 240

10F

Now you try

Example 11

a Numerical discrete **b** Categorical nominal
c Numerical continuous **d** Categorical ordinal

Example 12

a i Stratified **ii** Systematic
iii Simple random
b Yes. Wasram's team-mates at the soccer club are unlikely to represent the views of the wider community.

Exercise 10F

1 a Survey **b** Bias
c Population **d** Sample
2 a Categorical **b** Numerical
3 a Stratified **b** Simple random **c** Systematic
4 a Numerical **b** Categorical
c Categorical **d** Numerical
5 a Categorical nominal **b** Categorical ordinal
c Numerical continuous **d** Numerical discrete
6 a Categorical nominal **b** Numerical continuous
c Categorical ordinal **d** Numerical discrete
7 a i Simple random **ii** Stratified
iii Systematic
b Yes. Only the people who are confident in answering the questions will raise their hand.
8 a i Systematic **ii** Stratified
iii Simple random
b Yes. There is a good chance that more water birds are in that area.
9 Sales (3), Administration (7)
10 Length: Numerical continuous
Weight: Numerical continuous
Favourite food: Categorical nominal
Number of toys: Numerical discrete
Type of scratching post: Categorical ordinal
11 a Expand his sample to include other member of the class
b Survey other vets in the practice too
c Consider the price of bread in other types of shops
d Survey other students not in the extension program
12 The data might be only relevant if sorted into age groups
13 Answers may vary

10G

Now you try

Example 13

a i 5 **ii** 4 **iii** 7 **iv** 6
b i 0.75 **ii** 0.5 **iii** 0 **iv** 9

Example 14

a 87 **b** 97

Exercise 10G

1 a mean
b i 7.8 **ii** 9.9
2 a common (or frequent)
b i 2 **ii** 10
3 a middle
b i 5 **ii** 8
4 a range
b i 21 **ii** 22
5 a outlier
b i 36 **ii** 3
6

	Mean	Median	Mode	Range
a	6	7	8	7
b	8	6	5, 10	13
c	6	6	2	10
d	11	12	none	13
e	4	3.5	2.1	6.2
f	5	4.5	none	9.7
g	3	3.5	−3	12
h	0	2	3	11

7 a Outlier = 33, mean = 12, median = 7.5
b Outlier = 2, mean = 35.2, median = 42
c Outlier = −1.1, mean = 1.075, median = 1.4
d Outlier = −4, mean = −49, median = −59
8 a Yes **b** No
c No **d** Yes
9 a 12 **b** 5.5
c 6 **d** Bimodal; 4, 7
10 a 24.67 s
b 24.8 s
11 a $1 700 000
b ($354 500 and $324 000) drops $30 500
c ($570 667 and $344 800) drops $225 867
12 a 15 **b** 26
13 a 90 **b** 60
14 a 76.75
b i 71.4, B^+ **ii** 75, B^+ **iii** 80.2, A
c 81.4, he cannot get an A^+
d i 43 **ii** 93

10H

Now you try

Example 15

a i 125 **ii** 200 **iii** 175
b 75
c 144°

Exercise 10H

1 Mean, range
2 A skewed, B symmetrical
3 A, the gap looks larger because the vertical scale does not start at zero.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

4 **a** 17 **b** 3, 5, 6, 6, 7, 8, 8, 8, 9, 9, 9, 9, 9, 10, 10, 10, 10
c 8 **d** 9 **e** 7
f Skewed **g** $\frac{12}{17}$

5 **a** 25 **b** 30 cm **c** 173 cm
d 170 cm **e** 168.88 cm
f 186 cm

6 **a** 44% **b** 6600/year **c** 18/day
d 9.5/day **e** 108°

7 **a** 1.42 m **b** 0847, 8 : 47 a.m. **c** 1 hour 21 minutes
d 31 cm or 0.31 m **e** 11 hours 55 minutes

8 **a** 41 years **b** 24 years
c 16 years **d** 10 years, 14.5%

9 **a** 18°C
b 43 mm
c 5°C
d 47 mm
e January, February, March, December
f February, March, May
g August, September
h 9°C

10 **a** 45 m **b** 25 m **c** 20 m **d** 69 m
e 35 m − 14 m = 21 m more


11 **a** Channel 15, 27; channel 16, 24; channel 17, 26; channel 18, 24
b Graph A
c Graph B
d The scale on graph A starts at 23 but the scale on graph B starts at 0.
e Graph A is misleading as the scale expands the difference in column heights.

12 **a** Energy and Yum-crunch equal, Chunky, Sunshine
b No, it was equal to Energy.
c The Yum-crunch column has wider dimensions, a different colour and the cereal name on the column.
d All columns should be the same width, same colour and either all or none have the name.

13 **a** Rugby
b Rugby, soccer, basketball, cricket, tennis
c Rugby: 700 000; Soccer: 600 000
d Cricket, 800 000
e Cricket, rugby, soccer, basketball and tennis equal
f The length of each row is misleading because the ball sizes are not equal.
g Each picture should be an equal size.

14 **a** 10
b $560 000, $560 000, $750 000, $750 000, $750 000, $750 000, $870 000, $870 000, $870 000, $2 300 000
c $903 000
d $750 000
e Mean
f 9 out of 10 house prices are less than the mean of $903 000.
g The median is a better measure of 'average' as it is the middle value. The mean is increased by one very large value.

15 **a**



b 26 inch, 30 inch, 34 inch
c 42 inch, 47 inch, 50 inch, 55 inch, 60 inch, 65 inch

10I

Now you try

Example 16

a

Stem	Leaf
0	9
1	1 3 6
2	1 4 6 9 9
3	4 5 7
4	2 6
5	0 2

3 | 5 means 35

b 29
c 29
d Almost symmetrical

Example 17

a

Class A		Class B
	0	3
9 9 7 5	1	1 2 3 4 6 8
9 4 1	2	0
6 1	3	

2 | 4 means 24 pages

b The number of pages read by class A is between 15 and 36 while for class B it is between 11 and 20 except for a lower value of 3. In general class A is reading more.

Exercise 10I

1 **a** 3.5, 3.7, 4.1, 4.3, 4.8
b 52, 60, 61, 67, 73, 75

2 **a** 9
b **i** 8 min **ii** 35 min
c 21 min
d 21 min

3 **a** 26
b **i** 0 mm **ii** 6 mm
c **i** 21 mm **ii** 24 mm
d **i** 8 mm **ii** 15 mm
e Skewed
f Symmetrical

4 **a** **i**

Stem	Leaf
1	9 9
2	3 4 6 6 8 8 8
3	2 2 3 5
4	1
5	4

2 | 3 means 23

ii 28 **iii** 28 **iv** Skewed

b **i**

Stem	Leaf
1	5 5
2	1 3 3 3 6
3	0 1 1 3 4 5 5 7 9
4	2 2 5 5 5 8
5	0 0 1

4 | 2 means 42

ii 35 **iii** 23, 45 **iv** Almost symmetrical

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

c i

Stem	Leaf
32	0
33	3 7 8
34	3 4 5 5 7 8 9
35	2 2 4 5 8
36	1 3 5
37	0

34 | 3 means 34.3

ii 34.85 **iii** 34.5, 35.2 **iv** Almost symmetrical

d i

Stem	Leaf
15	7 8 9 9 9 9
16	1 4 7 7 7
17	3 5 7 7
18	5 5 7 9
19	3 8
20	0 2

17 | 7 means 177

ii 173 **iii** 159 **iv** Skewed

5 a

Stem	Leaf
0	5 6 6 8 8 9
1	0 1 1 2 2 2 4 4 5 6 7
2	0 1 2

1 | 2 means 12

b 9 **c** 12

6 a i

Set A		Set B
9 3 2	3	1
8 7 6 5 3 3 0 0	4	0 1 3 4 4 6 7 8 8 9 9
4 4 3	5	1 3

3 | 2 means 32

ii Set A has values spread between 32 and 54 while set B has most of its values between 40 and 53 with an outlier at 31.

b i

Set A		Set B
8 7 4 3	0	1 1 1 2 3 6 6 9 9
9 9 7 6 4	1	2 3 5 8 9
6 5 3 1	2	1 5 6
9 9 6 4 3 2	3	3 4 9
7 3 1	4	3 7 8
7 3 2	5	2 3 7
2 1	6	1 2
8 3	8	3 8
1	9	

4 | 1 means 4.1

ii Set A and B are similar. Set A has values between 0.3 and 9.1 and set B has values between 0.1 and 8.8. Both sets have many small numbers and fewer large numbers.

7 a

Collingwood		St Kilda
8 3	6	6 8 8
8 7 2 1 0	7	8
9 9 8 2 0	8	0 0 2 2 3 4 7 8
8 7 5 1	9	0 4
4 3 3 3	10	6 9
9 8 5	11	1 3 7 8
7	12	2 5 6
	13	8

10 | 6 means 106

b Collingwood $33\frac{1}{3}\%$ St Kilda $41\frac{2}{3}\%$

c Collingwood is almost symmetrical data. St Kilda is not symmetrical.

d Collingwood results seem to be consistent. St Kilda has groups of similar scores and, while less consistent, that team has higher scores.

8 a 16.1 s

b 2.3 s

c Yes, 0.05 s lower

9 a

Battery lifetime

Brand A		Brand B
3	7	2 3 4
	7	5 6 8 9 9
4 2	8	0 1 3
9 8 7 5	8	
4 4 3 2 1 0	9	0 1 2 3 4
9 9 8 7 6 5 5	9	5 6 8 8

8 | 3 represents 8.3 hours

b Brand A, 12; brand B, 8

c Brand A consistently performs better than brand B.

10 a 52

b 17.8

11 a 48%

b 15%

c In general, birth weights of babies are lower for mothers who smoke.

10J


Now you try

Example 18

a

Class interval	Frequency	Percentage frequency
10 –	3	25
20 –	2	$16\frac{2}{3}$
30 –	2	$16\frac{2}{3}$
40 –	2	$16\frac{2}{3}$
50–60	3	25
Total	12	100

b



c 3 **d** 25%

Exercise 10J

1 a class interval **b** frequency
c histogram **d** percentage frequency

2 a 20 **b** 20

3 a 3 **b** 360 000–370 000

4 a 2

b 20


c i 30% **ii** 65%

5 a

Class	Frequency	Percentage frequency
0–	23	46
10–	10	20
20–	6	12
30–	7	14
40–	4	8
	50	100

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

b




c **i** 33 **ii** 11

d 34%

6 **a**

Class	Frequency	Percentage frequency
50–	8	$13\frac{1}{3}$
60–	14	$23\frac{1}{3}$
70–	9	15
80–	25	$41\frac{2}{3}$
90–	4	$6\frac{2}{3}$
	60	100

b




c **i** 22 **ii** $63\frac{1}{3}\%$

7 **a**

Number of goals	Frequency
0 – 2	1
3 – 5	8
6 – 8	7
9 – 11	7
12 – 14	7
	30

b



c 9 **d** 7

8 **a** Symmetrical data **b** Skewed data

9 **a** $a = 15$ $b = 2$ $c = 30$ $d = 40$ $e = 20$ $f = 100$

b $a = 50$ $b = 28$ $c = 12$ $d = 40$ $e = 100$

10 85.5%

11 **a** **i** 20% **ii** 55% **iii** 80%
iv 75% **v** 50%

b **i** 30 **ii** 45

c **i** 2 **ii** 22

12 **a** Student A


b Make the intervals for their groups of data smaller so that the graph conveys more information.

13 **a** Minimum wage: \$204; maximum wage: \$940

b **i**

Weekly wages (\$)	Frequency
200–	2
300–	7
400–	6
500–	6
600–	11
700–	11
800–	5
900–	2
Total	50


ii



c **i**

Weekly wages	Frequency
200–	9
400–	12
600–	22
800–	7
Total	50

ii



d More intervals shows greater detail. Since first graph has each pair of intervals quite similar, these two graphs are quite similar.

Maths@Work: Personal trainer

1 **a** 88 bpm **b** 84 bpm
c 98 bpm **d** 119 bpm


2 **a** Yes **b** No **c** Yes **d** No
e Yes

3 Mean 94.6 bpm, mode 90 bpm, median 93.5 bpm, range 35 bpm

4 Mean 105.8 bpm, mode 98 bpm, median 104.5 bpm, range 59 bpm
The data has a higher mean, median and mode, and is more spread out (range is much higher).


ISBN 978-1-009-59315-1
© Greenwood et al. 2025
Photocopying is restricted under law and this material must not be transferred to another party.

5 Graph should look like the following.



Puzzles and games

1 IT'S RANDOM
2 1 – C, 2 – B, 3 – E, 4 – D, 5 – A
3 6 own both cat and a dog
4 2, 3, 4, 8, 8
5 7, 8, 9, 10 and 11
6 **a**




b 12 **c** $\frac{1}{4}$

Chapter checklist

1 **a** $\frac{1}{5}$ **b** $\frac{7}{10}$ **c** $\frac{9}{10}$

2



a 10 **b** $\frac{14}{15}$

3

	Petrol	No petrol	totals
Electric	4	3	7
No electric	11	2	13
Totals	15	5	20

a 2 **b** $\frac{11}{20}$

4

	1	2	3	4
1	2	3	4	5
2	3	4	5	6
3	4	5	6	7
4	5	6	7	8


a $\frac{1}{4}$ **b** $\frac{5}{8}$ **c** $\frac{7}{8}$

5

	K	E	Y
K		K, E	K, Y
E	E, K		E, Y
Y	Y, K	Y, E	


a $\frac{1}{6}$ **b** $\frac{1}{3}$

6



a $\frac{1}{2}$ **b** $\frac{3}{4}$

7



a $\frac{3}{5}$ **b** $\frac{7}{10}$

8 **a** $\frac{1}{4}$ **b** 625

9 **a** Numerical discrete **b** Categorical ordinal
10 **a** Systematic **b** Stratified
11 Yes, as data collected at the airports is likely to be biased towards airline travel.
12 Set 1 : mean = 3, mode = 3, median = 3, range = 4
Set 2 : mean = 18.5, mode = 21, median = 19.5, range = 13
13 Current mean = 68, new score = 78
14 **a** 4 **b** 2.5 **c** 70%
15 **b**: It has an incorrect scale on the vertical axis.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

16

Stem	Leaf
1	5 6 9
2	2 5 6 6 8 9
3	2 5 7
4	2 9

2|5 means 25 Mode = 26 Median = 27

17


Team A		Team B
2	1	2 6
8	2	3 5 7 9 9
1	3	1 4
9 5 0	4	
5	5	
9	6	
6	7	

2|3 means 23

Team A: mean 45, median 45, range 64
Team B: mean 25.1, median 27, range 25
Team B is more conslstent than team A

18

Class interval	frequency	Percentage frequency
30–	2	10
40–	3	15
50–	3	15
60–	5	25
70–	2	10
80–	3	15
90–	2	10
Total	20	100




Percentage of turtles older than 50 years = 75%

Short-answer questions

1 **a** $\frac{2}{3}$ **b** $\frac{5}{9}$ **c** $\frac{3}{5}$

2 **a**




b 15

c **i** $\frac{1}{5}$ **ii** $\frac{2}{5}$ **iii** $\frac{3}{10}$

3 **a**

	A	Not A	Total
B	9	16	25
Not B	8	12	20
Total	17	28	45

b




c **i** $\frac{4}{9}$ **ii** $\frac{1}{5}$ **iii** 8 **iv** 33

4 **a** 12 outcomes

	1	2	3	4
red	(red, 1)	(red, 2)	(red, 3)	(red, 4)
green	(green, 1)	(green, 2)	(green, 3)	(green, 4)
blue	(blue, 1)	(blue, 2)	(blue, 3)	(blue, 4)

b **i** $\frac{1}{6}$ **ii** $\frac{1}{6}$ **iii** $\frac{2}{3}$

5 **a**



b $\frac{2}{3}$

6 **a** $\frac{48}{120} = \frac{2}{5}$ **b** 14

7 **a** Categorical nominal **b** **i** Simple random **ii** systematic
c Yes, her extension science class may have different interests to others in the year level.

8 **a** 26 **b** 26.5 **c** 23, 31 **d** 18

9 **a**

Employee 1		Employee 2
9	1	7 7
5 3 1	2	0 4 8 9
9 8 6 5 5 4	3	2 7 7 8
6 5 0	4	0 1 2 5 8
3 1	5	

2|4 means 24


b **i** Employee 1 : 36, employee 2 : 37
ii Employee 1 : 36, employee 2 : 33
c Employee 1; they have a higher mean and more sales at the high end.
d Employee 1 approximately symmetrical, employee 2 skewed

10 **a**

Class interval	Frequency	Percentage frequency
120–	1	4
130–	3	12
140–	4	16
150–	8	32
160–	7	28
170–	2	8
Total	25	100

ISBN 978-1-009-59315-1
© Greenwood et al. 2025
Photocopying is restricted under law and this material must not be transferred to another party.

b



c **i** 4 **ii** 60%

11 **a** Mean = \$760 000

Median = \$590 000

b The median is better because the mean has been inflated by the outlier of \$2 900 000.

Multiple-choice questions

1 B
2 A
3 **i** A **ii** B **iii** D **iv** C
4 B
5 D
6 E
7 **i** B **ii** C
8 E
9 C


Extended-response questions

1 **a** **i**

		1st spin				
		1	2	3	4	5
2nd spin	1	(1, 1)	(2, 1)	(3, 1)	(4, 1)	(5, 1)
	2	(1, 2)	(2, 2)	(3, 2)	(4, 2)	(5, 2)
	3	(1, 3)	(2, 3)	(3, 3)	(4, 3)	(5, 3)
	4	(1, 4)	(2, 4)	(3, 4)	(4, 4)	(5, 4)
	5	(1, 5)	(2, 5)	(3, 5)	(4, 5)	(5, 5)

ii $\frac{12}{25}$ **iii** $\frac{4}{25}$ **iv** 24 **v** \$68

b **i**



ii 5 **iii** 10 **iv** $\frac{1}{3}$ **v** $\frac{4}{9}$

2 **a**

Airline A		Airline B
4 4 3 2 2 1 1 0	0	1 2 4 4
9 8 7 7 6	0*	5 5 6 7 7 8 9
4 4 4 3 3 3 2 2 1	1	0 0 0 0 1 2 2
9 9 8 8 7	1*	5 5 8 8 8 9 9
4 3	2	1 2 2
	2*	5 6
	3	
	3*	
	4	
	4*	
2	5	

1 | 2 means 12 1* | 5 means 15

b Yes, 52 min

c **i** Airline A: 12, airline B: 10.5
ii Airline A: 10.6, airline B: 12.4

d No, the median time is 12 min so half the flights have less than 12 min delay.

Chapter 11

Warm-up quiz

1 **a** **i** 0 **ii** 4 **iii** 1 **iv** 9
b **i** 0 **ii** 2 **iii** 5 **iv** 27
c **i** 3 **ii** −1 **iii** 2 **iv** −6
d **i** −1 **ii** 7 **iii** −2 **iv** 2

2 **a** Yes **b** No **c** Yes

3 **a** 0 **b** 3 **c** −5
d −3 **e** 4 **f** $\frac{4}{3}$

4 **a** $a^2 - 2a = 0$ **b** $a^2 - 3a + 1 = 0$

5 **a** $2x$ **b** x **c** $3x$
d $2x$ **e** 4 **f** 5

6 **a** $2x - 6$ **b** $2x - x^2$ **c** $x^2 - x - 2$
d $x^2 - 7x + 12$ **e** $x^2 - 4$ **f** $x^2 - 25$

7 **a**

x	−3	−2	−1	0	1	2	3
y	5	0	−3	−4	−3	0	5

b

x	−3	−2	−1	0	1	2	3
y	1	0	1	4	9	16	25

8 **a** $x = 5$ **b** $x = 3$ **c** $x = 1$ **d** $x = 5.5$

11A

Now you try

Example 1

a $x^2 - 3x - 5 = 0$ **b** $2x^2 + 8x - 7 = 0$
c $2x^2 + 3x - 4 = 0$

Example 2

a Not a solution **b** Is a solution

Example 3

a $x = 0, -5$ **b** $x = -2, 3$

Example 4

$x = \frac{2}{3}, -\frac{5}{2}$

Exercise 11A

1 **a** −4 **b** −1 **c** 72
2 **a** No **b** Yes **c** Yes
d No **e** No **f** Yes
3 **a** 1 **b** −3 **c** −2
4 **a** $x^2 - 7x - 2 = 0$ **b** $x^2 - 5x + 2 = 0$
c $x^2 - 4x + 1 = 0$ **d** $3x^2 - 6x - 4 = 0$
e $2x^2 + 2x + 1 = 0$ **f** $3x^2 - 3x + 4 = 0$
g $3x^2 + 4 = 0$ **h** $x^2 - 3x - 1 = 0$
5 **a** Yes **b** Yes **c** No **d** No
e Yes **f** No **g** Yes **h** No
6 Both are solutions
7 Both are solutions
8 **a** 0, −1 **b** 0, −5 **c** 0, 2 **d** 0, 7
e 0, 1 **f** 0, −5 **g** −1, 3 **h** 4, −2
i −7, 3 **j** 5, −4 **k** −7, 6 **l** −12, 9
9 **a** $\frac{1}{2}, -2$ **b** $-2, \frac{1}{3}$ **c** $-\frac{2}{5}, -4$

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

d $1, \frac{1}{3}$ **e** $-5, -\frac{2}{7}$ **f** $\frac{2}{3}, -\frac{1}{5}$

g $\frac{7}{11}, \frac{13}{2}$ **h** $-\frac{9}{4}, \frac{7}{2}$ **i** $\frac{4}{3}, -\frac{1}{7}$

10 a $0, -3$ **b** $0, 7$ **c** $1, -4$ **d** $\frac{1}{2}, -6$

11 a $-1, 2$ **b** $1, 3$ **c** $0, 4$ **d** $0, -3$
e $-4, 1$ **f** $-4, 4$

12 a $(x+2)(x+2) = 0$
b Both solutions are the same, $x = -2$
c **i** -3 **ii** 5

13 a $(x-1)(x+2) = 0$ **b** $x = 1, x = -2$
c Multiplying by a constant doesn't change a zero value.
d **i** $-2, 3$ **ii** $0, -2$

14 a Linear **b** Quadratic **c** Quintic
d Cubic **e** Quartic **f** Quintic

15 a $-2, -1, 3$ **b** $-11, 2, 5$

11B

Now you try

Example 5

$x = 0, -6$

Example 6

$x = 0, 5$

Example 7

$x = -7, 7$

Example 8

$x = -\sqrt{7}, \sqrt{7}$

Exercise 11B

1 a 8 **b** x **c** $3x$

2 a $x(x-3)$ **b** $2x(3x+2)$
c $4x(1-4x)$ **d** $2(x-2)(x+2)$

3 a $x = 0, x = 3$ **b** $x = 0, x = -1$

4 $3^2 = 3 \times 3 = 9$ and $(-3)^2 = -3 \times (-3) = 9$

5 a $x = 0, x = -3$ **b** $x = 0, x = -7$
c $x = 0, x = -4$ **d** $x = 0, x = 5$
e $x = 0, x = 8$ **f** $x = 0, x = 2$
g $x = 0, x = -\frac{1}{3}$ **h** $x = 0, x = \frac{1}{2}$

6 a $x = 0, x = 3$ **b** $x = 0, x = 4$
c $x = 0, x = -5$ **d** $x = 0, x = 3$
e $x = 0, x = 3$ **f** $x = 0, x = -4$

7 a $x = 0, x = 3$ **b** $x = 0, x = 2$
c $x = 0, x = 4$ **d** $x = 0, x = -3$
e $x = 0, x = -4$ **f** $x = 0, x = -3$

8 a $x = 3, x = -3$ **b** $x = 6, x = -6$
c $x = 5, x = -5$ **d** $x = 12, x = -12$
e $x = 9, x = -9$ **f** $x = 20, x = -20$

9 a $x = 2, x = -2$ **b** $x = 3, x = -3$
c $x = 5, x = -5$ **d** $x = 2, x = -2$
e $x = \sqrt{6}, x = -\sqrt{6}$ **f** $x = \sqrt{5}, x = -\sqrt{5}$
g $x = \sqrt{7}, x = -\sqrt{7}$ **h** $x = \sqrt{3}, x = -\sqrt{3}$

10 a 0 or 7 **b** 8 or -8 **c** 0 or -4

11 a $(x+2)(x-2) = x^2 - 4$ not $x^2 + 4$
b No, $x^2 \geqslant 0$, so $x^2 + 4 \geqslant 4$

12 a $x = 2, x = -2$ **b** $x = 5, x = -5$
c $x = 10, x = -10$ **d** $x = 0, x = -7$
e $x = 0, x = 7$ **f** $x = -1, x = 1$

13 a $x = -2, x = 2$ **b** $x = -1, x = 1$
c $x = -1, x = 1$ **d** $x = 0, x = -7$

14 a $x = -\frac{4}{3}, x = \frac{4}{3}$ **b** $x = -\frac{6}{5}, x = \frac{6}{5}$
c $x = -\frac{1}{5}, x = \frac{1}{5}$ **d** $x = -\frac{9}{5}, x = \frac{9}{5}$

11C

Now you try

Example 9

$x = -2, -4$

Example 10

a $x = 5, -2$ **b** $x = 6, 2$

Example 11

$x = 5$

Example 12

$x = 8, -5$

Exercise 11C

1 a $3, 2$ **b** $-5, -2$ **c** $5, -1$ **d** $-4, 3$

2 a $(x+5)(x+7)$ **b** $(x+9)(x-5)$
c $(x-2)(x-8)$

3 a $(x+5)(x+4) = 0$, $x+5 = 0$ or $x+4 = 0$, $x = -5$ or $x = -4$
b $(x-6)(x+4) = 0$, $x-6 = 0$ or $x+4 = 0$, $x = 6$ or $x = -4$

4 a $-6, -2$ **b** $-8, -3$ **c** $-5, -2$ **d** $-4, -6$
e $-6, -3$ **f** $-3, -1$

5 a $4, 8$ **b** $3, 6$ **c** $3, 7$ **d** $-3, 5$
e $-2, 8$ **f** $-5, 9$ **g** $4, 6$ **h** $-6, 7$
i $-12, 7$

6 a -3 **b** -2 **c** -7 **d** -12 **e** 5
f 8 **g** 6 **h** 9

7 a $-2, 5$ **b** $2, 5$ **c** 3 **d** $-4, 1$
e $-7, 2$ **f** 4 **g** $-6, 2$ **h** $-6, 1$
i $3, 5$ **j** $-8, 2$ **k** $-4, -2$ **l** $-9, 2$

8 11 am, 6 pm

9 a $x^2 - 3x + 2 = 0$ **b** $x^2 - x - 6 = 0$
c $x^2 + 3x - 4 = 0$ **d** $x^2 - 7x - 30 = 0$
e $x^2 - 10x + 25 = 0$ **f** $x^2 + 22x + 121 = 0$

10 a Equation is not written in standard form $x^2 + bx + c = 0$ so cannot apply Null Factor Law in this form.
b $x = -2, x = 3$

11 It is a perfect square: $(x-1)(x-1), (x-1)^2 = 0, x = 1$

12 a $-2, 3$ **b** $-5, -3$ **c** $-2, 8$ **d** $2, 3$ **e** 2
f -1 **g** 5 **h** -3 **i** $-7, 1$

11D

Now you try

Example 13

-8 or 5

Example 14

Width = 20 cm and length = 32 cm

Exercise 11D

1 a $-6, 3$ **b** $5, -4$ **c** $2, -5$

2 a $x(x+2) = 8$ **b** $2, -4$
c 2, width > 0 **d** $l = 4$ cm, $w = 2$ cm

3 $-8, 6$

4 $-5, 12$

5 $-2, 15$

6 $l = 20$ cm, $w = 12$ cm

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

7 $l = 9$ cm, $w = 4$ cm
8 $l = 3$ m, $w = 23$ m
9 $h = 5$
10 a $x = 3$ (-4 not valid)
b $x = 12$ **c** $x = 25$
11 -8 not valid because dimensions must be > 0
12 $x = -2$ or $x = 3$; both valid as both are integers
13 a $A = 100 - x^2$ **b** $x = 6$
14 a $A = x^2 + 5x + 15$ **b** $x = 7$

Progress quiz

1 a $x^2 - 3x - 8 = 0$
b $3x^2 - 4x - 6 = 0$
c $2x^2 - 2x - 3 = 0$
2 a $x = 2$ is not a solution **b** $x = -2$ is a solution
3 a $x = 0$ or $x = 7$ **b** $x = 5$ or $x = -2$
c $x = \frac{1}{3}$ or $x = \frac{3}{4}$ **d** $x = 4$ or $x = 3$
4 a $x = 0$ or $x = -11$ **b** $x = 0$ or $x = 6$
c $x = 0$ or $x = -5$ **d** $x = 0$ or $x = 4$
5 a $x = 7$ or $x = -7$ **b** $x = 1$ or $x = -1$
c $x = 2$ or $x = -2$ **d** $x = 3$ or $x = -3$
6 a $x = -3$ or $x = -8$ **b** $x = 3$ or $x = 12$
c $x = -5$ or $x = 7$
7 a $x = -4$ **b** $x = 7$ **c** $x = -3$ or $x = 8$
8 Length 9 m; width 6 m
9 Number is -12 or 7
10 a $x^2 - 5x + 6 = 0$
b $x^2 - 5x - 6 = 0$
c $x^2 + 6x + 9 = 0$

11E

Now you try

Example 15


a $x = 4$
b Maximum
c (4, 6)
d (2, 0), (6, 0)
e (0, −18)

Example 16

a

x	−3	−2	−1	0	1	2	3
y	0	−5	−8	−9	−8	−5	0

b



c i Minimum **ii** $x = 0$ **iii** (0, −9)
iv (0, −9) **v** (−3, 0) and (3, 0)

Exercise 11E

1 a highest **b** parabola **c** intercepts
d vertex **e** lowest **f** zero
2 a $x = 3$ **b** $x = 1$ **c** $x = -2$


3

	i	ii	iii	iv	v
a	$x = 2$	minimum	(2, −1)	(1, 0), (3, 0)	(0, 3)
b	$x = 0$	minimum	(0, −4)	(−2, 0), (2, 0)	(0, −4)
c	$x = 0$	maximum	(0, 3)	(−1, 0), (1, 0)	(0, 3)
d	$x = 0$	maximum	(0, 4)	(−2, 0), (2, 0)	(0, 4)
e	$x = 2$	minimum	(2, 1)	none	(0, 4)
f	$x = -1$	maximum	(−1, 7)	(−4, 0), (2, 0)	(0, 6)
g	$x = 0$	minimum	(0, 0)	(0, 0)	(0, 0)
h	$x = 0$	minimum	(0, −4)	(−2, 0), (2, 0)	(0, −4)
i	$x = 3$	maximum	(3, 4)	(1, 0), (5, 0)	(0, −5)
j	$x = 3$	minimum	$\left(3, -\frac{1}{2}\right)$	(2, 0), (4, 0)	(0, 4)
k	$x = 0$	maximum	(0, 2)	(−1, 0), (1, 0)	(0, 2)
l	$x = -2$	maximum	(−2, −1)	none	(0, −4)

4 a $y = x^2 - 1$

x	−2	−1	0	1	2
y	3	0	−1	0	3


b, c



5 a $y = x^2 + 2x - 3$

x	−4	−3	−2	−1	0	1	2
y	5	0	−3	−4	−3	0	5


b, c



6 a $y = 9 - x^2$

x	−3	−2	−1	0	1	2	3
y	0	5	8	9	8	5	0


b, c



7 a i 1 s, 3 s
ii One time is on the way up, the other is on the way down
b i 2 s **ii** 12 m **iii** 4 s


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

8 **a** 100 m **b** 155 m **c** 5 s
d Journey takes 1 s longer to go down to the ground
9 **a** $x = 1$ **b** $(1, -3)$
10 **a** $y = x^2$
b $y = x^2 - 4$
c $y = 1 - x^2$
d $y = x^2 - 2x + 1$
e $y = -x^2 - 2x$
f $y = x^2 - 3x - 4$
11 **a** Yes




Two x-intercepts

b Yes



One x-intercept

c Yes



Zero x-intercepts

d No
12 **a** $-2^2 = -1 \times 2^2 = -4$. It is not $(-2)^2 = 4$, so correct solution is $y = -2^2 + 2 \times 2 = -4 + 4 = 0$
b $-(-3)^2 = -1 \times 9 = -9$. It is not $-(-3)^2 = +9$, so correct solution is $y = -3 - (-3)^2 = -3 - 9 = -12$
13 **a** $x = -2, x = 2$
b $x = 0$
c **i** Infinite **ii** One **iii** None
14 **a** **i** $x = 0, x = 2$ **ii** $x = -1, x = 3$
b Two; a parabola is symmetrical
c One; this is the minimum turning point
d None; -1 is the minimum y-value so there are no values of y less than -1

11F


Now you try

Example 17

a

x	-2	-1	0	1	2
$y = x^2$	4	1	0	1	4
$y = 4x^2$	16	4	0	4	16
$y = \frac{1}{4}x^2$	1	$\frac{1}{4}$	0	$\frac{1}{4}$	1

b




c Axis of symmetry: $x = 0$, turning point $(0, 0)$
d **i** Narrower
ii Wider

Example 18

a

x	-2	-1	0	1	2
$y = -x^2$	-4	-1	0	-1	-4
$y = -4x^2$	-16	-4	0	-4	-16
$y = -\frac{1}{4}x^2$	-1	$-\frac{1}{4}$	0	$-\frac{1}{4}$	-1

b




c Axis of symmetry: $x = 0$, turning point $(0, 0)$
d **i** Narrower
ii Wider

Exercise 11F

1 **a** $y = 3x^2, y = x^2, y = \frac{1}{2}x^2$
b $y = -3x^2, y = -x^2, y = -\frac{1}{2}x^2$
c $(0, 0)$ **d** $x = 0$
e **i** $y = 3x^2$ **ii** $y = \frac{1}{2}x^2$
iii $y = -3x^2$ **iv** $y = -\frac{1}{2}x^2$
f **i** $y = -x^2$ **ii** $y = -3x^2$
iii $y = \frac{1}{2}x^2$
2 **a** positive
b negative
3 **a**

x	-2	-1	0	1	2
$y = x^2$	4	1	0	1	4
$y = 3x^2$	12	3	0	3	12
$y = \frac{1}{3}x^2$	$\frac{4}{3}$	$\frac{1}{3}$	0	$\frac{1}{3}$	$\frac{4}{3}$

b




c For all three graphs, the turning point is a minimum at $(0, 0)$ and the axis of symmetry is $x = 0$.
d **i** Narrower
ii Wider

ISBN 978-1-009-59315-1
© Greenwood et al. 2025
Photocopying is restricted under law and this material must not be transferred to another party.

4 a i

x	−2	−1	0	1	2
y	24	6	0	6	24

ii



iii Axis of symmetry: $x = 0$; turning point: $(0, 0)$

iv Narrower

b i

x	−2	−1	0	1	2
y	20	5	0	5	20

ii See part **b**.

iii Axis of symmetry: $x = 0$; turning point: $(0, 0)$

iv Narrower

c i

x	−2	−1	0	1	2
y	$\frac{2}{3}$	$\frac{1}{6}$	0	$\frac{1}{6}$	$\frac{2}{3}$

ii See part **a**.

iii Axis of symmetry: $x = 0$; turning point: $(0, 0)$

iv Wider

d i

x	−2	−1	0	1	2
y	$\frac{4}{5}$	$\frac{1}{5}$	0	$\frac{1}{5}$	$\frac{4}{5}$

ii See part **a**.


iii Axis of symmetry: $x = 0$; turning point: $(0, 0)$

iv Wider

5 a i

x	−2	−1	0	1	2
y	−8	−2	0	−2	−8

ii



iii Axis of symmetry: $x = 0$; turning point: $(0, 0)$; x- and y-intercept: $(0, 0)$

iv Narrower

b i

x	−2	−1	0	1	2
y	−12	−3	0	−3	−12

ii See part **a**.

iii Axis of symmetry: $x = 0$; turning point: $(0, 0)$; x- and y-intercept: $(0, 0)$

iv Narrower

c i

x	−2	−1	0	1	2
y	−2	$-\frac{1}{2}$	0	$-\frac{1}{2}$	−2

ii See part **a**.

iii Axis of symmetry: $x = 0$; turning point: $(0, 0)$; x- and y-intercept: $(0, 0)$

iv Wider

d i

x	−2	−1	0	1	2
y	$-\frac{4}{3}$	$-\frac{1}{3}$	0	$-\frac{1}{3}$	$-\frac{4}{3}$

ii See part **a**.

iii Axis of symmetry: $x = 0$; turning point: $(0, 0)$; x- and y-intercept: $(0, 0)$

iv Wider

6 a A **b** H

7 a D **b** C **c** A
d E **e** F **f** B

8 No, because both transformations are multiplying to 'a' and multiplication is commutative: $bc = cb$

9 $y = \frac{67}{242\,064}x^2$

10 a Reflection in the x-axis, dilation by a factor of 3 from the x-axis

b Reflection in the x-axis, dilation by a factor of 6 from the x-axis

c Reflection in the x-axis, dilation by a factor of $\frac{1}{2}$ from the x-axis

d Reflection in the x-axis, dilation by a factor of 2 from the x-axis

e Reflection in the x-axis, dilation by a factor of 3 from the x-axis

f Reflection in the x-axis, dilation by a factor of $\frac{1}{3}$ from the x-axis

11 a $y = -4x^2$ **b** $y = \frac{1}{3}x^2$

c $y = -4x^2$ **d** $y = -\frac{4}{3}x^2$


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

11G


Now you try

Example 19

a




b




Example 20

a




b




Example 21

a b




Exercise 11G


1 **a** **i** $(-2, 0)$ **ii** $(3, 0)$
b Left **c** Right
2 **a** **i** $(0, -3)$ **ii** $(0, 2)$
b Down **c** Up
3 **a** $(0, 3)$ **b** $(0, -4)$ **c** $(0, -4)$ **d** $(0, 25)$
4 **a** $y = x^2 + 1$ **b** $y = x^2 + 3$
c $y = x^2 - 2$




d $y = -x^2 + 4$ **e** $y = -x^2 + 1$
f $y = -x^2 - 5$




5 **a** $y = (x - 2)^2$ **b** $y = (x - 4)^2$
c $y = (x + 3)^2$



d $y = -(x - 3)^2$ **e** $y = -(x + 6)^2$
f $y = -(x + 2)^2$




6 **a** **i** $(2, 3)$ **ii** $(-1, -1)$
b **i** one, left, one, down **ii** two, right, three, up
7 **a** $y = (x - 3)^2 + 2$




ISBN 978-1-009-59315-1
© Greenwood et al. 2025
Photocopying is restricted under law and this material must not be transferred to another party.


b $y = (x-1)^2 - 1$




c $y = (x+2)^2 - 3$




d $y = (x+1)^2 + 7$




e $y = -(x-2)^2 + 1$




f $y = -(x-5)^2 + 3$




g $y = -(x+3)^2 - 4$



h $y = -(x+1)^2 - 5$



i $y = -(x-3)^2 - 6$




8 **a** J **b** D **c** B **d** H **e** K
f G **g** A **h** E **i** L **j** I
k F **l** C

9 **a** $y = (x-1)^2 + 1$ **b** $y = (x+2)^2 + 2$
c $y = (x+1)^2 - 3$ **d** $y = -(x-1)^2 + 4$
e $y = -(x-2)^2 - 2$ **f** $y = -(x+2)^2 + 4$

10 **a** (h, k) **b** $y = ah^2 + k$

11 **a** $y = -(x-2)^2 + 9$, turning point (2, 9)
$y = (x-7)^2 - 4$, turning point (7, −4)

b



11H

Now you try

Example 22


a i (0, 0), (−5, 0) **ii** (0, 0)
b i (−1, 0), (4, 0) **ii** (0, −12)

Example 23

a (0, 0) **b** $y = x(x-4)$ **c** (0, 0), (4, 0)
d $x = 2$ **e** (2, −4)

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


f



Example 24

a (0, −12) **b** $y = (x - 6)(x + 2)$
c (6, 0), (−2, 0) **d** $x = 2$ **e** (2, −16)

f



Exercise 11H


1 a $x(x + 2)$ **b** $x(x - 3)$
c $(x + 3)(x - 3)$ **d** $(x + 7)(x - 7)$
e $(x - 4)(x + 3)$ **f** $(x - 2)(x - 2) = (x - 2)^2$

2 a 0 **b** −3 **c** −1 **d** −0.5


3 a −4 **b** −9 **c** −1

4 a i (0, 0), (−7, 0) **ii** (0, 0)
b i (0, 0), (−3, 0) **ii** (0, 0)
c i (0, 0), (−4, 0) **ii** (0, 0)
d i (4, 0), (−2, 0) **ii** (0, −8)
e i (−2, 0), (5, 0) **ii** (0, −10)
f i (−3, 0), (7, 0) **ii** (0, −21)
g i (−3, 0), (1, 0) **ii** (0, −6)
h i (−4, 0), (−1, 0) **ii** (0, 12)
i i (2, 0), (3, 0) **ii** (0, 6)


5 a $y = x(x - 5)$




b $y = x(x + 1)$




c $y = x(x - 3)$




d $y = x(2 + x)$




e $y = x(5 + x)$



f $y = x(3 - x)$




g $y = -x(x + 8)$




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


h $y = -x(2 + x)$




i $y = -x(x - 1)$




6 a $y = (x - 2)(x - 1)$




b $y = (x - 3)(x - 1)$




c $y = (x + 3)(x - 1)$




d $y = (x + 2)^2$



e $y = (x + 1)^2$



f $y = (x + 4)(x - 2)$




7 a -3 **b** 2 **c** 3
d -4 **e** -5 **f** 6

8 a 30 m **b** 225 m

9 a 200 m **b** 1000 m


10 There is only one x-intercept so this must be the turning point.

11 a $y = x^2 - 1$




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


b $y = 9 - x^2$




c $y = x^2 + 5x$




d $y = 2x^2 - 6x$




e $y = 4x^2 - 8x$




f $y = x^2 - 3x - 4$




g $y = x^2 + x - 12$



h $y = x^2 + 2x + 1$



i $y = x^2 - 6x + 9$



Maths@Work: Fencing Crown land

1 a i 300 m **ii** 160 m **iii** 40 m

b i 15 000 m^2 **ii** 19 200 m^2 **iii** 7200 m^2

c when the width is 120 m gives the largest area. Small and large values at the width result in very thin rectangles, resulting in a small area.

2 a $L = 400 - 2x$

b $A = x(400 - 2x)$


c i 7200 m^2 **ii** 19 200 m^2 **iii** 12 800 m^2

3 a i (0, 0)

ii (0, 0) and (200, 0)

iii $x = 100$


iv (100, 20 000)




b $x = 100$
This is the turning point of the graph.
Maximum area = 20 000 m^2

c width = 100 m, Length = 200 m, the length is double the width.

d $A = x(600 - 2x)$
Yes, the length is double the width for the maximum area.



4



ISBN 978-1-009-59315-1
© Greenwood et al. 2025
Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Puzzles and games

1 **a** They are square numbers.
b $100^2 = 10\,000$

2 **a** $\frac{5}{3}, -\frac{7}{2}$ **b** $2, -1$ **c** $2^{\frac{2}{3}} + 1$

3 6 seconds

4 $\frac{n(n+1)}{2}$; $n = 11$

5 $x = 2$

6 **a** 20 units, 21 units, 29 units
b 17 units

7 **a** $k < 0$ **b** $k = 0$ **c** $k > 0$

Chapter checklist

1 $x^2 - 3x - 6 = 0$

2 Yes, $x = -2$ is a solution

3 $x = -2$ or $x = 5$

4 $x = 0$ or $x = 5$

5 $x = \pm 2$

6 $x = 10$ or $x = -7$


7 $x = 4$ or $x = 5$

8 Width = 7 cm, length = 12 cm

9 **i** $x = 1$
ii Minimum turning point $(1, -4)$
iii x-intercepts: $(-1, 0)$ and $(3, 0)$; y-intercept: $(0, -3)$


10

x	−3	−2	−1	0	1	2	3
y	25	16	9	4	1	0	1



11


x	−2	−1	0	1	2
$y = x^2$	4	1	0	1	4
$y = 3x^2$	12	3	0	3	12
$y = \frac{1}{3}x^2$	$1\frac{1}{3}$	$\frac{1}{3}$	0	$\frac{1}{3}$	$1\frac{1}{3}$



Turning point of each graph: minimum at $(0, 0)$
The graph of $y = 3x^2$ appears narrower than the graph of $y = x^2$
The graph of $y = \frac{1}{3}x^2$ appears wider than the graph of $y = x^2$


12

x	−2	−1	0	1	2
$y = -x^2$	−4	−1	0	−1	−4
$y = -4x^2$	−16	−4	0	−4	−16
$y = -\frac{1}{4}x^2$	−1	$-\frac{1}{4}$	0	$-\frac{1}{4}$	−1




Turning point of each graph: maximum at $(0, 0)$
The graph of $y = -4x^2$ appears narrower than the graph of $y = -x^2$
The graph of $y = -\frac{1}{4}x^2$ appears wider than the graph of $y = -x^2$


13



14




15




ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

16 x-intercepts: (0, 0) and (6, 0)
y-intercept: (0, 0)
turning point: (3, −9)



17 x-intercepts: (−2, 0) and (6, 0)
y-intercept: (0, −12)
turning point: (2, −16)




Short-answer questions

1 a $y = x^2 - 2x - 3$

x	−3	−2	−1	0	1	2	3
y	12	5	0	−3	−4	−3	0

b



2 a 0, −2 **b** 0, 4 **c** −3, 7 **d** 2, −2
e $-1, \frac{2}{5}$ **f** $\frac{1}{2}, \frac{4}{3}$

3 a 0, −3 **b** 0, 4 **c** −5, 5
d −9, 9 **e** −2, 2 **f** $-\sqrt{10}, \sqrt{10}$
g −3, −7 **h** −5, 8 **i** 4


4 a $x^2 - 5x = 0$; $x = 0, 5$
b $3x^2 - 18x = 0$; $x = 0, 6$
c $x^2 + 8x + 12 = 0$; $x = -6, -2$
d $x^2 - 2x - 15 = 0$; $x = -3, 5$
e $x^2 - 8x + 15 = 0$; $x = 3, 5$
f $x^2 + 3x - 4 = 0$; $x = -4, 1$

5 a $x^2 + 2x - 80 = 0$; $x = 8$ units
b $x^2 + 5x - 24 = 0$; $x = 3$ units


6 a i $x = 3$ **ii** minimum (3, −2)
b i $x = -1$ **ii** maximum (−1, 3)

7 a $y = 2x^2$ **b** $y = -x^2 + 2$
c $y = (x + 1)^2$ **d** $y = (x - 3)^2$
e $y = x^2 - 4$


8 a $y = x^2 + 2$




b $y = -x^2 - 5$




c $y = (x + 2)^2$



d $y = -(x - 3)^2$




e $y = (x - 1)^2 + 3$




ISBN 978-1-009-59315-1
© Greenwood et al. 2025
Photocopying is restricted under law and this material must not be transferred to another party.


f $y = 2(x+2)^2 - 4$




9 a $y = x^2 - 8x + 12$




b $y = x^2 + 10x + 16$



c $y = x^2 + 2x - 15$



d $y = x^2 + 4x - 5$




Multiple-choice questions

1 D	**2** B	**3** C	**4** A	**5** D
6 B	**7** E	**8** C	**9** A	**10** B


Extended-response questions

1 a $x + 5$ **b** $A = x^2 + 5x$
c $x = 2$ ($x = -7$ not valid as $x > 0$)
d Base = 4 m, height = 7 m

2 a i (0, 0), (6, 0) **ii** Maximum at (3, 18)
iii



b i (0, 0)
ii (4, 16)
iii (0, 0), (8, 0)
iv



c i Connor's rocket, 8 s
ii Sam's rocket, 2 m higher
iii 16 m, the same height

Semester review 2

Measurement

Short-answer questions

1 a 6.25% **b** 6.35 kg, 6.45 kg
2 a 22 cm^2 **b** 12 cm^2 **c** 25.92 m^2
3 a 25.13 m, 50.27 m^2 **b** 62.83 cm, 314.16 cm^2
c 13.19 cm, 13.85 cm^2
4 a 54.82 cm^2 **b** 26 m^2 **c** 24.57 m^2 **d** 36 cm^2
5 a 220 m^2 **b** 216 m^2 **c** 25.45 m^2
6 a 12 m^3 **b** 42 m^3 **c** 100.53 cm^3

Multiple-choice questions

1 C **2** C **3** D **4** C **5** A

Extended-response question

a 12.57 m^2
b 77.43 m^2
c $628

Indices

Short-answer questions

1 a $15p^3q^2$ **b** $\frac{a^2b}{2}$ **c** $54x^9y^6$ **d** -2

2 a $3 \times 4^2 = 48$ **b** $\frac{4}{10^3} = 0.004$

3 a 2400 **b** 1 080 000
c 0.0071 **d** 0.0000206

4 a 6.03×10^4 **b** 2.7×10^6
c 4×10^{-3} **d** 7.03×10^{-4}

5 a 3.07×10^{-2} kg **b** 4.24×10^6 kg
c 1.22×10^4 s **d** 7.45×10^{-6} years

Multiple-choice questions

1 C **2** B **3** D **4** E **5** C

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Extended-response question

a i 74 000 000 000 **ii** 7.4×10^{10}
b 1.87×10^{17}
c 8.72×10^{-7}

Geometry

Short-answer questions

1 a 180° **b** 360° **c** 540°
d 900° **e** 1080° **f** 1440°
2 a $x = 70$ **b** $x = 70$ **c** $a = 100, b = 140$
d $a = 70, b = 110$ **e** $x = 100, y = 110$
f $x = 67, y = 98$ **g** $x = 120$
h $a = 135$
3 a Yes, RHS **b** $\Delta ABC \equiv \Delta DEF$
c AB
4 a SAS **b** 2
5 a AAA **b** 2.4

Multiple-choice questions

1 C **2** B **3** A **4** E **5** B

Extended-response question

a AAA **b** 6 m **c** $\frac{5}{3} = 1.\dot{6}$ **d** 7.5 m
e 3.75 m

Algebraic techniques

Short-answer questions

1 a $-3x - 6$ **b** $x - x^2$ **c** $x^2 - 1$
d $x^2 - 8x + 16$ **e** $25x^2 - 4$
f $9x^2 - 24x + 16$ **g** $8x^2 + 10x - 3$
h $2x + 3$ **i** $4 - 4x + x^2$
2 a $-6x^2y$ **b** $42a^2b^2$ **c** $\frac{a}{2}$ **d** $\frac{-6x}{y}$
e $\frac{a}{b}$ **f** $\frac{1}{2y}$
3 a $x^2 - 9$ **b** $x^2 + 4x + 4$
c $6x^2 - 17x + 12$
4 a $3(x - 4)$ **b** $-7(x + 2)$
c $x(5x + 2)$ **d** $7x(2x - 3)$
5 a $(x + 3)(x - 3)$ **b** $(5 + 2y)(5 - 2y)$
c $(x - 6)(x + 3)$ **d** $(x + 2)(3x - 5)$
6 a $(x + 4)(x + 5)$ **b** $(x + 4)(x - 2)$
c $(x - 8)(x + 7)$ **d** $2(x - 6)(x + 4)$
7 a $\frac{3x}{2}$ **b** $\frac{3}{4}$ **c** 4 **d** $-x$
e $\frac{3}{2}$ **f** $\frac{9a}{x}$
8 a $\frac{x}{12}$ **b** $\frac{53x}{20}$ **c** $\frac{7x + 5}{10}$ **d** $\frac{11x + 7}{15}$

Multiple-choice questions

1 E **2** E **3** A **4** B **5** D

Extended-response question


a $10 - 2x$ and $8 - 2x$
b $(10 - 2x)(8 - 2x) = 80 - 36x + 4x^2$
c 48 m^2

Statistics and probability

Short-answer questions

1 a Stratified **b** Systematic

2 a




b i 24 **ii** 16
c i $\frac{1}{5}$ **ii** $\frac{1}{3}$

3 a

	1	2	3	4
H	H1	H2	H3	H4
T	T1	T2	T3	T4


b 8 **c** $\frac{1}{8}$ **d** $\frac{1}{4}$

4 a



b $\frac{1}{9}$ **c** $\frac{1}{3}$ **d** $\frac{2}{9}$

5 a



b i $\frac{1}{3}$ **ii** $\frac{1}{6}$ **iii** $\frac{1}{2}$

6 a Numerical
b

Stem	Leaf
1	0 1 1 2 3 5 7 8
2	2 5 5 5 6
3	1 2 2

1 | 3 means 13 aces

c Mode = 25, median = 20
d Skewed

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

7 a

Class interval	Frequency	Percentage frequency
0–	2	6.7
50–	4	13.3
100–	5	16.7
150–	9	30
200–	7	23.3
250–	3	10
Total	30	100

b



c i 6 **ii** 60%

Multiple-choice questions

1 D
2 A
3 a C **b** D
4 B
5 B

Extended-response question

a

		Red		
		1	2	3
Green	1	(1, 1)	(1, 2)	(1, 3)
	2	(2, 1)	(2, 2)	(2, 3)
	3	(3, 1)	(3, 2)	(3, 3)

b $\frac{4}{9}$

c i $\frac{1}{3}$ **ii** 18

d i 17 years **ii** 92


Introduction to quadratic equations and graphs

Short-answer questions


1 a $x = -5$ or $x = 3$ **b** $x = \frac{1}{2}$ or $x - \frac{5}{3}$
c $x = 0$ or $x = -2$ **d** $x = -3$ or $x = 3$
e $x = 2$ or $x = 7$ **f** $x = -4$

2 15 m by 8 m


3 a translated 3 units up




b Reflected in the x-axis and translated 4 units left



c Translated 2 units right and 5 units up



4 a (0, −12) **b** (−2, 0), (6, 0)
c (2, −16) **d**




Multiple-choice questions

1 B **2** B **3** E **4** C **5** E

Extended-response question

a (0, 0) and (6, 0), lands after 6 seconds.
b i 180 cm **ii** 3 seconds
c After 2 seconds and 4 seconds
d



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Essential Mathematics Core for the Victorian Curriculum Year 9
ISBN 978-1-009-59315-1
© Greenwood et al. 2025
Photocopying is restricted under law and this material must not be transferred to another party.
Cambridge University Press

12

Algorithmic thinking



Algorithms and networks

Imagine trying to organise the efficient running of a busy rail network without access to special programs and algorithms to help synchronise all the components of the system. The associated field of study is called Networks and this has its roots in Mathematics, Computer Science and Engineering. When working with Networks, the aim is to look at all the possible ways in which different trains can utilise platforms and rail networks so that the mass movement of people can occur efficiently and safely.

People who work with networks use computer programs and algorithmic thinking to set up possible timetables that aim to maximise the capacity of the system. Algorithmic thinking is required to analyse the situation and consider all the possible factors, and then computer programs help to find the most suitable models.

Such algorithmic modelling is critical in the management of busy rail networks as well as other transport systems all around the world.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025
Photocopying is restricted under law and this material must not be transferred to another party.
Cambridge University Press



In this chapter

Activity 1: Algorithms for number patterns and financial mathematics

1.1 Algorithms in number patterns

1.2 Algorithms in financial mathematics

Activity 2: Minimising and maximising

2.1 Cardboard boxes

2.2 Cylindrical cans

Activity 3: Sorting, simulations and sampling

3.1 Ways of sorting data to find the median

3.2 Simulations in probability calculations

Victorian Curriculum 2.0

This chapter covers the following content descriptors in the victorian Curriculum 2.0:

SPACE

VC2M9A06, VC2M9SP03, VC2M9P03

Please refer to the curriculum support documentation in the teacher resources for a full and comprehensive mapping of this chapter to the related curriculum content descriptors.

© VCAA

Online resources

A host of additional online resources are included as part of your Interactive Textbook, including HOTmaths content, video demonstrations of all worked examples, auto-marked quizzes and much more.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025
Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Introduction

An **algorithm** is a sequence of steps that, when followed, lead to the solution of a problem. It has a defined set of inputs and delivers an output. Each step in the algorithm leads to another step or completes the algorithm.

Algorithms occur in mathematics and computing, as well as in simple areas of daily life such as following a recipe. Algorithmic thinking is a type of thinking that involves designing algorithms to solve problems. The algorithms we design can then be written in a way that a computer program will understand, so that the computer does the hard computational work.

In the following activities you will carry out some algorithms as well as think about the design, analysis and implementation of your own algorithms.


The algorithms in the following activities will be described through the use of spreadsheets, flow charts, pseudocode (an informal programming language) and simulations. The following symbols will be used in the flow charts with arrows used to connect each stage.




For input/output stages



For process stages



For decision stages



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


Activity 1: Algorithms for number patterns and financial maths

Number and Algebra

Algorithms can be used to generate number patterns as well as to carry out tasks in the financial world. Some examples are seen in the following parts.

1.1 Algorithms in number patterns

Consider the flow-chart algorithm shown on the right.




a Trace through the algorithm by completing a table like the one shown below, updating the variable values as you go. Add any output to the line at the bottom of the table. The first run through to 'Is count < 10?' is filled in for you.

num	*n*	*count*
1	1	1
4	2	2
.	.	.
.	.	.
.	.	.
Output: 1, 4, …		


b What is the sequence of numbers generated in the output of part **a**?

c Which part of the algorithm controls how many numbers are displayed?

The triangular numbers are another number sequence as shown below.



d Consider the pattern in the triangular numbers by filling in the numbers in the boxes below.



e Complete a flow chart that generates the first 10 triangular numbers. Copy the flow chart from part **a**. The only box you will need to change is:

$num = num + n + (n + 1)$
$n = n + 1$

Test it on a friend using a table to see if they get the correct list of numbers as they trace through the algorithm.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


1.2 Algorithms in financial mathematics



a *Payslip calculations*

The weekly payslip for workers can be automatically generated once their hourly pay rate, their number of hours worked and their overtime hours are entered.


i Create the spreadsheet below using the given formulas. Recall that time-and-a-half is 1.5 times the normal rate and double time is two times the normal rate.
The workers' hours need to be entered; then in cell G3 enter the formula for the total pay. This formula can then be filled down for each worker.



	A	B	C	D	E	F	G
1	**Workers Inc. Payslip**						
2	**Employee**		**Hourly Rate**	Hours at normal rate	Time and a half hours	Double time hours	**Total pay**
3	Brown	Kai	18.7	18	5	2	=D3*C3+E3*1.5*C3+F3*2*C3
4	Clark	Emily	32.2	20	4	4	=D4*C4+E4*1.5*C4+F4*2*C4
5	Fredericks	Gwen	38.7	24	0	0	=D5*C5+E5*1.5*C5+F5*2*C5
6	Martino	Jack	21.3	12	6	6	=D6*C6+E6*1.5*C6+F6*2*C6
7	Perkins	Felicity	21.3	0	6	6	=D7*C7+E7*1.5*C7+F7*2*C7
8	Thomas	Harry	32.2	18	2	2	=D8*C8+E8*1.5*C8+F8*2*C8
9	Xu	Tony	21.3	10	8	8	=D9*C9+E9*1.5*C9+F9*2*C9
10							
11							
12							
13							
14							
15							
16							
17							
18							

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

ii In some work situations, overtime depends on the number of hours worked in a week. Consider the scenario where the number of hours worked over 38 hours earn time-and-a-half while the number of hours worked over 44 hours earn double time. Fill in the blanks in the flow-chart algorithm shown that would calculate an employee's earnings in a week under this system. The hourly rate (*rate*) and number of hours worked in a week (*hours*) are required as inputs.



iii Test your algorithm from part **ii** with the following inputs:

- *rate* = \$22 and *hours* = 32
- *rate* = \$22 and *hours* = 40
- *rate* = \$22 and *hours* = 48

b *Income tax calculation algorithm*

In section 2F you saw tax tables used to calculate how much tax a person must pay based on their taxable income.

The taxable income is taken as the gross income minus deductions. An example of a tax table with different tax brackets is shown below. Here you are taxed at a higher rate for dollars earned over certain amounts.

Taxable income	Tax on this income
0 – \$18 200	Nil
\$18 201 – \$37 000	19c for each \$1 over \$18 200
\$37 001 – \$80 000	\$3572 plus 32.5c for each \$1 over \$37 000
\$80 001 – \$180 000	\$17 547 plus 37c for each \$1 over \$80 000
\$180 001 and over	\$54 547 plus 45c for each \$1 over \$180 000

For example, for a taxable income of \$40 000 you are in the \$37 001 – \$80 000 tax bracket:

$$\begin{aligned}\text{Tax} &= \$3572 + 0.325 \times (\$40\,000 - \$37\,000) \\ &= \$4547\end{aligned}$$

i Using the above table, find the tax payable for these taxable incomes.

I \$10 000 **II** \$25 000 **III** \$50 000
IV \$100 000 **V** \$200 000

ii Complete an algorithm flow chart so that it determines a person's tax payable based on input of their gross income. Use the flow chart in part **a ii** as a guide.

iii Have a friend trace through your algorithm using a gross income of \$85 000. Do they complete the algorithm with the correct answer?

iv List four other values, chosen from different tax brackets, to test that your algorithm works correctly.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.


c *Simple interest*

In section 2G you studied simple interest. Recall that this is the interest (I) calculated at a set rate ($r\%$), over a certain period of time (t) on a principal amount (P). When calculating simple interest, the interest is the same for each period.

The formula is: $I = \frac{Prt}{100}$.


Consider the case where a sum of $\$2000$ is invested in a simple interest bank account with an interest rate of 4% p.a. for 5 years.

i Set up a spreadsheet like the one shown in Figure 1 below to display the total amount of interest earned after each year and the account balance. Once the initial formulas are entered, fill down the columns until $t = 5$. The formula in cell B5 is the simple interest formula using the given values.



	A	B	C	D
1	Principal	2000		
2	Interest rate % p.a.	4		
3				
4	Number of Years	Interest Earned	Amount	
5	1	=($B$1*$B$2*A5)/100	=$B$1+B5	
6	=A5+1	=($B$1*$B$2*A6)/100	=$B$1+B6	
7	=A6+1	=($B$1*$B$2*A7)/100	=$B$1+B7	
8	=A7+1	=($B$1*$B$2*A8)/100	=$B$1+B8	
9	=A8+1	=($B$1*$B$2*A9)/100	=$B$1+B9	
10				
11				
12				
13				

Figure 1 showing formulas



	A	B	C	D
1	Principal	$2,000		
2	Interest rate % p.a.	4		
3				
4	Number of Years	Interest Earned	Amount	
5	1	80	$2,080	
6	2	160	$2,160	
7	3	240	$2,240	
8	4	320	$2,320	
9	5	400	$2,400	
10				
11				
12				
13				

Figure 2 showing calculated values

The table in Figure 2 could also be generated in programming languages. The first stage is to design the flow-chart algorithm.

ii The steps of the algorithm are listed below. Use these steps to construct a flow chart for the algorithm.

Step 1 Receive inputs P, r and t.
Step 2 Set the counter n to $n = 1$ to represent the end of the first year.
Step 3 Calculate the simple interest and the current balance for year n.
Step 4 Display the current year n, the interest earned and the balance.
Step 5 Increase n by 1.
Step 6 If $n \leq t$ repeat steps 3 to 6, else End.

iii Use your algorithm in part **ii** above to find the total amount after 5 years if $P = 5000$ and $r = 6$.


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Activity 2: Minimising and maximising

Measurement and Geometry

In the following we will consider two common scenarios. The first is for a fixed surface area (amount of material), looking at the dimensions that maximise the volume. The second for a fixed volume of an object looking at the dimensions that minimise the surface area (i.e. the material required to form it).

2.1 Cardboard boxes



A packaging company has cardboard sheets 320 cm^2 in area to be used to form boxes. The boxes are required to have square bases. The box will use all the cardboard, so the 320 cm^2 will be the surface area.

The company wishes to set the dimensions of the box so that the box has the largest possible volume.




a For the square-based rectangular prism shown above, write down the formula for:

i its volume **ii** its surface area

b Using the surface area of 320 cm^2, the surface area formula can be rearranged to show that $h = \frac{320 - 2x^2}{4x}$.

In a spreadsheet, make a column for x and create columns to calculate h (using the formula above) and the volume. Start at $x = 0.1$ and increment (increase) your x values by 0.1. Using a smaller increment will give greater accuracy.



	A	B	C	D
1	Fixed surface area, cm^2:	320		
2				
3	base side length, x cm	height, h cm	volume, cm^3	
4	0.1	=($B$1-2*A4^2)/(4*A4)	=A4^2*B4	
5	=A4+0.1			
6				
7				
8				
9				

Fill down each column.

c From your spreadsheet determine the box dimensions (x and h) that maximise the box's volume.

d Use your spreadsheet from part **b** for the following different fixed surface areas and find the dimensions that deliver a maximum volume. You will only need to change the value in cell B1.

i 660 cm^2 **ii** 850 cm^2

e In conclusion, what shaped box maximises the volume of the box for a given surface area?


ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

2.2 Cylindrical cans

A company packages tinned food in cylindrical cans. It requires a fixed volume for these cans and wishes to use dimensions that give the minimum surface area to minimise the cost of production.

Recall the following formulas for a cylinder:


Volume: $V = \pi r^2 h$
Surface area: TSA $= 2\pi r^2 + 2\pi rh$



Let the required volume of the cans be 340 cm^3.

By letting $V = 340$ in $V = \pi r^2 h$, the formula can be rearranged to make h the subject to give $h = \dfrac{340}{\pi r^2}$.

a In a spreadsheet make a column for r and create columns to calculate h (using the formula above). Also create a column for the surface area. Start at $r = 0.1$ and increment (increase) your r values by 0.1.
In Microsoft Excel, the PI() function accesses π.



	A	B	C	D
1	Fixed Volume, V cm^3:	340		
2				
3	**radius, r cm**	**height, h cm**	**surface area, cm^2**	
4	0.1	=$B$1/(PI()*A4^2)	=2*PI()*A4^2+2*PI()*A4*B4	
5	=A4+0.1			
6				

Fill down each column.

b Use the graphing features of your spreadsheet to sketch a graph of surface area against radius. You should notice a minimum (lowest point) on your graph. Refer to your spreadsheet and write down the closest r and h values that give the minimum surface area.

c Repeat parts **a** and **b** using the following different volumes. (In your spreadsheet you will only need to change cell B1).
i $V = 400$ cm^3 **ii** $V = 500$ cm^3

d What do you notice about the relationship between the r and h values of a cylinder that minimise its surface area? Do you see many tin cans like this?

e *Extension*: Alter your increment size in your spreadsheet to see if you can find the r and h values that deliver the minimum surface area to a greater level of accuracy.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Activity 3: Sorting, simulations and sampling

Statistics and Probability

3.1 Ways of sorting data to find the median

Many statistical calculations have a set process or algorithm for calculating them. For example, the mean is found by adding the data values and dividing by the number of values. The range is found by finding the maximum and minimum values and calculating the difference between them.

As you have worked through in the past, the process for finding the median of a set of data is:

Step 1 Sort the data in ascending order.

Step 2 Locate the middle value. If there is an odd number of data values it will be the middle value; if an even number of data values, average the two middle values.

1 2 ④ 7 10
median

3 5 8 10 15 20
$\frac{8+10}{2}$ = ⑨ median

From an algorithmic point of view the most involved part of this process is sorting the list of data.

There are a number of different sorting algorithms already in existence. Here we will compare two different types: selection sort (which you may have seen in Year 8) and bubble sort.

3.2 Selection sort

This method involves finding the smallest (or largest) element in a list and swapping it with the first (or last) item in the list and then moving along the list until it is sorted.

For example, the data set ⑦ 8 6 12 ② 3 9 after one pass through becomes 2 8 6 12 7 3 9 where the minimum value 2 moves to the start of the list and swaps with the 7. The next pass through becomes 2 3 6 12 7 8 9 and so on.

a Use the data sets listed below to trace through the selection sort algorithm. Complete a table like the one shown to show the algorithm in action. Record the number of comparisons (comparing data elements to find the minimum) and number of swaps made along the way. The first row for the first data set is completed for you.

i 7 3 9 5 **ii** 6 2 8 5 4

Pass number	List order	Number of comparisons made	Number of swaps made
1	3 7 9 5	3	1
. . .			
		Total:	**Total:**

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

Bubble sort

This method works through the list by comparing each adjacent pair of numbers along the way and swapping them if necessary so that the smaller number is on the left. One pass through of the list ensures the biggest number is at the end of the list. The process continues until the list is sorted.

For example, for the data set 3 8 6 4 7 the first pass involves the following stages:

3	8	6	4	7
3	8	6	4	7
3	6	8	4	7
3	6	4	8	7
3	6	4	7	8

Each adjacent pair was compared (3 and 8 stayed where they were as they were already in order), which involved 4 comparisons and 3 swaps. Pass 2 will then work on the list 3 6 4 7.

b Use the bubble sort algorithm for the two data sets in part **a**, completing the table.

c Compare your tables from parts **a** and **b**.


i What do you notice about the number of comparisons required in the two algorithms? How does this compare to the number of data elements?

ii What do you notice about the number of swaps between the two algorithms?

iii Hence, in general which algorithm do you think is considered more efficient (runs faster or requires less processing)?

Now that we have algorithms for sorting a list of data we can call upon these to help us find the median of a set of data.

d Copy and complete the median flow-chart algorithm on the left. Use the box shapes as a guide. Include how you will use the number of elements in the list to locate the position of the median value in the list.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

3.3 Simulations in probability calculations

Simulations to determine good approximations for probabilities can be carried out using different techniques. For experiments with two equally likely outcomes, a coin is a good device to carry out the simulation. Other devices include dice, spinners, random number generators such as calculators and computers, and random number tables.

The following tasks will explore the use of some of these.

a *Simulation devices*

Consider the following scenarios and describe how you could use the listed device to carry out the simulation.

i Simulate the gender of the next baby born, using a coin.

ii Simulate the situation '1 in 3 chocolate bars wins a prize', using a 6-sided die.

iii Simulate getting the correct answer from a random guess on a multiple-choice question with 4 options, using a deck of cards. How could you simulate this using a 6-sided die?



b *Technology and random numbers*

You may have noticed in part **a** that some probabilities were easier to simulate than others with certain devices. The use of random numbers on calculators and computers makes this more achievable. For example, to represent a 2 in 5 chance you can generate a random integer between 1 and 5 and assign the numbers 1 and 2 to represent a win and 3, 4 and 5 a loss.
Many scientific calculators have a randint() function that can generate a random integer in a specified range. In Microsoft Excel RANDBETWEEN(1, 5) will generate a random integer between 1 and 5.

Consider a basketball player who gets on average 75% of her free throws in.

i Consider how you could use a random number generator to simulate her next free throw.

Technology would be helpful in the following but coins or a die could also be used effectively.

ii Using your chosen device, estimate the probability that she makes 5 or 6 of the 6 free throw shots she takes in a game by completing the following.

- Generate 6 random numbers and count if 5 or 6 of them are in. Use a table like the one below. Repeat for 50 trials.

Trial number	Number of free throw shots in out of 6	5 or 6 shots in? (Yes/No)
1		
2		
. . .		
50		

- Calculate the probability estimate: $\frac{\text{number of times 5 or 6 in (Yes)}}{\text{number of trials}}$

iii The theoretical probability is 0.534. How does your answer compare? Compare with other members of the class.

ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

c *Random number tables*

The simulation in part **b** could also be carried out using a random number table.

The random number table shown on the next page groups digits in groups of five but is generated with a random digit between 0 and 9 at each place. These numbers can then be grouped together to represent 1, 2, 3 digit numbers and more.

Consider the following scenario.

Over his career, an AFL player has a set shot for goal record of 68% goals, 24% behinds and 8% out on the full.
Since we have percentages we can use random numbers between 0 (i.e. 00) and 99. For a goal we could use the numbers 00–67 (this covers 68 out of 100 numbers, i.e. a 68% chance).

i Decide which sets of two digit numbers could then be used to represent a behind and out on the full.

Two digit numbers can be drawn from the random number table by selecting a starting point in the table and reading off two digits at a time to form a number.
For example, for the block of numbers 58315 10578 you would get the numbers 58, 31, 51, 05 (i.e. 5) and 78.

ii Use the random number table on the next page (starting from any location) to carry out 20 trials to estimate the probability that in 5 set shots on goal he scores at least 3 goals.

- Record your results in a table like the one below.

Trial number	Number of goals	At least 3 goals? (Yes/No)
1		
2		
. . .		
20		

- Estimate the probability: $\frac{\text{number of yes}}{\text{number of trials}}$
- Compare your result with other members of the class.



ISBN 978-1-009-59315-1
© Greenwood et al. 2025 Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.

58315	10578	77473	16526	53775
22646	82056	42313	50814	60650
07419	77083	11543	26629	08313
76940	62101	86568	08456	53641
15963	82704	06272	58036	61078
80644	86510	78615	06079	60154
30571	92400	07305	64811	03054
09182	14662	10472	97400	65696
34361	67837	16869	00904	79928
56845	84009	10030	28001	64238
64794	74684	75213	38693	27053
78586	70912	18697	55949	39557
92444	18934	51892	24300	20247
10287	04605	29245	52500	90501
75233	47520	50251	75778	69504
89599	84143	14821	26191	18076
35287	40045	24635	45782	12217
07320	03379	80797	26408	75542
43560	20875	53010	23045	23634
90623	75405	18139	31992	64709
06655	97645	90074	57757	48091
59959	72048	57584	39509	43253
88358	78732	45246	00345	02690
11524	56474	20503	02944	86411
45850	83072	27083	65098	92990
05013	21373	93138	21196	91294
80820	27119	29971	73672	21843
30859	16362	07037	82057	67908
19037	36352	26371	72254	33515
94869	80499	48086	43199	55744
02516	67531	73014	46866	52298
46238	37370	63515	37083	33247
26103	81339	93391	43856	95475
07664	85034	46581	88772	93372
18305	71127	91648	96303	65869
04887	10435	77400	30370	20995
08597	14871	08080	99425	73733
01876	18260	04657	87735	07273
16680	12966	75383	87195	86948
43454	22639	45772	62461	67602
89229	90868	03485	85955	73123
70283	78014	64377	40020	73714
19609	05831	80438	76003	50046
87442	56988	25210	21541	81928
43409	93065	52495	77536	81227

ISBN 978-1-009-59315-1
© Greenwood et al. 2025
Cambridge University Press
Photocopying is restricted under law and this material must not be transferred to another party.